Feasibility of an Intervention to Support Hearing and Vision in Dementia

DOI:

[10.1111/jgs.15936](https://doi.org/10.1111/jgs.15936)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Hooper, E., Simkin, Z., Abrams, H., Camacho, E., Charalambous, A. P., Collin, F., Constantinidou, F., Dawes, P., Elliott, R., Falkingham, S., Frison, E., Hann, M., Helmer, C., Himmelsbach, I., Hussain, H., Marié, S., Montecelo, S., Reeves, D., Regan, J., ... Leroi, I. (2019). Feasibility of an Intervention to Support Hearing and Vision in Dementia: The SENSE-Cog Field Trial. *Journal of the American Geriatrics Society*.
<https://doi.org/10.1111/jgs.15936>

Published in:

Journal of the American Geriatrics Society

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

**Feasibility of an intervention to support hearing and vision in dementia:
The SENSE-Cog Field Trial**

Emma Hooper,¹ Zoe Simkin¹, Harvey Abrams², Elizabeth Camacho³, Anna Pavlina
Charalambous⁴, Fideline Collin⁵, Fofi Constantinidou⁴, Piers Dawes⁶, Rachel Elliott³, Sue
Falkingham⁷, Eric Frison⁵, Mark Hann⁸, Catherine Helmer⁹, Ines Himmelsbach¹⁰, Hannah
Hussain³, Sarah Marié¹¹, Susana Montecelo¹¹, David Reeves⁸, Jemma Regan¹², Chryssoula
Thodi⁴, Lucas Wolski⁹, Iracema Leroi^{1,13, 14}

¹ Division of Neuroscience and Experimental Psychology, University of Manchester, U.K.

² University of South Florida, Tampa, Florida, USA

³ Manchester Centre for Health Economics, University of Manchester, U.K.

⁴ Department of Health Sciences, School of Sciences, European University Cyprus, Nicosia,
Cyprus

⁵ University of Bordeaux, INSERM, EUCLID/F-CRIN Clinical Trials Platform, Bordeaux
Population Health Center, France

⁶ Manchester Centre for Audiology and Deafness, University of Manchester, U.K.

⁷ Starkey Hearing technologies, U.K.

⁸ Centre for Biostatistics, University of Manchester, U.K.

⁹ University of Bordeaux, INSERM, Bordeaux Population Health Research Center, team LEHA,
France

¹⁰ Catholic University of Applied Sciences Freiburg, Freiburg, Germany

¹¹ Essilor International, Research & Development, Paris, France

¹² Research Associate, University of Manchester, UK

¹³ Global Brain Health Institute, Trinity College Dublin, Ireland

¹⁴ Greater Manchester Mental Health Foundation Trust, UK

(Please note that the first two names are joint first authors; the final author is the senior author; all other authors are in alphabetical order)

***Corresponding author:**

Iracema Leroi

Global Brain Health Institute

Room 0.60, Lloyd Building

Trinity College Dublin

Dublin 2, Ireland

+353 (0) 1 896 8500

Email: iracema.leroi@manchester.ac.uk

Running head: *Feasibility of sensory support in dementia*

This paper was first presented as part of a symposium at the British Society of Gerontology Annual meeting in Manchester, 5th July 2018.

Word count:

Abstract: 300; Main text: 2333; 2 tables; 1 figure; 1 supplementary table; 28 references.

Keywords: Dementia, hearing impairment, vision impairment, feasibility, acceptability, tolerability.

45 *Trial registration number:* The trial is a psychosocial intervention with an allocated ISRCTN
46 number 35019114 16th January 2018

47 **Impact statement:** We certify that this work is entirely novel and is the first study of hearing
48 and vision enhancement in people living with dementia. This interdisciplinary approach
49 makes a significant contribution to the literature and sets the stage for further full scale
50 evaluations of hearing and vision interventions to improve outcomes for people with
51 dementia. This is the first part of a two-part report.

52 **Abstract**

53 **Background:** People living with dementia (PwD) frequently experience hearing and vision
54 impairment that is under-recognised and under-treated, resulting in reduced quality of life.
55 Managing these impairments may be an important strategy to improve outcomes in PwD.

56 **Objective:** To field trial a multi-faceted ‘Sensory Intervention’ (SI) to enhance hearing and
57 vision in PwD.

58 **Design:** An international single arm, open label, feasibility, acceptability and tolerability
59 study.

60 **Setting:** Home-based, in the United Kingdom, France, and Cyprus.

61 **Participants:** Adults aged ≥ 60 with mild-moderate dementia and uncorrected or sub-
62 optimally corrected hearing and/or vision impairment, and their study partners (n=19
63 dyads).

64 **Intervention:** A ‘Sensory Intervention’ (SI), comprising assessment of hearing and vision,
65 fitting of corrective devices (glasses, hearing aids), and home-based support from a ‘sensory
66 support therapist’ for device adherence and maintenance, communication training, referral
67 to support services, environmental sensory modification and optimisation of social
68 inclusion.

69 **Measurements:** Ratings of study procedure feasibility, and intervention
70 acceptability/tolerability, ascertained through questionnaires, participant diaries, therapist
71 logbooks and semi-structured interviews.

72

Results: We successfully delivered all intervention components, and these were received and enacted as intended in all those who completed the intervention. No serious adverse events were reported. Acceptability (i.e. understanding, motivation, sense of achievement) and tolerability (i.e. effort, fatigue) ratings of the intervention were within *a priori* target ranges. We met recruitment and retention (93.8%) targets in two of the three sites. Participants completed >95% of diary entries, representing minimal missing data. Delays in the logistics circuit for the assessment and delivery of hearing aids and glasses were identified, requiring modification. The need for minor modifications to some outcome measures and the inclusion criteria were identified.

Conclusion: This is the first study combining home-based hearing and vision remediation in PwD and the positive feasibility, acceptability and tolerability findings suggest that a full-scale efficacy trial, with certain modifications, is achievable.

Introduction

People with dementia (PwD) are more likely to experience vision and hearing impairment than their healthy counterparts^{1,2}, and such impairments, particularly in combination, may impact negatively on quality of life³ and other outcomes^{4,5}, as well as imposing an additional burden on health, social and informal care^{6,7}. Importantly, there is some evidence that managing vision and hearing impairments with glasses and hearing aids respectively may improve outcomes⁸ but the evidence is still equivocal and represents a gap in understanding. Unfortunately, in the context of dementia, adherence to hearing aids and other devices is often low⁹. Thus, simply correcting the sensory impairment may be insufficient to have a positive impact. In contrast, an intervention targeting the wider issue of sensory impairment and adherence with corrective devices may have a role. To address this, we iteratively developed a multi-faceted 'sensory intervention' (SI) which includes assessment and management of hearing and vision deficits and additional support to aid adoption of the corrective devices into everyday life as well other components to support sensory function³.

A first step in evaluating a complex psychosocial intervention should be a field trial of the study design, components and implementation of the intervention¹⁰. Thus, the primary aim of our field trial was to evaluate: (1) the feasibility of the operational aspects of an evaluation trial of the intervention; and (2) the acceptability and tolerability of the intervention. Our secondary aim was to explore a signal of clinical and cost effectiveness, which we report elsewhere (in preparation). The results of this study have informed the

design and conduct of a full-scale randomised controlled trial (RCT) in five European sites (ISRCTN 17056211)¹¹.

Methods

Study design and participants

This was an international single-arm, open-label field study of a newly developed ‘sensory intervention’ to improve the hearing and/or vision of PwD in three sites: Bordeaux, France (Site B), Manchester, UK (Site M) and Nicosia, Cyprus (Site N). The study received favourable ethical opinion at each site. All participants provided written informed consent prior to their inclusion. The planned sample size was n=24 dyads (PwD and study partner), with 8 dyads per site. All dyads received the *basic* version of the SI, with a sub-set of 4 receiving a 12-week *extended* version. We recruited participants from memory assessment clinics, and dementia research registries such as Join Dementia Research in the UK¹². Detailed inclusion and exclusion criteria have been described elsewhere¹³. Briefly, these included people over the age of 60, living at home with a formal diagnosis of mild-moderate stage dementia (Alzheimer disease, vascular dementia or ‘mixed’ Alzheimer and vascular dementia) and with capacity to consent (as per the UK’s Mental Capacity Act, 2005)¹⁴. All had a clinically significant uncorrected or partially corrected (e.g. outdated prescription for sensory aids) hearing and/or vision problem, ascertained using a brief objective screening procedure. The inclusion threshold for hearing was >35 dB HL over 1-3 kHz and above in the better ear, and for vision was binocular corrected visual acuity of $\leq 6/9$, 5 Snellen metric or $\geq +0.2$ LogMAR and a visual field of $\geq 10^\circ$. We did not include people with congenital hearing and/or vision impairments. Study partners were informal carers in regular contact with the PwD.

131

132

[Insert Table 1 here]

133

We have detailed participants' demographic and clinical characteristics in Table 1. Briefly, all

134

PwD were above age 62 years and all study partners were above age 42. Of the PwD, 42%

135

(n=8) had hearing impairment only; 58% (n=11) had both vision and hearing impairment;

136

and none had vision impairment alone. There was an equal proportion of PwD due to

137

Alzheimer disease and vascular dementia; and one individual had 'mixed' dementia.

138

Description of the intervention

139

The basic SI comprised: a clinical vision and/or hearing assessment with prescription and

140

fitting of corrective lenses, provided by Essilor International ¹⁵, and/or hearing aids ('behind

141

the ear' Muse Mini i2400), provided by Starkey Hearing Technologies¹⁶, and information

142

about device maintenance. The extended SI comprised additional components, delivered by

143

a Sensory Support Therapist (SST) in the participant's own home: (1) individualised

144

adherence support; (2) communication training; (3) functional assessment and goal-setting;

145

(4) referral to health and social care services; (5) supplementary sensory aids to enhance the

146

home environment; and (6) fostering social inclusion. The SST was an occupational therapist

147

skilled in dementia who received additional training in hearing and vision rehabilitation.

148

149

Study procedures

150

The detailed study protocol and schedule of events are described elsewhere¹³ and shown in

151

Figure 1 in abbreviated form. Briefly, after informed consent, we screened PwD for hearing,

152

vision and cognitive impairment using the Sivantos Siemens HearCheck screener¹⁷, Peek

153

Acuity app¹⁸, and MoCA ¹⁹, followed by a baseline assessment and the intervention. The

basic SI was delivered over 4 weeks at all three sites to enable us to evaluate feasibility of study procedures. At Site M, the extended SI, delivered over 12 weeks in participants' homes, enabled us to evaluate further study procedures, feasibility of the intervention delivery, and its acceptability and tolerability.

[add Figure 1 here]

Evaluation framework

We based our evaluation on a modified version of the ACCEPTANCE framework for feasibility studies²⁰. Data were captured at baseline and within one week of the last intervention visit. At each visit for the extended SI, PwD and study partners completed diaries with in-house Likert-type scales (rating each aspect of acceptability and tolerability on a scale of 1=strongly disagree to 5= strongly agree) and space for free text, and the SST completed a log book and field notes. We conducted semi-structured interviews with a sub-sample of dyads at sites M and N who received either the basic (n=8 dyads) or extended (n=2 dyads) SI. The focus of the interviews was on participants' perception, experiences and acceptance of the SI.

Feasibility of trial procedures: These included our recruitment strategy, suitability of eligibility criteria, execution of the 'logistics circuit' for assessment and supply of hearing aids and glasses, feasibility of the participant diaries, data collection methods, suitability of the battery of effectiveness measures, and retention.

Described in detail elsewhere¹³, effectiveness measures for the PwD were: quality of life, mental wellbeing, neuropsychiatric symptoms, functional ability (dementia-, hearing- and vision-related), and relationship satisfaction. Effectiveness measures for the study partner

were: wellbeing, mental health, caregiving-related burden and stress, and relationship satisfaction. Health care resource use questionnaires were included. Since this was an open-label study, we did not evaluate randomisation and blinding procedures.

Feasibility of the intervention components and implementation: To determine whether the intervention was delivered, received and enacted as intended²¹, we obtained SST visit completion rates, visit duration and SST logbook feedback.

Acceptability of the intervention: The appropriateness of the delivery and receipt of the intervention²² was determined by: percentage dropouts due to non-acceptability and rate of serious adverse events. The 'acceptability' criterion for the extended SI was 100% of participants scoring within the *a priori* target ranges on a five point Likert-type scale: $\geq 3/5$ for '*understanding*', '*interest*', '*emotional response*', '*motivation*' and '*sense of achievement*'.

Tolerability of the intervention: This was operationalised by percentage dropouts due to intolerance of the intervention and diary ratings of '*effort*' and '*fatigue*' for the extended SI. The criterion for 'tolerability' was 75% of participants scoring the intervention with the *a priori* target ranges: $\geq 3/5$ for '*effort*' and '*fatigue*'.

Data analysis

We used descriptive statistics for the quantitative analysis since the study was not formally powered to detect specific post-intervention effect sizes. The small sample size increases the likelihood of a Type II error when using inferential statistics. We applied content

analysis²³, a reliable method of analysing of qualitative data using ‘coding units’, to the non-quantitative data from the semi-structured interviews, participant dyad diaries, researcher field notes and SST logbooks.

Results

Details of the feasibility of trial procedures and acceptability and tolerability of the intervention are outlined in Supplementary Table S1.

Feasibility of the trial procedures

(a) Recruitment and retention

Recruitment was successful in Sites M and N, but slower in Site N (2.6 dyads per month for 3 months and 1.3 dyads per month for 6 months, respectively) and did not reach target in Site B, which recruited 3 dyads. This resulted in a total sample size of 19 dyads from an intended sample of 24 dyads. The retention rate at Site M was 87.5% (one participant dyad withdrew due to study-related burden) and at Site N was 100%. All three dyads at Site B did not complete the study. Non-completion and failure to recruit at Site B was due to the lack of a pathway between the study site and the necessary referral sources and lack of infrastructure to support the logistics circuit. Screening and baseline visits were conducted according to protocol in all sites.

(b) Suitability of eligibility criteria

Investigators at all sites perceived that the cognitive score cut-off threshold (MoCA ≥ 12) was too high and would potentially exclude PwD who could meaningfully participate.

221 Additionally, of the 19 PwD who screened positive for hearing impairment, the assessing
222 audiologist did not prescribe hearing aids for five of the participants due to mildness of
223 impairment. None of these PwD received the extended intervention. All other
224 inclusion/exclusion criteria were considered appropriate by investigators.

225
226 (c) Execution of the service and device logistics circuit

227 Referrals to vision and/or audiology assessments post-baseline visit were successful
228 although we experienced some delays and variation across study sites, with delivery of
229 glasses ranging from 7-9 weeks and hearing aids 3-20 weeks post-baseline. Delays in the
230 logistics circuit impacted on the study timeline, with post-intervention assessments being
231 conducted 7-25 weeks post-baseline. Reasons for delay were clearly identified, including
232 difficulties in arranging study visits, inadequate communication among assessing clinicians
233 and the study team, and delays in delivery of devices from suppliers.

234
235 (d) Usability of study materials and suitability of effectiveness battery

236 Diary use by dyads was feasible and acceptable, with a 95% completion rate of entries for
237 PwD and 97% for the study partners. The battery of effectiveness measures was feasible and
238 well-tolerated, except for the self-efficacy and self-reported hearing and vision impairment
239 scales, which were difficult for the PwD to report on due to deteriorating insight. Missing
240 data on effectiveness scales for study completers was minimal (<10%) and within the *a*
241 *priori* acceptability threshold (see Supplementary Table S1).

242 *Feasibility of the intervention components and implementation*

We achieved 100% adherence to the study protocol for the basic SI at Sites M and N for study completers. At Site B, study procedures were not completed due to problems with the study team, thus we could not evaluate feasibility at this site. At Site M, 100% of components of the extended SI were delivered, received and enacted as intended, over a range of 7-12 sessions (median 9), and a median session duration of 95 minutes (range 45-135). This included certain iterative changes to the intervention recorded in the SST logbook. This number of sessions, together with the need to schedule vision and hearing assessments and wait for delivery of sensory aids, required 20 weeks for full intervention package to be delivered.

Acceptability and tolerability of the intervention

At Sites M and N there were no withdrawals due to lack of acceptability of the basic or extended SI. At site M, one dyad withdrew due poor tolerability of the extended SI (Table 2, participant 4). All adverse events were classified as 'mild', including poor fit or discomfort from corrective devices. This included expressions of concern about the potential to lose or damage the corrective device, resulting in anxiety of a mild level. No serious adverse events were experienced. For the extended SI, Likert-style mean acceptability ratings of '*understanding*', '*motivation*', '*emotional response*', '*interest*' and '*sense of achievement*' all fell within the target range, as did tolerability ratings of '*effort*' and '*fatigue*' (Supplementary Table S1 and Table 2). Themes emerging from the post-intervention semi-structured interviews were: (1) good acceptability of session duration; (2) home-based delivery was acceptable, convenient and desirable; (3) additional SST support was 'extremely helpful' in encouraging the introduction of the corrective devices and optimising activity engagement;

and (4) study evaluation procedures were burdensome for some dyads because it was challenging for the PwD to distinguish between their different impairments.

[Insert Table 2 here]

Discussion

This is the first reported study of a hearing and vision intervention in PwD, demonstrating that such an intervention is feasible as a home-based therapy, with slight modifications, in two of the three study sites. We ascertained that the intervention itself is acceptable to and tolerated by PwD and their study partners. We identified the need for modifications to the study design for a full clinical trial, including: tightening the logistics circuit, widening the recruitment pool, replacing the under-recruiting site, changing certain effectiveness measures and altering the inclusion criteria for level of cognitive impairment to MoCA ≥ 10 . Since most of the outcome measures are informant-rated or proxy-rated, it will be possible to capture accurate data for this group of participants. Diary feedback on participant fatigue, effort and motivation and other parameters allowed fine-tuning of the intervention, and underscored the need for careful tailoring to individualised requirements, an approach consistent with the conduct of pragmatic trials²⁴. We have incorporated all modifications into a final protocol for a full RCT. We have addressed the recruitment and retention problems at Site B by replacing it with a new site in Dublin, which has a dedicated dementia service a proven record of successful recruitment to non-pharmacologic RCTs. Furthermore, using the experience of this feasibility study, we have selected a further two European dementia services (Athens and Nice) with similarly strong research experience to participate in the full SENSE-Cog RCT (ISRCTN 17056211)¹¹, making five sites in total. The experience in this study enabled us to develop robust site selection criteria for the additional sites. Finally,

a limitation of this study was the extended SI was only delivered in one of the field trial sites, but this gave us rich data from which to develop the final extended SI for the RCT.

In summary, this is the first study combining hearing and vision remediation in PwD and the positive feasibility, acceptability and tolerability findings suggest that a full-scale efficacy trial with certain modifications is achievable.

Acknowledgements

Conflict of interest

HA and SF are employed by Starkey Hearing Technologies, SMa and SMO are employed by Essilor International. There are no other conflicts of interest.

Authors' contributions

IL and PD are the programme leads and conceptualised and designed the field trial. EH is the Senior Sensory Support Therapist. ZS and APC are research assistants. JR was study coordinator for the field trial. RE and EC provided health economic input. MH and DR provided statistical input for the study. IH and LW led the qualitative analysis. CH and FoC oversaw study delivery in their sites. FiC and EF were involved in the study design and interpretation of study results. CT, HA, SF, SMa and SMO provided professional input to the design and conduct of the trial. EH, ZS and IL took primary responsibility for writing the paper; all authors were involved in critical revision of the article.

We thank Christine Dickinson, University of Manchester for assisting with the design of the study. We thank our industry collaborators at The Outside Clinic, Starkey Hearing

Technologies, Essilor International, Siemens Hearing Aids, Sivantos Limited (HearCheck) and PEEK Vision Limited (PEEK Acuity). We thank the Greater Manchester NIHR Clinical Research Network and Greater Manchester Mental Health Trust (GMMH) for supporting the study. We thank the Research User Group for input on the design and development of the intervention and participants and their families for taking part in the study.

Sponsor's role

This work was supported by the European Union's Horizon 2020 research and innovation programme under grant agreement No 668648.

References

1. Bowen M, Edgar DF, Hancock B, et al. The Prevalence of Visual Impairment in People with Dementia (the ProVIDe study): a cross-sectional study of people aged 60–89 years with dementia and qualitative exploration of individual, carer and professional perspectives. *Heal Serv Deliv Res.* 2016;4(21):1-200. doi:10.3310/hsdr04210
2. Lin FR, Metter EJ, O'Brien RJ, Resnick SM, Zonderman AB, Ferrucci L. Hearing loss and incident dementia. *Arch Neurol.* 2011;68(2):214-220.
3. Leroi I, Pye A, Armitage CJ, et al. Research protocol for a complex intervention to support hearing and vision function to improve the lives of people with dementia. *Pilot Feasibility Stud.* 2017;3(1):1-16. doi:10.1186/s40814-017-0176-1
4. Guthrie DM, Davidson JGS, Williams N, et al. Combined impairments in vision, hearing and cognition are associated with greater levels of functional and communication

334 difficulties than cognitive impairment alone: Analysis of interRAI data for home care
 335 and long-term care residents in Ontario. *PLoS One*. 2018;13(2):e0192971.
 336 doi:10.1371/journal.pone.0192971

337 5. Lawrence V, Murray J, Ffytche D, Banerjee S. The experiences and needs of people
 338 with dementia and serious visual impairment: a qualitative study. *Int*
 339 *Psychogeriatrics*. 2009;21(3):511-518. doi:10.1017/S1041610209008424

340 6. Nyman SR, Innes A, Heward M. Social care and support needs of community-dwelling
 341 people with dementia and concurrent visual impairment. *Aging Ment Health*.
 342 2017;21(9):961-967. doi:10.1080/13607863.2016.1186151

343 7. Ke KM, Montgomery A-M, Stevenson M, O'Neill C, Chakravarthy U. Formal and
 344 informal care utilisation amongst elderly persons with visual impairment. *Br J*
 345 *Ophthalmol*. 2007;91(10):1279-1281. doi:10.1136/bjo.2006.113142

346 8. Dawes P, Wolski L, Himmelsbach I, Regan J, Leroi I. Interventions for hearing and
 347 vision impairment to improve outcomes for people with dementia: A scoping review.
 348 *Int Psychogeriatrics*. 2018:1-19. doi:10.1017/S1041610218000728

349 9. Hartley D, Rochtchina E, Newall P, Golding M, Mitchell P. Use of hearing AIDS and
 350 assistive listening devices in an older Australian population. *J Am Acad Audiol*.
 351 2010;21(10):642. doi:10.3766/jaaa.21.10.4

352 10. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and
 353 evaluating complex interventions: the new Medical Research Council guidance. *Br*
 354 *Med J*. 2008;337:a1655. doi:10.1136/bmj.a1655

- 355 11. Regan J, Frison E, Collin F, et al. Individualised sensory intervention to improve quality
356 of life in people with dementia and their companions (SENSE-Cog trial): study
357 protocol for a randomised controlled trial. *Trials*. 2019;20:80(1):1-15.
358 doi:10.1186/s13063-018-2973-0
- 359 12. Join Dementia Research. www.joindementiaresearch.nihr.ac.uk.
- 360 13. Regan J, Dawes P, Pye A, et al. Improving hearing and vision in dementia: Protocol for
361 a field trial of a new intervention. *BMJ Open*. 2017;7(11):1-12. doi:10.1136/bmjopen-
362 2017-018744
- 363 14. Department of Health. *Mental Capacity Act*. London; 2005.
- 364 15. Essilor International. <https://www.essilor.com/en/>.
- 365 16. Starkey Hearing Technologies. <https://www.starkey.com/>.
- 366 17. Sivantos Siemens Hearcheck. [https://www.bestsound-
367 technology.co.uk/nhs/equipment/hear-check/](https://www.bestsound-technology.co.uk/nhs/equipment/hear-check/).
- 368 18. Peek Vision. <https://www.peekvision.org/>.
- 369 19. Nasreddine ZS, Phillips NA, Bédirian V, et al. The Montreal Cognitive Assessment,
370 MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc*.
371 2005;53(4):695-699.
- 372 20. Charlesworth G, Burnell K, Hoe J, Orrell M, Russell I. Acceptance checklist for clinical
373 effectiveness pilot trials: a systematic approach. *BMC Med Res Methodol*.
374 2013;13(78):1-7. doi:10.1186/1471-2288-13-78

- 375 21. Burgio L, Corcoran M, Lichstein KL, et al. Judging outcomes in psychosocial
376 interventions for dementia caregivers: The problem of treatment implementation.
377 *Gerontologist*. 2001;41(4):481-489. doi:10.1093/geront/41.4.481
- 378 22. Sekhon M, Cartwright M, Francis JJ. Acceptability of healthcare interventions: An
379 overview of reviews and development of a theoretical framework. *BMC Health Serv*
380 *Res*. 2017;17(1):1-13. doi:10.1186/s12913-017-2031-8
- 381 23. Hsieh H-F, Shannon SE. Three approaches to qualitative content analysis. *Qual Health*
382 *Res*. 2005;15(9):1277-1288.
- 383 24. Hawe P, Shiell A, Riley T. Complex interventions: how “out of control” can a
384 randomised controlled trial be? *Br Med J*. 2004;328(June):1561-1563.
- 385 25. Chen G, Gully S, Eden D. Validation of a New General Self-Efficacy Scale. *Organ Res*
386 *Methods*. 2001;4(1):62-83. doi:10.1177/109442810141004
- 387 26. Yesavage JA. Geriatric depression scale. *Psychopharmacol Bull*. 1988;24(4):709-711.
- 388 27. Burns DD. *Ten Days to Self-Esteem: The Leader’s Manual*. Quill/HarperCollins
389 Publishers; 1993.
- 390 28. Zigmond A, Snaith R. The Hospital Anxiety and Depression Scale. *Acta Psychiatr*
391 *Scand*. 1983;67:361-370.

392

Legends

Figure 1: Flowchart of study procedures (submitted separately as a TIF file)

Table 1 Description of the baseline demographic and clinical variables in participants with dementia and their study partners

Variable	Category	Participants with Dementia	Study partner Participants
N		19	19
Age (Years)	Median (IQR) Range	76 (11) 63 to 88	67 (13) 43 to 82
Gender	Female Male	7 (36.8%) 12 (63.2%)	16 (84.2%) 3 (15.8%)
Duration of Cognitive Impairment (Months)	Median (IQR) Range	60 (54) 6 to 120	NA
Level of Cognitive Impairment (MoCA Total Score)	Mean (SD) Range	17.3 (3.7) 12 to 23	NA
Dementia Sub-Type	Alzheimer's Vascular Mixed	9 (47.4%) 9 (47.4%) 1 (5.3%)	NA
Sensory Impairment	Hearing only Vision only Hearing & Vision	8 (42.1%) 0 11 (57.9%)	NA
Relationship to PwD	Spouse/ Partner Son/ Daughter Other Relative	NA	13 (68.4%) 5 (26.3%) 1 (5.3%)
Hours per Week spent with PwD	Median (IQR) Range	NA	100 (115) 3 to 168

SD: standard deviation; IQR: interquartile range

405 **Table 2 Acceptability and tolerability of the extended Sensory Intervention***

		Ratings of SI visits by PwD, study partner and SST: Mean score (range)			
		Participant 1	Participant 2	Participant 3	Participant 4
Acceptability	Understanding ^{PwD}	4.7 (4-5)	4.6 (4-5)	3.1 (2-4)	3.3 (2-5)
	Motivation ^{PwD}	4.6 (4-5)	4.9 (4-5)	3.9 (3-5)	3.3 (2-4)
	Motivation ^{SP}	4.4 (4-5)	5 (5-5)	3.3 (2-4)	3.8 (3-4)
	Motivation ^{SST}	4 (4-4)	4.8 (4-5)	3.3 (2-4)	4.5 (4-5)
	Sense of achievement ^{SP}	4.4 (4-5)	4.7 (4-5)	3 (2-5)	3.5 (3-4)
	Sense of achievement ^{SST}	3.8 (3-4)	4.6 (4-5)	3.1 (2-4)	4 (3-5)
	Interest ^{SP}	4.7 (4-5)	5 (5-5)	3.6 (3-4)	3.8 (3-4)
	Interest ^{SST}	4 (4-4)	4.8 (4-5)	3.8 (2-4)	4.8 (4-5)
	Emotional response ^{SP}	4.1 (4-5)	4.4 (4-5)	3.1 (2-4)	3.3 (3-4)
Tolerability	Effort ^{PwD}	4.7 (4-5)	4.5 (4-5)	3.1 (2-5)	2.3 (1-4)
	Fatigue ^{PwD}	5 (5-5)	3.2 (2-5)	3.4 (1-4)	1.5 (1-2)

406 Key: ^{PwD} PwD rating of response; ^{SP} Study partner rating of PwD's response; ^{SST} SST rating of
 407 PwD's response.

408 * Rated by participants on a 5-point Likert-type scale: 1=strongly disagree; 2=disagree;
 409 3=neutral; 4=agree; 5=strongly agree (reverse rating for 'effort' and 'fatigue').

410

411 **Supplementary Table S1: Feasibility of trial procedures and intervention feasibility,**
412 **acceptability and tolerability**

Parameter and <i>a priori</i> evaluation criteria (if applicable)	Findings	Evidence to support finding	Changes implemented for RCT
Feasibility of study procedures			
Eligibility criteria: ≥75% screened meet study criteria	Criteria are acceptable except: (1) cognitive score cut-offs may be set too high and exclude PwD who may be appropriate; (2) HearCheck screening cut-off may not be stringent enough.	100% of those screened met inclusion criteria ^a . 5 participants who screened positive on hearing impairment using the HearCheck were deemed not clinically suitable for hearing aids on full assessment ^a . There was an imbalance of sensory diagnostic groupings across the sites ^a .	Inclusion criteria adjusted to MoCA ≥10. Remaining components of the SI will continue for any PwD not prescribed sensory aids following clinical assessment.
Recruitment: Total target number Rate	Successful at 2 of 3 sites. Slower than required for a larger trial.	100% at Site M and N; 38% at Site B ^a . Rate was 2.7 dyads per month at Site M and 1.3 dyads per month at Site N ^a . Incomplete recruitment at Site B.	Site B replaced with an alternative. Recruitment pool widened.
Retention: ≥60% completed all study procedures	Successful in 2 of 3 sites.	93.8% completed the study in Sites M and N; 0% completed in Site B ^a .	Site B replaced with an alternative.
Screening & baseline process:	Appropriate due to the length of assessment battery.	9 dyads had one visit; 10 had two visits ^a .	No changes indicated.
Outcome battery administration and suitability: ≥10% missing data suggests scale is not acceptable	Outcome rating scales are generally acceptable. Some scales were not suitable for the study population and require	<10% missing data from outcome rating scales at baseline and follow-up ^a . Missing items within given scales included gender-specific physiological	General Self Efficacy Scale ²⁵ dropped. Geriatric Depression Scale ²⁶ replaced with the Hospital Anxiety and Depression Scale ²⁸ .

	revision.	<p>items^a.</p> <p>Minimal or no concerns were noted on battery duration and level of difficulty, other than all 3 sites reporting problems with:</p> <p>PwD understanding the General Self Efficacy Scale²⁵ items^b;</p> <p>The Geriatric Depression Scale²⁶ was not appropriate for younger study partners^a;</p> <p>PwD self-report of hearing and vision impairment was not valid^a.</p> <p>The Relationship Satisfaction Scale²⁷ was difficult to administer in presence of the study partner^b</p>	<p>Caregiver reports of hearing and vision impairment introduced alongside PwD's self-report.</p> <p>Relationship Satisfaction Scale²⁷ administration procedure amended.</p>
Device logistics circuit:	Broadly feasible; areas for improvement identified.	<p>All prescribed hearing aids and glasses were received by participants^a.</p> <p>Delays in assessment for and receipt of corrective devices impacted on overall study timelines^a.</p>	<p>Logistics circuit tightened through training and identification of dedicated clinicians.</p> <p>Timeframe for SI delivery extended.</p>
Participant diary: ≥80% completion	Diary activity was feasible for both PwD and study partner.	95% of diary entries completed by both members of the dyad ^c .	No changes indicated.
<p><i>Feasibility of the Sensory Intervention (SI) components and implementation:</i></p> <p>Basic SI: Basic intervention (Sites M, N and B)</p> <p>Extended SI: Extended intervention (Site M)</p>			
Basic SI: Was the basic SI delivered, received and enacted as intended?	It is feasible, although timeline deviations were evident.	100% of participants received a vision and / or hearing assessment and prescription of	<p>Logistics circuit tightened up.</p> <p>Window for vision /</p>

		corrective devices (if indicated) within 20 weeks of baseline ^a . 100% of participants completed measures of device skills and knowledge (hearing aids / glasses) ^a .	hearing assessment specified as 1-8 weeks from randomisation.
Extended SI: Completion of extended SI within 12 weeks	It is feasible to complete the SI within 12 visits. The timeline of 12 weeks was not feasible due to logistics circuit delays and participant / SST availability. Successful delivery of each component is possible. It is viable to introduce the SI components in a flexible manner to account for delays in receiving hearing aids / glasses.	SI was completed over a mean of 9 visits (range 7-12) ^b . Time from baseline to follow-up was mean 18 weeks (range 17-20) ^a . 100% of participants completed functional assessment and set study-related goals ^b ; of those that continued the SI to completion, 100% of components were addressed ^b . Elements of the extended SI were successfully introduced prior to device delivery ^b .	Timeframe extended from 12 weeks to 18 weeks for SI delivery.
Acceptability of the intervention:			
Basic SI: Was the Sensory Intervention appropriate?	The basic intervention is acceptable	100% of participants were willing to receive their prescribed aids ^a . No participant withdrawals due to lack of acceptability ^a .	No changes indicated.
Extended SI: 100% of: Score ≥ 3 on PwD scales for understanding and motivation Score ≥ 3 on SP and SST scales for motivation and sense	The intervention is broadly acceptable. PwD may not demonstrate anticipated levels of sense of achievement; however there were no withdrawals due to lack of acceptability.	100% of mean scores are within range for PwD ^c . 100% of mean scores are within range for SP and SST ^{b,c} .	No changes indicated.

of achievement Score ≥ 3 on SP and SST scales for interest and emotional response			
<i>Tolerability of the intervention by participants:</i>			
Basic SI:	The basic intervention is tolerable	100% of participants were able to complete their vision and / or hearing assessment ^a . The basic intervention was completed over maximum 3 visits ^a .	No changes indicated.
Extended SI: 75% of: Score ≥ 3 on PwD scale for effort and fatigue	The intervention is broadly tolerable but the SST needs to be mindful that lower tolerability ratings could indicate withdrawal risk.	One participant withdrew after 4 SI visits due to perceived burden (Participant 4). This is reflected in their effort and fatigue scores ^c . 75% of scores were ≥ 3 . This is within the <i>a priori</i> range for tolerability.	SST to monitor diary responses and tailor the SI to the PwD's needs.

413

414 Key PwD = Person with dementia; SP= Study Partner; SST= Sensory Support Therapist

415 ^a Quantitative data; ^b SST logbook; ^c Participant dyad diaries

416

417

418