



Epsilon factors as algebraic characters on the smooth dual of GL_n

DOI:

[10.4064/bc120-1](https://doi.org/10.4064/bc120-1)

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Plymen, R. (2020). Epsilon factors as algebraic characters on the smooth dual of GL_n . In *Proceedings of conference on Quantum Dynamics* (Vol. 120, pp. 11-21). Banach Center Publications. <https://doi.org/10.4064/bc120-1>

Published in:

Proceedings of conference on Quantum Dynamics

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



EPSILON FACTORS AS ALGEBRAIC CHARACTERS ON THE SMOOTH DUAL OF GL_n

ROGER PLYMEN

ABSTRACT. Let K be a non-archimedean local field and let $G = GL_n(K)$. We have shown in previous work that the smooth dual $\mathbf{Irr}(G)$ admits a complex structure: in this article we show how the epsilon factors interface with this complex structure. The epsilon factors, up to a constant term, factor as invariant characters through the corresponding complex tori. For the arithmetically unramified smooth dual of GL_n , we provide explicit formulas for the invariant characters.

1. INTRODUCTION

Let K be a non-archimedean local field and let $G = GL_n(K)$. In the spirit of the Langlands program, this short paper provides a link between number theory and geometry. The arithmetic comes from ε -factors, which are naturally associated to representations of Galois groups, or more generally to representations of the Weil-Deligne group WD_K of K .

The smooth dual $\mathbf{Irr}(G)$ possesses a nice geometry: it is the disjoint union of smooth complex algebraic varieties, each of which is the quotient of a complex torus by a product of symmetric groups, see [BP].

In this article we show how the epsilon factors interface with this complex structure.

We continue with some background on the epsilon factors [L], [D]. The epsilon factors are very important and central in the theory of Artin L -functions. If E denotes a global field, then we have the absolute Weil group W_E , see [MP, 6.2.6]. The completed L -function $L(s, V)$ of a representation $W_E \rightarrow GL(V)$ of the Weil group W_E of the global field E defines a meromorphic function in the complex plane satisfying the functional equation

$$L(s, V) = \varepsilon(s, V)L(1 - s, V^*)$$

where V^* is the dual of the representation V of W_E , and the epsilon factor $\varepsilon(s, V)$ is defined by the product

$$(1) \quad \varepsilon(s, V) = \prod \varepsilon_{E_\nu}(s, V_\nu, \psi_\nu).$$

Here, ψ_ν is the local component at a place ν of a non-trivial additive character ψ of \mathbb{A}_E/E . Then ψ_ν is a non-trivial additive character of the local field E_ν , see [D, 5.11]. As usual, \mathbb{A}_E denotes the adèles of E .

Date: June 20, 2019.

From now on, we will focus on the non-archimedean places of the global field E , and we will write K for the non-archimedean local field E_ν . An elementary substitution, see §2, allows on to replace the epsilon factor $\varepsilon_K(s, V, \psi)$ with three variables, by the epsilon factor $\varepsilon_K(V, \psi)$ with two variables. From now on, we will be concerned with the epsilon factor $\varepsilon_K(V, \psi)$.

From the point of view of the local Langlands correspondence for GL_n , the relevant representations are the Weil-Deligne representations, see section 2. The set $\mathcal{G}_n(K)$ of equivalence classes of n -dimensional Weil-Deligne representations can be organised as a disjoint union of complex algebraic varieties:

$$\mathcal{G}_n(K) = \bigsqcup \mathfrak{X}$$

Each variety \mathfrak{X} arises in the following way. Let ρ' denote a Weil-Deligne representation of the Weil group W_K , and let m denote the number of indecomposable summands in ρ' . Then \mathfrak{X} is the quotient of a complex torus \mathfrak{T} of dimension m by a certain finite group \mathfrak{S} :

$$(2) \quad \mathfrak{X} = \mathfrak{T}/\mathfrak{S}.$$

By a *rational character*, or *algebraic character*, or simply *character*, we shall mean a morphism of algebraic groups

$$\mathfrak{T} \rightarrow \mathbb{C}^\times.$$

Such a character has the form

$$(z_1, \dots, z_m) \mapsto z_1^{\beta_1} \dots z_m^{\beta_m}$$

where the β_j are all integers.

Theorem 1.1. *Up to a constant $e(\mathfrak{X}, \psi)$, each epsilon factor $\varepsilon_K(V, \psi)$ factors through a rational character $\chi(\mathfrak{X}, \psi)$ of \mathfrak{T} . Quite specifically, we have*

$$(3) \quad (z_1, \dots, z_k) \mapsto (z_1^{\beta_1}, \dots, z_k^{\beta_k})$$

where the z_j are torus coordinates, \mathfrak{X} is the orbit of the Weil-Deligne representation

$$V_1 \otimes \mathrm{Sp}(d_1) \oplus \dots \oplus V_k \otimes \mathrm{Sp}(d_k)$$

and

$$\beta_j = (d_j - 1) \dim V_j^I + d_j [a(V_j) + n(\psi) \dim(V_j)]$$

where V_1, \dots, V_k are irreducible representations of the local Weil group W_K , $a(V_j)$ denotes the Artin conductor exponent of V_j , $n(\psi)$ denotes the conductor of ψ , and I denotes the inertia subgroup of W_K .

Since the conductors are integers, the number β_j is an integer. So the map (3) is a rational character of \mathfrak{T} .

The rather lengthy formula for the constant $e(\mathfrak{X}, \psi)$ appears later in this article, see (21) and (22).

We emphasize that the character $\chi(\mathfrak{X}, \psi)$ and the constant $e(\mathfrak{X}, \psi)$ depend only on the connected component \mathfrak{X} , once a base point has been chosen in \mathfrak{X} , and the additive character ψ has been chosen and fixed.

In §4, we focus on a part of the smooth dual of $\mathrm{GL}_n(K)$, namely the arithmetically unramified smooth dual. The extended quotient $T//W$ is a model for this part of the dual, where T is a maximal torus in the Langlands dual group $\mathrm{GL}_n(\mathbb{C})$ and W is the Weyl group of $\mathrm{GL}_n(\mathbb{C})$. We calculate explicitly the epsilon factors.

This article is an expanded account of a talk given at the conference *Geometry, representation theory and the Baum-Connes conjecture*, Fields Institute, July 2016, which was part of the EU Quantum Dynamics network activities. We thank Paul Baum for several valuable conversations, which led to major changes in the exposition of this article.

We take this opportunity to thank Paul Baum for an inspiring collaboration, which has withstood the test of time, and which has enabled us to discover and explore several new mathematical vistas. Paul observed that the article [BP] was the tip of a large iceberg. Further exploration led to several articles, of which [ABPS] is the most recent.

In writing this article, we were greatly influenced by the preprint of Ikeda [Ikeda]. The main background reference is Deligne [D], but we prefer to use the notation in Langlands[L]. We thank the referee for providing many insightful comments which influenced the final version of this Note.

2. WEIL-DELIGNE REPRESENTATIONS

We need to recall some material, following closely the exposition in [BP]. Let K be a non-archimedean local field. The normalized valuation will be denoted val_K , the norm of $x \in K$ will be denoted $\|x\|_K$, and the cardinality of the residue field of K will be denoted q_K . We have the local Artin reciprocity map

$$\mathrm{Art}_K : W_K \rightarrow K^\times.$$

We will write

$$\|w\| = \|\mathrm{Art}_K(w)\|_K = q_K^{-\mathrm{val}_K(\mathrm{Art}_K(w))}$$

for all $w \in W_K$.

The homomorphism $d : W_K \rightarrow \mathbb{Z}$ is defined by

$$d(w) = \mathrm{val}_K(\mathrm{Art}_K(w)).$$

The Weil group W_K fits into a short exact sequence

$$0 \rightarrow I_K \rightarrow W_K \xrightarrow{d} \mathbb{Z} \rightarrow 0$$

where I_K is the inertia group of K . A *Weil-Deligne representation* is a pair (ρ, N) consisting of a continuous representation $\rho : W_K \rightarrow \mathrm{GL}_n(V)$, $\dim_{\mathbb{C}}(V) = n$, together with a nilpotent endomorphism $N \in \mathrm{End}(V)$ such that

$$\rho(w)N\rho(w)^{-1} = \|w\|N.$$

For any $n \geq 1$, the representation $\mathrm{Sp}(n)$ is defined by

$$V = \mathbb{C}^n = \mathbb{C}e_0 + \cdots + \mathbb{C}e_{n-1}$$

with $\rho(w)e_i = \|w\|^i e_i$ and $Ne_i = e_{i+1}$ ($0 \leq i \leq n-1$), $Ne_{n-1} = 0$.

Let $\mathcal{G}_n(K)$ be the set of equivalence classes of semisimple n -dimensional Weil-Deligne representations. Let $\mathbf{Irr}(\mathrm{GL}_n(K))$ be the set of equivalence classes of irreducible smooth representations of $\mathrm{GL}_n(K)$.

We recall the local Langlands correspondence

$$\mathrm{rec}_K : \mathbf{Irr}(\mathrm{GL}_n(K)) \rightarrow \mathcal{G}_n(K)$$

which is unique subject to the conditions listed in [HT, p.2]. See also the single trace condition of Scholze in [Sch, Theorem 1.2(a)].

We identify the elements of the set $\mathcal{G}_1(K)$, the quasicharacters of W_K , with quasicharacters of K^\times via the local Artin reciprocity map Art_K . The local Langlands correspondence is compatible with twisting by quasicharacters [HT, p.2].

A quasicharacter $\eta : W_K \rightarrow \mathbb{C}^\times$ is (arithmetically) *unramified* if η is trivial on the inertia group I_K . In that case we have $\eta(w) = z^{d(w)}$ with $z \in \mathbb{C}^\times$. The group of unramified quasicharacters of W_K is denoted $\Psi(W_K)$. Let $\Phi = \Phi_K$ denote a geometric Frobenius element in W_K . The isomorphism $\Psi(W_K) \simeq \mathbb{C}^\times$ is secured by the map $\eta \mapsto \eta(\Phi_K)$.

Let now

$$\rho' = \rho_1 \otimes \mathrm{Sp}(d_1) \oplus \cdots \oplus \rho_m \otimes \mathrm{Sp}(d_m)$$

be a Weil-Deligne representation. In this formula, ρ_1, \dots, ρ_m are irreducible representations of the Weil group W_K . The set

$$\{\eta_1 \rho_1 \otimes \mathrm{Sp}(d_1) \oplus \cdots \oplus \eta_m \rho_m \otimes \mathrm{Sp}(d_m) : \eta_1, \dots, \eta_m \in \Psi(W_K)\}$$

will be called the *orbit* of ρ' under the action of

$$\Psi(W_K) \times \cdots \times \Psi(W_K)$$

(m factors). This orbit will be denoted $\mathfrak{X} = \mathfrak{X}(\rho')$. The orbits create a partition of $\mathcal{G}_n(K)$. The set $\mathcal{G}_n(K)$ is a disjoint union of orbits:

$$\mathcal{G}_n(K) = \bigsqcup \mathfrak{X}$$

We note that $\Psi(W_K)^m \simeq (\mathbb{C}^\times)^m$, a complex torus.

Each irreducible representation ρ of W_K has a *torsion number*: the order of the cyclic group of all those unramified characters η of W_K for which $\rho \otimes \eta \cong \rho$. The torsion number of ρ_i will be denoted τ_i . To determine the structure of each orbit, we have to pay attention to the torsion numbers of ρ_1, \dots, ρ_m and to the action of $\mathrm{GL}_n(\mathbb{C})$ by conjugation.

Let $\mu_\tau \subset \mathbb{C}$ denote the cyclic group of order τ . The orbit $\mathcal{O}(\rho')$ will have the structure of the *quotient* torus

$$(4) \quad \Psi(W_K) \times \cdots \times \Psi(W_K) / (\mu_{\tau_1} \times \cdots \times \mu_{\tau_m})$$

modulo a product of symmetric groups. We will write z_1, \dots, z_m for coordinates on this quotient torus. To be precise, we are writing z_j as a coset:

$$(5) \quad z_j = \eta_j(\Phi_K) \cdot \mu_{\tau_j}$$

In this way, the set $\mathcal{G}_n(K)$ acquires (locally) the structure of complex algebraic variety. Each connected component in $\mathcal{G}_n(K)$ is the quotient of a complex torus by a product of symmetric groups.

We recall that, given an irreducible representation V of W_K , there exists an irreducible representation V^{Gal} of Galois type such that $V = V^{\text{Gal}} \otimes \omega_s$ for some $s \in \mathbb{C}$, see [Tate, (2.2.1)].

This allows us to view the orbit $\mathfrak{X}(\rho')$ as a pointed complex algebraic variety, with base point

$$\rho' = \rho_1 \otimes \text{Sp}(d_1) \oplus \cdots \oplus \rho_m \otimes \text{Sp}(d_m).$$

where each ρ_j is of Galois type.

3. THE FORMULAS

The elementary substitution referred to in the Introduction is as follows. Let

$$\varepsilon_K(s, V, \psi) = \varepsilon_K(V \otimes \omega_{s-1/2}, \psi)$$

for all $s \in \mathbb{C}$, see [Tate, (3.6.4)], [L, p.6]. For $s \in \mathbb{C}$, $\omega_s : W_K \rightarrow \mathbb{C}^\times$ is the unramified quasicharacter defined by $\omega_s(w) = \|w\|_K^s$ for all $w \in W_K$. To compare this quasicharacter with those in §2, note the following:

$$\begin{aligned} \omega_s(w) &= \|w\|_K^s \\ &= q_K^{-s \cdot \text{val}_K(\text{Art}_K(w))} \\ &= z^{d(w)} \end{aligned}$$

with $z = q_K^{-s} \in \mathbb{C}^\times$.

If V is a 1-dimensional continuous complex representation of W_K , and $\chi : W_K \rightarrow \mathbb{C}^\times$ is the corresponding quasicharacter, then $\varepsilon_K(\chi, \psi)$ is the abelian local constant of Tate, see [Tate, (3.6.3)].

We note that $\varepsilon_K(V, \psi)$ is denoted $\varepsilon_K^{\text{Langlands}}(V, \psi)$ in [Ikeda] and $\varepsilon_L(V, \psi)$ in [Tate, 3.6].

We recall that, if (V, N) is any Φ -semisimple Weil-Deligne representation, then we have a finite direct sum decomposition of (V, N) into indecomposable Weil-Deligne representations as follows:

$$(6) \quad (V, N) = V_1 \otimes \text{Sp}(d_1) \oplus \cdots \oplus V_m \otimes \text{Sp}(d_m)$$

We will write

$$V_j = V_j^{\text{Gal}} \otimes \omega_{s_j}.$$

Lemma 3.1. *Let $a(V)$ denote the Artin conductor exponent of V . Then we have $a(V \otimes \omega_s) = a(V)$.*

Proof. The definition is

$$(7) \quad a(V) = \dim V - \dim V^I + \sum_{k \geq 1} \frac{1}{[I : I_k]} \cdot \dim V/V^{I_k}$$

where $I = I_0 \supset I_1 \supset \dots \supset I_k \supset \dots$ are the ramification subgroups of the inertia group I .

We have

$$\dim(V \otimes \omega_s) = \dim V$$

Now ω_s is an unramified quasi-character of W_K :

$$\omega_s(I) = \|\mathrm{Art}_K(I)\|^s = \|U_K\|^s = 1$$

and so

$$(V \otimes \omega_s)^{I_k} = V^{I_k}$$

for all $k \geq 0$. The result now follows from (7). \square

In particular, we have

$$a(V_j) = a(V_j^{\mathrm{Gal}}).$$

We need the following three items in order to compute epsilon factors.

3.1. Additivity. Additivity with respect to V , see [Tate, 3.4.2], [L, Theorem A (ii)]:

$$(8) \quad \varepsilon_K(V_1 \oplus \dots \oplus V_k, \psi) = \varepsilon_K(V_1, \psi) \dots \varepsilon_K(V_k, \psi)$$

3.2. Unramified twist. Behaviour under unramified twist, see [Tate, 3.4.5], [L, Lemma 22.4]:

$$(9) \quad \varepsilon_K(V \otimes \omega_s, \psi) = \varepsilon_K(V, \psi) q^{-s[a(V)+n(\psi) \dim V]}$$

where $a(V)$ is the Artin conductor exponent of V , and $n(\psi)$ is the conductor of ψ .

3.3. The extension formula. The extension to Weil-Deligne representations is as follows [Tate, 4.1.6]:

$$(10) \quad \varepsilon_K((V, N), \psi) := \varepsilon_K(V, \psi) \det(-\Phi|V^I/V_N^I)$$

3.4. The term $\varepsilon_K(V, \psi)$. A typical direct summand of (6) as a representation of W_K is

$$V_j^{\mathrm{Gal}} \otimes \omega_{s_j} \otimes \omega_k$$

with $1 \leq j \leq m$, $0 \leq k \leq d_j - 1$. We have

$$V_j^{\mathrm{Gal}} \otimes \omega_{s_j} \otimes \omega_k = V_j^{\mathrm{Gal}} \otimes \omega_{s_j+k}$$

For this summand, we have by (9)

$$(11) \quad \varepsilon_K(V_j^{\mathrm{Gal}} \otimes \omega_{s_j} \otimes \omega_k, \psi) = \varepsilon_K(V_j^{\mathrm{Gal}}, \psi) q^{-(s_j+k)[a(V_j^{\mathrm{Gal}})+n(\psi) \dim V_j^{\mathrm{Gal}}]}$$

and then the formula for $\varepsilon_K(V, \psi)$ follows from (8). Applying Lemma 3.1, we obtain

$$(12) \quad \varepsilon_K(V, \psi) = \prod_{j=1}^m \left(\varepsilon_K(V_j^{\mathrm{Gal}}, \psi) \right)^{d_j} \cdot q^{-[s_j d_j + \frac{(d_j-1)d_j}{2}][a(V_j)+n(\psi) \dim(V_j)]}$$

Note that Ikeda succeeds in describing the numbers $\varepsilon_K(V_j^{\mathrm{Gal}}, \psi)$ in terms of the non-abelian local class field theory of K , see [Ikeda, Theorem 5.4].

3.5. **The determinant.** The determinant is additive

$$\det(A \oplus B) = (\det A)(\det B)$$

and so it suffices to consider a typical factor in (10), namely

$$(13) \quad \det(-\Phi|E_j^I/(E_j^I)_{N_j})$$

where E_j is the W_K -module given by

$$\begin{aligned} E_j &= V_j \otimes (\omega_0 \oplus \omega_1 \oplus \cdots \oplus \omega_{d_j-1}) \\ &= (V_j^{\text{Gal}} \otimes \omega_{s_j}) \otimes (\omega_0 \oplus \omega_1 \oplus \cdots \oplus \omega_{d_j-1}) \\ &= V_j^{\text{Gal}} \otimes (\omega_{s_j} \oplus \omega_{s_j+1} \oplus \cdots \oplus \omega_{s_j+d_j-1}) \end{aligned}$$

We note that $V_j^I = (V_j^{\text{Gal}})^I$. Then the W_K -submodule fixed by the inertia group I is

$$E_j^I := V_j^I \otimes (\omega_{s_j} \oplus \omega_{s_j+1} \oplus \cdots \oplus \omega_{s_j+d_j-1})$$

The W_K -submodule of E_j annihilated by N_j is

$$(E_j)_{N_j} = V_j^{\text{Gal}} \otimes \omega_{s_j+d_j-1}$$

from which it follows that

$$(E_j)_{N_j}^I = V_j^I \otimes \omega_{s_j+d_j-1}$$

For the quotient we have the following W_K -module:

$$(14) \quad E_j^I/(E_j)_{N_j}^I \simeq V_j^I \otimes (\omega_{s_j} \oplus \cdots \oplus \omega_{s_j+d_j-2})$$

Recall that

$$\omega_s(\Phi) = \|\varpi_K\|^s = q_K^{-s}$$

It is enough to compute the action of $-\Phi$ on the W_K -module $V_j^I \otimes \omega_{s_j} \otimes \omega_k$ with $0 \leq k \leq d_j - 1$. On this W_K -module, $-\Phi$ will act as

$$q^{-(s_j+k)}(-\Phi|V_j^I)$$

and the determinant will be

$$q^{-(s_j+k) \dim V_j^I} \det(-\Phi|V_j^I)$$

There are $d_j - 1$ direct summands in (14) so the resulting determinant will be the product

$$(15) \quad \prod_{k=0}^{d_j-2} q^{-(s_j+k) \dim V_j^I} \cdot \det(-\rho_j(\Phi)|V_j^I)$$

$$(16) \quad = \det(-\rho_j(\Phi)|V_j^I)^{d_j-1} \cdot q^{-s_j(d_j-1) \dim V_j^I} \cdot q^{-(1+2+\dots+d_j-2) \dim V_j^I}$$

$$(17) \quad = \det(-\rho_j(\Phi)|V_j^I)^{d_j-1} \cdot q^{-s_j(d_j-1) \dim V_j^I} \cdot q^{-\frac{1}{2}(d_j-2)(d_j-1) \dim V_j^I}$$

provided that $d_j \geq 3$. By inspection, this formula is also valid for $d_j = 1$ or 2.

3.6. The term $\varepsilon_K((V, N), \psi)$. We recall the discussion in §2 of the quotient torus (4), especially the definition (5) of the torus coordinates z_1, \dots, z_m :

$$\begin{aligned} z_j &= \omega_{s_j}(\Phi_K) \cdot \mu_{\tau_j} \\ &= q_K^{-s_j} \cdot \mu_{\tau_j} \end{aligned}$$

where τ_j is the torsion number of V_j^{Gal} and $\mu_{\tau_j} \subset \mathbb{C}$ is the cyclic group of order τ_j .

From the extension formula (10) we infer that $\varepsilon((V, N, \psi))$ is the product of (12) and (15). This product is of the form

$$(18) \quad \text{const} \cdot z_1^{\beta_1} \cdots z_m^{\beta_m}$$

where

$$\beta_j = (d_j - 1) \dim V_j^I + d_j[a(V_j) + n(\psi) \dim(V_j)]$$

for all $1 \leq j \leq m$. Note that β_j is an *integer*:

$$\beta_j \in \mathbb{Z}.$$

The constant can be read off from (12) and (15). Apart from the constant term, the formula (18) for the epsilon factor is a *rational character* of the quotient torus (a complex torus of dimension m):

$$(z_1, \dots, z_m) \mapsto z_1^{\beta_1} \cdots z_m^{\beta_m}$$

Consider the following set:

$$(19) \quad \{\omega_{s_1} \otimes V_1^{\text{Gal}} \otimes \text{Sp}(d_1) \oplus \cdots \oplus \omega_{s_m} \otimes V_m^{\text{Gal}} \otimes \text{Sp}(d_m) : s_1, \dots, s_m \in \mathbb{C}\}.$$

After allowing for conjugacy in the Langlands dual group $\text{GL}_n(\mathbb{C})$, this set has the structure of a complex algebraic variety \mathfrak{X} in \mathcal{G}_n . In fact \mathfrak{X} is a connected component in $\mathcal{G}_n(K)$:

$$\mathfrak{X} \subset \mathcal{G}_n(K)$$

Applying the local Langlands correspondence, we have, by transport of structure, a connected component in the smooth dual:

$$\text{rec}_K^{-1}(\mathfrak{X}) \subset \mathbf{Irr}(\text{GL}_n(K))$$

We emphasize that this imposes a topology on $\mathbf{Irr}(\text{GL}_n(K))$ which is finer than the standard (Zariski) topology in representation theory. Our topology on the smooth dual is well-adapted to the study of the Hecke algebra $\mathcal{H}(\text{GL}_n(K))$ (see Theorem 3 in [BP]), and is, indeed, well-adapted to the study of epsilon factors, the subject of this article.

Looking carefully at the formulas (12) and (15), we see that the constant in (18) depends on the variety \mathfrak{X} , the choice of base point of \mathfrak{X} , and the additive character ψ . We will denote this constant by $e(\mathfrak{X}, \psi)$, so that (18) can be re-written

$$(20) \quad e(\mathfrak{X}, \psi) \cdot z_1^{\beta_1} \cdots z_m^{\beta_m}$$

which, up to the constant $e(\mathfrak{X}, \psi)$, factors as a rational character through \mathfrak{T} , in the notation of (2). The constant $e(\mathfrak{X}, \psi)$ is itself the product of

$$(21) \quad \prod_{j=1}^m \left(\varepsilon_K(V_j^{\text{Gal}}, \psi) \right)^{d_j} \cdot q^{-\frac{1}{2}(d_j-1)d_j[a(V_j)+n(\psi)\dim(V_j)]}$$

with

$$(22) \quad \prod_{j=1}^m \det(-\rho_j(\Phi)|V_j^I)^{d_j-1} \cdot q^{-\frac{1}{2}(d_j-2)(d_j-1)\dim V_j^I}$$

This establishes our main result, Theorem 1.1, which could perhaps be useful in the geometric Langlands program.

The terms $\varepsilon_K(V_j^{\text{Gal}}, \psi)$ are the epsilon factors attached to irreducible representations of the local absolute Galois group G_K . These terms are defined in [Ikeda, p.15]. There is one case where they are readily computed.

Lemma 3.2. *Let ψ be an additive character $K \rightarrow \mathbb{C}^\times$. Then we have $\varepsilon_K(1, \psi) = 1$.*

Proof. We start with the classical formula in [Tate, 3.6.3]:

$$\varepsilon_K(\chi, \psi) = \chi(c) \frac{\int_{\mathcal{O}^\times} \chi^{-1}(u)\psi(u/c) du}{\left| \int_{\mathcal{O}^\times} \chi^{-1}(u)\psi(u/c) du \right|}$$

where c is an element of K^\times of valuation $a(\chi) + n(\psi)$. Now set $\chi = 1$ and let $n(\psi) = k$. Then we take $c = \varpi^k$. Then $u \in \mathcal{O}^\times \implies u/c \in \varpi^{-k}\mathcal{O}^\times$. But we have $\psi(\varpi^{-k}\mathcal{O}) = 1$ since ψ has conductor k . Therefore we have

$$\begin{aligned} \varepsilon_K(1, \psi) &= \frac{\int_{\mathcal{O}^\times} \psi(u/c) du}{\left| \int_{\mathcal{O}^\times} \psi(u/c) du \right|} \\ &= \frac{\text{vol}(\mathcal{O}^\times)}{|\text{vol}(\mathcal{O}^\times)|} \\ &= 1 \end{aligned}$$

□

4. THE ARITHMETICALLY UNRAMIFIED REPRESENTATIONS OF $\text{GL}_n(K)$

Here, the underlying representation of the Weil group is the trivial n -dimensional representation $\rho : W_K \rightarrow \text{GL}_n(\mathbb{C})$. So we have $V_j^{\text{Gal}} = 1, 1 \leq j \leq n$.

Let W be the Weyl group \mathfrak{S}_n . The arithmetically unramified representations of $\text{GL}_n(K)$ have, by definition, the following set of Langlands parameters (Weil-Deligne representations):

$$(23) \quad \{\omega_{s_1} \otimes \text{Sp}(d_1) \oplus \cdots \oplus \omega_{s_k} \otimes \text{Sp}(d_k) : s_j \in \mathbb{C}\}$$

where $d_1 + \cdots + d_k = n$. This set determines a complex algebraic variety \mathfrak{X} in $\mathcal{G}_n(K)$.

We choose ψ to have conductor 0, and now apply Lemma 3.2. In this case $\beta_j = d_j - 1$. We have

$$\begin{aligned} \varepsilon_K((V, N, \psi)) &= e(\mathfrak{X}, \psi) \prod_{j=1}^m q^{-(d_j-1)s_j} \\ &= e(\mathfrak{X}, \psi) \prod_{j=1}^m z_j^{d_j-1} \end{aligned}$$

where

$$e(\mathfrak{X}, \psi) = \prod_{j=1}^m (-1)^{d_j-1} q^{-(d_j-1)(d_j-2)/2}$$

and $z_j := q^{-s_j}$.

The epsilon factor records the dimensions d_j of the special representations $\mathrm{Sp}(d_j)$ which occur in the Weil-Deligne representation (V, N) .

We will now re-organise the partition $d_1 + \dots + d_k = n$. Suppose that this partition has distinct parts t_1, \dots, t_m with $t_1 < t_2 < \dots < t_m$ and that t_j is repeated r_j times so that

$$(24) \quad r_1 t_1 + \dots + r_m t_m = n.$$

Then, as a function on the complex torus $(\mathbb{C}^\times)^{r_1 + \dots + r_m}$, the epsilon factor is *invariant under the following product of symmetric groups*:

$$\mathfrak{S}_{r_1} \times \mathfrak{S}_{r_2} \times \dots \times \mathfrak{S}_{r_m}$$

and therefore factors through the following quotient variety

$$(\mathbb{C}^\times)^{r_1} / \mathfrak{S}_{r_1} \times \dots \times (\mathbb{C}^\times)^{r_m} / \mathfrak{S}_{r_m}$$

Let T denote the standard maximal torus in the Langlands dual group $\mathrm{GL}_n(\mathbb{C})$, and let W be the Weyl group of $\mathrm{GL}_n(\mathbb{C})$. We have the *inertia space*

$$\mathcal{I}_W(T) := \{(w, t) : w \in W, t \in T, wt = t\}$$

Then $\mathcal{I}_W(T)$ admits an action of W , namely $\alpha \cdot (w, t) = (\alpha w \alpha^{-1}, \alpha \cdot t)$. We define $T//W$ to be the quotient:

$$T//W := \mathcal{I}_W(T)/W$$

This is the *extended quotient* of T by W , sometimes called the *inertia stack*, see [JM, §1.1].

Let T^w denote the w -fixed set, and let $Z(w)$ be the W -centralizer of w . Choose one w in each W -conjugacy class, then we have

$$T//W = \bigsqcup T^w / Z(w).$$

The conjugacy classes in the symmetric group W are in bijection with the partitions of n . The partition in (24) determines the permutation comprising

r_1 cycles of length t_1, \dots, r_m cycles of length t_m . If this permutation is denoted w , then we have

$$T^w = (\mathbb{C}^\times)^{r_1} \times \dots \times (\mathbb{C}^\times)^{r_m}$$

The centralizer $Z(w)$ is a product of wreath products

$$\mathbb{Z}/t_1\mathbb{Z} \wr \mathfrak{S}_{r_1} \times \dots \times \mathbb{Z}/t_m\mathbb{Z} \wr \mathfrak{S}_{r_m}$$

but the cyclic groups act trivially and so we have

$$T^w/Z(w) = (\mathbb{C}^\times)^{r_1}/\mathfrak{S}_{r_1} \times \dots \times (\mathbb{C}^\times)^{r_m}/\mathfrak{S}_{r_m}$$

Every irreducible component in $T//W$ is accounted for in this way. The epsilon factors have precisely the amount of symmetry required to factor through these quotient varieties.

EXAMPLE. Here, we consider the following Weil-Deligne representation of $\mathrm{GL}_{19}(K)$:

$$\omega_{s_1}\mathrm{Sp}(2) \oplus \omega_{s_2}\mathrm{Sp}(2) \oplus \omega_{s_3}\mathrm{Sp}(2) \oplus \omega_{s_4}\mathrm{Sp}(3) \oplus \omega_{s_5}\mathrm{Sp}(3) \oplus \omega_{s_6}\mathrm{Sp}(7)$$

The epsilon factor of this representation is

$$\mathrm{const} \cdot z_1 z_2 z_3 z_4^2 z_5^2 z_6^6$$

which will factor through the following irreducible component of the extended quotient $T//W$:

$$\mathrm{Sym}^3(\mathbb{C}^\times) \times \mathrm{Sym}^2(\mathbb{C}^\times) \times \mathbb{C}^\times$$

This perfectly illustrates the symmetry properties of the epsilon factors. Each epsilon factor has precisely the symmetry, *neither more nor less*, of the corresponding irreducible component in the extended quotient $T//W$. Each epsilon factor will therefore factor through the corresponding irreducible component in $T//W$.

5. THE GENERAL CASE

Let G be a reductive p -adic group and let ${}^L G$ be the L -group

$${}^L G = G^\vee \rtimes W_K$$

where G^\vee is the complex dual group of G . Let

$$r : {}^L G \rightarrow \mathrm{GL}(V)$$

be a representation of ${}^L G$ on the complex vector space V , as in [Bor, 2.6]. Let W'_K denote the Weil-Deligne group $W_K \rtimes \mathbb{C}$, defined by $w.z = ||w||z$. As in [Bor, 8.2], let ϕ be an L -parameter for G :

$$\phi : W'_K \rightarrow {}^L G$$

Following [Bor, 12.1], we have the composite

$$r \circ \phi : W'_K \rightarrow \mathrm{GL}(V)$$

The representation $r \circ \phi$ determines, and is determined by, a Weil-Deligne representation (V, N) . So we obtain an epsilon factor $\varepsilon_K((V, N), \psi)$ to which we may apply all the preceding material.

REFERENCES

- [ABPS] A.-M. Aubert, P. Baum, R.J. Plymen, M. Solleveld, Smooth duals of inner forms of GL_n and SL_n , *Doc. Math.* 24 (2019), to appear.
- [Bor] A. Borel, Automorphic L -functions, *Automorphic Forms, Representations and L -functions*, Proc. Symp. Pure Math. 33 part 2 (1979) 27– 61.
- [BP] J. Brodzki, R.J. Plymen, Geometry of the smooth dual of $GL(n)$, *C.R. Acad. Sci. Paris* 331 (2000) 213 - 218.
- [D] P. Deligne, Les constantes des équations fonctionnelles des fonctions L , *Springer Lecture Notes* 349 (1973) 501 - 595.
- [HT] M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Ann. Math. Study, Princeton, 2001.
- [Ikeda] K.I. Ikeda, On the ϵ -factors of Weil-Deligne representations, preprint 2016.
- [L] R. P. Langlands, On the functional equation of the Artin L -functions, unpublished article, <https://publications.ias.edu/sites/default/files/a-ps.pdf>.
- [MP] Yu.I. Manin, A.A. Panchishkin, *Introduction to modern number theory*, Encyclopaedia of Mathematical Sciences 49, Springer, Berlin 2007.
- [JM] J. Morava, HKR characters and higher twisted sectors, *Contemporary Math.* 403 (2006) 143–152.
- [Sch] P. Scholze, The local Langlands correspondence for GL_n over p -adic fields, *Invent. Math.* 192 (2013) 663–715.
- [Tate] J. Tate, Number Theoretic Background, *Automorphic Forms, Representations and L -functions*, Proc. Symp. Pure Math. 33 part 2 (1979) 3 - 26.

SCHOOL OF MATHEMATICS, MANCHESTER UNIVERSITY, MANCHESTER M13 9PL, UK
E-mail address: roger.j.plymen@manchester.ac.uk