Permafrost and periglacial processes in mid- and low-latitude mountains regions

DOI:
[10.1002/ppp.2034](https://doi.org/10.1002/ppp.2034)

Document Version
Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Ruiz-fernández, J., Oliva, M., & Hughes, P. (2019). Permafrost and periglacial processes in mid- and low-latitude mountains regions. *Permafrost and Periglacial Processes*. <https://doi.org/10.1002/ppp.2034>

Published in:
Permafrost and Periglacial Processes

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Permafrost and periglacial processes in mid- and low-latitude mountains regions

Journal:	<i>Permafrost and Periglacial Processes</i>
Manuscript ID	PPP-19-0068
Wiley - Manuscript type:	Editorial
Date Submitted by the Author:	04-Oct-2019
Complete List of Authors:	Ruiz-Fernández, Jesús; Department of Geography, University of Oviedo Oliva, Marc; University of Barcelona, Physical Geography Hughes, Phil; University of Manchester, Geography
Keywords:	Periglacial processes, mid- and low-latitude mountains, Permafrost

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 WILEY

2 *Permafrost and Periglacial Processes (PPP)*

3 Permafrost and
4 Periglacial
5 Processes

6 **Permafrost and periglacial processes in mid- and low-latitude mountains regions**

7 Jesús Ruiz-Fernández¹ | Marc Oliva² | Philip Hughes³

8 ¹ Department of Geography, University of Oviedo, Spain

9 ² Department of Geography, Universitat de Barcelona, Catalonia, Spain

10 ³ Department of Geography, School of Environment, Education and Development, University of
11 Manchester, Manchester, UK

12 **Short title:** Periglacial processes in mid- and low-latitude mountains

13
14 **Correspondence:** Jesús Ruiz-Fernández, Department of Geography, University of
15 Oviedo. C/ Amparo Pedregal s/n, 33011, Oviedo, Asturias, Spain.

16 Email: ruizjesus@uniovi.es

17
18 **ABSTRACT**

19 The Special Issue that justifies this Editorial has been designed to commemorate the 25th
20 anniversary of the 1st Spanish scientific meeting of the International Permafrost
21 Association (IPA) organized in 1994 in Madrid. This Special Issue entitled “Permafrost
22 and periglacial processes in mid- and low-latitude mountains regions” includes nine
23 papers from various mountain regions of the globe, such as the Pyrenees, Sierra Nevada,
24 Galician and Cantabrian Mountains in Iberian Peninsula, Atacama and Central Andes in
25 South America, Absaroka ranges in North America, and Seckauer Tauern Range in the
26 European Alps. These articles provide new methodologies and approaches focusing on a

wide range of periglacial phenomena, such as past cryogenic environments, active periglacial processes, permafrost and ground thermal regime, as well as rock glaciers. As a result of this Special Issue, we have detected several knowledge gaps that should be addressed in the future by the scientific community studying permafrost and periglacial processes in mid- and low-latitude mountains regions, namely: i) improving the geochronology of past periglacial environments and associated paleoclimatic implications; ii) the study of periglacial features existing in low-altitude mountain environments and/or in the lowlands; iii) improve our current knowledge of active periglacial processes, permafrost distribution and ground thermal regime in mid-altitude mountains; iv) the monitoring of periglacial mass wasting processes and mechanisms of sediment transfer; v) the interaction between glaciers and periglacial processes, and vi) geoeological dynamics in response to climate scenarios anticipating significant changes of temperature and moisture in periglacial regions.

KEYWORDS

Permafrost, periglacial processes, mid- and low-latitude mountains

1 | INTRODUCTION

Climate and environmental changes condition Earth's ecosystems in a great variety of ways in cold-climate regions. The most commonly used records to reconstruct environmental and climatic conditions are marine and lake sediments and ice cores¹⁻⁴. The evolution of glaciers, which also constitute an important source of climate and environmental information, has also been extensively studied at various scales and perspectives⁵⁻⁹. Conversely, the study of periglacial environments has received, in general, less attention from the scientific community, except in the case of some specific

1
2
3 52 topics (such as permafrost¹⁰⁻¹³ and rock glaciers^{10,14-16}). However, their study is crucial to
4
5 53 understand past and present-day dynamics of large areas of the Earth's surface as 25% of
6
7 54 the land surface is currently affected by periglacial conditions¹⁰. The geomorphology,
8
9
10 55 hydrology, soil formation and ecology of these areas is strongly conditioned by the
11
12 56 presence of seasonal or permanent ground ice conditions^{17,18}.
13
14
15 57
16
17 58 Research on permafrost and periglacial geomorphology has mainly focused on polar
18
19 59 regions (mainly in the Arctic^{15,19}, but also in Antarctica^{11,13,20}), as well as in the world's
20
21 60 major mountain ranges, many of them located in mid-latitude mountain regions (Alps^{12,21-}
22
23 61 ²², Andes^{23,24}, Himalaya²⁵, Pyrenees²⁶, etc.). The magnitude of periglacial processes and
24
25 62 associated landforms that exist in these areas have led to a greater development of
26
27 63 periglacial research in these sectors, compared to other environments. This has also been
28
29 64 encouraged because of the pressing problems that occur in many of these regions in
30
31 65 relation to the degradation of permafrost and related mass wasting processes^{19,27}, as well
32
33 66 as the logistical facilities provided by polar and alpine research bases.
34
35
36
37 67
38
39 68 However, to fully understand the dynamics of periglacial environments, it is necessary to
40
41 69 extend research to other mountainous areas that are much less studied. This is often due
42
43 70 to their lower altitude/latitude and more marginal character compared to the classic
44
45 71 periglacial areas of the world. Likewise, it is not only necessary to examine the present-
46
47 72 day periglacial processes and associated landforms, but it is also important to improve
48
49 73 our understanding of the periglacial paleoenvironments.
50
51
52 74
53
54
55 75 This Special Issue has been designed to commemorate 25 years since the first Spanish
56
57 76 scientific meeting of the International Permafrost Association (IPA). Dr D. Palacios and
58
59
60

Dr A. Gómez-Ortiz organized this meeting in 1994 in Madrid. In the same year, the IPA-Spain group was created and integrated as an associate member of the IPA^{28,29}. Since then, six IPA-Spain meetings have been organized and, from 2007 (with the creation of the IPA-Portugal group in 2005²⁹), the IPA sections of both countries have jointly organized seven Iberian congresses^{28,29}. The last congress was held in Jaca (Spanish Pyrenees) in June 2019. Throughout these 25 years a thriving network of scientific collaboration has been developed amongst the researchers integrated in the Iberian section of the IPA²⁹. In addition to the aforementioned meetings, this scientific collaboration has led to the publication of several monographic books and special issues in scientific journals²⁸. For example, between 2000 and 2017, 230 scientific publications have been counted within the Iberian network of the IPA²⁹. These publications involved 198 researchers belonging to 64 different institutions (29 Spanish, 5 Portuguese and 34 from other countries), allowing a considerable advance in the knowledge of the Cryosphere of the Iberian Peninsula²⁹⁻³¹, but also of the polar regions (where Iberian researchers are making significant contributions^{13,18,20,27}), and in many other mountainous of the Earth^{24,32-34}.

Thus, with the purpose of better understanding permafrost and periglacial processes in mid- and low-latitude mountains regions, this Special Issue includes nine studies focusing on a wide and diverse range of specific topics that we have synthesized in three main sections: i) periglacial paleoenvironments, ii) active periglacial processes, permafrost and ground thermal regime, and iii) rock glaciers.

2 | PERIGLACIAL PALEOENVIRONMENTS

1
2
3 101 The study of paleoenvironments from different perspectives represents one of the classic
4
5 102 topics of periglacial research. Two articles focused on this topic have been included in
6
7 103 this Special Issue. Rodríguez-Ochoa et al.³⁵ study relict periglacial soils on Quaternary
8
9 104 terraces in the NE sector of the Iberian peninsula, through geochronological,
10
11 105 micromorphological and physico-chemical methods. The main results show the existence
12
13 106 of various features related to cryogenic processes in the soil profiles.
14
15
16
17
18

19 108 Viana-Soto and Alberti³⁶ reconstruct paleotemperatures in northwest Iberia based on
20
21 109 periglacial geomorphological evidence, such as block fields, block slopes, rock glaciers
22
23 110 and stratified slope deposits. During the coldest part of the Late Pleistocene between 21
24
25 111 and 25 ka cal BP, permafrost and genesis of block fields, block slopes and rock glaciers
26
27 112 occurred at elevations above 700 m. Prior to this, before 30 ka cal BP, the prevailing snow
28
29 113 conditions favored the formation of stratified slope deposits and genesis of glaciers in all
30
31 114 of the mountain systems, although without the presence of permafrost.
32
33
34
35
36

37
38 116 **3 | ACTIVE PERIGLACIAL PROCESSES, PERMAFROST AND GROUND**
39
40 117 **THERMAL REGIME**

41
42 118 The relationship between periglacial processes and soil thermal regime is one of the topics
43
44 119 that has been more intensely studied over the last decades. In this Special issue, a number
45
46 120 of works have also focused on the monitoring of periglacial dynamics and the activity of
47
48 121 periglacial landforms. This is the case of the research article by Gómez-Ortiz et al.³⁷,
49
50 122 focusing on Sierra Nevada, one of the best studied massifs with regards to its past
51
52 123 environmental history and current periglacial dynamics. The authors show evidence
53
54 124 widespread periglacial conditions in this massif above 2500 m, although permafrost is
55
56 125 only present in glaciated environments during the Little Ice Age.
57
58
59
60

Gjorup et al.³⁸ studied the local variability of soil temperature and moisture at different depths in three sites along a altitudinal gradient in the Licancabur Volcano (Atacama, northern Chile). Temperature profiles of all study sites are characterized by absence of permafrost, abundant freeze-thaw cycles, greater thermal variations close to soil surface year round, and most soil temperatures readings close to 0°C.

In the eastern margin of the European Alps Kellerer-Pirklbauer³⁹ investigated the characteristics and distribution of sporadic permafrost. The main results of this work indicate sporadic permafrost occurrence in the summit sector of the study area, with mean annual temperatures close to 0°C at the soil surface, and -1.4°C at 2.5 m depth, as well as the existence of permafrost patches in the transition area between rock glaciers and talus cones. From the projection of statistically significant trends, this work highlights the presumably disappearance of permafrost in the next decades in part of the sites studied.

Tapia Baldis and Liaudat⁴⁰ examine the evidence of rockslides and rock avalanches in the Central Andes of Argentina and investigate their possible association with permafrost degradation. Over the long-term permafrost degradation favors a deeper failure process, whereas in the shorter term shallow active layer detachment is favored by shear-displacement along pre-existing joints, as a result of short-term periods of climate warming. It is expected that the magnitude and frequency of rockslide hazards will increase during the 21st century.

Serrano et al.⁴¹ provide a comprehensive review of the periglacial environments and frozen ground in four massifs of the Central Pyrenean high mountain area (Infierno-

Argualas, Posets, Maladeta and Monte Perdido). In these areas, the lower limit of frozen ground is at ~2,650 m a.s.l. and permafrost is dominant above 2,900 m a.s.l. Future projected changes in temperature and precipitation are likely to have a significant influence on high mountain the snow cover and mountain permafrost in this areas.

4 | ROCK GLACIERS

Rock glaciers are probably the best indicators of permafrost conditions in mid- and low-latitude mountains regions³⁰. In relict features, the analysis of their morphometric dimensions can reveal patterns of past activity³¹. In this Special Issue, González-Gutiérrez et al.⁴² examined the surface macro-fabrics in relict rock glaciers in the Cantabrian Mountains (NW Spain) and shown evidence of the topographical and morphostructure controls on their formation and activity.

The scale of periglacial landscapes in the mountains of North America is highlighted in the paper by Seligman et al.⁴³, who identify 661 rock glaciers in the Beartooth and northern Absaroka ranges, Montana, USA. Landscapes in these areas are shifting from predominantly glacial to periglacial regimes. As glaciers retreat in response to climate warming, the authors show that rock glaciers could soon become the most important source of ice in the region.

5 | PERSPECTIVES

The nine papers included in this Special Issue cover different areas of mid- and low-latitude mountains regions, both of the Northern and Southern Hemispheres, namely Pyrenees, Sierra Nevada, Ebro Basin, Galician and Cantabrian Mountains in Iberian Peninsula, Atacama and Central Andes in South America, Absaroka ranges in Montana

(North America), and Seckauer Tauern Range in the European Alps. This Special Issue brings together important new information to help judge the effect of past and present-day periglacial morphodynamics in various mountains regions around the globe. Individual studies published here address wide-ranging topics in periglacial paleoenvironments, active periglacial processes, permafrost and ground thermal regime, and also rock glaciers.

These studies have also highlighted knowledge gaps, and many issues still require more attention, such as the need to:

- Substantially improve the knowledge of past periglacial environments, especially regarding geochronology and paleoclimatic implications.
- Examine the periglacial features developed in the lowlands and/or low altitude mountain environments of mid-latitude ranges, which are currently still poorly known.
- Improve our current knowledge of active periglacial processes, permafrost distribution and ground thermal regime in intermediate elevation belts of mid-latitude mountain areas, as well as in the high intertropical mountain ranges, including large stratovolcanoes.
- Increase the monitoring of periglacial mass wasting processes and mechanisms of sediment transfer in mid- and low-latitude mountain areas.
- Further understand the dynamic relationship between glaciers and the periglacial environment, which directly affect paraglacial processes and mechanisms of sediment transfer, especially under a changing future climate.

- Improve understanding of wider geocological dynamics in response to the climate scenarios anticipating higher temperatures and lower moisture conditions in most of these regions⁴⁴.

ACKNOWLEDGEMENTS

The guest editors appreciate the opportunity given by the journal Permafrost and Periglacial Processes, particularly by the editor-in-chief Prof Dr Mauro Guglielmin, to organize this Special Issue on Permafrost and Periglacial Processes in Mid- and Low-latitude Mountains Regions. We would like to congratulate all authors for the research papers submitted and acknowledge all reviewers for their work in reviewing the manuscripts. This work fits within the research topics of the CRONOANTAR (CTM2016-77878-P) and PALEOGREEN (CTM2017-87976-P) projects of the Spanish Ministry of Economy, Industry and Competitiveness, and the NUNANTAR project (02/SAICT/2017 - 32002) of the Fundação para a Ciência e a Tecnologia of Portugal. Marc Oliva is supported by the Ramón y Cajal Program (RYC-2015-17597) and the Research Group ANTALP (Antarctic, Arctic, Alpine Environments; 2017-SGR-1102) funded by the Government of Catalonia through the AGAUR agency.

REFERENCES

1. Petit JR, Jouzel J, Raynaud D, Barkov NI, Barnola JM, Basile I, Bender M, Chappellaz J, Davis M, Delaygue G, Delmotte M, Kotlyakov VM, Legrand M, Lipenkov VY, Lorius C, Pépin L, Ritz C, Saltzman E, Stievenard M. Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica. *Nature*. 1999;399:429-436.

- 224 2. de Vernal A, Hillaire-Marcel C. Natural Variability of Greenland Climate,
 225 Vegetation, and Ice Volume During the Past Million Years. *Science*.
 226 2008;320(5883):1622-1625. DOI: 10.1126/science.1153929
- 227 3. Oliva M, Antoniades D, Giralt S, Granados I, Pla-Rabes S, Toro M, Liu EJ,
 228 Sanjurjo J, Vieira G. The Holocene deglaciation of the Byers Peninsula
 229 (Livingston Island, Antarctica) based on the dating of lake sedimentary
 230 records. *Geomorphology*. 2016;261:89-102. DOI:
 231 10.1016/j.geomorph.2016.02.029
- 232 4. Antoniades D, Giralt S, Geyer A, Álvarez-Valero AM, Pla-Rabes S,
 233 Granados I, Liu EJ, Toro M, Smellie JL, Oliva M. The timing and widespread
 234 effects of the largest Holocene volcanic eruption in Antarctica. *Scientific*
 235 *Reports*. 2018;8:17279. DOI: 10.1038/s41598-018-35460-x
- 236 5. Hughes PD, Woodward JC, Gibbard PL. Quaternary glacial history of the
 237 Mediterranean mountains. *Progress in Physical Geography*.
 238 2006;30(3):334-364. DOI: 10.1191/0309133306pp481ra
- 239 6. Mackintosh A, Golledge N, Domack E, Dunbar R, Leventer A, White D,
 240 Pollard D, DeConto R, Fink D, Zwartz D, Gore D, Lavoie C. Retreat of the
 241 East Antarctic ice sheet during the last glacial termination. *Nature*
 242 *Geoscience*. 2011;4:195-202. DOI: 10.1038/ngeo1061
- 243 7. Nick FM, Vieli A, Howat IM, Joughin I. Large-scale changes in Greenland
 244 outlet glacier dynamics triggered at the terminus. *Nature Geoscience*.
 245 2009;2:110-114. DOI: 10.1038/ngeo394
- 246 8. Navarro FJ, Jonsell U, Corcuera Labrado MI, Martín Español A. Decelerated
 247 mass loss of Hurd and Johnsons Glaciers, Livingston Island, Antarctic

- 248 Peninsula. *Journal of Glaciology*. 2013;59(213):115-128. DOI:
249 10.3189/2013JoG12J144.
- 250 9. Palacios D, Gómez-Ortiz A, Andrés N, Salvador F, Oliva M. Timing and
251 new geomorphologic evidence of the last deglaciation stages in Sierra
252 Nevada (southern Spain). *Quaternary Science Reviews*. 2016;150:110-129.
253 DOI: 10.1016/j.quascirev.2016.08.012.
- 254 10. French H. *The Periglacial Environment*. 3rd edition. Wiley, Chichester, UK;
255 2007.
- 256 11. Bockheim JG. Permafrost distribution in the southern circumpolar region
257 and its relation to the environment: A review and recommendations for
258 further research. *Permafrost and Periglacial Processes*. 1995;6:27-45. DOI:
259 10.1002/ppp.3430060105
- 260 12. Boeckli L, Brenning A, Gruber S, Noetzli J. Permafrost distribution in the
261 European Alps: calculation and evaluation of an index map and summary
262 statistics. *The Cryosphere*. 2012;6(4):807-820.
- 263 13. Vieira G, Bockheim J, Guglielmin M, Balks M, Abramov AA, Boelhouwers
264 J, Cannone N, Ganzert L, Gilichinsky DA, Goryachkin S, López-Martínez J,
265 Meiklejohn I, Raffi R, Ramos M, Schaefer C, Serrano E, Simas F, Sletten
266 R, Wagner D. Thermal state of permafrost and active-layer monitoring in the
267 antarctic: Advances during the international polar year 2007–2009.
268 *Permafrost and Periglacial Processes*. 2010;21:182-197. DOI:
269 10.1002/ppp.685
- 270 14. Humlum O. The climatic significance of rock glaciers. *Permafrost and*
271 *Periglacial Processes*. 1998;9:375-395.

15. Humlum O. The geomorphic significance of rock glaciers: estimates of rock glacier debris volumes and headwall recession rates in West Greenland. *Geomorphology*. 2000;35(1-2):41-67.
16. Guglielmin M, Ponti S, Forte E. The origins of Antarctic rock glaciers: periglacial or glacial features? *Earth Surface Processes and Landforms*. 2018;43:1390-1402. DOI: 10.1002/esp.4320
17. Oliva M, Pereira P, Ruiz-Fernández J, Nieuwendam A. Recent advances in the study of active layer thermal regime and seasonal frost dynamics in cold climate environments. *Catena*. 2017;149:515-518. DOI: 10.1016/j.catena.2016.08.030
18. Ruiz-Fernández J, Oliva M, Nývlt D, Cannone N, García-Hernández C, Guglielmin M, Hrbáček F, Roman M, Fernández S, López-Martínez J, Antoniades D. Patterns of spatio-temporal paraglacial response in the Antarctic Peninsula region and associated ecological implications. *Earth-Science Reviews*. 2019;192:379-402. DOI: 10.1016/j.earscirev.2019.03.014
19. Fritz M, Vonk JE, Lantuit H. Collapsing Arctic coastlines. *Nature Climate Change*. 2017;7(1):6-7. DOI: 10.1038/nclimate3188.
20. López-Martínez J, Serrano E, Schmid T, Mink S, Linés C. Periglacial processes and landforms in the South Shetland Islands (northern Antarctic Peninsula region). *Geomorphology*. 2012;155-156:62-79.
21. Matsuoka N, Ikeda A, Date T. Morphometric analysis of solifluction lobes and rock glaciers in the Swiss Alps. *Permafrost and Periglacial Processes*. 2005;16:99-113.

22. Kellerer-Pirklbauer A, Lieb Gk, Kleinfierchner H. A new rock glacier inventory of the eastern European Alps. *Austrian Journal of Earth Sciences*. 2012;105(2):78-93.
23. Trombotto DT. Survey of cryogenic processes, periglacial forms and permafrost conditions in South America. *Revista Do Instituto Geologico*. 2000;21(1-2):33-55.
24. Valcárcel-Díaz M, Carrera-Gómez P, Coronato A, Castillo-Rodríguez F, Rabassa J, Pérez-Alberti A. Cryogenic landforms in the Sierras de Alvear, Fuegian Andes, subantarctic Argentina. *Permafrost and Periglacial Processes*. 2006;17:371-376.
25. Barsch D. Mass transport by active rock glaciers in the Khumbu Himalaya. *Geomorphology*. 1998;26(1-2):215-222. DOI: 10.1016/S0169-555X(98)00060-9
26. Lugon R, Delaloye R, Serrano E, Reynard E, Lambiel C, González-Trueba JJ. Permafrost and Little Ice Age glacier relationships, Posets Massif, Central Pyrenees, Spain. *Permafrost and Periglacial Processes*. 2004;15:207-220. DOI: 10.1002/ppp.494
27. Oliva M, Ruiz-Fernández J. Coupling patterns between paraglacial and permafrost degradation responses in Antarctica. *Earth Surface Processes and Landforms*. 2015;40(9):1227-1238. DOI: 10.1002/esp.3716
28. Serrano Cañadas E. Periglacialismo y permafrost. *Polígonos. Revista de Geografía*. 2016;28:15-29.
29. García-Hernández C, Ruiz-Fernández J, Serrano-Cañadas E. Social network analysis in Geosciences: scientific collaboration between periglacial scholars

- in the Iberian Peninsula. *Cuadernos de Investigación Geográfica*. DOI: 10.18172/cig.3939
30. Oliva M, Serrano E, Gómez-Ortiz A, González-Amuchastegui MJ, Nieuwendam A, Palacios D, Pérez-Alberti A, Pellitero-Ondicol R, Ruiz-Fernández J, Valcárcel M, Vieira G, Antoniades D. Spatial and temporal variability of periglaciation of the Iberian Peninsula. *Quaternary Science Reviews*. 2016;137:176-199. DOI: 10.1016/j.quascirev.2016.02.017
31. Fernandes, M., Palma, P., Lopes, L., Ruiz-Fernández, J., Pereira, P., Oliva, M. Spatial distribution and morphometry of permafrost-related landforms in the Central Pyrenees and associated paleoclimatic implications. *Quaternary International*. 2018;470(A):96-108.
32. Vieira G, Mora C, Ali F. New observations indicate the possible presence of permafrost in North Africa (Djebel Toubkal, High Atlas, Morocco). *Cryosphere*. 2017;11(4):1691-1705. DOI: 10.5194/tc-11-1691-2017
33. Oliva M, Žebre M., Guglielmin M., Hughes PD, Çiner A, Vieira G, Bodin X, Andrés N, Colucci RR, García-Hernández C, Mora C, Nofre J, Palacios D, Pérez-Alberti A, Ribolini A, Ruiz-Fernández J, Sarıkaya MA, Serrano E, Urdea P, Valcárcel M, Woodward JC, Yıldırım C. Permafrost conditions in the Mediterranean region since the Last Glaciation. *Earth-Science Reviews*. 2018;185:397-436. DOI: 10.1016/j.earscirev.2018.06.018
34. Oliva M, Ruiz-Fernández J. Late Quaternary environmental dynamics in Lenin Peak area (Pamir Mountains, Kyrgyzstan). *Science of the Total Environment*. 2018;645:603-614. DOI: 10.1016/j.scitotenv.2018.07.178
35. Rodríguez-Ochoa R, Olarieta JR, Santana A, Castañeda C, Calle M, Rhodes E, Bartolomé M, Peña-Monné JL, Sancho C. Relict periglacial soils on

1
2
3 344 Quaternary terraces in the Central Ebro Basin (NE Spain). *Permafrost and*
4
5 345 *Periglacial Processes*. 2019. DOI: 10.1002/ppp.2005 (in this issue).
6
7
8 346 36. Viana Soto A, Pérez-Alberti A. Periglacial deposits as indicators of
9
10 347 paleotemperatures. A case study in the Iberian Peninsula: the mountains of
11
12 348 Galicia. *Permafrost and Periglacial Processes*. 2019. DOI:
13
14 349 10.1002/ppp.2026 (in this issue).
15
16
17 350 37. Gómez A, Oliva M, Salvador-Franch S, Palacios D, Tanarro, LM.
18
19 351 Monitoring permafrost and periglacial processes in Sierra Nevada (Spain)
20
21 352 from 2001 to 2016. *Permafrost and Periglacial Processes*. 2019 (in this
22
23 353 issue), doi: 10.1002/PPP.2002.
24
25
26 354 38. Gjorup DF, Francelino MR, Michel RFM, Senra EO, Schaefer CEGR.
27
28 355 Pedoclimate monitoring in the periglacial high mountain soils of Atacama,
29
30 356 northern Chile. *Permafrost and Periglacial Processes*. 2019 (in this issue).
31
32
33 357 39. Kellerer-Pirklbauer A. Long-term monitoring of sporadic permafrost at the
34
35 358 eastern margin of the European Alps (Hochreichart, Seckauer Tauern Range,
36
37 359 Austria). *Permafrost and Periglacial Processes*. 2019. DOI:
38
39 360 10.1002/ppp.2021 (in this issue).
40
41
42 361 40. Tapia Baldis C, Trombotto D. Rockslides and rock avalanches in the Central
43
44 362 Andes of Argentina and their possible association with permafrost
45
46 363 degradation. *Permafrost and Periglacial Processes*. 2019. DOI:
47
48 364 10.1002/ppp.2024 (in this issue).
49
50
51 365 41. Serrano E, González González M, Gómez Lende M, López Moreno I,
52
53 366 Sanjosé Blasco JJ, González-Trueba JJ, Pisabarro A. Periglacial
54
55 367 environments and frozen ground in the Central Pyrenean high mountain area.
56
57
58
59
60

- 368 Ground thermal regime and distribution of landforms and processes.
369 *Permafrost and Periglacial Processes*. 2019 (in this issue).
- 370 42. González-Gutiérrez RB, Santos-González J, Gómez-Villar A, Redondo-
371 Vega JM. Surface Macro-Fabric Analysis of Relict Rock Glaciers in the
372 Cantabrian Mountains (NW Spain). *Permafrost and Periglacial Processes*.
373 2019 (in this issue).
- 374 43. Seligman ZM, Klene AE, Nelson FE. Rock glaciers of the Beartooth and
375 northern Absaroka ranges, Montana, USA. *Permafrost and Periglacial*
376 *Processes*. 2019. DOI: 10.1002/ppp.2019 (in this issue).
- 377 44. IPCC Climate Change 2013. The Physical Science Basis: Contribution of
378 Working Group I to the Fifth Assessment Report of the Intergovernmental
379 Panel on Climate Change. Cambridge University Press, Cambridge; 2013.