Transforming clinical trials in rheumatology

DOI:

[10.1038/s41584-020-0491-4](https://doi.org/10.1038/s41584-020-0491-4)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Pitzalis, C., Choy, E. H. S., & Buch, M. (2020). Transforming clinical trials in rheumatology: towards patient-centric precision medicine. *Nature Reviews Rheumatology*, 16(10), 590-599. <https://doi.org/10.1038/s41584-020-0491-4>

Published in:

Nature Reviews Rheumatology

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

CONSIDERATIONS FOR IMPROVING QUALITY OF CARE OF PATIENTS WITH RHEUMATOID ARTHRITIS AND ASSOCIATED COMORBIDITIES

Authors: Tore K. Kvien^{*1}, Alejandro Balsa², Neil Betteridge³, Maya H Buch⁴, Patrick Durez⁵, Ennio Favalli⁶, Guillaume Favier⁷, Cem Gabay⁸, Rinie Geenen⁹, Ioanna Gouni-Berthold¹⁰, Frank van den Hoogen¹¹, Alison Kent¹², Lars Klareskog¹³, Mikkel Østergaard¹⁴, Karel Pavelka¹⁵, Joaquim Polido-Pereira¹⁶, Anne Grete Semb¹, Magnus Sköld^{17,18}, Maxime Dougados¹⁹

Author affiliations: ¹Diakonhjemmet Hospital, Oslo, Norway, ²Hospital Universitario La Paz, IdiPAZ, Institute for Health Research, Madrid, Spain, ³Betteridge Associates, London, United Kingdom, ⁴Chapel Allerton Hospital, University of Leeds, Leeds, United Kingdom (current affiliation: Centre for Musculoskeletal Research, School of Biological Sciences, University of Manchester, United Kingdom), ⁵Rheumatology, Cliniques universitaires Saint-Luc, Université Catholique de Louvain, Institut de Recherche Expérimentale et Clinique (IREC), Brussels, Belgium, ⁶Department of Rheumatology, ASST Gaetano Pini-CTO Institute, Milan, Italy, ⁷Global Strategy Group, KPMG LLP, London, United Kingdom, ⁸Geneva University Hospital, Geneva, Switzerland, ⁹Department of Psychology, Utrecht University, Utrecht, Netherlands, ¹⁰Polyclinic for Endocrinology, Diabetes and Preventive Medicine, University of Cologne, Cologne, Germany, ¹¹Sint Maartenskliniek, Nijmegen, Netherlands, ¹²Salisbury NHS Foundation Trust Hospital, Salisbury, United Kingdom, ¹³Division of Rheumatology, Department of Medicine, Karolinska Institutet and Karolinska University Hospital, Stockholm, Sweden, ¹⁴Copenhagen Center for Arthritis Research, Center for Rheumatology and Spine Diseases, Rigshospitalet, University of Copenhagen, Denmark, ¹⁵Institute of Rheumatology, Prague, Czech Republic, ¹⁶Hospital Santa Maria, Lisbon, Portugal, ¹⁷Respiratory Medicine Unit, Department of Medicine Solna and Center for Molecular Medicine, Karolinska Institutet, Stockholm, Sweden, ¹⁸Department of Respiratory Medicine and Allergy, Karolinska University Hospital Solna, Stockholm, Sweden, ¹⁹Cochin Hospital, Paris, France

Corresponding author:

Tore K. Kvien

Dept. of Rheumatology, Diakonhjemmet Hospital

Box 23 Vindern

N-0319 Oslo

t.k.kvien@medisin.uio.no

Abstract

Objective: Rheumatoid arthritis (RA) is a chronic autoimmune inflammatory disorder with a global prevalence of approximately 0.5-1%. Patients with RA are at an increased risk of developing comorbidities (e.g. cardiovascular disease, pulmonary disease, diabetes and depression). Despite this, there are limited recommendations for the management and implementation of associated comorbidities. This study aimed to identify good practice interventions in the care of RA and associated comorbidities.

Methods: A combination of primary research (180+ interviews with specialists across 12 European rheumatology centres) and secondary research (literature review of existing publications and guidelines / recommendations) were used to identify challenges in management and corresponding good practice interventions. Findings were prioritised and reviewed by a group of 18 rheumatology experts including rheumatologists, comorbidity experts, a patient representative and a highly specialised nurse.

Results: Challenges throughout the patient pathway (including delays in diagnosis and referral, shortage of rheumatologists, limited awareness of primary care professionals) and 18 good practice interventions were identified in the study. The expert group segmented and prioritised interventions according to three distinct stages of the disease (1. suspected RA, 2. recent diagnosis of RA and 3. established RA). Examples of good practice interventions included: enabling self-management (self-monitoring and disease management support e.g. lifestyle adaptations); early arthritis clinic; rapid access to care (online referral, triage, ultrasound-guided diagnosis); dedicated comorbidity specialists; enhanced communication with primary care (hotline, education sessions); and integrating patient registries into daily clinical practice.

Conclusion: Learning from implementation of good practice interventions in centres across Europe provides an opportunity to more widely improved care for patients with RA and associated comorbidities.

Key words: Rheumatoid arthritis, Comorbidities, Quality of care, Early arthritis clinic, Primary care

Key messages

What is already known about this subject?

- RA has a global prevalence of about 0.5-1%. The global burden of RA has risen, despite no significant change in prevalence.
- Despite publication of international guidelines and recommendations on RA management, there are limited international recommendations on the management of the several RA associated comorbidities.
- There are many challenges in the delivery for RA care and associated comorbidities which exist at the level of the healthcare system, health care providers and patient, across the patient pathways.

What does this initiative add?

- This initiative highlights good practice interventions which are in place at selected centres across Europe and addresses some of the challenges as well as opportunities in delivery of care.
- An expert multi-disciplinary panel consisting of rheumatologists, specialists in associated comorbidities, a patient representative and a highly specialist nurse prioritised, by consensus, the good practice interventions at three different disease stages; suspected RA, recent diagnosis of RA and established RA.
- The study findings inform European considerations on good practice management of RA and associated comorbidities.

How might this impact on clinical practice on future developments?

- The interventions highlighted in this report could be potentially implemented by and adapted to varying healthcare settings to improve the quality of care of patients with RA and their associated comorbidities.

Introduction

Rheumatoid arthritis (RA) is one of the most common autoimmune inflammatory diseases worldwide¹, affecting between 0.5 - 1% of the global population². Although life expectancy of patients with RA has increased over the last decades³, they are still at increased risk of developing comorbid conditions⁴.

Comorbidities discussed in this paper refer to conditions that may either be directly or indirectly caused by RA, RA treatment, or by RA outcomes, such as decreased mobility and functional impairment. Multi-morbidities related to increasing age and lifestyles of patients with RA are also of note and included in this paper's definition of comorbidities⁵.

When compared to the general population, patients with RA have the following adjusted lifetime hazard ratios: 1.41 for the development of cardiovascular disease (40-70% increased risk)⁶, 8.96 for interstitial lung disease (ILD) (7% increased risk)⁷ and 1.5 for diabetes⁸. Estimates of the prevalence of depressive disorder in patients with RA range between 13 and 20%^{9,10} which is 2-3 times higher than for the general population whilst the disability adjusted life years of patients with RA has increased from 3.3 million in 1990 to 4.8 million in 2010¹¹.

Guidelines and recommendations, such as the 2016 EULAR points to consider for reporting, screening for and preventing selected comorbidities¹², primarily address screening and prevention of comorbidities. However there is a clear need for recommendations on management of RA associated comorbidities¹³.

The objectives of this study were to 1) understand the current challenges in the management of patients living with RA and associated comorbidities, 2) identify and collate approaches undertaken by rheumatology teams to address these challenges and 3) discuss potential strategies to replicate and implement these interventions across healthcare systems and care settings to improve patient outcomes.

Materials and Methods

The study followed a three-step approach:

1. A literature review using an integrative approach to identify reported and evidence-based challenges and good practice examples in the management of RA and its associated comorbidities. The review included:
 - Academic literature search. A database search for articles in PubMed using a combination of prioritized search terms was conducted. The search terms included: rheumatoid arthritis, comorbidities, cardiovascular risk, diabetes, depression, interstitial lung disease, management, early identification, diagnosis, challenges in care, patient outcomes, guidelines, recommendation, quality of care, screening, diagnosis, delay referral, treat to target, disease burden, multidisciplinary approach to care, self-management, personalised care etc. Additional articles were retrieved through the citation-tracking of original publications and were based on the recommendation from the expert panel.
 - Grey literature search. The publicly available grey literature was investigated by applying the same prioritized search terms in search engines. Non-English-language references were excluded unless there was sufficient explanatory text in English. The time allotted for the grey literature search was one hour or until saturation was reached, whichever came first. Saturation was defined as not identifying new literature to include in analysis for 40 min or 5 consecutive search pages, whichever came first.
2. Visits to 12 RA centres across Europe, representing a multinational approach to RA management with each centre being located in a different European country. The centres were selected based on a set of criteria which included:
 - Geography: the centres, each located in a different European country, were selected to represent a multinational approach to RA management

- Focus: the centre's experience with RA was considered; both specialised and more generalist centres were selected to ensure the results of the study would be applicable to the wider clinical community
- Centre type: the centres' models and involvement in the wider healthcare ecosystem (e.g. private versus public funding, size, partnerships within the community etc.) were assessed

Participating centres included Diakonhjemmet Hospital (Oslo, Norway), Cochin Hospital (Paris, France), Institute of Rheumatology (Prague, Czech Republic), Hospital Santa Maria (Lisbon, Portugal), University Hospital La Paz (Madrid, Spain), Chapel Allerton Hospital and University of Leeds (Leeds, United Kingdom), University Clinic Saint-Luc (Brussels, Belgium), Geneva University Hospital (Geneva, Switzerland), Sint Maartenskliniek (Nijmegen, Netherlands), Karolinska University Hospital and the Centre for Rheumatology (Stockholm, Sweden), Rigshospitalet (Copenhagen, Denmark) and ASST Gaetano Pini-CTO Institute (Milan, Italy).

Approximately 180 qualitative semi-structured interviews were conducted to gain a robust understanding of the management of RA and associated comorbidities through the visits. An interview guide was created to facilitate structured collection of quantitative and qualitative insights. The questions were designed to capture the RA service set-up and patient pathway design within each centre and assessing the key challenges, which were categorised as "gaps in care" and "drivers" (i.e. reasons for gaps in care).

All the interviews were conducted on an individual basis over a duration of approximately one hour. Before the start of the interview, all participants were given an overview of the project objectives.

The individuals interviewed included a wide range of healthcare professionals (HCPs) such as rheumatologists, nurses, physiotherapists, occupational therapists and RA-associated comorbidity specialists, pharmacists, practice management.

Interview responses were systematically analysed for insights and patterns. Thematic analysis was undertaken to identify all challenges and interventions, which were subsequently reviewed by the expert panel.

3. The findings from the above secondary and primary research were tested with an expert panel. This panel of European experts was set up to ensure relevance in a range of healthcare systems. It consisted of 18 members including 16 clinicians of which there were 12 rheumatologists (one from each centre visited), one cardiologist, one pulmonologist, one diabetologist and one psychologist (i.e. one representative for the four preselected common comorbidities). The panel also included one rheumatology specialist nurse, acknowledging evidence that highlights the importance of the role of nurses in practice and one patient representative to capture the views and perspectives of patients. The expert panel were involved throughout the study and met on three occasions to guide the project objectives and discuss project findings.

Ethical approval was not required for this study since it did not include data collection from patients or other individuals.

Results

Several challenges in the management of RA and associated comorbidities were identified. Gaps in the care of patients with RA included: delays in securing a diagnosis consisting of seeking medical advice (>12 weeks¹⁴); referral delays (with a median of 4 visits to the general practitioner before patients are referred to a rheumatologist¹⁵) leading to an overall delay in diagnosis (in several countries, it can take up to a year from symptom onset¹⁶); and upon diagnosis and treatment initiation, poor patient adherence to therapy (varying between 30-80%¹⁷) and lack of monitoring (up to 50% of patients do not have access to an regular review in a secondary care setting¹⁸). Drivers of these gaps in care were evident at a healthcare system, patient and HCP level [Table 1].

Challenges were also evident throughout the patient pathway for the care of RA associated comorbidities. One pertinent challenge was the limited guidance for the care of comorbidities in patients with RA [table 2].

Through secondary research several guidelines and recommendations were identified for some comorbidities but the presence of guidelines differed depending on the comorbidity [Table 3].

The expert panel recommended to frame the findings of the study against three different stages of disease [Table 4].

These disease stages were used to segment and prioritise good practice interventions. In total, 18 good practice interventions were identified, through the centre visits, which span across the whole patient pathway. Interventions were highlighted as applicable to one, multiple or all disease stages [Table 5].

All 18 interventions were reviewed by the expert panel and prioritised by ranking the highest impact on patient outcomes, against each stage of the disease: suspicion of RA; recent diagnosis of RA; established disease (whilst recognising that for each one of them patients have distinct needs and requirements). The top three interventions for each disease stage are shown in Table 6 and are described below.

Interventions: Diagnosing RA and early intervention

Rapid access to specialised care. Patients with suspected RA must be seen rapidly to facilitate the effective prevention and management of disease progression and joint damage¹⁹. Delays in diagnosis and referral from primary care mean patients can arrive at secondary centres with irreversible damage to their joints and other organs. It is vital that these patients are triaged rapidly into care and start treatment straight away²⁰. Rapid access to care may include online referral systems (patient self-enrollment) and access to diagnostic services (e.g. blood tests and ultrasound). Hospital Universitario La Paz has implemented an online direct communication system between the rheumatology department and primary care practitioners to allow two-way messaging system for rapid patient referrals within 24 hours. A number of centres run rapid access clinics, such as the 'clinically suspect arthralgia' (CSA) clinic for undifferentiated arthralgia at the Institute of Rheumatology in the Czech Republic, the pre-RA CCP clinic Leeds Hospital in the UK and the sophisticated triaging system at Hospital de Santa Maria which aims to increase accessibility for new patients and rapidly assess the presence and severity of RA. These services have reduced waiting times, improved the experience for patients and facilitated two-way communication between primary and secondary care.

Early arthritis clinic. Early treatment reduces overall disease progression and therefore prevents the accumulation of damage that causes irreversible joint deformities. Clinics are dedicated to ensuring timely clinical assessment and diagnosis of patients with suspected RA, including initial triage, to which patients can be referred by primary care physicians (PCPs) and other specialists. The Institute of Rheumatology, Hospital de Santa Maria, Cliniques Universitaires Saint-Luc, Sint Maartenskliniek, Diakonhjemmet Hospital, Leeds Teaching Hospitals, Karolinska University Hospital, Hospital Universitario La Paz, ASST Gaetano Pini-CTO Institute and Rigshospitalet have all implemented early arthritis clinics leading to more patients being diagnosed quickly and starting early disease-modifying therapy combined with a treat-to-target strategy; with follow-up maintained in the early arthritis clinic until target is reached and/or directed to escalated treatment pathways and service as indicated.

Enhanced communication with primary care. Often patients are not transitioned between different levels of the healthcare system in a timely manner²¹ which can be due to a number of factors including lack of awareness of referral pathways and systems, suboptimal communication between specialists and PCPs, and sharing of up-to-date patient information²². Enhanced communication strategies identified include a hotline for PCPs to access specialist advice within 48 hours, linked information systems leading to collaboration between primary and secondary care, educational and training sessions. Hôpital Cochin

implemented the “Hospital and City Rheumatology Network” and Diakonhjemmet Hospital have a dedicated primary care coordinator. ASST Gaetano Pini-CTO Institute delivers education and training for PCPs in Italy regarding RA, developing a network with primary care. Enhanced communication facilitates improved integration of services and patients accessing specialist treatment in a timely manner.

Interventions: Management of comorbidities

Comprehensive comorbidity assessment. Certain patients with RA are at increased risk of morbidity and mortality due to existence of comorbidities such as cardiovascular disease, ILD, diabetes and depression^{23,24,25,26}. Systems for cross-specialty care and follow-up can be fragmented²³. Centres may implement regular (typically 6-monthly) full comorbidity assessment as a standalone service or in conjunction with RA appointments. Hôpital Cochin has a Comorbidities Education in Rheumatoid Arthritis (COMEDRA) service which is a doctor-led programme on RA comorbidity management and supported by a programme coordinator, and Hospital Universitario La Paz has a systematic inflammatory osteoporosis screening. Comorbidity assessments may lead to reduced patient mortality due to comorbidity complications in patients with RA (e.g. cardiovascular disease and ILD).

Dedicated comorbidity specialist. Comorbid diseases may not always be given appropriate focus and attention because rheumatologists are not specialists in these areas²⁷. To overcome this, centres may deploy an in-house comorbidity specialist e.g. a cardiologist dedicated to the cardiovascular manifestations of rheumatic disorders. Diakonhjemmet Hospital has a preventive cardio-rheuma clinic run by a cardiologist employed in the Dept. of Rheumatology, and Hospital de Santa Maria has a dedicated psychologist for patients with RA. The input of comorbidity specialists may raise awareness regarding comorbidities amongst rheumatology HCPs. At the Karolinska University Hospital for instance, pulmonary complications can be presented during discussion rounds with the participation of pulmonologists, rheumatologists and radiologists.

Combined clinics. To effectively and more efficiently manage comorbidities associated with RA, with minimum burden on patient hospital visits, centres may implement combined clinics with both rheumatologists and the respective comorbidity specialists. Hospital de Santa Maria, ASST Gaetano Pini-CTO Institute and Hospital Universitario La Paz have combined clinics with rheumatologists and pulmonologists to manage patients with RA with or at risk of ILD. Hospital de Santa Maria has a joint rheuma-obstetrics clinic to manage the potential problems that can arise during pregnancy in rheumatological patients. Leeds Teaching Hospitals has several joint comorbidity services with dermatology, gastroenterology, immunology, neurology, respiratory, renal and obstetrics. These combined or stand-alone clinics may cover the wider aspects of RA management, including lifestyle factors. For example, Rigshospitalet in Copenhagen offers a sleep clinic and smoking cessation clinic for patients with RA helping patients to live healthier lives. Joint clinics may lead to a greater focus on comorbidities and involvement of specialist comorbidity expertise leading to more effective decisions and better patient outcomes.

Interventions: Encouraging patient self-management and empowering health professionals

Effectively utilizing the skill mix of the multidisciplinary care team. The findings of this study indicated an increasing number of patients with RA requiring comprehensive care and management. Rheumatologists may not have enough time, capacity or specific skills to address these important problems outside diagnosis, inflammatory assessment and pharmacological treatment. Other health professionals (e.g. nurses, physio- and occupational therapists, care coordinators and administrative staff) often have the competencies that are well suited for some of the tasks required to ensure best practice. By facilitating a more active role of these healthcare professionals in the management of patients, they are able to bring a different dimension which adds to a holistic and patient-centred approach to care. For example, at Sint Maartenskliniek, the pharmacist is a key member of the care team, delivering medication support for patients. At Diakonhjemmet Hospital, Cliniques Universitaires Saint-Luc, Rigshospitalet and Leeds Teaching Hospitals the enhanced role of the nurse helps support rheumatologists in care delivery (e.g. in Leeds the nurse-delivered services at the centre include outpatient clinics and a day care unit where

rheumatology nurses manage the delivery of biologics to patients). Such activities can be done in various ways between countries and health systems as shown in online supplementary Table S1.

Enabling self-management. Patients may not always have long enough consultation sessions with rheumatologists and rheumatology nurses and other HCPs to fully educate themselves about the impact of the disease and how to manage it²². As part of the COMEDRA initiative at Hôpital Cochin, patients are taught to self-assess and self-monitor their disease activity and more importantly to feel comfortable to undertake this. At Diakonhjemmet Hospital a 'learning and coping' centre is run by a multidisciplinary team led by an experienced nurse. Cliniques Universitaires Saint-Luc enable self-management through personalising care delivered to patients. All these interventions have been put in place not only to improve adherence to treatment but also to help patients manage their lives and not just their disease activity; it is about empowerment and feeling in control.

Interventions: Care delivery process optimisation

Integrating patient registries into daily clinical practice. Several registries have been implemented at different centres to collect longitudinal real life data, for example at the Institute of Rheumatology; Swiss Clinical Quality Management in Rheumatic Diseases (SCQM) at the Hôpitaux Universitaires de Genève; Swedish Rheumatology Quality Registry (SRQ) at the Karolinska University Hospital and the Centre for Rheumatology in Stockholm; DANBIO registry at Rigshospitalet; and NOR-DMARD at Diakonhjemmet Hospital. Hospital de Santa Maria uses the Portuguese Reuma.pt registry to support research and clinical practice and Leeds Teaching Hospitals has a focus on research to support clinical practice. Registries may inform real life results of clinical management from different participating centres and thereby also provide information which may stimulate the different centres to improve their quality of care. Further, linkages between registries may help the understanding of the relationships between RA and comorbidities. Karolinska University Hospital has had a strong focus on this type of research^{28,29}

Patient-centred care journey. Patients need to be satisfied and confident with their care to be fully engaged with their care^{30,31}. Patient-centred care can be defined as when patients are empowered throughout the duration of their journey whether it relates to the physical environment or the experience of the services received. At the Sint Maartenskliniek, patient-centred design reduces time spent travelling between each care station, and provides more time with the HCPs; all of which improves the overall patient experience. This intervention enables HCPs to improve patient engagement and helps empower and support patients throughout their care journey.

Key considerations in selected comorbidity care for patients with RA

The visits to study centres (primary research) and the input from the expert panel provided a number of key considerations for the care of comorbidities (cardiology, pulmonology, diabetes and depression) in patients with RA.

Cardiovascular disease (CVD): Key elements to consider for improvement of the quality of care of CVD in patients with RA are listed below:

- Communication about CVD risk factors between medical professionals as well as screening of CVD risk factors. It was noted that CVD risk prediction in patients with RA using risk calculators developed for the general population are generally inaccurate in prediction of future CVD.^{32,33} Modifying the predicted risk by a 1.5 multiplier as per recommendation²⁴² may also not necessarily reclassify patients to their appropriate risk class.^{34,35} However adding information on carotid plaque/atherosclerosis may reclassify patients to a more appropriate risk class in up to 30 to 60% of cases.³⁴
- Organisation and responsibility of CVD risk management through cardiologists working alongside or within rheumatology departments e.g. preventive cardio-rheumatology clinics. It was noted that lipid-lowering, antihypertensive therapies and non-pharmacological recommendation (e.g. focusing on life-style factors) can be implemented safely and effectively with recommended goal achievement over approximately 3 consultations in 80% - 90% of the patients.³⁶ Systemic

inflammation or lipid levels at baseline or anti-rheumatic medication do not have an impact on statin dose needed to obtain LDL goals.³⁷

- Availability of data regarding CVD risk management.

Pulmonary disease: Interventions and results from the study highlight the importance of early screening and detection. This can include chest x-ray, computed tomography (CT) and functional assessment with spirometry and six-minute walking test (6MWT). Regular follow-up in outpatient clinics are also important. Management of adverse respiratory events, side effects from medications and smoking cessation advice should be included. The expert panel emphasized the importance of the fieldwork findings from the visits of study centres and especially the collaboration between rheumatologists and pulmonologists. ILD should be diagnosed in multidisciplinary discussions where the presence of rheumatologists is encouraged. These conferences can also enable pulmonologists to consider the possibility of an underlying systemic disease including RA in patients with ILD.

Diabetes: Study results highlight the importance of screening for and detecting diabetes, coordination of care between RA and diabetologists and regular follow-ups. The expert panel advised that the effective treatment of RA with targeted therapies can improve long-term glycaemic control in patients with diabetes and RA.³⁸

Depression: The results of the study highlight the importance of screening and detection, periodic follow-up, guided self-help and targeted depression history. These can be achieved through dedicated psychological and psychiatric services as well as focusing on impacts of depression e.g. sleep disturbance and socioeconomic consequences. The expert panel recommended a strong focus on prevention through paying attention to individual factors (e.g. sleep, physical activity, interest in undertaking any activities), encouraging a healthy lifestyle and treatment of depression (moderate – severe) if guided self-help and exercise fail (e.g. treatment with cognitive-behavioural treatment [CBT] or psychotherapy). The expert panel noted that the ideal model to achieve the above activities would be the integration of psychology resources as part of the wider multidisciplinary team although the experts also recognised local constraints.

Discussion

Challenges in the care of RA and associated comorbidities are well documented and results of the interviews confirmed the findings from the literature review. However, the study highlighted the different types of challenges experienced by the centres. In order to improve patient care, it is recommended that centres identify and prioritise interventions most suited to their population, patient pathways and current challenges in delivery of care. The need for comprehensive and consensus-approved guidelines and recommendations to improve the care of patients with RA and the several associated comorbidities was clear in the findings. The experts acknowledged that this requires further exploration and expert input with more robust methodology in order to develop guidelines and recommendations.

It was recognised that there were limitations to the study methodology and several actions were taken to mitigate these. One drawback was the lack of patient interviews at each individual centre which limited the patient perspective on challenges in the patient pathways and intervention benefits (e.g. benefits of self-management). The role of the very experienced patient representative on the expert committee (NB) helped to overcome this by providing a collective patient perspective, on behalf of patients. Secondly, the inclusion of one centre per country limited the findings with respect to the representativeness of the wider healthcare system in which they reside. This was taken into consideration when conducting research which did not solely focus on centres themselves but rather their entire ecosystems; including the various institutions they collaborate with, whether formally or informally. However, the goal of this study was rather to find centres that had developed good quality of care practices which could serve as models if implemented in other centres.

Thirdly, centre selection was biased towards larger teaching centres. This was mitigated by gathering and documenting interviewees' perspectives on how to replicate interventions in smaller and less resource rich centres; although it was accepted that this was incomplete.

The project group decided to focus on four main comorbidities; CVD, ILD, diabetes and depression. The literature review and also centre visits highlighted other comorbidities of importance. For example it has been shown that osteoporosis is twice as common in patients with RA compared with controls of the same age^{39,40} and that fractures are also more common.^{41,42} Services to support primary and secondary prevention are of value in improving patient morbidity.

Infections are other important comorbidities which can partly be related to the disease, partly to treatment. Several studies have described treatment related occurrence of infections. Importantly, EULAR has recently published updated recommendations on vaccinations⁴³ and editorials have addressed the important aspects of hepatitis and tuberculosis in this population.⁴⁴

Guidelines and / or recommendations on periodontitis, malignancy and gastrointestinal disorders were not assessed but it was noted that these conditions lead to poor outcomes for patients with RA^{45,46,47}

We believe that the most important message from this paper is the description of examples of good clinical practices and care models. One of the main findings was the success of early arthritis clinics. Impressively, many clinics reported to see patients within 1-2 days since they focused on the importance of early diagnosis and treatment. This may appear more aspirational for the majority of departments with limitations in resource and funding to deliver such a service. Nevertheless, these examples illustrate the importance of prioritizing early RA pathways of care within rheumatology services to optimize efficient diagnosis and management. However, we also recognize that not all recommendations are feasible in every setting since different clinical structures and workforce challenges may limit the implementation of for example multidisciplinary care approaches. However, the recommendation may hopefully serve as a benchmark to support efforts to obtain more resources to enhance quality of care.

The strength and impact of this paper would have been improved if data had been available on longitudinal clinical outcomes to demonstrate the benefit of the good practice recommendations shown in Table 5. Long-term clinical data are available for some of the recommendations and are included in the reference list.

It is envisaged that rheumatology departments could utilise this study as an initial platform to draw upon models appropriate to local services and pathways for adaptation and implementation. Interested centres are welcome to contact the centres who took part in this study for further knowledge transfer.

The panel of experts recognise that data capture should also be done in the community. Community-based physicians could propose a list of items which, if accepted by rheumatologists, would lead to harmonization of standards of care across all care settings along the patient journey.

The next steps for the project group are to do additional work on dissemination of our findings. A comprehensive study report will be published on the web-site of the project accompanied by an appendix with single centres reports. More importantly, we also plan to support implementation of the examples of models of good clinical care into other settings.

In conclusion, we identified 18 models of good clinical practice which are linked to three different disease stages. Further work is needed to explore the ability to implement each of the interventions (e.g. the development of tools and manuals detailing their implementation) and the results achieved.

Acknowledgements: The authors thank Sanofi Genzyme who funded this study. The authors are also grateful to Olivia Bryan, Emma Hanson, Jessica O'Neil, Zoe Philips and Thishi Surendranathan from the KPMG team for their input and contribution to the work.

Previous publications: The main results were previously published as abstracts in ARD related to presentations at EULAR 2019: Kvien TK, Pavelka K, Polido-Pereira J, et al. THU0638 Considerations for improving quality of care in Rheumatoid Arthritis and associated comorbidities. [Ann Rheum Dis](#). 2019;**78**:613-6; Betteridge N, Balsa A, Buch M, et al. PARE0018 Enabling patient-centred care in Rheumatoid Arthritis and associated comorbidities. [Ann Rheum Dis](#). 2019;**78**:2185; Kent A, Gabay C, Geenen R, et al. AB1368-HPR Enhanced role of nurses and other healthcare

professionals (HCPs) in the care of Rheumatoid Arthritis and associated comorbidities. [Ann Rheum Dis](#) 2019;**78**:2147-8. Further, the full report of the project has been published online: <https://home.kpmg/content/dam/kpmg/uk/pdf/2020/02/improving-quality-of-care.pdf>.

Disclosures:

Tore K Kvien has received fees for speaking and/or consulting from AbbVie, Biogen, Celltrion, Egis, Eli Lilly, Hikma, MSD, Mylan, Novartis, Oktal, Orion Pharma, Pfizer, Roche, Sandoz, Sanofi and UCB and received research funding to Diakonhjemmet Hospital from AbbVie, BMS, MSD, Pfizer, Roche and UCB.

Alejandro Balsa has received Grant/research support, fees for consultancies or as a speaker for Abbvie, Pfizer, Novartis, BMS, Nordic, Sanofi-Genzyme, Sandoz, Lilly, UCB and Roche.

Neil Betteridge has received fees for speaking and/or consulting from Amgen, Eli Lilly, Grunenthal, GSK, Heart Vlavle Voice, Janssen, Roche, Sanofi Genzyme and Sanofi Regeron.

Maya H Buch has received fees for speaking and/or consulting from AbbVie, AstraZeneca, Bristol-Myers-Squibb, Chugai, Eli Lilly, Merck-Serono, Pfizer, Roche, Sandoz and Sanofi, and research funding to University of Leeds from Pfizer, Roche and UCB.

Patrick Durez has received fees for speaking and/or consulting from Bristol-Myers-Squibb, Celltrion, Lilly and Sanofi Genzyme.

Ennio G. Favalli has received fees for speaking and/or consulting from AbbVie, BMS, Eli Lilly, Gilead, MSD, Novartis, Pfizer, Roche, Sanofi-Genzyme and UCB.

Cem Gabay has received fees for consulting from Roche, Sanofi Genzyme, Regeneron, Pfizer, Lilly and Ab2 Bio Ltd.

Rinie Geenen has received fees for speaking from Sanofi Genzyme.

Ioanna Gouni-Berthold has received fees for consulting from Amgen, Akcea, Sanofi Genzyme and Regeneron.

Frank van den Hoogen has received fees for consulting from AbbVie, Biogen, Celltrion, Roche, Sanofi Genzyme, Pfizer and Munidpharma.

Alison Kent has received fees for speaking and/or consulting from UCB, Bristol-Myers-Squibb, MSD, Amgen, Abbvie, Pfizer, Novartis and Sanofi.

Lars Klareskog has received research grants to Karolinska Institutet from Janssen, Pfizer, BMS, GSK, UCB.

Mikkel Østergaard has received fees for speaking and/or consulting from Abbvie, Bristol-Myers-Squibb, Boehringer-Ingelheim, Celgene, Eli-Lilly, Hospira, Janssen, Merk, Novartis, Novo, Orion, Pfizer, Regeneron, Roche, Sanofi and UCB.

Karel Pavelka has received fees for speaking and/or consulting for AbbVie, Amgen, Biogen, Bristol-Myers-Squibb, Egis, MSD, UCB.

Joaquim Polido-Pereira has received fees for speaking and/or consulting from AbbVie, MSD, Pfizer, Roche and Tecnimed.

Anne Grete Semb has received fees for speaking and/or consulting from AbbVie, Novartis, Sanofi and Bayer and have an unrestricted research collaboration with Eli Lilly which includes transfer of funds to Diakonhjemmet Hospital from Eli Lilly.

Magnus Sköld has received research grants from Boehringer Ingelheim and Roche, speakers fee/consultancy from AstraZeneca, Boehringer Ingelheim, GlaxoSmithKline, Novartis, Mundipharma, Sandoz and Roche.

Maxime Dougados has received fees for speaking and/or consulting from AbbVie, Biogen, Eli Lilly, BMS , MSD, Novartis, Pfizer, Roche, Sandoz, Sanofi and UCB and his department has received research grants from AbbVie, BMS, MSD, Pfizer, Roche, Lilly, Janssen, Novartis, UCB.

Table 1. Gaps in care and their associated drivers evident across the journey of patients with RA

Stage of Patient Journey	Gaps in care	Drivers ⁽¹⁾
Awareness and prevention	Delays in patients seeking medical advice	Low public awareness ¹⁴ Lack of patient education ⁴⁸
Referral	Delayed referrals from PCPs to rheumatologist	Limited awareness of PCPs on signs and symptoms of RA ⁴⁹ Long waiting times for specialists ²¹ Joint stiffness and inflammation are common for many other conditions Lack of knowledge regarding the referral pathway or the importance of rapid referral ²¹
Diagnosis	Delay in diagnosis	Complex diagnostic requirements ¹⁴ Limited availability of imaging facilities ²¹²² Shortage of rheumatologists ²² Limited number of HCPs trained to interpret diagnostic results (including imaging) ¹⁴
Treatment and management (pharmacological and non-pharmacological)	Delayed treatment initiation Poor patient adherence to therapy Lack of coordinated treatment	Budgetary cap and restrictive clinical recommendations regarding choice of therapy ²² Disease management not aimed at optimising quality of life ⁵⁰ Deviation from recommendations, or difficulty in implementing them into clinical practice ²² Lack of regular review and engagement with patients' needs ²⁷ Treatment adherence ⁵⁰ Competing lifestyle factors ⁵⁰
Follow-up	Lack of monitoring Suboptimal patient outcomes	Capacity constraint – leading to long waiting time for examinations with imaging modalities ²² Lack of regularity in follow-up care ²² Access to rheumatologists ²²

Notes: (1) List not exhaustive

Abbreviations: RA: rheumatoid arthritis; HCP: healthcare professional; PCP: primary care professional

Table 2. Gaps in care and their associated drivers evident across the journey of patients with RA associated comorbidities

Stage of Patient Journey	Gaps in care	Drivers ⁽¹⁾
Screening	Lack of screening for comorbidity risk factors	Limited screening pathways and frameworks in place to detect comorbidities amongst patients with RA ²³ Lack of comprehensive comorbidity-specific recommendations Limited comorbidity screening ²⁵
Referral	Delay in referral	Lack of HCP education on referral pathways ⁵¹ Logistical and ambulatory difficulty can negatively impact outpatient attendance ²³⁵
Diagnosis	Delay in diagnosis	Comorbidities are often underdiagnosed and more likely to be diagnosed when severe ²⁵ Symptoms of RA/RA treatment can mask comorbidity signs and symptoms ²⁶
Treatment and management (pharmacological and non-pharmacological)	Inadequate management of comorbidities	Inadequate communication across multidisciplinary team ²³ Limited recording of drug-drug interactions Patients with RA who have comorbidities do not always receive the recommended comorbidity treatment ²³
Follow-up	Suboptimal patient outcomes	Issues with patient data inconsistencies in capturing and sharing across different systems ²⁶ Poor follow-up appointment attendance ⁵¹ Increased morbidity and mortality is seen in patients with complex therapeutic needs ⁵¹
Notes: (1) List of drivers not exhaustive Abbreviations: RA: rheumatoid arthritis; HCP: healthcare professional; PCP: primary care professional		

Table 3. Guidelines and / or recommendations present for comorbidities in RA

Comorbidity	Authors	Guidelines / recommendations
Cardiovascular disease	Regulatory bodies or disease associations	<ol style="list-style-type: none"> 1. 2016 Update EULAR recommendations for CVD risk management⁵² 2. 2016 EULAR points to consider for reporting, screening for and preventing selected comorbidities¹² 3. 2019 European Society of Cardiology / European Atherosclerosis Society guidelines for the management of dyslipidaemias⁵³ 4. Spanish Society of Rheumatology (2011). Update of the clinical practice guideline for the management of rheumatoid arthritis in Spain ⁵⁴ 7. Haute Autorite De Sante (2017): Clinical practice guidelines: Rheumatoid Arthritis⁵⁵
Pulmonary disease	Academic literature	<ol style="list-style-type: none"> 1. Diagnosis of Idiopathic Pulmonary Fibrosis. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline⁵⁶ (associated rheumatological diseases are discussed from a pulmonary perspective) 2. 2019 Swedish Respiratory Society Guidelines for Idiopathic Pulmonary Fibrosis⁵⁷
Diabetes	No existing guidelines or recommendations were found at a regional or national level for diabetes	
Depression	Existing recommendations / points to consider published by regulatory bodies or disease associations	<ol style="list-style-type: none"> 1. 2016 EULAR points to consider for reporting, screening for and preventing selected comorbidities¹² 2. NICE (2009): Depression in adults with a chronic physical health problem: recognition and management⁵⁸

Table 4. Disease stages

Disease stage	Description
Suspicion of RA	Patients who have not yet been given a diagnosis but may have interacted with primary care or had a first contact with a rheumatologist, and who may have suspected RA (inflammatory manifestations not diagnosed)
Recent diagnosis of RA	Patients have been given a diagnosis of RA, and usually have been started on treatment with regular follow-up
Established disease / structural damage	Patients have had a diagnosis of RA for a number of years or may have presented late (most likely from less developed healthcare economies). At this stage treatments may have become progressively less effective at reducing inflammation and preventing further joint damage

Table 5. Good practice interventions relevant to the three disease stages of patients with RA

Intervention	Definition	Suspected RA	Recently diagnosed with RA	Established RA / structural damage
Rapid access to care	Fast-track access to care for patients with RA done via online referral form reviewed every 24 hours, hotline leading to appointments within 48 hours for diagnostic services including blood tests and joint imaging	X	X	X
Enhanced communication across wider care team	Availability of reliable communication channels (e.g. emails, online forms) enabling easy dialogue between specialists and PCPs; and providing and coordinating education programmes to ensure the wider care team are kept up-to-date with developments in best practice care	X	X	X
Early arthritic clinic	Clinic dedicated to ensuring timely clinical assessment and diagnosis of patients with suspected RA	X	X	
Comprehensive comorbidity assessment	Comorbidity assessment in patient baseline assessment and follow-up of newly diagnosed patients	X	X	X
Tailored education to patients and family members	Programmes to increase understanding of diagnosis, treatment plans, and how to live with the disease that is sensitive to individual patient needs		X	
Role of the care coordinator	Care coordinator role to help to manage the burden of navigating contacts across multiple HCPs		X	X
Dedicated comorbidity specialist HCP	Specific role or clinic to support the management of comorbidities in the context of RA		X	X
Enabling self-management	Provision of tools and resources to patients to monitor and manage their RA and reduce dependence on healthcare services		X	X
Enhanced therapy services	Additional care centred around non-physician-led management including care led by therapists to promote rehabilitation and enablement			X
Day clinic services	The coordination of services enabling provision of stacked outpatient appointments across specialities and disciplines over one day or session	X	X	X
Virtual engagement with patients	Digital enablement of autonomy, self-management and empowerment by providing a channel of direct communication with attending physicians in addition to online access to education around the disease, networks and peer support	X	X	X
Integrating patient registries into daily clinical practice	Employment of evidence-based practice, informed by clinical research and supported by systematic capture and monitoring of data, in order to improve the quality of clinical care delivery and promote evolution of care models	X	X	X
Patient-centred care journey	Curation of care processes and physical environment that enables the patient to feel empowered and supported during their journey through care	X	X	X

Effectively utilizing the skill mix of the multidisciplinary care team	Enablement of non-physician HCPs to taking on greater responsibility in the assessment and management of patients	X	X	X
Integrative and shared care solutions	Ensuring regular communication between all attending physicians and other HCPs as part of a holistic and integrated approach to care	X	X	X
Collaborating with PAGs	Increasing communication with local and regional patient advocacy groups through working group sessions, conference attendance, newsletters and patient liaisons	X	X	X
Developing care networks	Developing networks with community-based RA services to maintain high-quality 'joined-up' care past the point of discharge	X	X	X
Quality management programmes	A coordinated approach and a robust system for measuring, tracking and improving the quality of care	X	X	X
Abbreviations: RA: rheumatoid arthritis; HCPs: healthcare professionals; PAGs: patient advocacy groups; PCPs: primary care professionals				

Table 6. Prioritised interventions per disease stage

Disease stage	Prioritised interventions ⁽¹⁾
Suspicion of RA	<ol style="list-style-type: none"> 1. Rapid access to care 2. Enhanced communication with primary care 3. Early arthritis clinic
Recent diagnosis of RA	<ol style="list-style-type: none"> 1. Enabling self-management 2. Early arthritis clinic 3. Comprehensive comorbidity assessment
Established disease / structural damage	<ol style="list-style-type: none"> 1. Dedicated comorbidity specialist 2. Integrating patient registries into daily clinical practice 3. Enabling self-management
Notes: (1) The top three priority interventions are listed	

Table S1. Examples of extended roles of the multidisciplinary team members across different disease stages

Examples identified	Suspected RA	Recently diagnosed with RA	Established RA / structural damage
Providing education on RA and comorbidities		X	X
Conducting history and joint examination	X	X	X
Coordinating care		X	X
Delivering preliminary comorbidity screening		X	
Providing treatment		X	X
Being first point-of-contact		X	X
Collaborating with community-based teams		X	X
Providing holistic support	X	X	X
Empowering self-management		X	X

References

- ¹ Song L, Yangsheng Y, Yinshi Y, et al. Microbial Infection and Rheumatoid Arthritis. [J Clin Cell Immunol](#) 2013; 174.
- ² Angel J, Akesson K, Amadio P.C, et al. The burden of musculoskeletal conditions at the start of the new millennium. [World Health Organization](#) 2003; 919:1-218.
- ³ Norton S, Carpenter L, Nikiphorou E, et al. Excess mortality in rheumatoid arthritis: Gains in life expectancy over 25 years. [Rheumatology](#). 2014; Abstract O34.
- ⁴ Dougados M, Soubrier M, Antunez A, et al. Prevalence of comorbidities in rheumatoid arthritis and evaluation of their monitoring: results of an international, cross-sectional study (COMORA). [Ann Rheum Dis](#) 2014; 73:62-8.
- ⁵ Radner H, Yoshida K, Smolen J, et al. Multimorbidity and rheumatic conditions - enhancing the concept of comorbidity. [Nat. Rev. Rheumatol](#) 2014; 10:252–6.
- ⁶ Holmqvist M, Ljung L, Askling J. Acute coronary syndrome in new-onset rheumatoid arthritis: a population-based nationwide cohort study of time trends in risks and excess risks. [Ann Rheum Dis](#) 2017; 76:1642-7.
- ⁷ Bongartz T, Nannini C, Medina-Velasquez Y, et al. Incidence and mortality of interstitial lung disease in rheumatoid arthritis: a population-based study. [Arthritis Rheum](#) 2010; 62:1583-91.
- ⁸ Solomon DH, Love TJ, Canning C, et al. Risk of diabetes among patients with rheumatoid arthritis, psoriatic arthritis and psoriasis. [Ann Rheum Dis](#) 2010; 69:2114-7.
- ⁹ Sheehy C, Murphy E, Barry M. Depression in rheumatoid arthritis--underscoring the problem. [Rheumatology](#) 2006; 45:1325-7.
- ¹⁰ Matcham F, Rayner L, Steer S, et al. The prevalence of depression in rheumatoid arthritis: a systematic review and meta-analysis. [Rheumatology \(Oxford\)](#) 2013; 52:2136-48.
- ¹¹ Cross M, Smith E, Hoy D, et al. The global burden of rheumatoid arthritis: estimates from the global burden of disease 2010 study. [Ann Rheum Dis](#) 2014; 73:1316-22.
- ¹² Baillet A, Gossec L, Carmona L, et al. Points to consider for reporting, screening for and preventing selected comorbidities in chronic inflammatory rheumatic diseases in daily practice: a EULAR initiative. [Ann Rheum Dis](#) 2016; 75:965-73.

-
- ¹³ Winthrop K, Weinblatt M, Crow M, et al. Unmet need in rheumatology: reports from the Targeted Therapies meeting 2018. [Ann Rheum Dis](#) 2019; 78:872-8.
- ¹⁴ Bosworth A, Hancock T, Hewitt J, et al. Breaking Down Barriers: Rheumatoid Arthritis and Public Awareness. [National Rheumatoid Arthritis Society \(NRAS\)](#) 2013; 1-39.
- ¹⁵ Stack R, Nightingale P, Jinks C, et al. Delays between the onset of symptoms and first rheumatology consultation in patients with rheumatoid arthritis in the UK: an observational study. [BMJ Open](#) 2019; 9:e024361.
- ¹⁶ Aletaha D, Neogi T, Silman AJ, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. [Ann Rheum Dis](#) 2010; 69:1580-8
- ¹⁷ [National Rheumatoid Arthritis Society \(NRAS\)](#). Keep taking the pills – the critical importance of adherence in the management of rheumatoid arthritis. 2018.
- ¹⁸ Hider SL, Blagojevic-Bucknall M, Whittle R, et al. 2016 What does a primary care annual review for RA include? A national GP survey. [Clin Rheumatol](#) 2016; 35:2137-8.
- ¹⁹ Smolen JS, Landewé R, Bijlsma J, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. [Ann Rheum Dis](#) 2017; 76:960-77.
- ²⁰ Villeneuve E, Nam JL, Bell MJ, et al. A systematic literature review of strategies promoting early referral and reducing delays in the diagnosis and management of inflammatory arthritis. [Ann Rheum Dis](#) 2013; 72:13-22.
- ²¹ Garneau KL, Iversen MD, Tsao H, et al. Primary care physicians' perspectives towards managing rheumatoid arthritis: room for improvement. [Arthritis Res Ther](#) 2011; 13(6).
- ²² Miltenburger C, Karampampa K, Richard BT, et al. A survey of Barriers to Treatment Access in Rheumatoid Arthritis. [i3 innovus](#) 2009; 05:1-29.
- ²³ Peters MJL, Symmons DPM, McCarey D, et al. EULAR evidence-based recommendations for cardiovascular risk management in patients with rheumatoid arthritis and other forms of inflammatory arthritis. [Ann Rheum Dis](#) 2010; 69:325-31.
- ²⁴ Dixon WG, Hyrich KL, Watson KD on behalf of the British Society for Rheumatology Biologics Register, et al. Influence of anti-TNF therapy on mortality in patients with rheumatoid arthritis-associated interstitial lung disease: results from the British Society for Rheumatology Biologics Register. [Ann Rheum Dis](#) 2010; 69:1086-91.
- ²⁵ Bartels CM, Saucier JM, Thorpe CT, et al. Monitoring diabetes in patients with and without rheumatoid arthritis: a Medicare study. [Arthritis Res Ther](#) 2012; 14(4):1-9.
- ²⁶ Nakajima A, Inoue E, Shimizu Y, et al. Presence of comorbidity affects both treatment strategies and outcomes in disease activity, physical function, and quality of life in patients with rheumatoid arthritis. [Clin Rheumatol](#) 2015; 34:441-9.
- ²⁷ [National Rheumatoid Arthritis Society \(NRAS\)](#). The Impact of Rheumatoid Arthritis Comorbidities. 2012; 1-29.
- ²⁸ Radner H, Chatzidionysiou K, Nikiphorou E, et al. 2017 EULAR recommendations for a core data set to support observational research and clinical care in rheumatoid arthritis. [Ann Rheum Dis](#) 2018; 77:476-9.
- ²⁹ Eriksson J, Askling J, Arkema E. The Swedish Rheumatology Quality Register: optimisation of rheumatic disease assessments using register-enriched data [Clin Exp Rheumatol](#) 2014; 32:S-147-9.
- ³⁰ Doyle C, Lennox L, Bell D. A systematic review of evidence on the links between patient experience and clinical safety and effectiveness. [BMJ Open](#) 2013; 3:1-18.
- ³¹ Vare P et al. Delivering a one-stop, integrated, and patient-centered service for patients with rheumatic diseases. [SAGE Open Med](#) 2016; 4:2050312116654404.
- ³² Arts EE, Popa C, Den Broeder AA, et al. Performance of four current risk algorithms in predicting cardiovascular events in patients with early rheumatoid arthritis. [Ann Rheum Dis](#). 2015; 74(4):668-74.
- ³³ Crowson C, Matteson EL, Roger VL, et al. Usefulness of Risk Scores to Estimate the Risk of Cardiovascular Disease in Patients With Rheumatoid Arthritis. [Am J Cardiology](#) 2012; 110:420-4.

-
- ³⁴ Semb AG, Ikdahl E, Hisdal J, et al. Exploring cardiovascular disease risk evaluation in patients with inflammatory joint diseases. *Int J Cardiol* 2016; 223:331-6.
- ³⁵ Corrales A, González-Juanatey C, Peiró ME, et al. Carotid ultrasound is useful for the cardiovascular risk stratification of patients with rheumatoid arthritis: results of a population-based study. *Ann Rheum Dis*. 2014; 73:722-7.
- ³⁶ Rollefstad S, Kvien TK, Holme I, et al. Treatment to lipid targets in patients with inflammatory joint diseases in a preventive cardio-rheuma clinic. *Ann Rheum Dis* 2013; 72:1968-74.
- ³⁷ Rollefstad S, Ikdahl E, Hisdal J, et al. Systemic inflammation in patients with inflammatory joint diseases does not influence statin dose needed to obtain LDL cholesterol goal in cardiovascular prevention. *Ann Rheum Dis*. 2015; 74:1544-50.
- ³⁸ Otsuka Y, Kiyohara C, Kashiwado Y, et al. Effects of tumor necrosis factor inhibitors and tocilizumab on the glycosylated hemoglobin levels in patients with rheumatoid arthritis; an observational study. *PLoS One* 2018; 13(4).
- ³⁹ Haugeberg G, Uhlig T, Falch JA, et al. Bone mineral density and frequency of osteoporosis in female patients with rheumatoid arthritis: results from 394 patients in the Oslo County Rheumatoid Arthritis register. *Arthritis Rheum* 2000 43(3):522-30.
- ⁴⁰ Haugeberg G, Uhlig T, Falch JA, et al. Reduced bone mineral density in male rheumatoid arthritis patients: frequencies and associations with demographic and disease variables in ninety-four patients in the Oslo County Rheumatoid Arthritis Register. *Arthritis Rheum* 2000; 43:2776-84.
- ⁴¹ Ørstavik RE, Haugeberg G, Uhlig T, et al. Self-reported non-vertebral fractures in rheumatoid arthritis and population based controls: incidence and relationship with bone mineral density and clinical variables. *Ann Rheum Dis* 2004; 63:177-82.
- ⁴² Ørstavik RE, Haugeberg G, Mowinckel P, et al. Vertebral deformities in rheumatoid arthritis: a comparison with population-based controls. *Arch Intern Med* 2004; 23; 164:420-5.
- ⁴³ Furer V, Rondaan C, Heijstek M, et al. 2019 update of EULAR recommendations for vaccination in adult patients with autoimmune inflammatory rheumatic diseases. *Ann Rheum Dis* 2020;79:39-52.
- ⁴⁴ Winthrop L. Update on tuberculosis and other opportunistic infections associated with drugs blocking tumour necrosis factor α . *Ann Rheum Dis* 2005; 64 (suppl 4): 29-30.
- ⁴⁵ Rodríguez-Lozano B, González-Febles J, Garnier-Rodríguez J, et al. Association between severity of periodontitis and clinical activity in rheumatoid arthritis patients: a case– control study. *Arthritis Res Ther* 2019; 21:27.
- ⁴⁶ Nayak P, Luo R, Elting L, et al. Impact of Rheumatoid Arthritis on the Mortality of Elderly Patients Who Develop Cancer: A Population-Based Study. *Arthritis Care Res* 2017; 69:75-83.
- ⁴⁷ Van Den Hoek J, Boshuizen HC, Roorda LD, et al. Association of Somatic Comorbidities and Comorbid Depression With Mortality in Patients With Rheumatoid Arthritis: A 14-Year Prospective Cohort Study. *Arthritis Care Res* 2016; 68:1055-60.
- ⁴⁸ [National Rheumatoid Arthritis Society \(NRAS\)](#). Rheumatology in Wales, The State of Play, 2016.
- ⁴⁹ [UK Parliament](#). Early identification and diagnosis of rheumatoid arthritis. 2010.
- ⁵⁰ Marengo M and Suarez-Almazor M. Improving treatment adherence in patients with rheumatoid arthritis; what are the options *Int J Clin Rheumtol* 2015; 10:345–56.
- ⁵¹ Selmi C, Santis M, Gershwin M. Liver involvement in subjects with rheumatic disease. *Arthritis Res Ther*. 2011; 13:226.
- ⁵² Agca R, Heslinga S, Rollefstad S, et al. EULAR recommendations for cardiovascular disease risk management in patients with rheumatoid arthritis and other forms of inflammatory joint disorders: 2015/2016 update. *Ann Rheum Dis* 2017; 76:17-28.
- ⁵³ Mach F, Baigent C, Catapano A, et al. 2019 European Society of Cardiology / European Atherosclerosis Society Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J* 2019; 00:1-78.

-
- ⁵⁴ Andreu Sanchez JL, Balsa A, Gualda E, et al. GUIPCAR Group. Update of the clinical practice guideline for the management of rheumatoid arthritis in Spain. [Spanish Society of Rheumatology](#). 2011.
- ⁵⁵ André-Vert J, Guillez P, Scémama O. Clinical practice guidelines: Rheumatoid Arthritis - Medical, social and organisational aspects of treatment (excluding surgery and drugs). [Haute Autorite De Sante](#). 2017.
- ⁵⁶ Raghu G, Remy-Jardin M, Myerset J, et al. Diagnosis of Idiopathic Pulmonary Fibrosis. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline. [Am J Respir Crit Care Med](#). 2018; 198:44–68.
- ⁵⁷ Skold M. Idiopatisk Lungfibros, Vårdprogram. [Swedish Respiratory Society](#). 2019.
- ⁵⁸ Pilling S, Anderson I, Goldberg D, et al. Depression in adults with a chronic physical health problem: recognition and management. [NICE](#). 2009.