Sex-dependent Effects of Developmental Hypoxia on Cardiac Mitochondria from Adult Murine Offspring

DOI:

[10.1016/j.freeradbiomed.2020.11.004](https://doi.org/10.1016/j.freeradbiomed.2020.11.004)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Hellgren, K., Premanandhan, H., Quinn, C., Trafford, A., & Galli, G. (2020). Sex-dependent Effects of Developmental Hypoxia on Cardiac Mitochondria from Adult Murine Offspring. *Free Radical Biology and Medicine*. <https://doi.org/10.1016/j.freeradbiomed.2020.11.004>

Published in:

Free Radical Biology and Medicine

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Sex-dependent Effects of Developmental Hypoxia on Cardiac Mitochondria from Adult Murine Offspring

Kim T. Hellgren¹, Hajani Premanandhan¹, Callum J. Quinn¹, Andrew W Trafford¹
and Gina L. J. Galli*¹

¹Division of Cardiovascular Sciences, School of Medical Sciences, University of Manchester, Manchester M13 9NT, UK

*Corresponding Author

Dr Gina Galli

Division of Cardiovascular Sciences,

School of Medical Sciences,

University of Manchester,

3.09 Core Technology Facility,

46 Grafton St,

Manchester M13 9NT, UK

gina.galli@manchester.ac.uk

Abstract

Insufficient oxygen supply (hypoxia) during fetal and embryonic development can lead to latent phenotypical changes in the adult cardiovascular system, including altered cardiac function and increased susceptibility to ischemia reperfusion injury. While the cellular mechanisms underlying this phenomenon are largely unknown, several studies have pointed towards metabolic disturbances in the heart of offspring from hypoxic pregnancies. To this end, we investigated mitochondrial function in the offspring of a mouse model of prenatal hypoxia. Pregnant C57 mice were subjected to either normoxia (21%) or hypoxia (14%) during gestational days 6-18. Offspring were reared in normoxia for up to 8 months and mitochondrial biology was assessed with electron microscopy (ultrastructure), spectrophotometry (enzymatic activity of electron transport chain complexes), microrespirometry (oxidative phosphorylation and H_2O_2 production) and Western Blot (protein expression). Our data showed that male adult offspring from hypoxic pregnancies possessed mitochondria with increased H_2O_2 production and lower respiratory capacity that was associated with reduced protein expression of complex I, II and IV. In contrast, females from hypoxic pregnancies had a higher respiratory capacity and lower H_2O_2 production that was associated with increased enzymatic activity of complex IV. From these results, we speculate that early exposure to hypoxia has long term, sex-dependent effects on cardiac metabolic function, which may have implications for cardiovascular health and disease in adulthood.

Graphical Abstract

Keywords

Heart
Mitochondria
Developmental programming
Fetal hypoxia
Reactive oxygen species

67 **Abbreviations**

- 68 ADP - adenosine diphosphate
- 69 CS - citrate synthase
- 70 ETC – electron transport chain
- 71 ET_{CI+CII} - electron transfer capacity with complex I and II substrates
- 72 ET_{CII} - electron transfer capacity with complex II substrates
- 73 FCCP - carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone
- 74 GD – gestational day
- 75 H_2O_2 – hydrogen peroxide
- 76 HRP - horseradish peroxidase
- 77 I/R – ischemia reperfusion
- 78 $LEAK_{CI}$ - Leak respiratory state with complex I substrates
- 79 MPG - malate pyruvate and glutamate
- 80 O_2^- - superoxide
- 81 OXPHOS – oxidative phosphorylation
- 82 $OXPHOS_{CI}$ - oxidative phosphorylation with complex I substrates
- 83 $OXPHOS_{CI+CII}$ - oxidative phosphorylation with complex I and II substrates
- 84 PKC ϵ - protein kinase C epsilon
- 85 RCR - respiratory control ratio
- 86 ROS – reactive oxygen species
- 87 ROX - residual non-mitochondrial oxygen consumption
- 88 SOD - superoxide dismutase
- 89 TMPD - TM/AS, N,N,N,N-tetramethyl-p-phenylenediamin
- 90

1.0. Introduction

Heart disease remains the leading cause of death worldwide ¹. Despite significant advances in treatment options, prevention strategies are the most cost-effective way to reduce the socioeconomic burden of these diseases. Such strategies have traditionally targeted behavioral and lifestyle risk factors, such as smoking and obesity. However, the seminal work of Barker ² demonstrated that adverse events during pregnancy can predispose offspring to heart disease in adulthood. This phenomenon, termed developmental programming, provides a window of opportunity to prevent the development of heart disease in early life. Nevertheless, before effective treatments can be designed, it is crucial to understand the mechanisms leading to cardiac dysfunction in offspring from high-risk pregnancies.

Insufficient oxygen supply to an embryo or foetus, termed developmental hypoxia, occurs in a wide range of high-risk pregnancies, including preeclampsia, placental insufficiency, placental infection, maternal anaemia, gestational diabetes and high altitude pregnancy ^{3,4}. Animal models have shown the foetus initially responds to hypoxia by preferentially distributing blood flow to vital organs, such as the heart and brain ⁴⁻⁶. While this strategy is protective in the short term, a sustained redistribution of blood flow is associated with asymmetric fetal growth restriction, increased peripheral resistance and cardiac abnormalities, including hypertrophy ⁴. In adulthood, the hearts of offspring from hypoxic pregnancies continue to express abnormal phenotypes, including diastolic dysfunction, sustained increases in myocardial contractility and enhanced responsiveness to β -adrenoreceptor stimulation ⁷⁻⁹. Additionally, prenatal hypoxia appears to sensitise the adult heart to ischaemia and reperfusion (I/R) injury ¹⁰⁻¹². Interestingly, females from hypoxic pregnancies appear to be partially protected

from cardiac dysfunction, suggesting that the effects of developmental hypoxia are gender-specific ^{11,13,14}. In aggregate, these studies suggest developmental hypoxia programmes a dysfunctional cardiac phenotype in offspring that cannot be reversed by normalising oxygen availability after birth.

Given that most of these experiments have been conducted in isolated hearts, the cardiac dysfunction in offspring from hypoxic pregnancies cannot be explained by autonomic influences, altered peripheral resistance (cardiac afterload) or circulating catecholamines. Therefore, the intrinsic properties of the myocardial cells have been altered. While multiple cellular mechanisms may account for cardiac dysfunction, recent evidence suggests intrauterine stress can alter offspring cardiac metabolism, particularly at the level of the mitochondria. For example, nutritional stress during development leads to a range of cardiac mitochondrial abnormalities in fetal and adult offspring, including structural disorganization, impaired mitophagy, reduced oxygen consumption, decreased proton leak and altered fission/fusion dynamics ¹⁵⁻¹⁷. Similarly, exposure to excess glucocorticoids during development can programme cardiac mitochondrial dysfunction in adult offspring, leading to increased sensitivity to ischemia/reperfusion injury, higher levels of H₂O₂ production and a reduced capacity to produce ATP ¹⁸. Collectively, these studies suggest mitochondrial remodeling represents a major mechanism underlying the developmental programming of heart disease.

To our knowledge, only one group has investigated the long-term effects of developmental hypoxia on offspring mitochondrial function. Thompson and colleagues has shown pregnant guinea pigs exposed to 10.5% atmospheric oxygen between gestational days (GD) 28-65 (term 65-72 days) led to sex-specific alterations in

mitochondrial enzymatic complex activities, mitochondrial DNA content, protein expression and respiration¹⁹⁻²². However, the moderate level of hypoxia used in these studies (10.5%) caused a significant reduction in maternal weight, which is indicative of nutritional stress. It is therefore difficult to conclude whether mitochondrial function is being altered by developmental hypoxia or nutritional stress, or a combination of the two. To this end, we have undertaken a comprehensive investigation into the effects of developmental hypoxia on adult offspring mitochondrial morphology, respiratory capacity, reactive oxygen species production, enzymatic activity and protein expression. Our results suggest developmental hypoxia has long-term, sex-dependent effects on cardiac mitochondrial function; in particular, males from hypoxic pregnancies have a lower mitochondrial respiratory capacity and generate more H₂O₂ under basal conditions, and females from hypoxic pregnancies have greater respiratory scope and produce less H₂O₂ under basal conditions, compared to their normoxic counterparts. We speculate that these phenotypes have long-term implications for metabolic health and the susceptibility to heart disease.

2.0. Methods

2.1. Animal model

There are several strategies to model chronic fetal hypoxia, but most (e.g. placental embolisation, reduced uterine blood flow) cause impaired placental perfusion, thereby decreasing the delivery of nutrients as well as oxygen to the foetus^{23,24}. In these instances, it is difficult to separate the effects of fetal nutrient restriction versus fetal hypoxia in programming cardiovascular dysfunction in the offspring. To this end, we have utilised a rodent model of prenatal hypoxia developed by the Giussani laboratory

that does not affect maternal food intake, thereby allowing the effects of developmental hypoxia to be studied in isolation ^{8,25-27}.

2.2. Oxygen incubation protocols

All procedures comply with The UK Animals (Scientific Procedures) Act 1986 and EU directive 2010/63. The ARRIVE guidelines were followed for reporting the use of animals in scientific experiments. Local ethical approval was granted by The University of Manchester Animal Welfare Ethical and Review Board. Pregnant C57BL/J6 mice (aged 12 weeks, 24.1 ± 6.0 g), were bred, mated and maintained at The University of Manchester (UK). Mice were housed in standard Individually Ventilated Cages (IVC) with normal oxygen levels and a 12:12 light cycle with ad libitum food and water. The pregnant mice were randomly assigned to two groups; normoxia (N, 21% O₂) or hypoxia (H, 14% O₂). For hypoxic incubations, mice were transferred to an environmental chamber (Coy O₂ In Vivo Glove Box, Coy Laboratory Products, Grass Lake, MI) at gestational day (GD) 6 where they were subjected to 14% oxygen. Levels of humidity (60%), CO₂ (<1%) and temperature (22°C) were controlled throughout chamber incubation. Maternal food intake and water intake was monitored at regular intervals throughout the procedure, and maternal body weight was measured before and after chamber incubation. Mice were removed from the chamber at GD 18 and allowed to litter in normoxia (GD 21 $\pm$ 1 day). Litters were culled down to six (3 males and 3 females) to assure standardized maternal care and feeding. Sexing was done by visual determination of the presence or absence of dark pigmentation on the perineum ^{28,29}. Offspring were weaned from the mother at 3 weeks of age, and group housed in normal conditions. Experiments were performed on mouse offspring aged between 25-32 weeks.

2.3. *Electron microscopy*

We used electron microscopy to assess left ventricular mitochondrial morphology in adult mice aged between 30-32 weeks. 5 images were analyzed from each animal (n = 2 males and 2 females from normoxic pregnancies, and n = 5 males and 5 females from hypoxic pregnancies). In brief, hearts were removed from the animal, the atria were discarded, and the ventricles were cut longitudinally to separate the left and right chambers. A section of the left ventricular free wall was isolated and 2mm horizontal slices were taken from the mid myocardial layer; care was taken to isolate the same area from each animal. The slices were immediately fixed by immersion with 4% formaldehyde and 2.5% glutaraldehyde in 0.1M HEPES. The tissue was then removed from the fixation solution and prepared according to the Elisman protocol³⁰. Briefly, the tissue was stained with heavy metals, dehydrated (stepwise), infiltrated with Taab 812 Hard Resin (stepwise), embedded in silicon wells and finally polymerized at 60°C for 24 hours³¹. A single block was randomly selected from a bag and muscle orientation was determined in semithin sections. Ultrathin sections (up to 1mm², Reichert Ultracut ultramicrotome) were cut longitudinally in relation to the muscle fibres. Samples were then attached to 200 mesh copper EM grids and imaged at x890 magnification with a FEI Tecnai 12 Biotwin microscope at 80kV acceleration voltage with a Gatan Orius SC1000 CCD camera. A grid was randomly selected, and squares were given numerical values. A random number generator was used to select 5 grid squares per animal and the images were analyzed using the free hand tool in Image J (version 1.52k, National Institute of Health, MD). Mitochondria were traced and total mitochondrial area was expressed relative to the total area of the cell (students were blinded to the experimental sample).

2.4. Mitochondrial function

Mitochondrial function was investigated in male (n = 7 normoxic and 9 hypoxic) and female (n = 7 normoxic and 8 hypoxic) offspring aged between 26-31 weeks.

2.4.1. Mitochondrial oxygen consumption and H₂O₂ production

Mitochondrial respiration was assessed by high-resolution respirometry using an Oroboros Oxygraph-2k (Oroboros Instruments, Innsbruck, Austria) coupled to a fluorescent LED2-Module, allowing simultaneous measurement of O₂ consumption and H₂O₂ production, respectively. Mice were killed by cervical dislocation and the heart was immediately excised. ~50mg of fresh left ventricular tissue was homogenized (IKA Ultra-turrax T25) in mitochondrial respiration media (MiR05; containing: EGTA 0.5mM, MgCl₂ 3mM, K-MES 60mM, KH₂PO₄ 10mM, HEPES 20mM, sucrose 110mM and 1% BSA). 0.16 ± 0.04 mg of homogenised tissue was injected into each of the two chambers of the Oxygraph-2K for measurement of mitochondrial respiration and H₂O₂ production. The rest of the homogenate was frozen at -80°C for analysis of enzymatic function (see section 2.4.2). To measure H₂O₂ production, 10µM Amplex® UltraRed and 1U/ml horseradish peroxidase (HRP) were added to each chamber. Amplex® UltraRed oxidizes in the presence of H₂O₂ and forms resorufin, using HRP as a catalyst. Amplex® UltraRed was excited at 563nm and emission was read at 587nm. 5U/ml superoxide dismutase (SOD) was also added to the chambers to convert any extramitochondrial superoxide (O₂⁻) to H₂O₂.

Substrate inhibitor titration protocols (SUIT protocols) were designed according to Pesta et al ³² (see Fig. 1). Firstly, pyruvate (5mM), malate (2 mM) and glutamate (10mM) were added to achieve LEAK respiratory state with complex I (CI) substrates in the absence of

adenylates (LEAK_{CI}). When oxygen consumption was stable, saturating ADP (5mM) was injected to activate oxidative phosphorylation with CI substrates (OXPHOS_{CI}). Succinate (10mM) was then added to assess the additive effects of complex II (CII) substrates on oxidative phosphorylation (OXPHOS_{CI+CII}).

Figure 1: Original trace of simultaneous measurement of mitochondrial oxygen consumption and H₂O₂ production in mouse ventricular homogenate. Data is from a male offspring from a hypoxic pregnancy. Ventricular homogenate (0.1 mg ml⁻¹) was added to the chamber and a range of substrates and inhibitors were injected to investigate the electron transport chain (see methods section for details). Abbreviations; MPG, malate pyruvate and glutamate; ADP, adenosine diphosphate; SUC, succinate; FC, Carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone (FCCP); ROT, rotenone; AA, antimycin-A; O₂,

the chamber was opened for reoxygenation; TM/AS, N,N,N,N-tetramethyl-p-phenylenediamine (TMPD) and ascorbate; AZ, azide.

To uncouple mitochondria and assess maximum electron transfer capacity with CI+CII substrates (ET_{CI+CII}), carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone (FCCP) was titrated in steps to a final concentration of 0.1-0.3 μ M. Next, the CI inhibitor rotenone (0.5 μ M) was added to assess ET_{CI+CII} with CII substrates only. To block the ET-pathway and assess residual non-mitochondrial oxygen consumption (ROX), the complex III (CIII) inhibitor, antimycin A, was added (2.5 μ M). To assess complex IV (CIV) activity in isolation, the electron donor N,N,N',N'-tetramethyl-p-phenylenediamine (TMPD, 0.5 mM) was added in combination with ascorbate (2mM) to avoid autooxidation of TMPD. Lastly, the CIV inhibitor sodium azide (50mM) was added to assess background non-mitochondrial oxygen consumption from the addition of TMPD.

A separate protocol was used to assess the impact of CIV inhibition on H_2O_2 production in control adult mice ($n = 4$). Ventricular homogenates were incubated with substrates to achieve a steady-state OXPHOS_{CI+CII} (pyruvate, malate, glutamate, succinate and ADP) whilst measuring H_2O_2 (HRP, amplex red and SOD). Next, sodium azide was titrated in 0.25mM steps from 0 to 1.75 mM to selectively inhibit CIV.

All O_2 consumption and H_2O_2 production data were normalized to a marker of mitochondrial density, citrate synthase (CS) activity. When analyzing the effect of sodium azide on H_2O_2 production, H_2O_2 was expressed as a percentage of the amount of O_2 consumed (H_2O_2/O_2). To estimate mitochondrial efficiency of ATP production, the OXPHOS-coupling efficiency ratio was calculated as $1 - (LEAK_{CI} / OXPHOS_{CI})$, and the respiratory control ratio (RCR) was calculated as $OXPHOS_{CI} / LEAK_{CI}$ ³³.

294
295 2.4.2. *Spectrophotometric analysis of enzymatic activities and antioxidant capacity*

296 Spectrophotometric assays were undertaken on ventricular homogenates from male (n =
297 5 normoxic and 5 hypoxic) and female (n = 6 normoxic and 6 hypoxic) frozen tissue
298 samples from (aged between 25-29 weeks). Homogenates were assayed for protein
299 content, citrate synthase activity and enzymatic activity of electron transport chain
300 complexes. For all assays, absorbance was measured with a BioTek Synergy HTX
301 multimode reader (BioTek, Swindon, UK). For enzymatic assays, values are expressed
302 as maximum enzymatic activity per min divided by protein content. Protein content of
303 ventricular homogenates were assessed using the Quick start Bradford dye reagent kit
304 (Bio-Rad laboratories, Watford, UK); absorption was read at 550nm at 25°C.

305
306 Protocols for measuring CI, CII and CIV enzyme activities, as well as citrate synthase
307 activity, were designed according to Spinazzi et al. ³⁴. In brief, approximately 20-50mg of
308 frozen ventricular tissue was homogenized using a FastPrep-24™ 5G instrument (MP
309 Biomedicals, Santa Ana, CA) in solution containing 20mM TRIZMA-base, 40mM KCl,
310 2mM EGTA and 250mM Sucrose (pH 7.4). The tissue was homogenized in two cycles of
311 30 seconds with a 180-second pause in between each trial at 4°C. Samples were then
312 spun at 600g for 10 minutes at 4°C and the supernatant was stored in -80°C until the
313 day of the assay. Maximal enzymatic activity rate (V_{max}) was assessed over a ten-minute
314 period with a BioTek Synergy HTX multimode reader (BioTek, Swindon, UK) at 37°C.
315 The buffer components for each individual assay are given in Supplementary Table S1.

2.5. Mitochondrial protein expression of electron transfer chain complexes

Protein expression of complexes in the electron transport chain was measured with Western Blot in males from normoxic (n = 5) and hypoxic (n = 5) pregnancies, and females from normoxic (n = 5) and hypoxic (n = 5) pregnancies, aged between 26-32 weeks. The protocols have been described previously in detail ³⁵⁻³⁷. Following cardiac excision, a ~0.5 cm³ region of the left ventricular free wall was removed, snap frozen and stored in liquid nitrogen until analysis. Whole homogenates (~50 mg starting material) were prepared in RIPA buffer with protease and phosphatase inhibitors (0.1mg/ml phenylmethanesulphonylfuroide, 100 mmol/l sodium orthovanadate, 1 mg/ml aprotinin, 1 mg/ml leupeptin) and protein content was determined (DC Protein Assay, BioRad, UK). Proteins were separated by PAGE and transferred to nitrocellulose membranes. 10µg of protein was used for each sample, and membranes were blocked with 5% milk in TBS-T and incubated with the primary antibody cocktail (Abcam-110413, Cambridge, UK: 1:1000 concentration) and the secondary antibody IRDye® 800CW IgG_{2a}-Specific (Licor, UK: 1:20000 concentration). Membranes were visualized by chemiluminescence (Syngene, UK) or IR-Dye labeled secondary antibodies (Licor, UK). As the 'classical' housekeeping proteins can prove problematic with experimental treatments we opted for a total protein stain to normalize for sample loading, as suggested by Li et. Al., ³⁸, and an internal control to normalize between gels. Total protein transferred to the membrane was determined by REVERT total protein stain (Licor, UK). Blots were repeated in triplicate on separate occasions and data was averaged.

The antibody generated five separate bands, one for each protein corresponding to the five complexes of the ETC pathway (Fig. 1 supplementary material). The value for each separate protein in each sample was divided by the total value for the total protein as well as the internal control.

2.6. Calculations and statistics

Maternal BW, FI, WI are expressed as scatter plots (including all measured points) and statistically compared by fitting linear regression curves to compare slopes. Offspring BW is presented as scatter plots expressing means $\pm$ SEM and compared with mixed-effect analysis using maternal oxygen level as the random effect and the Tukey's correction for multiple comparisons test. For CIV inhibition, a linear regression curve was fitted to the data to assess if the slope was non-zero. For mitochondrial density where multiple observations (n) have been obtained from the same animal (N), linear mixed modeling (SPSS Statistics. IBM, USA) was performed thus accounting for the nested (clustered) design of the experiment e.g. multiple observations from the same heart (with treatment (hypoxic/normoxic), sex and animal as co-factors). All remaining data were statistically analyzed using a 2-way analysis of covariance (ANCOVA), with sex and intrauterine oxygen levels as independent variables, and age as a covariate (n-values and p-values are given in the figure legends). For mitochondrial H₂O₂ production, data were log-transformed to obtain a normal distribution prior to performing the ANCOVA.

3.0. Results

3.1. Maternal and offspring biometry

There were no differences in maternal body weight, water intake or food intake between normoxic and hypoxic dams (Fig. 2A-C), confirming that any programmed effects in the offspring are most likely due to hypoxia alone, rather than differences in maternal food consumption. There were no differences in offspring body weight between hypoxic and normoxic groups at any of the ages tested (Fig. 2D-E).

Figure 2: Effects of developmental hypoxia on maternal and fetal biometry. Maternal body weight (A), food (B) and water intake (C) were monitored in normoxic (green squares, $n = 16$) and hypoxic dams (blue squares, $n = 11$). Due to space restrictions within the environmental chamber, the body weight of pregnant dams could not be measured during the hypoxic incubation period (gestational days 6-18). Offspring body weight was also monitored in males (D, circles) from normoxic and hypoxic pregnancies ($n = 24$ offspring from 9 pregnancies and 23 offspring from 8 pregnancies, respectively) and females (E, triangles) from normoxic and hypoxic pregnancies ($n = 20$ offspring from 8 pregnancies and 20 offspring from 7 pregnancies, respectively) from birth to 15 weeks. Data are presented as scatter points for measurements of an individual animal. Statistics were run using a mixed-model effects with Tukey's correction for multiple comparisons.

3.2. Offspring mitochondrial oxygen consumption

Mitochondrial homogenate preparations were of good quality, as attested by high respiratory control ratios (9.2 ± 0.3) and OXPHOS-coupling efficiency ratios (0.89 ± 0.006) with complex I substrates (malate, pyruvate and glutamate). Mitochondrial oxygen consumption and H_2O_2 production responded to substrates and inhibitors in the expected manner³⁹ (Fig. 1), but FCCP had modest effects on mitochondrial respiration with only 9.3% of preparations responding positively. In the other preparations, even very low concentrations of FCCP ($<0.1 \mu M$) had no effect on oxygen consumption or caused a small inhibition (see Fig. 1A), suggesting that mouse cardiac homogenates are already operating at or near their theoretical maximum rate of oxygen consumption.

In male mice, developmental hypoxia reduced mitochondrial oxygen consumption in the $Leak_{CI}$ and $OXPHOS_{CI}$ states, as well as flux through CIV alone (Fig. 3, circular symbols). In contrast, developmental hypoxia increased mitochondrial oxygen consumption in female mice in all states, except for $LEAK_{CI}$ (Fig. 3, triangular symbols). Within the normoxic group, male mitochondrial oxygen consumption was higher than females in all respiratory states (Fig. 3, green symbols), but there were no sex-dependent differences in the hypoxic group (Fig. 3, blue symbols). There was no effect of treatment or gender on the OXPHOS-coupling efficiency ratio (Fig. 3F) or the RCR (data not shown).

Figure 3: Effect of developmental hypoxia on mitochondrial oxygen consumption. Mitochondrial oxygen consumption was measured in males from normoxic (green circles, $n = 6-7$) and hypoxic (blue circles, $n = 8-9$) pregnancies, and females from normoxic (green triangles, $n = 7$) and hypoxic (blue triangles, $n = 8$) pregnancies. Each panel represents a different respiratory state; (A) Leak respiration with substrates for complex I (LEAK_{CI}), (B) oxidative phosphorylation with substrates for complex I (OXPHOS_{CI}), (C) oxidative phosphorylation with substrates for Complex I and II (OXPHOS_{CI+II}), (D) electron transfer capacity with substrates for Complex II (ET_{CII}), (E) electron donation to Complex IV (CIV) and (F) OXPHOS-coupling efficiency ratios (1-(L/P)). Data are mean $\pm$ SEM. Asterix denotes a difference between mice from normoxic and hypoxic pregnancies, and Ψ denotes a difference between male and female mice (Two-way ANCOVA, one symbol = $p < 0.05$, two symbols = $p < 0.01$).

3.3. Offspring mitochondrial H_2O_2 production

In male mice, developmental hypoxia increased H_2O_2 production when it was measured in the LEAK_{CI} state and when CIII was inhibited with antimycin A (Fig. 4A and F, circular symbols). In contrast, females from hypoxic pregnancies produced less H_2O_2 compared to their normoxic counterparts in the OXPHOS_{CI+CII} state, and in the ET states (Fig. 4, triangular symbols). With regard to sex-dependent differences, normoxic male H_2O_2 production was higher than normoxic females in all respiratory states (Fig. 4, green symbols), but there were no differences in H_2O_2 production between hypoxic males and hypoxic females (Fig. 4, blue symbols).

Having discovered differences in mitochondrial oxygen consumption and H_2O_2 production between treatment groups, we next sought to determine the underlying mechanisms. In principle, changes in CIV oxygen consumption (Fig. 3E) may account for differences in basal H_2O_2 production between experimental groups. To explore this possibility, we performed experiments in control adult mice where we partially inhibited CIV with sodium azide and simultaneously measured H_2O_2 production (Fig. 5). We found that a dose-dependent inhibition of CIV caused a stepwise increase in H_2O_2 production.

Figure 4: Effects of developmental hypoxia on mitochondrial H_2O_2 production. H_2O_2 production was measured in males from normoxic (green circles, $n = 5$) and hypoxic (blue circles, $n = 5$) pregnancies, and females from normoxic (green triangles, $n = 5$) and hypoxic (blue triangles, $n = 6$) pregnancies. Each panel represents a different respiratory state; (A) Leak respiration with substrates for complex I ($LEAK_{CI}$), (B) oxidative phosphorylation with substrates for complex I ($OXPHOS_{CI}$), (C) oxidative phosphorylation with substrates for complex I and II ($OXPHOS_{CI+II}$), (D) electron transfer capacity with substrates for complex II (ET_{CII}), (E) electron transfer capacity with substrates for complex I+II (ET_{CI+II}), and (F) residual oxygen consumption (ROX) in the presence of antimycin A. Data are mean ± SEM. Asterisk denotes a statistically significant difference between mice from normoxic and hypoxic pregnancies, and Ψ denotes a difference between male and female mice (Two-way ANCOVA, one symbol = $p < 0.05$, two symbols = $p < 0.01$).

Figure 5: Relationship between complex IV activity and H_2O_2 production.

Oxygen consumption (A) was measured simultaneously with H_2O_2 production (B) during dose-dependent inhibition of complex IV with sodium azide. Data are presented as mean $\pm$ SEM, $n=4$ adult females from normoxic pregnancies in both panels. A linear regression curve was fitted to the data and was found to be significantly different from zero.

3.4. Offspring mitochondrial morphology

Differences in mitochondrial oxygen consumption and H_2O_2 production between experimental groups may also be explained by variable mitochondrial densities. Therefore, we investigated mitochondrial morphology with electron microscopy and measured the enzymatic activity of a common marker for mitochondrial content, citrate synthase. Developmental hypoxia had no effect on mitochondrial density or citrate synthase activity in any of the treatment groups (Fig. 6) and there were no differences between genders.

Figure 6: Effects of developmental hypoxia on mitochondrial density and citrate synthase activity. Transmission electron microscope images of left ventricular mitochondria in male offspring from normoxic (A, N = 2) and hypoxic (B, N = 5) pregnancies, and female offspring from normoxic (C, N = 2) and hypoxic (D, N = 5) pregnancies. Labels in image represent; M: mitochondria; Sr: sarcoplasmic reticulum; Scale bar: 1 μ m. Panel E-F show mean data $\pm$ SEM for mitochondrial density (n = 5 images per mouse) and citrate synthase activity (n = 5 offspring per group), respectively. For mitochondrial density, linear mixed modeling was performed to account for the nested (clustered) design of the experiment e.g. multiple observations from the same animal. For citrate synthase activity, a 2-way ANCOVA was performed. No statistical differences were found between any of the experimental groups.

3.5. Offspring mitochondrial enzymatic activity

Next, we investigated the possibility that differences in mitochondrial oxygen consumption and H₂O₂ production were underlined by variable enzymatic activities of complexes in the electron transport chain. We found that male offspring from hypoxic pregnancies had higher CI enzymatic activity (Fig. 7A), compared to their normoxic counterparts, and females from hypoxic pregnancies had lower CII activity (Fig. 7B) and higher CIV activity (Fig. 7C), compared to their normoxic counterparts.

Figure 7: Effects of developmental hypoxia on the activity of key mitochondrial enzymes. Enzymatic activity of complex I (A), complex II (B) and complex IV (C) were measured in males from normoxic (green circles, $n = 5$) and hypoxic (blue circles, $n = 5$) pregnancies, and females from normoxic (green triangles, $n = 5$) and hypoxic (blue triangles, $n = 6$) pregnancies. Data are presented as mean $\pm$ SEM. Asterisk signifies a statistically significant difference between mice from normoxic and hypoxic pregnancies (Two-way ANCOVA, one symbol = $p < 0.05$, two symbols = $p < 0.01$).

With regards to gender-specific effects, there were no differences in enzymatic activity between males and females in offspring from normoxic pregnancies, but females from hypoxic pregnancies had lower CI and CII activity, and higher CIV activity, compared to their male counterparts (Fig. 7A-C).

3.6. Offspring mitochondrial complex I-V protein expression

Lastly, we investigated the possibility that differences we observed in mitochondrial function could be explained by differential protein expression of electron transport chain complexes. CI, CII and CIV protein expression were lower in males from hypoxic pregnancies, compared to their normoxic counterparts (Fig. 8A, B and D), but there were no differences in the female group. Within the normoxic group, males had higher CI, CII and CIV protein expression compared to females (Fig. 8A, B and D), but there were no sex-specific effects within the hypoxic group.

Figure 8: Effect of developmental hypoxia on protein expression of respiratory chain Complexes. Protein expression was measured with Western Blot in males from normoxic (green circles, $n = 5$) and hypoxic (blue circles, $n = 5$) pregnancies, and females from normoxic (green triangles, $n = 5$) and hypoxic (blue triangles, $n = 6$) pregnancies. Each panel represents a different complex in the respiratory chain; complex I (A), II (B), III (C), IV (D) and V (E). Data are mean $\pm$ SEM. Asterisk denotes a difference between mice from normoxic and hypoxic pregnancies, and Ψ denotes a statistically significant difference between male and female mice (Two-way ANCOVA, one symbol = $p < 0.05$, two symbols = $p < 0.01$). The full Blot is given in supplementary Figure S1.

4.0. Discussion

This is the first comprehensive study to address the effects of developmental hypoxia on mitochondrial function in adult offspring. We show that male adult offspring from hypoxic pregnancies possess mitochondria with a reduced respiratory capacity, increased H_2O_2 production and a lower protein expression of CI, CII and CIV. In contrast, females from hypoxic pregnancies had a higher respiratory capacity, greater CIV enzymatic activity, and reduced H_2O_2 production, despite lower enzymatic activity of CII. From these results, we speculate that early exposure to hypoxia has long term sex-dependent effects on metabolic function, which may have implications for susceptibility to cardiac disease in adulthood.

4.2. Effects of developmental hypoxia on male offspring mitochondrial function

The first major and novel finding from this study is that male offspring from hypoxic pregnancies had greater levels of basal H_2O_2 production, compared to their normoxic counterparts. Previous studies have shown developmental hypoxia can cause oxidative stress in fetal²² and perinatal⁴⁰ mammals, but to our knowledge, this is the first study to show developmental hypoxia can programme ROS levels in adulthood. It is well-known that the over-production of mitochondrial ROS plays a major role in cardiac pathologies, particularly I/R injury⁴¹. Interestingly, several studies have shown developmental hypoxia programmes cardiac sensitivity to I/R injury in adult male offspring^{7,11,12}, partly due to the downregulation of protein kinase C epsilon (PKC ϵ) via DNA methylation¹³. In addition to differential expression of PKC ϵ , it is tempting to speculate that elevated basal H_2O_2 production sets a functional deficit in the hearts of males from hypoxic pregnancies, which may predispose them to postischemic oxidative stress.

Our data provides some information on the possible mechanism underlying the elevated basal H_2O_2 production in males from hypoxic pregnancies. H_2O_2 production in hypoxic males was higher than their normoxic counterparts in all respiratory states (although it only reached statistical significance in the $LEAK_{CI}$ state), and the difference was also apparent in the presence of rotenone and antimycin A, which block the CI_Q and $CIII_Q$ ROS sites, respectively ⁴². In the presence of these inhibitors, ROS production commonly occurs at the $CIII_{Qo}$ site, which is a major source of oxidative stress in the postischemic heart ⁴³. Nevertheless, pharmacological inhibition of the $CIII_{Qo}$ site (e.g. with myxothiazol or stigmatellin) would be necessary to confirm its involvement because other sites proximal to CIII can contribute to ROS production under these conditions ⁴². Furthermore, other factors may explain our results, including differential antioxidant profiles. Clearly, further research is necessary to confirm the site or source of elevated H_2O_2 production in males from hypoxic pregnancies.

In addition to H_2O_2 production, developmental hypoxia significantly reduced mitochondrial oxygen consumption in $LEAK_{CI}$ and $OXPHOS_{CI}$ states, as well as flux through CIV alone. The reduction in mitochondrial respiration was associated with a decrease in CI, CII and CIV protein expression, suggesting that these complexes are downregulated by developmental hypoxia, leading to lower respiratory capacity. These results are in agreement with data from Thompson's group ^{19,20} that showed developmental hypoxia reduces mitochondrial respiration rate and enzymatic activities of CI and CIV in adult male guinea pigs from hypoxic pregnancies. A reduction in mitochondrial respiratory capacity compromises ATP production, especially under situations of increased metabolic demand, such as exercise or disease. Therefore, in addition to elevated basal

H₂O₂ production, limitations in mitochondrial electron transport may contribute to cardiac dysfunction and sensitivity to I/R in males from hypoxic pregnancies.

The mitochondrial phenotype in males from hypoxic pregnancies may be explained by several factors. It is possible that developmental hypoxia directly altered fetal mitochondrial function and the phenotype persisted into adulthood. Indeed, several studies have shown intrauterine stress alters embryonic and fetal mitochondria, leading to the overproduction of ROS and a self-reinforcing cycle of mitochondrial dysfunction that persists into adulthood, and may even be inherited^{44,45}. In support of this idea, a previous study found evidence of oxidative stress in male fetal guinea pig hearts exposed to 10.5% oxygen, and this was associated with a reduction in the enzymatic activity of CIV²². Alternatively, developmental hypoxia may have altered fetal mitochondria indirectly due to its effects on cardiac structure. The foetus initially responds to hypoxia by preferentially distributing blood flow to the heart and brain, which eventually leads to increased peripheral resistance and cardiac remodeling⁴⁻⁶. This remodeling is apparent at the cellular level⁴⁶⁻⁴⁸, which may require mitochondrial alterations. Lastly, mitochondrial programming by intrauterine stress could also be achieved via stable epigenetic alterations to the nuclear genome⁴⁹. Indeed, recent studies have shown developmental hypoxia alters global DNA methylation patterns in rats leading to a reprogramming of the cardiac transcriptome, with a major focus on mitochondrial genes⁵⁰. Therefore, the observed differences in male mitochondrial protein expression in the present study may have epigenetic origins. Integrative longitudinal studies that monitor the effects of developmental hypoxia on mitochondrial function across the life course would help to discern between these possibilities.

4.2. Effects of developmental hypoxia on female offspring mitochondrial function

The second major and novel finding from this study is that adult female offspring from hypoxic pregnancies possess mitochondria with a higher respiratory capacity, increased mitochondrial efficiency and lower levels of basal H₂O₂ production. These results suggest developmental hypoxia programmes a mitochondrial phenotype in females that can generate ATP at a higher capacity and efficiency, while limiting ROS production. To our knowledge, this is the first study which has shown developmental hypoxia can programme a seemingly advantageous metabolic phenotype. From this perspective, it is interesting to note that several studies have shown female hearts are less affected by developmental hypoxia than males^{20,51,52}, and they also recover better from I/R injury^{7,11,12}. In theory, a greater mitochondrial respiratory capacity and lower H₂O₂ production could help to sustain ATP production and limit oxidative stress during oxygen and/or nutrient deprivation. Therefore, we propose that mitochondrial adaptations programmed by developmental hypoxia may protect the female heart from subsequent I/R stress in adulthood.

The increased respiratory capacity in females from hypoxic pregnancies was present under all respiratory states (except for Leak respiration), and with substrate combinations for CI and CII, as well as direct electron transfer to CIV with TMPD. Therefore, one mechanistic explanation is that developmental hypoxia increased mitochondrial density in female mice. However, analysis of cardiomyocyte EM images found no significant differences in mitochondrial content between females from normoxic or hypoxic pregnancies. While we acknowledge that our EM analysis has limitations and a more robust approach should be taken to confirm these morphological findings (i.e. using stereology or 3D reconstruction), we also found no differences in female citrate synthase activity, which is an excellent marker of mitochondrial content⁵³. Taken together, it seems unlikely that the differences in respiratory capacity between females

from normoxic and hypoxic pregnancies is related to mitochondrial content. An alternative explanation is that developmental hypoxia increases CIV activity in female mice. In support of this explanation, enzymatic activity of CIV was elevated in females from hypoxic pregnancies, compared to their normoxic counterparts, and this was associated with a trend towards higher CIV protein expression. In addition to increasing respiratory capacity, enhanced complex IV activity is also known to increase mitochondrial oxygen affinity which is beneficial for maintaining ATP production under conditions of low oxygen availability, such as I/R injury^{54,55}. Therefore, it would be interesting to investigate the effects of developmental hypoxia on female mitochondrial oxygen kinetics.

With regard to the mechanism underlying the reduced basal H₂O₂ production, enzymatic activity and protein expression of the main ROS producing complexes (CI and CIII) were not altered by developmental hypoxia in female mice. However, theoretically, the increase in CIV activity in females from hypoxic pregnancies could decrease the reduction of redox centers in CI or CIII, thereby reducing electron leak and ROS generation from these complexes⁵⁶. To explore this possibility, we performed experiments in control adult mice where we partially inhibited CIV with sodium azide and simultaneously measured H₂O₂ production. We found that a dose-dependent inhibition of CIV caused a stepwise increase in H₂O₂ production. This relationship has been demonstrated previously in isolated cardiomyocytes from embryonic chick cardiomyocytes where reduced CIV leads to enhanced ROS production⁵⁷. Therefore, greater CIV activity in females from hypoxic pregnancies may contribute to the observed reduction in H₂O₂ production in this experimental group. Nevertheless, there are other mechanistic explanations to explore, including differences in antioxidant capacity.

The mechanisms underlying mitochondrial programming in female mice are likely to be similar to males (discussed above), but the resultant phenotype is obviously very different. These results are strongly aligned to numerous studies that have demonstrated sex-dependent differences in the susceptibility of offspring to fetal stress (reviewed in^{45,58,59}). Identifying the underlying cause of these differences is an active area of research, but recent work suggests several fundamental differences between males and females may influence programming susceptibility, including; pattern of development (genetic, transcriptional and morphological), growth rate, sex hormones, placental plasticity, the regulation of epigenetic processes, metabolic hormones, the rate of ageing and lifespan^{45,58}. With regard to the cardiovascular system, several studies have shown oestrogen plays a protective role in the programming of hypertension, while testosterone is detrimental⁵⁹. Given that oestrogen is also known to modulate cardiac mitochondrial biogenesis, oxidative phosphorylation and ROS production⁶⁰, this hormone may have played a role in orchestrating the gender-specific responses in the present study. In this respect, it would be interesting to repeat the experiment in ovariectomized and castrated mice. There is also strong evidence that sex-dependent differences in placental plasticity play a major role in cardiovascular programming⁶¹. Indeed, numerous studies have shown developmental hypoxia leads to placental remodeling in rodents⁶²⁻⁶⁵, with female placentas adapting much better than males⁶⁶. Given the essential role that the placenta plays in the provision of fetal nutrients, sex-dependent placental remodeling may have important consequences for cardiac function and metabolism. Lastly, recent work has shown that epigenetic and transcriptomic signatures associated with developmental hypoxia are sex-dependent⁶⁷, and predominantly mitochondrial⁵⁰. Therefore, the gender-specific mitochondrial protein expression that we observed may represent persistent epigenetic marks.

4.3. Effects of gender on mitochondrial structure and function in offspring from normoxic pregnancies

Previous work has demonstrated substantial differences between males and females in mitochondrial structure and function, but the results are highly tissue-specific. In most tissues, including liver, brain, adipose and skeletal muscle, females exhibit a higher respiratory capacity than males, which is often associated with greater mitochondrial content⁶⁸. In contrast, two studies found no differences in cardiac mitochondrial respiratory capacity between male and female rats, despite a reduced mitochondrial content in females^{69,70}. Furthermore, a third study found mitochondrial respiration with glutamate and malate was higher in males compared to females⁷¹. In line with this latter study, we found male normoxic mice had a greater respiratory capacity than their female counterparts, and this difference was present in all respiratory states and substrate combinations, as well as CIV flux with TMPD. We go further to show the increase in male respiratory capacity was associated with higher male CI, CII and CIV protein expression, providing a mechanistic explanation for these sex-dependent differences.

In addition to respiratory capacity, previous work has shown sex-dependent differences in cardiac ROS production. In general, female cardiomyocytes generate less ROS than males at rest and after pathological stimuli, and they are also less prone to age-dependent oxidative stress^{69,72,73}. Some studies suggest the reduced ROS production in females is due to a superior antioxidant capacity⁷³, while others point towards reduced electron leak from CI and CIII⁶⁹. In contrast to these studies, normoxic males in our study had lower basal H₂O₂ production than their female counterparts in all respiratory states and substrate combinations. We cannot explain the discrepancy between these studies, but factors such as species, strain and age may play a role in determining sex-dependent differences.

777

778 **5.0. Conclusions and future directions**

779 In conclusion, our study has shown developmental hypoxia has long-term, sex-specific
780 implications for metabolism. We speculate that these differences may have implications
781 for disease susceptibility. For example, increased basal H₂O₂ production and lower
782 respiratory capacity in males from hypoxic pregnancies may predispose mitochondria to
783 cardiac dysfunction and I/R injury, a condition that is largely driven by oxidative stress ⁴¹.
784 In contrast, the greater respiratory capacity and lower H₂O₂ production in females from
785 hypoxic pregnancies may help to sustain ATP production and limit oxidative stress under
786 conditions of oxygen deprivation. These sex-specific differences in response to
787 developmental hypoxia may help to explain why females from hypoxic pregnancies are
788 less susceptible to I/R injury, compared to their male counterparts ^{11,12}. While purely
789 speculative at this stage, the present study provides the foundation to test this
790 hypothesis and design future gender-specific metabolic therapies to prevent cardiac
791 dysfunction in offspring from hypoxic pregnancies.

792

793 **6.0 Acknowledgements**

794 This study was funded by a PhD studentship awarded to Kim Hellgren by the Medical
795 Research Council Doctoral Training Partnerships. We would like to thank Aleksandr
796 Mironov, Dr. Christian Pinali and Samantha Forbes for their help in acquiring and
797 interpreting the EM images. In addition, we would like to thank Dr. Katherine Dibb, Dr
798 Javier Iglesias-Gonzalez and Professor David Eisner for their much-valued advice during
799 the study.

800 **Supplementary Tables and Figures**

801 **Table S1: Conditions for spectrophotometric assays of respiratory chain enzymes and citrate synthase activities in**
 802 **ventricular homogenates**

	CI	CII	CIV	CS
λ (nm)	600	600	550	412
Buffer	KP, 50mM	KP, 25 mM	KP, 50mM	TRIS, 100mM
pH	7.50	7.50	7.5	8.0
Substrates/electron acceptors	NADH 100 μM; DCPIP 80 μM; Ub-1 60 μM	Succinate 20 mM; DCPIP 80 μM; DUB 50 μM	Cyt c H ₂ ; 60 μM	DTNB 100 μM; Ac CoA 300 μM; Oxaloacetate 0.5mM
Other reagent(s)	BSA 3 mg ml ⁻¹ ; KCN 300 μM	BSA 1 mg ml ⁻¹ ; KCN 300 μM	KCN 300 μM	Triton X-100 0.1%;
Specific inhibitor	Rotenone 10 μM	Malonate 10 mM	KCN 300 μM	-

Abbreviations: CI, complex I; CII, complex II; CIV, complex IV; CS, citrate synthase; λ, selected wavelength for the assay; ε, extinction coefficient; Ac CoA, acetyl coenzyme A; BSA, fatty acid-free bovine serum albumin; Cyt c H₂, reduced cytochrome c; DCPIP, 2,6-dichlorophenolindophenol; DUB, decylubiquinol; DTNB, 5,5'-dithiobis(2-nitrobenzoic acid); KCN, potassium cyanide; KP, potassium phosphate buffer; Tris, Tris buffer; Ub1, biquinone1

Figure S1: Representative images of Western Blots for one replicate. (A) Each gel was loaded and imaged at two wavelengths; the samples and an internal control (historical left ventricular mouse sample that was used for all blots) were loaded with the antibody and imaged at 800nm (green bands: top to bottom; complex V, complex III, complex IV, complex II, and complex I). In addition, a ladder and the total protein stain, REVERT, was imaged on the same Blot at 700nm (red bands). (B-C) Original Blots for one replicate showing sample protein expression of Complexes V, III, IV, II and I (top to bottom) in an internal control (IC), normoxic female (NF, n = 5), hypoxic female (HF, n = 5), normoxic male (NM, n = 5) and hypoxic male (HM, n = 5). Averages for three replicates are given in Figure 8 of the main manuscript.

References

1. Dagenais, G.R., *et al.* Variations in common diseases, hospital admissions, and deaths in middle-aged adults in 21 countries from five continents (PURE): a prospective cohort study. *The Lancet*.
2. Barker, D.J.P., Osmond, C., Winter, P.D., Margetts, B. & Simmonds, S.J. WEIGHT IN INFANCY AND DEATH FROM ISCHAEMIC HEART DISEASE. *The Lancet* **334**, 577-580 (1989).
3. Hutter, D., Kingdom, J. & Jaeggi, E. Causes and Mechanisms of Intrauterine Hypoxia and Its Impact on the Fetal Cardiovascular System: A Review. *International Journal of Pediatrics* **2010**, 9 (2010).
4. Giussani, D.A. The fetal brain sparing response to hypoxia: physiological mechanisms. *The Journal of physiology* **594**, 1215-1230 (2016).
5. Cohn, H.E., Sacks, E.J., Heymann, M.A. & Rudolph, A.M. Cardiovascular responses to hypoxemia and acidemia in fetal lambs. *Am J Obstet Gynecol* **120**, 817-824 (1974).
6. Giussani, D.A., Spencer, J.A., Moore, P.J., Bennet, L. & Hanson, M.A. Afferent and efferent components of the cardiovascular reflex responses to acute hypoxia in term fetal sheep. *The Journal of physiology* **461**, 431-449 (1993).
7. Niu, Y., *et al.* Maternal Allopurinol Prevents Cardiac Dysfunction in Adult Male Offspring Programmed by Chronic Hypoxia During Pregnancy. *Hypertension* **72**, 971-978 (2018).
8. Giussani, D.A., *et al.* Developmental Programming of Cardiovascular Dysfunction by Prenatal Hypoxia and Oxidative Stress. *PloS one* **7**, e31017 (2012).
9. Zhang, L. Prenatal Hypoxia and Cardiac Programming. *Journal of the Society for Gynecologic Investigation* **12**, 2-13 (2005).
10. Rueda-Clausen, C.F., Morton, J.S., Lopaschuk, G.D. & Davidge, S.T. Long-term effects of intrauterine growth restriction on cardiac metabolism and susceptibility to ischaemia/reperfusion. *Cardiovascular research* **90**, 285-294 (2011).
11. Xue, Q. & Zhang, L. Prenatal Hypoxia Causes a Sex-Dependent Increase in Heart Susceptibility to Ischemia and Reperfusion Injury in Adult Male Offspring: Role of Protein Kinase C ϵ . *Journal of Pharmacology and Experimental Therapeutics* **330**, 624-632 (2009).
12. Li, G., *et al.* Effect of Fetal Hypoxia on Heart Susceptibility to Ischemia and Reperfusion Injury in the Adult Rat. *Journal of the Society for Gynecologic Investigation* **10**, 265-274 (2003).
13. Patterson, A.J., Chen, M., Xue, Q., Xiao, D. & Zhang, L. Chronic prenatal hypoxia induces epigenetic programming of PKC ϵ gene repression in rat hearts. *Circulation research* **107**, 365-373 (2010).
14. Shah, A., *et al.* Cardiovascular susceptibility to α -adrenergic stimulation in vivo; ischemic myocardial injury in male and female rat offspring exposed to prenatal hypoxia. *Clinical science* **131**, 2303 (2017).

- 865 15. Beauchamp, B., *et al.* Undernutrition during pregnancy in mice leads to
866 dysfunctional cardiac muscle respiration in adult offspring. *Bioscience*
867 *reports* **35**(2015).
- 868 16. Ferey, J.L.A., *et al.* A maternal high-fat, high-sucrose diet induces
869 transgenerational cardiac mitochondrial dysfunction independently of
870 maternal mitochondrial inheritance. *American journal of physiology. Heart*
871 *and circulatory physiology* **316**, H1202-h1210 (2019).
- 872 17. Larsen, T.D., *et al.* Diabetic Pregnancy and Maternal High-Fat Diet Impair
873 Mitochondrial Dynamism in the Developing Fetal Rat Heart by Sex-
874 Specific Mechanisms. *Int J Mol Sci* **20**, 3090 (2019).
- 875 18. Peng, J., *et al.* The detrimental effects of glucocorticoids exposure during
876 pregnancy on offspring's cardiac functions mediated by hypermethylation
877 of bone morphogenetic protein-4. *Cell Death & Disease* **9**, 834 (2018).
- 878 19. Thompson, L.P., Song, H. & Polster, B.M. Fetal Programming and Sexual
879 Dimorphism of Mitochondrial Protein Expression and Activity of Hearts of
880 Prenatally Hypoxic Guinea Pig Offspring. *Oxidative medicine and cellular*
881 *longevity* **2019**, 7210249 (2019).
- 882 20. Thompson, L.P., Chen, L., Polster, B.M., Pinkas, G. & Song, H. Prenatal
883 hypoxia impairs cardiac mitochondrial and ventricular function in guinea
884 pig offspring in a sex-related manner. *Am J Physiol Regul Integr Comp*
885 *Physiol* **315**, R1232-R1241 (2018).
- 886 21. Al-Hasan, Y.M., Pinkas, G.A. & Thompson, L.P. Prenatal Hypoxia
887 Reduces Mitochondrial Protein Levels and Cytochrome c Oxidase Activity
888 in Offspring Guinea Pig Hearts. *Reproductive Sciences* **21**, 883-891
889 (2014).
- 890 22. Al-Hasan, Y.M., *et al.* Chronic Hypoxia Impairs Cytochrome Oxidase
891 Activity Via Oxidative Stress in Selected Fetal Guinea Pig Organs.
892 *Reproductive Sciences* **20**, 299-307 (2013).
- 893 23. McMillen, I.C. & Robinson, J.S. Developmental origins of the metabolic
894 syndrome: prediction, plasticity, and programming. *Physiological reviews*
895 **85**, 571-633 (2005).
- 896 24. Morrison, J.L. Sheep models of intrauterine growth restriction: fetal
897 adaptations and consequences. *Clinical and experimental pharmacology*
898 *& physiology* **35**, 730-743 (2008).
- 899 25. Allison, B.J., *et al.* Divergence of mechanistic pathways mediating
900 cardiovascular aging and developmental programming of cardiovascular
901 disease. *FASEB journal : official publication of the Federation of American*
902 *Societies for Experimental Biology* **30**, 1968-1975 (2016).
- 903 26. Kane, A.D., Herrera, E.A., Camm, E.J. & Giussani, D.A. Vitamin C
904 prevents intrauterine programming of in vivo cardiovascular dysfunction in
905 the rat. *Circulation journal : official journal of the Japanese Circulation*
906 *Society* **77**, 2604-2611 (2013).
- 907 27. Giussani, D.A. & Davidge, S.T. Developmental programming of
908 cardiovascular disease by prenatal hypoxia. *Journal of Developmental*
909 *Origins of Health and Disease* **4**, 328-337 (2013).

- 910 28. Wolterink-Donselaar, I.G., Meeding, J.M. & Fernandes, C. A method for
911 gender determination in newborn dark pigmented mice. *Lab Anim (NY)* **38**,
912 35-38 (2009).
- 913 29. Deeney, S., Powers, K.N. & Crombleholme, T.M. A comparison of sexing
914 methods in fetal mice. *Lab Anim (NY)* **45**, 380-384 (2016).
- 915 30. Pinali, C., Bennett, H., Davenport, J.B., Trafford, A.W. & Kitmitto, A.
916 Three-Dimensional Reconstruction of Cardiac Sarcoplasmic Reticulum
917 Reveals a Continuous Network Linking Transverse-Tubules. *Circulation*
918 *Research* **113**, 1219-1230 (2013).
- 919 31. Deerinck, T.J., *et al.* Enhancing Serial Block-Face Scanning Electron
920 Microscopy to Enable High Resolution 3-D Nanohistology of Cells and
921 Tissues. *Microscopy and Microanalysis* **16**, 1138-1139 (2010).
- 922 32. Pesta, D. & Gnaiger, E. High-Resolution Respirometry: OXPHOS
923 Protocols for Human Cells and Permeabilized Fibers from Small Biopsies
924 of Human Muscle. Vol. 810 25-58 (Humana Press, Totowa, NJ, 2012).
- 925 33. Lemieux, H., *et al.* Impairment of mitochondrial respiratory function as an
926 early biomarker of apoptosis induced by growth factor removal. *bioRxiv*,
927 151480 (2017).
- 928 34. Spinazzi, M., Casarin, A., Pertegato, V., Salviati, L. & Angelini, C.
929 Assessment of mitochondrial respiratory chain enzymatic activities on
930 tissues and cultured cells. *Nature Protocols* **7**, 1235 (2012).
- 931 35. Lawless, M., *et al.* Phosphodiesterase 5 inhibition improves contractile
932 function and restores transverse tubule loss and catecholamine
933 responsiveness in heart failure. *Scientific Reports* **9**, 6801 (2019).
- 934 36. Briston, S.J., *et al.* Impaired beta-adrenergic responsiveness accentuates
935 dysfunctional excitation-contraction coupling in an ovine model of
936 tachypacing-induced heart failure. *J Physiol* **589**, 1367-1382 (2011).
- 937 37. Caldwell, J.L., *et al.* Dependence of cardiac transverse tubules on the
938 BAR domain protein amphiphysin II (BIN-1). *Circulation research* **115**,
939 986-996 (2014).
- 940 38. Li, R. & Shen, Y. An old method facing a new challenge: re-visiting
941 housekeeping proteins as internal reference control for neuroscience
942 research. *Life Sci* **92**, 747-751 (2013).
- 943 39. Makrecka-Kuka, M., Krumschnabel, G. & Gnaiger, E. High-Resolution
944 Respirometry for Simultaneous Measurement of Oxygen and Hydrogen
945 Peroxide Fluxes in Permeabilized Cells, Tissue Homogenate and Isolated
946 Mitochondria. *Biomolecules* **5**, 1319-1338 (2015).
- 947 40. Puente, B.N., *et al.* The oxygen-rich postnatal environment induces
948 cardiomyocyte cell-cycle arrest through DNA damage response. *Cell* **157**,
949 565-579 (2014).
- 950 41. Chouchani, E.T., *et al.* Ischaemic accumulation of succinate controls
951 reperfusion injury through mitochondrial ROS. *Nature* **515**, 431-435
952 (2014).
- 953 42. Brand, M.D. Mitochondrial generation of superoxide and hydrogen
954 peroxide as the source of mitochondrial redox signaling. *Free Radical*
955 *Biology and Medicine* **100**, 14-31 (2016).

- 956 43. Chen, Y.-R. & Zweier, J.L. Cardiac Mitochondria and Reactive Oxygen
957 Species Generation. *Circulation research* **114**, 524-537 (2014).
- 958 44. Gyllenhammer, L.E., Entringer, S., Buss, C. & Wadhwa, P.D.
959 Developmental programming of mitochondrial biology: a conceptual
960 framework and review. *Proceedings of the Royal Society B: Biological*
961 *Sciences* **287**, 20192713 (2020).
- 962 45. Dearden, L., Bouret, S.G. & Ozanne, S.E. Sex and gender differences in
963 developmental programming of metabolism. *Molecular Metabolism* **15**, 8-
964 19 (2018).
- 965 46. Botting, K.J., McMillen, I.C., Forbes, H., Nyengaard, J.R. & Morrison, J.L.
966 Chronic hypoxemia in late gestation decreases cardiomyocyte number but
967 does not change expression of hypoxia-responsive genes. *Journal of the*
968 *American Heart Association* **3**(2014).
- 969 47. Botting, K.J., *et al.* IUGR decreases cardiomyocyte endowment and alters
970 cardiac metabolism in a sex- and cause-of-IUGR-specific manner.
971 *American Journal of Physiology-Regulatory, Integrative and Comparative*
972 *Physiology* **315**, R48-R67 (2018).
- 973 48. Bae, S., Xiao, Y., Li, G., Casiano, C.A. & Zhang, L. Effect of maternal
974 chronic hypoxic exposure during gestation on apoptosis in fetal rat heart.
975 *Am J Physiol Heart Circ Physiol* **285**, H983-990 (2003).
- 976 49. Wadhwa, P.D., Buss, C., Entringer, S. & Swanson, J.M. Developmental
977 origins of health and disease: brief history of the approach and current
978 focus on epigenetic mechanisms. *Seminars in reproductive medicine* **27**,
979 358-368 (2009).
- 980 50. Gao, Y., *et al.* Multi-Omics Integration Reveals Short and Long-Term
981 Effects of Gestational Hypoxia on the Heart Development. *Cells* **8**, 1608
982 (2019).
- 983 51. Rueda-Clausen, C.F., Morton, J.S. & Davidge, S.T. Effects of hypoxia-
984 induced intrauterine growth restriction on cardiopulmonary structure and
985 function during adulthood. *Cardiovascular Research* **81**, 713-722 (2008).
- 986 52. Hemmings, D.G., Williams, S.J. & Davidge, S.T. Increased myogenic tone
987 in 7-month-old adult male but not female offspring from rat dams exposed
988 to hypoxia during pregnancy. *American Journal of Physiology-Heart and*
989 *Circulatory Physiology* **289**, H674-H682 (2005).
- 990 53. Larsen, S., *et al.* Biomarkers of mitochondrial content in skeletal muscle of
991 healthy young human subjects. *The Journal of physiology* **590**, 3349-3360
992 (2012).
- 993 54. Bienfait, H.F., Jacobs, J.M. & Slater, E.C. Mitochondrial oxygen affinity as
994 a function of redox and phosphate potentials. *Biochimica et biophysica*
995 *acta* **376**, 446-457 (1975).
- 996 55. Gnaiger, E. Bioenergetics at low oxygen: dependence of respiration and
997 phosphorylation on oxygen and adenosine diphosphate supply.
998 *Respiration Physiology* **128**, 277-297 (2001).
- 999 56. Dawson, T.L., Gores, G.J., Nieminen, A.L., Herman, B. & Lemasters, J.J.
1000 Mitochondria as a source of reactive oxygen species during reductive

- stress in rat hepatocytes. *The American journal of physiology* **264**, C961-967 (1993).
57. Duranteau, J., Chandel, N.S., Kulisz, A., Shao, Z. & Schumacker, P.T. Intracellular signaling by reactive oxygen species during hypoxia in cardiomyocytes. *J Biol Chem* **273**, 11619-11624 (1998).
 58. Aiken, C.E. & Ozanne, S.E. Sex differences in developmental programming models. *Reproduction (Cambridge, England)* **145**, R1-13 (2013).
 59. Intapad, S., Ojeda, N.B., Dasinger, J.H. & Alexander, B.T. Sex Differences in the Developmental Origins of Cardiovascular Disease. *Physiology* **29**, 122-132 (2014).
 60. Ventura-Clapier, R., Piquereau, J., Veksler, V. & Garnier, A. Estrogens, Estrogen Receptors Effects on Cardiac and Skeletal Muscle Mitochondria. *Front Endocrinol (Lausanne)* **10**, 557-557 (2019).
 61. Thornburg, K.L., O'Tierney, P.F. & Louey, S. Review: The placenta is a programming agent for cardiovascular disease. *Placenta* **31 Suppl**, S54-S59 (2010).
 62. Fisher, J.J., Bartho, L.A., Perkins, A.V. & Holland, O.J. Placental mitochondria and reactive oxygen species in the physiology and pathophysiology of pregnancy. *Clinical and Experimental Pharmacology and Physiology* **47**, 176-184 (2020).
 63. Nuzzo, A.M., *et al.* Placental Adaptation to Early-Onset Hypoxic Pregnancy and Mitochondria-Targeted Antioxidant Therapy in a Rodent Model. *The American Journal of Pathology* **188**, 2704-2716 (2018).
 64. Song, H., Telugu, B.P. & Thompson, L.P. Sexual dimorphism of mitochondrial function in the hypoxic guinea pig placenta†. *Biology of Reproduction* **100**, 208-216 (2018).
 65. Sferruzzi-Perri, A.N., Higgins, J.S., Vaughan, O.R., Murray, A.J. & Fowden, A.L. Placental mitochondria adapt developmentally and in response to hypoxia to support fetal growth. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 1621-1626 (2019).
 66. Cuffe, J.S.M., *et al.* Mid- to late term hypoxia in the mouse alters placental morphology, glucocorticoid regulatory pathways and nutrient transporters in a sex-specific manner. *The Journal of physiology* **592**, 3127-3141 (2014).
 67. Chen, X., Zhang, L. & Wang, C. Prenatal hypoxia-induced epigenomic and transcriptomic reprogramming in rat fetal and adult offspring hearts. *Scientific Data* **6**, 238 (2019).
 68. Ventura-Clapier, R., *et al.* Mitochondria: a central target for sex differences in pathologies. *Clinical science* **131**, 803-822 (2017).
 69. Colom, B., Oliver, J., Roca, P. & Garcia-Palmer, F.J. Caloric restriction and gender modulate cardiac muscle mitochondrial H₂O₂ production and oxidative damage. *Cardiovascular research* **74**, 456-465 (2007).
 70. Moulin, M., *et al.* Sexual Dimorphism of Doxorubicin-Mediated Cardiotoxicity. *Circulation: Heart Failure* **8**, 98-108 (2015).

- 1047 71. Ribeiro, R.F., *et al.* Sex differences in the regulation of spatially distinct
1048 cardiac mitochondrial subpopulations. *Mol Cell Biochem* **419**, 41-51
1049 (2016).
- 1050 72. Lagranha, C.J., Deschamps, A., Aponte, A., Steenbergen, C. & Murphy, E.
1051 Sex differences in the phosphorylation of mitochondrial proteins result in
1052 reduced production of reactive oxygen species and cardioprotection in
1053 females. *Circulation research* **106**, 1681-1691 (2010).
- 1054 73. Kander, M.C., Cui, Y. & Liu, Z. Gender difference in oxidative stress: a
1055 new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med*
1056 **21**, 1024-1032 (2017).
1057