Correlated pairs near the Fermi surface

DOI:

[10.1007/BFb0037545](https://doi.org/10.1007/BFb0037545)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Piechocki, W., Bishop, RF., & Stevens, GA. (1984). Correlated pairs near the Fermi surface. In H. Kümmel, & M. L. Ristig (Eds.), *Recent Progress in Many-Body Theories* (pp. 110-112). (Series on Lecture Notes in Physics; Vol. 198). Springer Nature. <https://doi.org/10.1007/BFb0037545>

Published in:

Recent Progress in Many-Body Theories

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

CORRELATED PAIRS NEAR THE FERMI SURFACE

W. Piechocki
Nuclear Theory Department
Institute of Nuclear Research
Hoza 69, PL-00-681 Warsaw, Poland

and

R.F. Bishop and G.A. Stevens
Department of Mathematics
UMIST, P.O.Box 88
Manchester M60 IQD, U.K.

One of the problems of the fifties, which occupied a lot of many body physicists dealing with strongly interacting Fermi systems of infinite extent like nucleon matter or liquid ^3He , was the occurrence of a singularity at the Fermi surface in the Bethe-Goldstone (BG) equation. It was generally believed, at that time, that this singularity was associated with the existence of a superfluid state¹.

In the next two decades we can observe a few developments². The approach I would like to discuss is the generalization of the BG equation. It includes, apart from particle-particle (pp) ladders, hole-hole (hh) ladders and all possible mixed pp and hh ladders. The generalized ladder equation of Mehta³ and the Galitskii-Feynman (GF) equation investigated by the Manchester group⁴ comprise the same type of ladder diagrams with the property that particles and holes are treated completely symmetrically. Mehta's equation does not lose the BG equation singularity. The Manchester group, on the other hand, has not noticed any problem near the Fermi surface but has discovered that the GF T-matrix possesses a pole in the bound-state region which corresponds to the formation of bound-state pairs in the medium.

Results obtained by using symmetrical ladder equation are interesting because we know that bound composite clusters of fermions comprised of an even number of fermions (particles) and/or an even number of their superfluid-like phase⁵. As an example we have the Cooper pairs which are responsible for superconductivity in weakly interacting Fermi systems.

We have decided to examine a generalized ladder equation again, but in the context of the coupled-cluster (CC) method; the CC method is flexible and general enough to enable investigation of the pairing from different points of view and with better and better approximation. We have solved⁶ the complete ladder (CLAD) equation of the CC method, corresponding to generalized ladder equations of the perturbation theory, for the separable two-body interaction. We have restricted ourselves to one-term S-wave interaction but the extension to both multi-term separable interactions and to arbitrary higher partial waves would

be straightforward. The solution for the states close to the Fermi surface contains both the bound-pairs and the singularity. To understand that result better we have examined excited states⁷. Using the Emrich's ansatz for the excited state, we have solved the ladder type equation corresponding to the CLAD equation of the ground state. The solution is characterized entirely by the quantities which describe the pairing of the ground state. For the attractive interaction, the excited state has lower energy than the ground state which means that our reference state chosen to be a filled Fermi-sphere plane-wave determinant is not the best possible point of departure. For the repulsive interaction, on the other hand, we observe a real excitation for densities above a certain critical density and no excitation below that density; in the latter range of density we also do not observe any pairing in the ground state.

Both results, for the ground state and for the excited state, suggest the existence of the pairing in a many-fermion strongly interacting system. The question arises whether the observed pairing exists in the exact solution of the Schrödinger equation. To examine that, we should find a suitable truncation scheme of the CC method which would give a possibility to see the pairing in its successive levels⁸. We cannot apply the Bochum truncation scheme⁹, because it was invented to describe the normal ground state of strongly interacting systems; by introducing a gap at k_F in the single-particle energy spectrum, this method relegates the pairing correlations near the Fermi surface to the terms of a very high order.

Presently we know how to treat the short-range repulsion of a two-body interaction, but we do not know how to describe properly the Fermi surface.

This research was supported in part by the Polish - U.S. Maria Sklodowska-Curie Fund and by the SERC of Great Britain.

References

- ¹J.S. Bell and E.J. Squires, Adv.Phys.10 (1961) 211 and references therein
- ²B.H. Brandow, Phys.Rev. 152 (1966) 863; G.A. Baker Jr. and J.L.Gammel, Phys.Rev. C6 (1972) 403; J.P. Jeukenne, A. Lejeune and C. Mahaux, Nucl.Phys. A 245 (1975) 411.
- ³M.L. Mehta, Nucl.Phys. 20 (1960) 533.

- ⁴R.F. Bishop and M.R. Strayer, J.Phys. G, L13 (1977) and reference therein.
- ⁵W. Kohn and D. Sherrington, Rev.Mod.Phys. 42 (1970) 1.
- ⁶R.F. Bishop, W. Piechocki and G.A. Stevens, to be published.
- ⁷R.F. Bishop, W. Piechocki and G.A. Stevens, to be published.
- ⁸W. Piechocki, Invited talk presented at 'The 7th (Pan-American) Workshop on Condensed Matter Theories', Altenberg, West-Germany, August 22 -27, 1983.
- ⁹B.D. Day and J.G. Zabolitzky, Nucl.Phys. A 366 (1981) 221.