Towards a coupled-cluster gauge-field approach to quantum hydrodynamics

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Arponen, J., Bishop, RF., Pajanne, E., & Robinson, NI. (1988). Towards a coupled-cluster gauge-field approach to quantum hydrodynamics. In J. S. Arponen, RF. Bishop, & M. Manninen (Eds.), *Condensed Matter Theories, Vol. 3* (pp. 51-66). Plenum Publishing Corporation.
[http://personalpages.manchester.ac.uk/staff/raymond.bishop/RFB_papers/\[055\] CMT_3\(1988\)51](http://personalpages.manchester.ac.uk/staff/raymond.bishop/RFB_papers/[055] CMT_3(1988)51)

Published in:

Condensed Matter Theories, Vol. 3

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Condensed **MATTER** **THEORIES**

VOLUME **3**

Edited by

Jouko S. Arponen

University of Helsinki
Helsinki, Finland

R. F. Bishop

University of Manchester Institute of Science and Technology
Manchester, United Kingdom

and

Matti Manninen

Helsinki University of Technology
Espoo, Finland

Plenum Press • New York and London

ISBN 0-306-42829-6
LC 87-656591

Proceedings of the 11th International Workshop on Condensed Matter Theories held
July 27-August 1, 1987, in Oulu, Finland

© 1988 Plenum Press, New York
A Division of Plenum Publishing Corporation
233 Spring Street, New York, N.Y. 10013

All rights reserved

No part of this book may be reproduced, stored in a retrieval system, or transmitted
in any form or by any means, electronic, mechanical, photocopying, microfilming,
recording, or otherwise, without written permission from the Publisher

Printed in the United States of America

TOWARDS A COUPLED CLUSTER GAUGE-FIELD APPROACH TO QUANTUM HYDRODYNAMICS

J. Arponen*, R.F. Bishop[†], E. Pajanne[‡] and N.I. Robinson[†]

*Department of Theoretical Physics, University of Helsinki
Siltavuorenpenger 20C, SF-00170 Helsinki, Finland

[†]Department of Mathematics
University of Manchester Institute of Science and Technology
P.O. Box 88, Manchester M60 1QD, England

[‡]Research Institute for Theoretical Physics
Siltavuorenpenger 20C, SF-00170 Helsinki, Finland

1. INTRODUCTION

The extended coupled cluster method (ECCM)^{1,2} has been discussed in an earlier volume of this series,³ where it was also applied to provide an effective gauge-field description of a charged impurity in a polarisable medium (e.g. positron in a metal).⁴ The purpose of the present paper is to apply the ECCM and comparable gauge field techniques to the zero-temperature condensed Bose fluid, in the most general case of arbitrary spatial inhomogeneity and arbitrary time dependence. Our principal aim is to derive the appropriate hydrodynamical balance equations for such local observables as the number density, current density and energy density. This is achieved by coupling the system to scalar and vector gauge fields, so that the theory may then be formulated in a completely gauge-invariant fashion to take fully into account the underlying $U(1)$ symmetry imposed by number conservation.

The ECCM formalism^{1-3,5,6} is based on the equations of motion for a set of linked-cluster amplitudes which completely characterise the many-body system. At the lowest level of truncation, we shall show explicitly that the formalism degenerates to the mean-field description of Gross⁷ and Pitaevskii⁸ for the condensate wavefunction or one-body order parameter. We particularly examine the decompositions of the general (off-diagonal) matrix elements of the one- and two-body density operators, in terms of these linked-cluster ECCM amplitudes. The various hydrodynamical balance equations are shown to follow straightforwardly. Furthermore, we show how these equations are also exactly satisfied by most practical approximation schemes which might be applied to the otherwise complete ECCM description. It is also of considerable practical importance that each of the ECCM amplitudes obeys the exact cluster property of vanishing when some subset of the cluster of particles described by the given amplitude becomes infinitely far removed from the remainder. The resulting "quasi-locality" of the basic ECCM amplitudes then implies that our gauge-invariant formalism is in principle capable of a complete hydrodynamical description of the system at zero temperature, including possible states of topological excitation or deformation, and of broken symmetry.

2. ELEMENTS OF THE ECCM

The $\exp(S)$ similarity transformation of the normal coupled cluster method (CCM)^{9,10} has proven to be extraordinarily powerful and versatile in the development of simple and finite algorithms for the ground-state energy of a wide variety of quantum many-body systems.¹¹ The desire to obtain approximations for the expectation values of other operators, which are properly compatible with the energy values, led to the introduction¹ of another operator S'' composed of a set of linked-cluster amplitudes. This correspondingly led to the consideration of a double similarity transformation and to the development of the so-called ECCM. We have shown⁶ how the method may be viewed as a biorthogonal formulation for a rather general quantal many-body problem. It is therefore best not to restrict ourselves from the start to Hamiltonians H or their relevant subsequent transforms that are necessarily manifestly hermitian.

Arbitrary time-dependent states $|\Psi(t)\rangle$ and $\langle\Psi'(t)|$ are now parametrised in the ECCM as,

$$\begin{aligned} |\Psi(t)\rangle &= e^{\phi(t)} e^{S(t)} |\Phi\rangle = e^{\phi(t)} e^{S(t)} e^{-S''(t)} |\Phi\rangle, \\ \langle\Psi'(t)| &= e^{-\phi(t)} \langle\Phi| e^{S''(t)} e^{-S(t)}, \end{aligned} \quad (1)$$

where $|\Phi\rangle$ is some suitable model state such that the algebra of all operators in the many-body Hilbert space is spanned by the two Abelian subalgebras of creation and destruction operators defined with respect to $|\Phi\rangle$. Thus, the subalgebra of creation (destruction) operators and the state $|\Phi\rangle$ ($\langle\Phi|$) are assumed to be cyclic in the sense that essentially all of the ket (bra) states of the space can be constructed by acting on $|\Phi\rangle$ ($\langle\Phi|$) with appropriate elements of the creation (destruction) operator subalgebra. The operators $S(t)$ and $S''(t)$ contain respectively only creation and only destruction operators (defined again with respect to the cyclic vector $|\Phi\rangle$),

$$S''(t)|\Phi\rangle = 0 = \langle\Phi|S(t), \quad (2)$$

and with corresponding amplitudes that are linked-cluster quantities. The factor $\phi(t)$ in Eq.(1) is a, largely inconsequential, c-number scale factor. The normalisation condition,

$$\langle\Psi'(t)|\Psi(t)\rangle = 1, \quad (3)$$

is manifestly preserved for all times, and the average-value functional $\langle A \rangle$ of an arbitrary operator A ,

$$\langle A \rangle \equiv \bar{A}(t) \equiv \langle\Psi'(t)|A|\Psi(t)\rangle \quad (4)$$

may evidently be written as the model-state expectation value of an operator $\hat{A}(t)$ which is the double similarity transform already mentioned of the operator A ,

$$\langle A \rangle = \langle\Phi|\hat{A}(t)|\Phi\rangle; \quad \hat{A}(t) \equiv e^{S''(t)} e^{-S(t)} A e^{S(t)} e^{-S''(t)}. \quad (5)$$

The linked-cluster amplitudes pertaining to the operators S and S'' are now regarded as independent variables which may be determined variationally by demanding stationarity of the action functional,

$$\mathcal{A} \equiv \int dt \langle\Phi| e^{S''(t)} e^{-S(t)} (i\partial/\partial t - H) e^{S(t)} |\Phi\rangle. \quad (6)$$

The stationarity conditions then yield the eigenvalue equations for the ground-state energy E_0 in the time-independent case,

$$\hat{H}|\Phi\rangle = E_0|\Phi\rangle \quad ; \quad \langle\Phi|\hat{H} = E_0\langle\Phi| \quad , \quad (7)$$

and the dynamical evolution equations for the operators S and S'' in the general case. The time-derivatives in Eq.(6) are easily taken, using the fact that the creation and destruction subalgebras are Abelian,

$$\dot{A} = \int dt [\dot{i}\langle\Phi|e^{S''}S|\Phi\rangle - \bar{H}] = \int dt [-i\langle\Phi|\dot{S}''e^{S''}S|\Phi\rangle - \bar{H}] \quad , \quad (8)$$

and where in the second form we have performed an integration by parts. The structure of Eq.(8) strongly suggests a change of independent operators from $S(t)$ and $S''(t)$ to a new pair $\Sigma(t)$ and $\tilde{\Sigma}(t)$, defined as

$$\Sigma(t)|\Phi\rangle \equiv (1 - |\Phi\rangle\langle\Phi|)e^{S''(t)}S(t)|\Phi\rangle \quad ; \quad \tilde{\Sigma}(t) \equiv S''(t) \quad , \quad (9)$$

in terms of which most of the resulting formalism has a more symmetric appearance. Thus, for example, the action takes the form,

$$A = \int dt [i\langle\Phi|\tilde{\Sigma}(t)\dot{\Sigma}(t)|\Phi\rangle - \bar{H}] = \int dt [-i\langle\Phi|\dot{\tilde{\Sigma}}(t)\Sigma(t)|\Phi\rangle - \bar{H}] \quad . \quad (10)$$

Henceforth we specialise to the case where the cyclic vector $|\Phi\rangle$ is the bare boson vacuum, and where the many-boson configuration space is parametrised by real-space coordinates $x \equiv (x^1, x^2, x^3)$. The basic single-boson creation and destruction field operators, a_x^+ and a_x respectively, obey the usual canonical commutation relations,

$$[a_x, a_y^+] = \delta^{(3)}(x-y) \equiv \delta(x-y) \quad . \quad (11)$$

We shall consistently denote coordinate-space three-vectors as $r, r', x_1, x_2, \dots$, and their corresponding infinitesimal volume elements as $dr, dr', dx_1, dx_2, \dots$. The basic ECCM operators $\Sigma(t)$ and $\tilde{\Sigma}(t)$ are now parametrised as,

$$\Sigma(t) = \sum_{n=1}^{\infty} \frac{1}{n!} \int dx_1 \dots \int dx_n \sigma_n(x_1 \dots x_n; t) a_{x_1}^+ \dots a_{x_n}^+ \quad (12)$$

$$\tilde{\Sigma}(t) = \sum_{n=1}^{\infty} \frac{1}{n!} \int dx_1 \dots \int dx_n \tilde{\sigma}_n(x_1 \dots x_n; t) a_{x_1} \dots a_{x_n} \quad ,$$

in which the fundamental amplitudes σ_n and $\tilde{\sigma}_n$ are all linked-cluster quantities, and which may, without loss of generality be assumed to be symmetric under arbitrary permutations of their coordinate-space arguments. We note that with these parametrisations, the states of Eq.(1) are not particle-number eigenstates, and we thus work from the outset in a number non-conserving formulation of the type introduced by Bogoliubov.¹² If we constrain ourselves to a Hilbert space of definite particle number, N , the identity operator has the following useful resolution

$$I = \sum_{n=0}^{\infty} \frac{1}{n!} \int dx_1 \dots \int dx_n a_{x_1}^+ \dots a_{x_n}^+ |\Phi\rangle\langle\Phi| a_{x_n} \dots a_{x_1} \quad , \quad (13)$$

wherein the $n = 0$ term is simply the model state projector $|\Phi\rangle\langle\Phi|$.

The ECCM formulation is now completely specified by the set of c-number amplitudes $\{\sigma(x_1 \dots x_n; t), \tilde{\sigma}(x_1 \dots x_n; t)\}$, which may be regarded as a complete set of dynamic variables. It is not difficult to see from Eqs.(5), (9) and (12) that the lowest (one-body) amplitudes σ_1 and $\tilde{\sigma}_1$ are precisely the usual "condensate wavefunctions", $\langle a_x \rangle = \sigma_1(x; t)$ and $\langle a_x^+ \rangle = \tilde{\sigma}_1(x; t)$, appropriate to a general non-uniform and temporally-varying macroscopically occupied condensate. All of the average-value functionals of Eqs.(4) and (5) may, for example, be fully specified in terms of the basic ECCM amplitudes, $A = A[\sigma_n, \tilde{\sigma}_n]$. Stationarity of the action of Eqs.(10) and (12) then also

easily leads to the fundamental evolution equations,

$$\begin{aligned} i\dot{\sigma}_n(x_1 \dots x_n; t) &= \delta \bar{H} / \delta \tilde{\sigma}_n(x_1 \dots x_n; t) \quad , \\ -i\dot{\tilde{\sigma}}_n(x_1 \dots x_n; t) &= \delta \bar{H} / \delta \sigma_n(x_1 \dots x_n; t) \quad ; \end{aligned} \quad (14)$$

where the functional derivatives of an arbitrary expectation value $\bar{A} = \bar{A}[\sigma_n, \tilde{\sigma}_n]$ are defined with respect to infinitesimal changes in accord with the earlier notation and the symmetry under permutations of the arguments of the basic ECCM amplitudes, as

$$\delta \bar{A} = \sum_{n=1}^{\infty} \frac{1}{n!} \int dx_1 \dots \int dx_n \left[\frac{\delta \bar{A}}{\delta \sigma_n(x_1 \dots x_n; t)} \delta \sigma_n(x_1 \dots x_n; t) + \frac{\delta \bar{A}}{\delta \tilde{\sigma}_n(x_1 \dots x_n; t)} \delta \tilde{\sigma}_n(x_1 \dots x_n; t) \right] \quad (15)$$

We discussed in an earlier volume in this series³ how the ECCM exactly maps an arbitrary quantal many-body problem into an equivalent classical Hamiltonian field theory in some well-defined complex symplectic phase space. The underlying classical fields in this formulation are just the c-number configuration-space amplitudes $\sigma(x_1 \dots x_n; t)$ and $\tilde{\sigma}(x_1 \dots x_n; t)$. It has been explicitly demonstrated that what underpins this equivalence is the exact ECCM result (which may be derived directly from the definitions already given³) that the average-value functional for the commutator of two arbitrary operators A and B , may be expressed in terms of a *generalised Poisson bracket* $\{ \bar{A}, \bar{B} \}$,

$$\langle \Psi'(t) | [A, B] | \Psi(t) \rangle = i \{ \bar{A}, \bar{B} \} \quad , \quad (16)$$

where

$$\begin{aligned} i \{ \bar{A}, \bar{B} \} \equiv \sum_{n=1}^{\infty} \frac{1}{n!} \int dx_1 \dots \int dx_n \left[\frac{\delta \bar{A}}{\delta \sigma_n(x_1 \dots x_n; t)} \frac{\delta \bar{B}}{\delta \tilde{\sigma}_n(x_1 \dots x_n; t)} - \frac{\delta \bar{A}}{\delta \tilde{\sigma}_n(x_1 \dots x_n; t)} \frac{\delta \bar{B}}{\delta \sigma_n(x_1 \dots x_n; t)} \right] \end{aligned} \quad (17)$$

We stress again that all of the basic amplitudes $\{ \sigma_n, \tilde{\sigma}_n \}$, which now completely specify the system, are linked-cluster quantities, and are hence quasi-local in the sense of obeying the cluster property.

Finally, in order to simplify the notation, we shall henceforth often drop the time label from, and thus leave implicit the time-dependence of, our basic amplitudes $\{ \sigma_n, \tilde{\sigma}_n \}$ and other dynamical variables constructed from them.

3. THE GAUGE-FIELD APPROACH

We now consider the application of the ECCM formalism to a Bose fluid composed of N identical uncharged bosons of mass m each, and interacting (in units with $\hbar = 1$) via two-body forces described by a Hamiltonian,

$$h_{(0)} = -\frac{1}{2m} \sum_{j=1}^N \nabla_j^2 + \sum_{i=1}^N \sum_{j=1}^{i-1} v(x_i - x_j) \quad (18)$$

Since the vacuum boson field theory already described does not conserve particle number, it is convenient to introduce a chemical potential, μ , and work

with the grand-canonical Hamiltonian $H_{(0)} \equiv h_{(0)} - \mu N$, with N the number operator. Furthermore, since the basic Hamiltonian of Eq.(18) does conserve particle number, we wish to formulate the theory in a *gauge-invariant* fashion so that the underlying $U(1)$ symmetry is preserved.

This is most naturally achieved from the outset by introducing *external gauge fields* and coupling the system to them.¹³ These fields are the usual scalar and vector potentials, $\phi(x,t) \equiv \phi(x)$ and $\vec{A}(x,t) \equiv \vec{A}(x)$ respectively, familiar from electrodynamics where they arise similarly due to charge conservation, and which couple respectively to the operators for the particle number density $\rho(x)$ and the canonical current density $\vec{j}(x)$,

$$\begin{aligned} \rho(r) &\equiv \rho(r,r) \quad ; \quad \rho(r,r') \equiv a_r^\dagger a_{r'} \\ \vec{j}(r) &\equiv \lim_{r' \rightarrow r} \frac{i}{2m} (\vec{\nabla}_r - \vec{\nabla}_{r'}) \rho(r,r') \end{aligned} \quad (19)$$

In this way we are led to study the Hamiltonian $H_{(0)} \rightarrow H$,

$$H = \sum_{j=1}^N \frac{1}{2m} [-i\vec{\nabla}_j - \vec{A}(x_j)]^2 + \sum_{j=1}^N [\phi(x_j) - \mu] + \sum_{i=1}^N \sum_{j=1}^{i-1} v(x_i - x_j) \quad , \quad (20)$$

which can be equivalently written in second-quantised form as

$$\begin{aligned} H &= H_{(0)} + \int dr \left[\left\{ \phi(r) + \frac{1}{2m} \vec{A}^2(r) \right\} \rho(r) - \vec{A}(r) \cdot \vec{j}(r) \right] \quad , \\ H_{(0)} &= \frac{1}{2m} \int dr (\vec{\nabla}_r a_r^\dagger) \cdot (\vec{\nabla}_r a_r) + \frac{1}{2} \int dr \left[v(r-r') a_r^\dagger a_{r'}^\dagger a_{r'} a_r - \mu \int dr a_r^\dagger a_r \right] \end{aligned} \quad (21)$$

The essence of the gauge-field approach is that by suitable choices of the gauge fields we can put the system and the observer into arbitrary relative motion. The coupling to the gauge fields may be thought of as inducing *local or differential Galilei transformations* in the system. Particularly simple examples are: (i) a global Galilei transformation to put the system into uniform motion with constant relative velocity $\vec{V}_0$ can be induced by the choice $\vec{A}(r) = m\vec{V}_0$, $\phi(r) = -\frac{1}{2}mV_0^2$; and this leads in the case of translationally-invariant systems to the wavefunction achieving an extra overall phase factor $\exp(i\vec{V}_0 \cdot \vec{P})$ where $\vec{P}$ is the total momentum operator which generates spatial translations; and (ii) the system can be put into uniform rotation with angular velocity $\vec{\Omega}$ by the choice $\vec{A}(r) = m\vec{\Omega} \times \vec{r}$, $\phi(r) = -\frac{1}{2}m|\vec{\Omega} \times \vec{r}|^2$; and this leads in the case of a rotationally-invariant system to the wavefunction acquiring an extra overall phase factor $\exp(i\vec{\Omega} \cdot \vec{L})$ where $\vec{L}$ is the total angular momentum operator which generates spatial rotations.

For the case of the more general local $U(1)$ gauge transformations, the Schrödinger wavefunction transforms in the usual way by acquiring an extra overall phase factor as,

$$\Psi(x_1 \dots x_N) \rightarrow \Psi'(x_1 \dots x_N) = \exp \left[-i \sum_{j=1}^N \Lambda(x_j, t) \right] \Psi(x_1 \dots x_N) \quad , \quad (22)$$

where $\Lambda(r,t) \equiv \Lambda(r)$ is some arbitrary gauge parameter or phase field. As usual, one easily sees from Eq.(20) that the transformed wavefunction Ψ' satisfies the usual time-dependent Schrödinger equation, with a Hamiltonian of the same form as Eq.(20) except with gauge fields which are correspondingly transformed as,

$$\vec{A} \rightarrow \vec{A}' = \vec{A} - \vec{\nabla} \Lambda \quad ; \quad \phi \rightarrow \phi' = \phi + \partial \Lambda / \partial t \quad . \quad (23)$$

It is clear that under these local Galilei transformations induced by the gauge fields, the (gauge-invariant) physical forces try to create local translational motions which vary both temporally and spatially throughout the system. A proper hydrodynamical description must then evidently be able to distinguish in a gauge-invariant fashion between, for example, the local hydrodynamical kinetic energy density of translation and the intrinsic kinetic energy density in the local rest frame. In the remainder of this paper we show how the ECCM can be used in this gauge-field approach to provide a complete (and, in principle, exact) hydrodynamical account of the system. Within the ECCM we have seen above that any aspect of the system can be described in terms of the linked-cluster amplitudes $\{\sigma_n, \tilde{\sigma}_n\}$. The gauge transformation of Eq.(22) may be equivalently specified in terms of them as,

$$\begin{aligned}\sigma_n(x_1 \dots x_n) &\rightarrow \sigma'_n(x_1 \dots x_n) = \exp \left[-i \sum_{j=1}^n \Lambda(x_j) \right] \sigma_n(x_1 \dots x_n), \\ \tilde{\sigma}_n(x_1 \dots x_n) &\rightarrow \tilde{\sigma}'_n(x_1 \dots x_n) = \exp \left[+i \sum_{j=1}^n \Lambda(x_j) \right] \tilde{\sigma}_n(x_1 \dots x_n).\end{aligned}\quad (24)$$

4. EVOLUTION OF THE ECCM AMPLITUDES: GENERALISED GROSS-PITAEVSKII THEORY

We now wish to examine more closely the dynamic equations (14). The average-value functional $\bar{H}$ for the Hamiltonian of Eqs.(20)-(21) is readily given in terms of the one- and two-body density matrices. The one-body density matrix is the expectation value of the one-body density operator of Eq.(19), $\rho(r, r') \equiv \langle \Psi(t) | \rho(r, r') | \Psi(t) \rangle$. Use of Eqs.(1) and (9) leads to the result,

$$\begin{aligned}\bar{\rho}(r, r') &= \langle \Phi | \tilde{\Sigma} a_r^\dagger I a_r, \Sigma | \Phi \rangle \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \int dx_1 \dots \int dx_n \tilde{\sigma}_{n+1}(rx_1 \dots x_n) \sigma_{n+1}(r'x_1 \dots x_n),\end{aligned}\quad (25)$$

where the second equality follows from the first by insertion of a resolution of the identity I of the form of Eq.(13) in the place so indicated.

The two-body density matrix is also needed to evaluate the expectation value $\bar{V}$ of the potential energy,

$$\bar{V}[\sigma_n, \tilde{\sigma}_n] = \frac{1}{2} \int dr \int dr' v(r-r') D(r, r') \quad ; \quad D(r, r') \equiv \langle a_r^\dagger a_r^\dagger a_r a_r \rangle. \quad (26)$$

For present purposes we shall not need the explicit functional form of $\bar{V} = \bar{V}[\sigma_n, \tilde{\sigma}_n]$, although for specific further applications the general two-body density matrix is probably most conveniently presented in terms of the *reduced subsystem amplitudes*^{1,14} χ_n and ϕ_n ,

$$\langle a_{r_1}^\dagger a_{r_2}^\dagger a_{r_3} a_{r_4} \rangle = \sum_{n=0}^{\infty} \frac{1}{n!} \int dx_1 \dots \int dx_n \phi_{n+2}(r_1 r_2; x_1 \dots x_n) \chi_{n+2}(r_3 r_4; x_1 \dots x_n), \quad (27)$$

where we have used a similar insertion of the unit operator as in Eq.(25), and where

$$\begin{aligned}\chi_{n+2}(rr'; x_1 \dots x_n) &\equiv \langle \Phi | a_{x_1} \dots a_{x_n} \hat{a}_r^\dagger \hat{a}_{r'}^\dagger | \Phi \rangle, \\ \phi_{n+2}(rr'; x_1 \dots x_n) &\equiv \langle \Phi | \hat{a}_r^\dagger \hat{a}_{r'}^\dagger a_{x_1} \dots a_{x_n} | \Phi \rangle.\end{aligned}\quad (28)$$

Explicit expressions for these reduced subsystem amplitudes in terms of the basic ECCM amplitudes $\{\sigma_n, \tilde{\sigma}_n\}$ are not needed here, but may be found using the functional derivative techniques described elsewhere^{3,5} to evaluate

various matrix elements involving double similarity-transformed operators of the form of Eq.(5).

After some straightforward manipulations, the equations of motion (14), with the Hamiltonian of Eqs.(20)-(21), can be more explicitly written as,

$$i \frac{d}{dt} \sigma_n(x_1 \dots x_n) = \sum_{j=1}^n h(x_j) \sigma_n(x_1 \dots x_n) + \frac{\delta \bar{V}}{\delta \tilde{\sigma}_n(x_1 \dots x_n)}, \quad (29)$$

$$-i \frac{d}{dt} \tilde{\sigma}_n(x_1 \dots x_n) = \sum_{j=1}^n \tilde{h}(x_j) \tilde{\sigma}_n(x_1 \dots x_n) + \frac{\delta \bar{V}}{\delta \sigma_n(x_1 \dots x_n)},$$

where the reduced one-body Hamiltonians are defined as,

$$h(r) \equiv -\frac{1}{2m} \nabla_r^2 + \frac{i}{m} \vec{A}(r) \cdot \vec{\nabla}_r + \frac{i}{2m} [\vec{\nabla}_r \cdot \vec{A}(r)] + \frac{1}{2m} \vec{A}^2(r) + \phi(r) - \mu, \quad (30)$$

$$\tilde{h}(r) \equiv -\frac{1}{2m} \nabla_r^2 - \frac{i}{m} \vec{A}(r) \cdot \vec{\nabla}_r - \frac{i}{2m} [\vec{\nabla}_r \cdot \vec{A}(r)] + \frac{1}{2m} \vec{A}^2(r) + \phi(r) - \mu.$$

We note that Eqs.(29)-(30) comprise a coupled set of nonlinear and non-local equations for the basic ECCM amplitudes. They may be regarded as a formally exact generalisation of the approximate nonlinear Gross-Pitaevskii equations^{7,8} for a weakly interacting condensed Bose fluid. The Gross-Pitaevskii equations are actually obtained as the lowest-order (SUB1) truncation of our equations (29), in which all of the amplitudes $\{\sigma_n, \tilde{\sigma}_n\}$ with $n > 1$ are set to zero. The resulting equations are precisely the self-consistent time-dependent Hartree equations for the amplitudes $\sigma_1(r)$ and $\tilde{\sigma}_1(r)$, which, as we have seen, are just the condensate wavefunctions. By approximating the two-body potential by a repulsive delta function form, the equilibrium (time-independent) version of the resulting local equation is the usual Gross-Pitaevskii equation with the well-known kink-soliton solutions in a one-dimensional geometry and comparable vortex solutions in a cylindrical geometry.¹⁵

5. THE ONE-BODY DENSITY MATRIX

The fundamental equations of motion (29) and (30) suffice to construct the time-evolution of the expectation value of an arbitrary operator. Special interest attaches to the one-body density matrix $\bar{\rho}(r, r')$. We note, for example, from Eq.(25) and the fact that each of the basic ECCM amplitudes $\{\sigma_n, \tilde{\sigma}_n\}$ obeys the cluster property, that $\bar{\rho}(r, r')$ behaves at large separations as,

$$\bar{\rho}(r, r') \xrightarrow{|\vec{r}-\vec{r}'| \rightarrow \infty} \tilde{\sigma}_1(r) \sigma_1(r'), \quad (31)$$

and hence exhibits the off-diagonal long-range order typical of superfluid systems.¹⁶ Its equation of motion is readily constructed, using the representation of Eq.(25), to give

$$\begin{aligned} \frac{\partial}{\partial t} \bar{\rho}(r, r') = & \frac{i}{2m} (\nabla_{r'}^2 - \nabla_r^2) \bar{\rho}(r, r') + \frac{1}{m} [\vec{A}(r) \cdot \vec{\nabla}_r + \vec{A}(r') \cdot \vec{\nabla}_{r'}] \bar{\rho}(r, r') \\ & + i [\phi(r) - \phi(r') + \frac{1}{2m} \{\vec{A}^2(r) - \vec{A}^2(r')\}] \bar{\rho}(r, r') \\ & + \frac{1}{2m} [\{\vec{\nabla}_r \cdot \vec{A}(r)\} + \{\vec{\nabla}_{r'} \cdot \vec{A}(r')\}] \bar{\rho}(r, r') + iC(r, r'), \end{aligned} \quad (32)$$

where the function $C(r, r')$ may be expressed in terms of the generalised

Poisson bracket defined in Eqs.(16) and (17) as,

$$C(r, r') = i\{\bar{V}, \bar{\rho}(r, r')\} \quad (33)$$

We also note that the result expressed in Eqs.(32) and (33) can be more directly derived by starting from the Heisenberg equation of motion for the operator $\rho(r, r')$, and taking its expectation value in the state $|\Psi(t)\rangle$. We can thus show either in this way or, more laboriously, by using the explicit form of the functional $\bar{V}[\sigma_n, \sigma_n]$, that the function $C(r, r')$ can be expressed in terms of the two-body density matrix as,

$$\begin{aligned} C(r, r') &= \frac{1}{2} \int dx_1 \int dx_2 v(x_1 - x_2) \langle [a_{x_1}^\dagger a_{x_2}^\dagger a_{x_2} a_{x_1}, a_{r'}^\dagger a_{r'}] \rangle \\ &= \int dx [v(r-x) - v(r'-x)] \langle a_x^\dagger a_{r'}^\dagger a_{r'} a_x \rangle \end{aligned} \quad (34)$$

We readily see from Eq.(25) that under the gauge transformation (24) previously discussed, the one-body density matrix transforms as,

$$\bar{\rho}(r, r') \rightarrow \bar{\rho}'(r, r') = e^{i[\Lambda(r) - \Lambda(r')]} \bar{\rho}(r, r') \quad (35)$$

and that both the diagonal term $\bar{\rho}(r, r)$ and, from Eqs.(26)-(28), the functional $\bar{V}[\sigma_n, \sigma_n]$ are gauge-invariant. It is now convenient to transform to relative and centre-of-mass coordinates,

$$\vec{\xi} \equiv \vec{r} - \vec{r}' \quad ; \quad \vec{R} \equiv \frac{1}{2}(\vec{r} + \vec{r}') \quad (36)$$

since our goal of deriving the hydrodynamical balance equations for the most important local physical field densities may most readily be attained by expanding the equation of motion (32) for $\rho(r, r') \equiv \bar{\rho}(R|\xi)$ in powers of ξ up to a given order, and with coefficients whose specific combinations in each order can be identified as being gauge-invariant.

The straightforward Taylor expansion,

$$\bar{\rho}(R|\xi) = \bar{\rho}(R) + \beta^a(R) \xi^a + \frac{1}{2} \gamma^{ab}(R) \xi^a \xi^b + \frac{1}{6} \delta^{abc}(R) \xi^a \xi^b \xi^c + O(\xi^4) \quad (37)$$

where $\bar{\rho}(R) \equiv \bar{\rho}(R, R)$ is the number density at space-point R , contains coefficients which are certainly not gauge-invariant. The roman superscripts $a, b, c, \dots$ in Eq.(37) are Cartesian three-vector indices; and the various coefficients $\gamma^{ab}, \delta^{abc}, \dots$ are assumed to be completely symmetric tensors, with no loss of generality. Summation over repeated indices is also assumed in Eq.(37) and henceforth. We note, for example, from Eq.(19) that the first order coefficient $\beta^a(R)$ may be written as,

$$\beta^a(R) = -imj^a(R) \quad (38)$$

in terms of the local canonical current density $j^a(R)$, which is definitely not gauge-invariant. By considering the gauge transformation properties expressed in Eqs.(35) and (23), it is not difficult to show that the following combinations of tensors, of first, second and third order respectively, are gauge-invariant,

$$J^a(R) = J^a \equiv j^a - (\bar{\rho}/m)A^a \quad (39a)$$

$$\Gamma^{ab}(R) = \Gamma^{ab} \equiv -(1/m^2)\gamma^{ab} - (1/\bar{\rho})j^a j^b \quad (39b)$$

$$\begin{aligned} \Delta^{abc}(R) = \Delta^{abc} &\equiv (1/2im^2)\delta^{abc} + (1/2m\bar{\rho})(j^a \gamma^{bc} + j^b \gamma^{ca} + j^c \gamma^{ab}) \\ &+ (m/\bar{\rho}^2)j^a j^b j^c + (\bar{\rho}/24m^2)(\partial^a \partial^b A^c + \partial^b \partial^c A^a + \partial^c \partial^a A^b) \end{aligned} \quad (39c)$$

where $\partial^a Z^b \equiv \partial Z^b / \partial R^a$ for an arbitrary vector $Z^a = Z^a(R)$. We note in particular that Eq.(39a) expresses the true (invariant) current density $\vec{J}(R)$ in terms of the (non-invariant) canonical current density $\vec{j}(R)$. In terms of these gauge-invariant combinations, the one-body density matrix has the convenient expansion,

$$\ln[\bar{\rho}(R|\xi)/\bar{\rho}(R)] = -i\xi^a [A^a_{mp} - \frac{1}{2}m^2 \rho^{-1} \xi^a \xi^b \Gamma^{ab}] - \frac{i}{3} \xi^a \xi^b \xi^c \left[-\frac{1}{24} (\partial^a \partial^b A^c + \partial^b \partial^c A^a + \partial^c \partial^a A^b) + m^2 \rho^{-1} \Delta^{abc} \right] + O(\xi^4) \quad (40)$$

Apart from the last term in Eq.(32) we are now ready to expand this equation of motion in powers of ξ^a . For this last term we may use the representation in Eq.(34). Expanding in gauge-invariant quantities, we may readily derive the expansion,

$$C(r, r') = -\xi^a \rho^{-1} F^a_{int} + \frac{1}{2} i \xi^a \xi^b \rho^{-1} \left[A^a_{F_{int}} + A^b_{F_{int}} + m W^{ab} \right] + O(\xi^3) \quad (41)$$

with the following definitions of the new gauge-invariant entities $\vec{F}_{int} = \vec{F}_{int}(R)$ and $W^{ab} = W^{ab}(R)$,

$$F_{int}(R) \equiv -\frac{1}{\rho(R)} \int dx \left[\vec{\nabla}_R v(R-x) \right] \langle a^\dagger_R a^\dagger_x a_x a_R \rangle \quad (42)$$

$$W^{ab}(R) \equiv -\frac{1}{\rho(R)} \int dx \left[\frac{\partial v(R-x)}{\partial R^a} \langle a^\dagger_x j^b(R) a_x \rangle + \frac{\partial v(R-x)}{\partial R^b} \langle a^\dagger_x j^a(R) a_x \rangle \right].$$

The operator $\vec{J}(R)$ in Eq.(42) is the gauge-invariant total current density operator, given by analogy with Eqs.(19) and (39a) as,

$$\vec{J}(R) = -\frac{i}{2m} \left[a^\dagger_R (\vec{\nabla}_R a_R) - (\vec{\nabla}_R a^\dagger_R) a_R \right] - \frac{1}{m} \vec{A}(R) a^\dagger_R a_R \quad (43)$$

It is useful for later purposes to consider the physical meaning of the above coefficients. From their definitions in Eq.(42) we see that: (i) $\vec{F}_{int}(R)$ represents the average force per particle due to the internal (pairwise) interactions in the presence of correlations; and (ii) since the gauge-invariant velocity field $\vec{u}(R)$ may be defined as,

$$\vec{u}(R) \equiv \vec{J}(R)/\rho(R) \quad (44)$$

then $W^{ab}(R)$ is the *symmetrised velocity-force correlation function*. Its trace, $W^{aa}(R)$, thus represents twice the average rate of increase of internal kinetic energy per particle at space-point R , due to the interparticle forces.

Finally, by inserting Eqs.(37) and (41) into the equation of motion (32) for the one-body density matrix, a comparison of the coefficients of the terms of zeroth, first and second order in ξ^a respectively, yields the following *local* evolution equations,

$$\frac{\partial \bar{\rho}}{\partial t} = -\partial^a j^a + \frac{1}{m} \bar{\rho} (\partial^a A^a) + \frac{1}{m} A^a (\partial^a \bar{\rho}) \quad (45a)$$

$$\frac{\partial j^a}{\partial t} = \frac{1}{m^2} \partial^b \gamma^{ab} + \frac{1}{m} \left[(\partial^a A^b) j^b + (\partial^b j^a) A^b \right] - \frac{\bar{\rho}}{m} \left[\partial^a \phi + \frac{1}{m} A^b (\partial^a A^b) \right]$$

$$+ \frac{1}{m} j^a (\partial^b A^b) + \frac{\bar{\rho}}{m} F^a_{int} \quad (45b)$$

and

$$\begin{aligned} \frac{\partial \gamma^{ab}}{\partial t} = & -\frac{i}{m} \partial^c \delta^{abc} + \frac{1}{m} [\gamma^{ac} (\partial^b A^c) + \gamma^{bc} (\partial^a A^c) + \gamma^{ab} (\partial^c A^c) + (\partial^c \gamma^{ab}) A^c] \\ & + j^a [m \partial^b \phi + A^c (\partial^b A^c)] + j^b [m \partial^a \phi + A^c (\partial^a A^c)] + \frac{1}{4m} \partial^c (\rho \partial^a \partial^b A^c) \\ & - \rho [A^a F_{int}^b + A^b F_{int}^a + m W^{ab}] \quad . \end{aligned} \quad (45c)$$

We shall now analyse Eqs.(45a-c) in some detail, and show that they lead to the *hydrodynamical balance equations* for the most important local physical observables.

6. HYDRODYNAMICS AND THE CONTINUITY EQUATIONS

In a hydrodynamic description, the most fundamental quantities that parametrise the system are the number, momentum and energy densities and the corresponding fluxes or currents that describe their flows. We show now how Eqs.(45a-c) relate respectively to the continuity or balance equations for these quantities.

6.1. Equation of Motion for the Number Density

Taking into account the definition (39a) of the total (gauge-invariant) current density $\vec{J}(R)$, we see easily that Eq.(45a) takes the usual form,

$$\partial \bar{\rho} / \partial t + \vec{\nabla} \cdot \vec{J} = 0 \quad , \quad (46)$$

of the *current continuity equation*. As is well known, this equation simply expresses the local form of the global number conservation law, $dN/dt = 0$. It is clearly gratifying to obtain this equation so readily since its correct imposition has been our primary objective in introducing the external gauge fields. We also stress that not only is Eq.(46) exactly true in the complete, untruncated formalism but, equally importantly, it also holds in most practical truncations that need to be made in real implementations of the ECCM. For example, a well-used approximation hierarchy is the so-called SUBn scheme in which the ECCM amplitudes $\{\sigma_i, \sigma_i^*\}$ with $i > n$ are set to zero, and where the remaining amplitudes with $1 \leq i \leq n$ are obtained as the solutions to the $2n$ coupled equations of motion (14) or (29)-(30). Since our derivation of Eq.(45a) proceeded via Eqs.(32) and (25), it should be immediately clear that Eq.(46) is also true in an arbitrary SUBn truncation.

6.2. Equation of Motion for the Momentum Density

It is not difficult to re-express Eq.(45b) in terms of gauge-invariant tensor quantities as,

$$\frac{\partial J^a}{\partial t} + \partial^b (J^a J^b / \bar{\rho}) = \frac{\bar{\rho}}{m} F^a \quad , \quad (47)$$

where we have also made use of the lower-order continuity equation (46). The vector field $\vec{F}$ in Eq.(47) is seen to comprise three separate pieces,

$$\vec{F}(R) = \vec{F} \equiv \vec{F}_{int} + \vec{F}_{ext} + \vec{F}_{kin} \quad , \quad (48)$$

where: (i) $\vec{F}_{int}$, defined in Eq.(42), has already been explained to be the average force per particle due to the internal interparticle forces of interaction; (ii) $\vec{F}_{ext}$ is the external force per particle due to the gauge fields, and which has the familiar (gauge-invariant) Lorentz form from the analogous and better-known application to electrodynamics,

$$\vec{F}_{ext} \equiv \vec{E} + \vec{u} \times \vec{B} \quad ; \quad \vec{B} \equiv \vec{\nabla} \times \vec{A} \quad , \quad \vec{E} \equiv -\vec{\nabla} \phi - \partial \vec{A} / \partial t \quad , \quad (49)$$

in terms of the velocity field $\vec{u} = \vec{u}(R)$ defined in Eq.(44); and (iii) $\vec{F}_{kin}$ is defined to be,

$$\vec{F}_{kin} = -m\rho^{-1}\partial^b\Gamma^{ab}, \quad (50)$$

and is hence proportional to the divergence of the (gauge-invariant) *kinetic stress tensor*, $\Gamma^{ab} = \Gamma^{ab}(R)$, which can be readily seen from Eq.(40) to be proportional to the second-order cumulant coefficient in the expansion for the one-body density matrix. The vector field $\vec{F} = \vec{F}(R)$ thus has the interpretation of being the total average force per particle at some space-point R in the fluid.

Equation (47) is clearly the equation of motion for the current or momentum density. Its physical content is perhaps more readily appreciated by rewriting it, using the usual definition of the convective derivative, d/dt ,

$$d/dt \equiv \partial/\partial t + \vec{u}^b\partial_b, \quad (51)$$

as

$$m[\rho^{-1}d\vec{J}/dt + \vec{u}(\vec{\nabla}\cdot\vec{u})] = \vec{F}. \quad (52)$$

A further combination of Eq.(52) with the lower-order continuity equation (46) then yields the very simple result,

$$m d\vec{u}/dt = \vec{F}, \quad (53)$$

which is precisely the correct Newtonian equation of motion for the system.

Before proceeding we point out again here that the equation of motion (47) or (53) is also true in SUBn approximation for the same basic reason as was discussed in the case of the lower-order continuity equation (46). Indeed both continuity equations remain true in even more drastic sub-approximations where some specific class of terms is further neglected from those which rightfully contribute to the potential energy expectation value $\bar{V} = \bar{V}[\sigma_n, \sigma_n]$. Further approximations of this type are often made in such realistic applications of the ECCM to physical systems with, for example, strongly repulsive (hard-core) interparticle forces. In these cases each such term contributing to $\bar{V}$ transforms under the gauge transformations in precisely the same way. The symmetry is thereby obeyed by each term separately, and the conservation laws hence continue to remain valid.

It is also worth pointing out that the equation of motion (47) can profitably be thought of as a conservation law which is the natural *generalisation of the well-known f-sum rule* away from the linear response regime. Thus, we can show by acting on the system at equilibrium with an infinitesimal sudden impulsive change in the external scalar potential, $\delta\phi(r,t) = \psi(r)\delta(t)$, that the corresponding infinitesimal change in the number density $\delta\rho(r,t)$ is continuous, but its time-derivative changes discontinuously at $t=0$ by an amount that can be calculated from Eqs.(46)-(50). By further relating the change in density, $\delta\rho$, as the system responds (linearly) from its equilibrium to an arbitrary small change $\delta\phi$ in the external scalar potential via the usual density-density response function, we can readily show that the linear response of our system obeys the f-sum rule. We point out again that the validity of the f-sum rule remains true even in the practical truncation or approximation schemes already discussed.

6.3. Equation of Motion for the Energy Density

We finally turn our attention to Eq.(45c), which we again wish first to express in terms of gauge-invariant tensors. It turns out to be especially

convenient to express the second-order equation of motion in terms of the particular gauge-invariant kinetic stress tensor K^{ab} , which is related to the kinetic energy density as we see below, and which is defined as

$$K^{ab}(R) = K^{ab} \equiv \frac{1}{2}m(\Gamma^{ab} - \rho u^a u^b) \quad (54)$$

A straightforward but rather lengthy calculation employing the lower-order equations of motion (46) - (47), yields the conservation law,

$$\begin{aligned} \frac{\partial K^{ab}}{\partial t} + \partial^c \left[\Delta^{abc} + u^a K^{bc} + u^b K^{ac} + u^c K^{ab} - m \rho u^a u^b u^c \right. \\ \left. + \frac{\bar{\rho}}{24m^2} (\epsilon^{bcd} \partial_a B^d + \epsilon^{acd} \partial_b B^d) \right] \\ = \frac{1}{2} \bar{\rho} (W^{ab} + u^a E^b + u^b E^a) + m^{-1} (\epsilon^{bcd} K^{ac} + \epsilon^{acd} K^{bc}) B^d, \end{aligned} \quad (55)$$

where ϵ^{abc} is the usual antisymmetric unit tensor of third rank, and $\vec{E}$ and $\vec{B}$ are the external force fields of Eq.(49).

We finally show how Eq.(55) can be used to study the conservation law of total energy in differential form. The global energy conservation law is easily obtained by calculating the time derivative of $\bar{H} = \bar{H}[\sigma_n, \sigma_n; t]$ from Eq.(15) as,

$$\begin{aligned} \frac{d\bar{H}}{dt} = \frac{\partial \bar{H}}{\partial t} + \sum_{n=1}^{\infty} \frac{1}{n!} \int dx_1 \dots \int dx_n \left[\frac{\delta \bar{H}}{\delta \sigma_n(x_1 \dots x_n)} \dot{\sigma}_n(x_1 \dots x_n) \right. \\ \left. + \frac{\delta \bar{H}}{\delta \tilde{\sigma}_n(x_1 \dots x_n)} \dot{\tilde{\sigma}}_n(x_1 \dots x_n) \right] \quad (56) \end{aligned}$$

From the basic equations of motion (14), it immediately follows that $d\bar{H}/dt = \partial \bar{H}/\partial t$, and hence that the total energy is conserved, $d\bar{H}/dt = 0$, in the case when the gauge fields are time-independent. From the corresponding local form which we now investigate, we shall identify the energy current density and the local energy source density in the general case when the external gauge fields are time-dependent.

The total Hamiltonian (20) or (21) is the sum, $H = H_1 + V$, of a one-body part H_1 and a two-body part V due to pairwise interactions. The one-body contribution to the expectation value $\bar{H}$ is expressed in terms of a local energy density $\epsilon_1(R)$,

$$\bar{H}_1 = \int dR \epsilon_1(R) \quad (57)$$

which may readily be evaluated to give the explicit form,

$$\epsilon_1(R) = \frac{1}{8m} \nabla_R^2 \bar{\rho}(R) + \frac{1}{2} m \Gamma^{aa}(R) + \frac{1}{2} m \bar{\rho}(R) \vec{u}^2(R) + \bar{\rho}(R) [\phi(R) - \mu] \quad (58)$$

The third term, $\frac{1}{2} m \rho u^2$, represents the hydrodynamical kinetic energy density of average translational motion; while the second term, $\frac{1}{2} m \text{Tr}(\Gamma)$, together with the first term, represents the kinetic energy density in the local rest frame. (We note that since the first term in Eq.(58) is a perfect divergence, it is strictly speaking redundant, since its contribution to $\bar{H}_1$ is zero. Its inclusion does however ensure the positivity of the kinetic energy density.) We note also that each of the terms in Eq.(58) is separately gauge-invariant, except for the potential energy term, $\rho\phi$, due to the external scalar potential.

Although the two-body energy $\bar{V}$ cannot simply be similarly expressed

in terms of a local energy density, $\epsilon_2(R)$, we now show that its time-derivative may be comparably decomposed. From Eq.(26) we first calculate,

$$\begin{aligned}\partial D(r, r') / \partial t &= -i \langle [a_{r'}^\dagger a_r^\dagger, H] \rangle \\ &= -\vec{\nabla}_r \cdot \langle a_{r'}^\dagger \vec{J}(r) a_r \rangle - \vec{\nabla}_{r'} \cdot \langle a_r^\dagger \vec{J}(r') a_{r'} \rangle, \quad (59)\end{aligned}$$

using the Heisenberg equation of motion. We thus find the result,

$$\partial \bar{V} / \partial t = \frac{1}{2} \int dr' v(r-r') \partial D(r, r') / \partial t = (\partial / \partial t) \int dr \epsilon_2(r), \quad (60)$$

where we have defined $\epsilon_2(r)$ in terms of its time-derivative as,

$$\begin{aligned}\frac{\partial \epsilon_2(r)}{\partial t} &\equiv - \int dr' v(r-r') \vec{\nabla}_r \cdot \langle a_{r'}^\dagger \vec{J}(r) a_r \rangle \\ &= -\frac{1}{2} \bar{\rho}(r) W^{aa}(r) - \vec{\nabla}_r \cdot \vec{P}(r), \quad (61)\end{aligned}$$

and where we have used the definition (42) and the new definition,

$$\vec{P}(r) \equiv \int dr' v(r-r') \langle a_{r'}^\dagger \vec{J}(r) a_r \rangle. \quad (62)$$

The first term in Eq.(61) represents the average rate of decrease of the kinetic energy density due to the interparticle interactions, as already discussed above; while the local field $\vec{P}$ in the second term is clearly an energy flux due to the internal forces. It is defined so that the interaction energy of any pair of particles is asymmetrically associated wholly with one of them.

We finally combine the above results as,

$$\partial \bar{H} / \partial t \equiv \int dr \partial \epsilon(r) / \partial t = \int dr [\partial \epsilon_1(r) / \partial t + \partial \epsilon_2(r) / \partial t]. \quad (63)$$

The total energy density $\epsilon(r)$ now satisfies a continuity equation which can readily be found from Eqs.(58) and (61), and by employing our previous gauge-invariant equations of motion (55) and (46), in the form,

$$\frac{\partial \epsilon}{\partial t} + \vec{\nabla} \cdot \vec{J}_\epsilon = S_\epsilon, \quad (64)$$

where the total energy flux vector, $\vec{J}_\epsilon$, is given by,

$$\begin{aligned}\vec{J}_\epsilon(r) = \vec{J}_\epsilon &\equiv \frac{1}{8m} \nabla^2 \vec{J} + \left[\frac{m}{2\bar{\rho}} \text{tr } \Gamma + \frac{1}{2} m u^2 + \phi - \mu \right] \vec{J} \\ &+ (\Delta^{aab} + m u \Gamma^{ab}) \hat{r}^b - \frac{\bar{\rho}}{12m^2} \vec{\nabla} \times \vec{B} + \vec{P}, \quad (65)\end{aligned}$$

with $\hat{r}^b$ a unit vector in the b-direction; and where the energy source density is given by,

$$\begin{aligned}S_\epsilon(r) = S_\epsilon &\equiv \bar{\rho} \frac{\partial \phi}{\partial t} - \vec{J} \cdot \frac{\partial \vec{A}}{\partial t} \\ &= \bar{\rho} \frac{d\phi}{dt} + \vec{J} \cdot \vec{E}. \quad (66)\end{aligned}$$

In the second form for S_ϵ above we have used the convective derivative of Eq.(51). We note finally that due to the appearance of terms involving the scalar potential ϕ , none of the quantities $\epsilon(r)$, $\vec{J}_\epsilon(r)$ and $S_\epsilon(r)$ is gauge-invariant. We may trivially rewrite the energy density balance equation in the equivalent form however,

$$\frac{\partial}{\partial t} (\epsilon - \bar{\rho} \phi) + \vec{\nabla} \cdot (\vec{J}_\epsilon - \phi \vec{J}) = \vec{J} \cdot \vec{E}, \quad (67)$$

by combining it with the current continuity equation (46), where now the subtracted energy density $(\epsilon - \rho\phi)$ and flux $(\vec{J}_\epsilon - \vec{J}\phi)$ are fully gauge-invariant, as is the source term $\vec{J} \cdot \vec{E}$.

7. SUMMARY AND DISCUSSION

The ECCM formally decomposes a quantum many-body problem into a non-local classical field theory for a set of interacting, non-local, n -body fields $\{\sigma_n, \sigma_n^*\}$, $n = 1, 2, \dots$, which in the coordinate-space representation considered here are n -point functions of arguments $(x_1, x_2, \dots, x_n)$.^{3,5} The lowest-order amplitudes, $\sigma_1(x) = \langle a^\dagger \rangle$ and $\sigma_1(x) = \langle a \rangle$, are just the condensate wavefunctions for the Bose fluid,¹⁵ and are clearly local by definition, since they depend on a single space coordinate. However, the higher-order amplitudes are also of a conceptually similar character due to their linked-cluster or connected property, which ensures that they vanish as the relative separation of any two spatial arguments approaches infinity. For example, $\sigma_2(x, y) = \langle a^\dagger a^\dagger \rangle_{\text{connected}} = \langle a^\dagger a^\dagger \rangle - \langle a^\dagger \rangle \langle a^\dagger \rangle$. Then, for sufficiently long-wavelength phenomena in the infra-red limit, the relevant values of the relative separations will be small in comparison with any natural scale lengths. In this way the basic ECCM amplitudes $\{\sigma_n, \sigma_n^*\}$ may be regarded as a set of *generalised quasi-local order parameters*, by analogy with the more conventional and strictly local (one-body) order parameters of condensed matter theory. The total energy expectation value, $\bar{H} = \bar{H}[\sigma_n, \sigma_n^*]$ is thus an almost local functional of these classical (c-number) quasi-local order parameters.

From the above viewpoint it is natural to compare the ECCM with both the *phenomenological Ginzburg-Landau theory*¹⁵ and the *density-functional theory*.^{17,18} In Ginzburg-Landau phenomenology, the total energy (or free energy at non-zero temperatures) is expressed as a strictly local fourth-order functional of the order parameters $\langle a^\dagger \rangle$ and $\langle a \rangle$. We stress however that whereas this approach is intended only to be valid close to the relevant phase transition where the order parameters are small, the present ECCM by contrast provides an exact microscopic description of a pure state at zero temperature. Similarly, in the density-functional theory of Hohenberg, Kohn and Sham,^{17,18} the total energy is again constructed as a functional of another local parameter, namely the particle number density $\rho(x)$. In its simplest manifestations, such as the Thomas-Fermi approximation, the total energy is taken to be a completely local functional of $\rho(x)$. We may thus view the ECCM as a rather different approach to incorporate in a very precise and exact manner the non-local corrections to these simple local approximations, and with which virtually all of the later work in density-functional theory has been concerned.

In the present work we have described an application of the ECCM to a condensed Bose fluid. We have concentrated on the properties of a single trajectory in the ECCM phase space, where a representative point in this symplectic ECCM phase space is fully characterised by the set of amplitudes $\{\sigma_n, \sigma_n^*\}$. The statistical mechanics of a physical system is nowadays often described in terms of the totality of all such possible trajectories -- namely the whole *phase portrait*. The present treatment may thus be viewed as the first step towards a more comprehensive statistical-mechanical description. In this first step however we have already shown how the individual trajectories fully comply with the pertinent gauge symmetries and the local hydrodynamical conservation laws for such physically relevant variables as the local densities for particle number, momentum and energy.

We have seen both how the ECCM in principle exactly generalises the earlier Gross-Pitaevskii formalism for the condensate wavefunctions, and how it provides a complete hydrodynamical description of the zero-temperature

condensed Bose fluid. In this latter regard, the locality features inherent to the ECCM make it powerful enough to incorporate a proper description of *topological excitations* as the vortex lines observed in liquid ^4He . Such excitations are created by specific topological boundary conditions on the amplitudes $\{\sigma_n, \sigma_n\}$ which prevent their decay by any quasi-local processes. These boundary conditions may only be properly imposed in such a theory as the ECCM in which the cluster property is exactly obeyed by all relevant amplitudes. The boundary condition is thus typically imposed at some sufficiently large distance -- from the vortex core in the case of a vortex line, for example, and where the phases of σ_n and σ_n increase or decrease respectively by $2\pi n$ when the centre-of-mass coordinate, $R \rightarrow \infty$, winds once about the core. In this way, all physical observables for the excitation are ultimately describable in terms of the lowest amplitudes σ_1 and σ_1 -- just as in Ginzburg-Landau theory -- since the higher amplitudes cannot obtain long-range contributions due to their observance of the cluster property.

In conclusion, we have demonstrated that the ECCM leads to a complete and correct hydrodynamical description of a zero-temperature condensed Bose liquid, which incorporates the possibility of describing topological excitations and non-equilibrium dynamics. The hydrodynamical continuity equations are valid not only in the complete description, but, most importantly, also in the usual practical truncation schemes necessary to implement it. Their gauge-invariant form has further enabled us to provide a description which, for example, correctly separates the average (hydrodynamical) kinetic energy of translation from the average kinetic energy in the local rest frame. By contrast, other microscopic treatments have often found real difficulty in obtaining this separation.

The present approach has clearly demonstrated the applicability of the ECCM into manifestly non-perturbative areas, far removed from its origins.^{9,10} We hope that further work along these lines will focus on the geometrical aspects of the underlying ECCM phase space in order to develop, for example, new approximation schemes that go beyond such truncations described here which still hark back to the perturbation-theoretical roots.

REFERENCES

1. J. Arponen, Ann.Phys. (NY) 151: 311 (1983).
2. J. Arponen and E. Pajanne, in: "Recent Progress in Many-Body Theories," H. Kummel and M. L. Ristig (eds.), Lecture Notes in Physics Vol. 198, Springer-Verlag, Berlin (1984), p.319.
3. J. Arponen, R. F. Bishop and E. Pajanne, in: "Condensed Matter Theories," Vol.2, P. Vashishta, R. Kalia and R. F. Bishop (eds.), Plenum, New York (1987), p. 357.
4. J. Arponen, R. F. Bishop and E. Pajanne, in: "Condensed Matter Theories," Vol. 2, P. Vashishta, R. Kalia and R. F. Bishop (eds.), Plenum, New York (1987), p. 373.
5. J. Arponen, R. F. Bishop and E. Pajanne, "Extended coupled cluster method: I. Generalised coherent bosonisation as a mapping of quantum theory into classical Hamiltonian mechanics," Phys.Rev. A36: 2519 (1987).
6. J. Arponen, R. F. Bishop and E. Pajanne, "Extended coupled cluster method: II. Excited states and generalised random phase approximation," Phys.Rev. A36: 2539 (1987).
7. E. P. Gross, Ann.Phys. (NY) 4: 57 (1958); Nuovo Cim. 20:454 (1961).
8. L. P. Pitaevskii, Sov. Phys.-JETP 13: 451 (1961).
9. F. Coester, Nucl.Phys. 7: 421 (1958).
10. F. Coester and H. Kummel, Nucl.Phys. 17: 477 (1960).
11. R. F. Bishop and H. Kummel, Physics Today 40 (No.3): 52 (1987).
12. N. N. Bogoliubov, J.Phys. USSR 11: 23 (1947).

13. Ta-Pei Cheng and Ling-Fong Li, "Gauge Theory of Elementary Particle Physics," Clarendon Press, Oxford (1984), p.229.
14. H. Kummel, K. H. Lührmann and J. G. Zabolitzky, Phys.Rep. 36C: 1 (1978).
15. A. L. Fetter and J. D. Walecka, "Quantum Theory of Many-Particle Systems," Academic Press, New York (1965).
16. G. Rickayzen in: "The Helium Liquids," J. G. M. Armitage and I. E. Farquhar (eds.), Academic Press, London (1975), p.95.
17. P. Hohenberg and W. Kohn, Phys.Rev. 136: B864 (1964).
18. W. Kohn and L. J. Sham, Phys.Rev. 140: A1133 (1965).