Schmidt Hammer exposure dating (SHED): Establishment and implications for the retreat of the last British Ice Sheet

DOI:

[10.1016/j.quageo.2016.02.002](https://doi.org/10.1016/j.quageo.2016.02.002)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Tomkins, M., Dortch, J., & Hughes, P. (2016). Schmidt Hammer exposure dating (SHED): Establishment and implications for the retreat of the last British Ice Sheet. *Quaternary Geochronology*, 33(46-60), 46-60. <https://doi.org/10.1016/j.quageo.2016.02.002>

Published in:

Quaternary Geochronology

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Schmidt Hammer exposure dating (SHED): Establishment and implications for the retreat of the last British Ice Sheet

M.D. Tomkins^a, J.M. Dortch^{a*}, P.D. Hughes^a

^a Department of Geography, School of Environment, Education and Development, University of Manchester, Cryosphere Research At Manchester, M13 9PL Manchester, United Kingdom

Corresponding author: Jason.Dortch@manchester.ac.uk (J.M. Dortch)

Keywords:

British Ice Sheet

TCN dating

Schmidt hammer exposure dating

Glacial chronology

Abstract

Ninety-eight surfaces related to the retreat of the British Ice Sheet (BIS) and dated using Terrestrial Cosmogenic Nuclide Dating (TCND) were sampled using the Schmidt Hammer to expand on relative dating techniques and establish Schmidt Hammer exposure dating (SHED) as an effective method for dating glacial landforms in the UK. The BIS is an effective analogue for contemporary glacial systems but our understanding of its retreat under changing climate conditions is constrained by a limited number of dates obtained from existing methods (¹⁴C, OSL). These methods are restricted in their application to glacial environments and while TCND has addressed this to some degree, its cost and potential for outliers encourage the establishment of new techniques. SHED fulfils this requirement by providing a cost-efficient method for obtaining numerous direct ages that are of comparable accuracy and precision to TCND. A multi-lithology approach has established that many rock types are unsuitable for numerical dating. However, a robust calibration curve was generated ($R^2 = 0.81$, $p = < 0.01$) for granite surfaces and applied to 31 undated granite erratics on Shap Fell, NW England. SHED indicates that BIS retreat occurred at 16.5 ± 0.5 ka, a conclusion which supports our current understanding of regional deglaciation and indicates that SHED can be a valuable and cost-effective geochronological tool.

1. Introduction

The British Ice Sheet (BIS), which reached its maximum extent by 27 ka (Sejrup *et al.*, 2005), is of particular importance as an analogue for contemporary glacial systems (Clark *et al.*, 2012). Our understanding of BIS extent is relatively robust (Figure 1) as recent research (Sejrup *et al.*,

2005; Bradwell *et al.*, 2009b) has demonstrated the persistence of extensive off-shore ice cover and the extension of the Fennoscandian Ice Sheet across the North Sea as first reconstructed by Geikie (1894) and Croll (1875) respectively. However, due to limited direct evidence for these early assessments (Clark *et al.*, 2012), most reconstructions proposed that off-shore ice was extremely restricted, an assessment popularised by Bowen *et al.*, (1986), although this has been challenged by recent research (e.g. Scourse *et al.*, 2009; Ballantyne, 2010). BIS retreat was highly dynamic and spatially variable, as retreat fundamentally shifted from marine to terrestrial terminating margins (Clark *et al.*, 2012) and was punctuated by major readvance events (Boulton *et al.*, 2002). As such, the geomorphological record is highly fragmented as ice expansion during readvance events such as the Wester Ross Readvance (WRR) (Ballantyne *et al.*, 2009a) and Loch Lomond Stadial (LLS) (Golledge, 2010) reworked and removed previous evidence of BIS retreat; thus preventing the straightforward coupling of glacial and climatic records. Even though climatic datasets are available, of sufficient resolution, and cover an appropriate timespan to facilitate analysis of the geomorphological record (e.g. Lowe *et al.*, 2008), robust comparison is not possible without precise and accurate geochronometric dating to constrain the nature and timing of BIS deglaciation.

In response to this, many relative and radiometric dating methods have been used to form the basis for reconstructions of BIS retreat (Ballantyne, 2010; Chiverrell and Thomas, 2010; Clark *et al.*, 2012). The broad agreement of these studies suggests that they are relatively robust assessments, although the research community is not without disagreement (Bowen *et al.*, 2002). While the number of dated landforms is increasing (cf. A. Hughes *et al.*, 2011), it is clear that further dating is still necessary (Ballantyne, 2010) as many sites remain undated and our understanding of retreat is often based on a limited number of samples using methods which can produce erroneous or uncertain results. It is clear that the limitations of existing methods must be addressed and new methods developed to better reconstruct past events and further constrain glacial response times.

The primary method for dating glacial retreat is Terrestrial Cosmogenic Nuclide Dating (TCND) which measures rock surface exhumation based on the spallation of upper atmospheric cosmic rays, which fragment target elements (Balco, 2011) and form radioactive isotopes (Briner, 2011). The application of TCND to boulders deposited on moraines (e.g. Everest *et al.*, 2006; Dortch *et al.*, 2010a; Wilson *et al.*, 2013b) and bedrock surfaces (e.g. Stone *et al.*, 1998; Dortch *et al.*, 2010a,b; Glasser *et al.*, 2012; Hughes *et al.*, 2016) can provide a direct age for glacial landforms. Despite the widespread application of this technique (Cockburn and Summerfield, 2004), there are numerous sources of error in calculating TCN exposure ages and many methodological uncertainties which are still to be addressed (Applegate *et al.*, 2012; Mackey and Lamb, 2013; Owen and Dortch 2014). In particular, nuclide production rate variation due to altitude and latitude necessitates the use of scaling schemes to calculate site specific production rates. Further research under the CRONUS-Earth Project (Phillips *et al.*, 2015) and the widespread application

of new exposure age calculators (Marrero *et al.*, 2015) may provide a consistent method for data generation although currently, the choice of appropriate scaling scheme remains uncertain.

The single largest source of uncertainty that leads to scattering of TCN ages is geologic uncertainty (Dortch *et al.*, 2013) which includes processes that affect sample stability (Bursik, 1991), moraine stabilisation (Ivy-Ochs *et al.*, 2008; Dortch *et al.*, 2009), transient or topographic shielding (Dunne *et al.*, 1999), inheritance (Putkonen and Swanson, 2003) and weathering rate variation (Colman, 1981; Sjöberg and Broadbent, 1991). Therefore, although the development and application of TCN dating has been effective in constraining the timing of BIS retreat, it is clear that researchers should be cautious in their interpretation of exposure ages and employ statistical analysis to identify outliers (Hallet & Putkonen, 1996; Briner *et al.*, 2005; Heyman *et al.*, 2013). The greatest barrier to isolating and correcting many of these uncertainties is the cost of TCN dating, which prevents high sample studies and statistically robust identification and rejection of erroneous results (Dortch *et al.*, 2013). Careful geomorphological analysis and the use of multiple isotopes can go some way to addressing these issues although this time-consuming approach is not always successful. TCN dating is a powerful geochronological tool but its associated uncertainties necessitate further empirical testing and refining of the method. High cost and limited facility resources demonstrate that the development of cost efficient methods is crucial to augment TCN dating studies in the pursuit of more detailed studies (for example, asynchronous ice sheet lobes).

Moving beyond relative dating, this study will establish Schmidt hammer exposure dating (SHED) as an effective method for numerical dating of glacial landforms in the UK by adopting a multi-site and multi-lithology calibration using TCN exposure ages. A lithology specific calibration curve is extended beyond 20 ka and applied to undated surfaces in a region well dated using other methods to demonstrate the validity of SHED. By combining a methodological and practical approach, this study not only provides the foundation for further application of SHED but can test and validate geomorphological interpretations and further our understanding of BIS retreat. Ultimately, we show that SHED can provide a direct age for glacial landforms and is of comparable accuracy and precision to TCN dating while providing immediate and cost efficient results.

2. Study area

TCN dated surfaces were identified and sampled based on 11 published articles, with 89 samples located in Scotland, the majority of these north of the Cairngorms including samples from Skye, Torridon and the Orkneys, Scotland. Two further sampling sites are located on Shap Fell, NW England and on the Aran Ridge, N Wales (Figure 1). Although climatic variation is evident between these regions, particularly for the drier sites in North East Scotland, the majority of sites are typically wet, often in excess of >3000mm/yr and certainly greater than 2000mm/yr (Met Office, 2015). Samples were obtained from a range of depositional and erosional environments

including moraines, roches moutonnées and summit and corrie blockfields, with contrasting topographic settings, between 21m a.s.l. in Slapin Bay, Skye (Small *et al.*, 2012) to 928m a.s.l. on Maol Chean Dearg, Torridon (Fabel *et al.*, 2012). Topographic variation may play a key role in localised precipitation (Ballantyne, 1983; Brundson *et al.*, 2001). The use of a statistically large dataset is necessary to analyse whether SHED can be applied in regions of similar climate or identify whether local climatic and topographic settings result in non-linear weathering over regional spatial scales. Sampling was focused on five lithologies; granite, sandstones, gneiss quartz and Ordovician felsic ash-flow tuffs. Adopting a multi-lithology and multi-landform approach is of particular value as this study is the first attempt to establish SHED in the UK.

3. Methods

3.1. Review of Schmidt Hammer methods

The ‘N’ type Schmidt hammer employed in this study was first applied to glacial environments in the 1960s (Goudie, 2006) and records a rebound value (R Value), a measure of how far a piston rebounds after being fired at a surface. Harder rocks will result in higher R values (Aydin and Basu, 2005). The use of the Schmidt Hammer as a relative dating tool was established by Matthews and Shakesby (1984) and is based on the principle that surfaces which have been uncovered for longer periods of time will be relatively softer than surfaces which have been uncovered recently. Relative dating has been largely limited to Holocene surfaces (e.g. Evans *et al.*, 1999; Shakesby *et al.*, 2011; Rode and Kellerer-Pirklbauer, 2012) with only a few attempts to extent the method beyond 10 ka. This restricted scope is of little value from the perspective of studying the BIS as retreat pre-dates the Holocene.

Lithology is an important precursor to Schmidt hammer testing as variance between weak and strong lithologies is highly significant (Selby, 1993). Given the importance of lithology in determining R values, relative dating can only be applied to single lithologies (Winkler, 2005) and has been limited to single site studies (*valley - mountain range scale*), as variation in climate may drive changes in physical and chemical weathering rates and result in non-linearity over larger spatial scales. The technique is relatively precise, able to differentiate between deposits at ± 300 yr (Matthews and Shakesby, 1984; McCarroll, 1987) and up to ± 100 yr intervals (Evans *et al.*, 1999).

However, the Schmidt Hammer is sensitive to surface discontinuities and rock moisture (Sumner & Nel, 2002) and surface moisture content (Viles *et al.*, 2011). Rock moisture is unlikely to have a significant impact within the confines of this study due to the ubiquitous >2000 mm/yr precipitation rates, as the 2-10 point R value variation identified by Sumner and Nel (2002) was between ‘oven dry’ and ‘fully saturated’ samples. The variation in the highly precipitous mountainous regions of the UK is unlikely to be as significant. However, surface moisture content may be of greater significance as it can alternate stochastically within short timescales and can result in an observable difference in R values (Viles *et al.*, 2011).

To avoid anomalous results, procedures as outlined by Viles *et al.*, (2011) must be followed. The Schmidt hammer should be used perpendicular to the surface (Aydin, 2015), avoiding rough, irregular (Williams and Robinson, 1983) and lichen covered surfaces (Matthews and Owen, 2008), which significantly reduce R values, while restricting sampling to horizontal surfaces and blocks of sufficient size, typically in excess of 25kg (Sumner and Nel, 2002; Demirdag *et al.*, 2009). However, there are contentious issues, particularly in regards to operator variance, deemed insignificant by Day and Goudie (1977). In contrast, Viles *et al.*, (2011) show that R value variation between researchers for the original ‘N’ type Schmidt Hammer can be significant but varies before ($p = 0.00$) but not after ($p = 0.10$) carborundum treatment, which removes surface irregularities. However, variation in this test was highly influenced by significantly higher readings by one operator (Viles *et al.*, 2011). In this study, all surfaces were sampled by one author (Tomkins) and therefore results are internally consistent. Clearly, further work on the impacts of operator variance needs to be undertaken.

Additionally, there is no consensus on how many R values are necessary on each surface (Table-1; Goktan and Gunes, 2005). Given the sensitivity of the tool to surface discontinuities, moisture content and surface texture, a sufficiently large dataset, typically greater than 25 R Values for weak and moderately strong rocks (Selby, 1980) is necessary to permit rapid identification of anomalous results (Shakesby *et al.*, 2006). We follow the methods of Niedzielski *et al.*, (2009) who provide statistical analysis for 14 key lithologies and show that the minimum sample size requirement varies significantly between lithologies. For this study, 30 R values were recorded for each sampled surface, irrespective of lithology, as this ensured that even coarse surfaces generated statistically significant results and eliminates the need for carborundum surface preparation while removal of the lowest values was not undertaken as this “produces different effects and is generally not recommended” (Niedzielski *et al.* 2009).

3.2. Schmidt hammer exposure dating (SHED)

SHED has been undertaken in New Zealand (Winkler, 2009; Stahl *et al.*, 2013), Portugal (Sánchez *et al.*, 2009) and Norway (Matthew and Owen, 2010; Matthews and Winkler, 2011) and requires independent dating controls to generate R value calibration curves. These previous studies have used a limited number ($n=9$) of TCN exposure ages as calibration points (Winkler, 2009) while Matthews and Owen (2010) based their calibration on two control points, one dated to the Little Ice Age at 100 ± 50 yr based on lichenometry (Matthews, 2005) and the other dated to 9700 yr based on ^{14}C (Dahl *et al.*, 2002; Matthews & Dresser, 2008). This study adopts and advocates a different approach by sampling many TCN dated surfaces from published research in order to develop a robust calibration curve (Table-2).

The majority of previous studies focus solely on the Holocene, although the study by Sánchez *et al.*, (2009) sampled surfaces beyond 100 ka. White and Brantley (2003) suggest that weathering

rates decline exponentially through time, thus glacial boulders from Holocene deglaciation would exhibit significantly lower R values compared to surfaces recently uncovered (< 100 yr). Although the R value difference between 10-20 or 90-100 ka surfaces is likely to be of lesser magnitude, Sánchez *et al.*, (2009) suggest that Schmidt Hammer methods may be applicable to surfaces beyond 100 ka. SHED can be of higher precision than TCN dating (Matthews and Owen, 2010) although it is not yet clear if this will be observed beyond the Holocene. Moreover, the precision of SHED will be, in part, limited by the precision of the technique used to develop the calibration curve.

3.3. Sampling Procedure

R values were collected for the following lithologies: granite ($n = 38$), sandstone ($n = 36$), gneiss ($n = 10$), quartz ($n = 9$) and Ordovician felsic ash-flow tuffs ($n = 5$). Sampling was limited to sites in which 2 or more surfaces had been dated as this ensured anomalous results could be more easily identified. Additionally, sampling was restricted to dates generated using Beryllium 10 (^{10}Be), the most commonly used isotope due to its straightforward decay chain.

A further 31 surfaces dated using Chlorine 36 (^{36}Cl) were sampled (Vincent *et al.*, 2010; Ballantyne *et al.*, 2009b; Wilson *et al.*, 2013b) but are not considered in this study as many of the surfaces, in particular those dated by Ballantyne *et al.*, (2009b) and Wilson *et al.*, (2013b), may be compromised by inheritance. Although the remaining ^{36}Cl surfaces (Vincent *et al.*, 2010) conform to our current understanding of regional deglaciation (Clark *et al.*, 2012), significant variation between scaling schemes for this isotope is possible, often in excess of 50% (Schimmelpfennig *et al.*, 2009) due to minor discrepancies in the calibration process (Swanson and Caffee, 2001, Licciardi *et al.*, 2008). Given the potential for these errors, ^{36}Cl samples are not included in this analysis but will be considered in future. As a result, three ^{10}Be surfaces from P. Hughes *et al.* (2012) and Wilson *et al.*, (2013b) are not included in the analysis.

Dated surfaces were located using maps from published papers, GPS data and sample photographs provided by researchers. However, some incorrect surfaces may have been sampled due to inaccurate GPS data recorded during the 1990s. The GPS Selective Availability (SA) feature, disabled in 2000, intentionally degraded civilian GPS accuracy by up to 100m. To compensate for this, the largest erratics which showed no sign of instability and were sufficiently removed from any nearby cliff faces were selected. Sampling surfaces in the correct location and from the correct lithology should be representative of the dated sample. In many cases, it was possible to identify direct evidence of sampling (Figure 2, particularly for Torridonian Sandstone surfaces where lichen growth was slowest (Ballantyne *et al.*, 2009a), but for other lithologies this was only possible for recent research (Kirkbride *et al.*, 2014).

3.4. R value and Age calibration

Having generated over 10,000 R values, the Schmidt Hammer may have lost condition (Goudie, 2006). Testing of a specific Sandstone boulder at the start and end of sampling indicated a four point difference (*June 2013*, 48.08 ± 0.82 - *December 2014*, 44 ± 1.05). The data were compiled chronologically and thus, it was possible to calibrate each individual R value to correct for temporal changes in spring tension. It was assumed that the tool deteriorated linearly through time which yields an average reduction per R value of 0.00083%.

The CRONUS Earth Calculator (Balco *et al.* 2008) was used to recalculate all ^{10}Be ages with reference to the Loch Lomond Local Production Rate (LL LPR) (Fabel, D., pers. comm. 2014) which is based on ^{10}Be concentration in samples from boulders on a Younger Dryas moraine and independently constrained by ^{14}C (Fabel *et al.*, 2012). This production rate (3.92 ± 0.18) minimises scaling uncertainties that are present in global scaling schemes (Balco, 2011). Although other local production rates are available, including the North West Highlands LPRs (NWH) of Ballantyne and Stone (2012), these have not been independently dated and assume retreat of Younger Dryas glaciers between 11.6-12.2 ka based on rapid warming observed in climate records (Brooks and Birks, 2000; Brooks *et al.*, 2012) at the onset of the Holocene. Ages referred to in this study are based on the St time-independent scheme (Lal, 1991; Stone, 2000) assuming 0 mm erosion. Recalibrated LL LPR ages are on average 14.3% older than ages based on global production rates.

3.5. Incomplete/incorrect data

Samples from Stone *et al.*, (1998), Everest and Kubik (2006) and Bradwell *et al.*, (2008b) did not incorporate or provide data for topographic shielding. Values were generated using a python code developed by Li (2013) based on an OS Terrain 50 DTM available from Edina Digimap and calculated using ArcGIS, with an azimuth interval of 10° and elevation interval of 5° . Additionally, some studies did not include reference to sample density or thickness. The latter was assumed to be 5cm (Darvill, 2013) as below this, cosmogenic nuclide concentrations rapidly decrease and nuclide production is quickly dominated by muonic reactions rather than nucleonic spallation (Gosse and Phillips, 2001). For sample density, lithology specific values from other publications were used although exposure ages are unlikely to change significantly due to these variables. Granitic rocks from Glen Eaniach (Everest and Kubik 2006) were assumed to be 2.7 g cm^{-3} based on the values in nearby Coire an Lochain (Kirkbride *et al.*, 2014) while Sandstone bedrock from An Teallach (Stone *et al.*, 1998) was assumed to be 2.6 g cm^{-3} based on the bedrock densities inferred by Ballantyne *et al.*, (2009a) near the Redpoint and Gairloch moraines. For Gneissic boulders on the Wester Ross peninsula, dated by Everest *et al.*, (2006), the nearest analogue is those found on the Orkneys by Phillips *et al.*, (2008), and the density of these samples, at 2.4 g cm^{-3} is applied here.

The CRONUS Earth recalibrated age for SHAP-03 ($20.05 \pm 1.15 \text{ ka}$) may be inaccurate. LL LPR ages are typically 6.8-7% older than exposure ages derived from the North West Highlands 11.6 ka LPR (Ballantyne *et al.*, 2014), which SHAP-03 is significantly older (13.4%). Additionally,

NWH 12.2 LPR and LL LPR ages are almost identical (Ballantyne, C., pers. comm. 2014). This is observed for the other SHAP samples (Maximum difference of 0.4%). The 8.2% increase for SHAP-03 is therefore a major anomaly. This error is unlikely to be associated with CRONUS Earth as it would be observed for all samples. Therefore, it is suggested that the ^{10}Be concentration may have been reported incorrectly. To address this, SHAP-03 has been calibrated by dividing the reported age by the NWH 11.9 production rate (4.09) and multiplying by the new production rate for the LL LPR (3.92) to generate an age of 18.78 ± 0.9 ka (Ballantyne, C., pers. comm. 2014). The increase from NWH 11.6 to LL LPR (6.1%) is in better agreement with Ballantyne *et al.*, (2014) although does differ more from the NWH 12.2 age (1.2%) than the other SHAP samples. Unfortunately, it is not possible to generate accurate uncertainties using this method, so the original uncertainty as reported by Wilson *et al.*, (2013c) is used. As such, rigorous calibration using the CRONUS Earth calculator is the recommended method for ^{10}Be age calibration.

3.6. Rejection of young/old ages

Of the 98 dated surfaces sampled in this study, 22 samples were not accepted as the results may have been compromised by inheritance ($n = 6$), overturning ($n = 3$), transient shielding or site disturbance ($n = 4$) or when Schmidt hammer testing was incorrectly conducted on unsuitable glacially polished or Quartz rich surfaces ($n = 8$). Additionally, sample WH1 from Phillips *et al.*, (2008) (19.67 ± 1.19 ka, R value: 27.64 ± 0.48) was not included in the sandstone lithology regression. This sample was anomalously soft compared to all other sandstones in excess of three standard deviations. The full dataset is presented in Table 1.

4. Results and discussion

4.1. R Value-Exposure Age Relationships

For many lithologies, the expected age-hardness relationship is not observed (Figure 3, Table 3). Sandstone, quartz, gneiss and Ordovician felsic ash-flow tuff do not have significant correlations, which indicates that ‘numerical’ dating using the Schmidt Hammer would be of little value. However, a statistically significant relationship is evident for granite surfaces ($R^2 = 0.81$, $p = <0.01$) and derived from a statistically acceptable dataset ($n = 25$). In order to test the validity of this calibration, SHED was applied to undated granite surfaces on Shap Fell, NW England with the results compared with other data sources (TCND, OSL) and discussed in the context of our understanding of regional deglaciation. However, before these results are presented, it is necessary to acknowledge some limitations of this study and discuss the suitability of each lithology to SHED.

4.2. Limitations and Sources of Error

4.2.1. Age Range

As this study relies on previously dated surfaces, the application of SHED to Holocene and pre-LGM deposits is limited. Given the extent of the BIS at its maximum (Bowen *et al.*, 1986, Serjup *et al.*, 2005), pre-LGM deposits are found only in the south-east and tend to be poorly preserved in the landscape (Rose, 2009). Additionally, much of the evidence for pre-LGM glacial stages is inferred from tills (Hamblin *et al.*, 2000), outwash deposits (Rose *et al.*, 2001) and offshore records (Holmes, 1997) which are not often dated using TCND, although studies by Balco and Rovey (2008) and Dehnert and Schlüchter (2008) demonstrate the value of applying this method to sedimentary deposits. However, there have been few attempts to constrain the southern limit of the BIS using this method (Ballantyne, 2010). While there are exposure ages that pre-date the LGM, they are either highly inaccessible (Everest *et al.*, 2013) or have such significant levels of uncertainty (Bowen *et al.*, 2002) that is not clear if they are truly representative of deglaciation. In this study, only seven samples are dated to older than 20 ka.

Additionally, no surfaces were sampled which post-date the Loch Lomond Stadial and only five samples were within this interval. This is despite significant evidence for LLS landforms and deposits (Sissons, 1979) although the number of TCN exposure ages related to this glacial readvance is remarkably low. LLS deposits have been dated in Cwm Idwal, North Wales (Phillips *et al.*, 1994), Keskadale, Lake District, (P. Hughes *et al.*, 2012) and in Scotland on Hoy (Ballantyne, 2007a), in the Grampians (Golledge *et al.*, 2007) and Galloway Hills (Ballantyne *et al.*, 2013) and in Coire an Lochain (Kirkbride *et al.*, 2014). However, the vast majority of published TCN exposure ages pre-date the LLS. Of the ~160 exposure ages reviewed in Ballantyne (2010), only 9 were directly attributed to the LLS, with 6 further samples from Lough Acorrymore, Ireland (Bowen *et al.*, 2002; Ballantyne *et al.*, 2008b) exhibiting a complex deglaciation history.

Recent publications have attempted to date Holocene glacial advances. In particular, the study by Kirkbride *et al.*, (2014) and its companion piece by Harrison *et al.*, (2014) infer ‘Little Ice Age’ (Mann *et al.*, 2009) glaciation in the Cairngorms. Further Holocene exposure ages have been reported for rock slope failures (RSF) throughout Scotland (Ballantyne *et al.*, 2014) although less than 1/3 of these are associated with debuitressing during deglaciation. These are the only studies which report accurate Holocene exposure ages in the UK. This contrasts markedly with other SHED studies in Norway (Matthews & Owen, 2010) and New Zealand (Winkler, 2009) which focus solely on Holocene deposition, and compare very recent deposits (< 0.1 ka) with early Holocene landforms. While the age range of this study and the previously mentioned publications is similar (~10-15 ka), the timeframe for that age range is key, given the exponential decline in weathering rates through time (White and Brantley, 2003). This is supported by evidence from Sanchez *et al.*, (2009) and Stahl *et al.*, (2013) as the magnitude of R value reductions decreases through time. While SHED is perhaps most effective at dating recent glacial deposits (< 10 ka), this study has shown that it can be extended beyond 20 ka for granite surfaces. It is clear that further testing of LLS equivalent and Holocene surfaces in northern

Europe is a priority for further research, but it is acknowledged that extending this method beyond 25 ka in the UK is unlikely given the extent of the BIS at its maximum. The testing and application of SHED in reasonably well-dated regions (>1800 TCN ages) with multiple glacial cycles (<400 ka) preserved in the landscape (e.g. the Himalaya) is an important step in establishing SHED as a key geochronological tool (Dortch *et al.*, 2013; Owen and Dortch, 2014).

4.2.2. Cosmogenic Uncertainty

This study also treats TCN exposure ages as ‘*absolute ages*’ but this infers a level of accuracy and precision which is often unwarranted. While samples affected by inheritance, site disturbance, reworking or transient shielding have been omitted, significant errors (1-2 ka) are possible. Thus, some exposure ages used in this study may be challenged by future research. For all dated samples used in this study TCN exposure ages have a mean uncertainty of ± 1.3 ka. The uncertainty associated with the clustering of ages is typically greater than individual age uncertainties. Significant uncertainty encourages researchers to informally match glacial deposits to high magnitude climatic drivers (Blaauw *et al.*, 2007) but this level of uncertainty is unacceptable for short term glacial events such as the WRR or LLS as the range of uncertainty exceeds the range of the climatic event. While the establishment of LPRs has generated more precise exposure ages, it is not clear if further improvements are possible. While TCND is an effective method for dating glacial landforms, it must be acknowledged that its moderate precision adds an element of uncertainty which is currently difficult to address.

4.3. Lithological Suitability

4.3.1. Granite

SHED of granitic surfaces is particularly promising as a clear R value-exposure relationship is evident (Figure 3A), indicating that this lithology weathers linearly over significant spatial scales and for regions of similar climate. Samples are relatively young (< 25 ka) and were free from deformities (e.g. cracks). Granite samples were also collected from a range of altitudes (21-928m a.s.l.). Furthermore, the impact of altitude is not evident in our data. For all granite surfaces, there is no R value-altitude relationship ($R^2 = 0.01$, $p = 0.71$). This contrasts with research by Ballantyne *et al.*, (1998) who found appreciable variation in R values above and below weathering limits for Torridon sandstone although this perhaps reflects the preservation of the landscape beneath cold-based ice and not R-value variation as a result of enhanced weathering with altitude. However, when 9 high altitude samples (895-928m) from Coire an Lochan (Kirkbride *et al.*, 2014) are removed, a clear relationship is evident ($R^2 = 0.82$, $p = <0.01$), implying a causal link between R values and altitude. However, a stronger relationship is observed between sample altitude and deglaciation age ($R^2 = 0.89$, $p = <0.01$). As such, this indicates that the potential link between altitude and R values is simply a function of deglaciation age, as low lying samples underwent deglaciation last, resulting in higher R values than mid-altitude samples, which underwent deglaciation earlier and are proportionally harder. Additionally, although granites may vary in ‘base’ hardness (Goudie, 2006), e.g. between

Cairngorm (Brook *et al.*, 2004) and Shap granite (Day and Goudie, 1977), these results suggest that granites are relatively homogenous in texture and mineral assemblage, which leads to time-dependent weathering rates which are consistent between granite source areas.

However, not all granites are suitable to SHED. In particular, the quartz-rich pegmatite erratics in Glen Eaniach (Everest and Kubik, 2006) are not internally consistent ($R^2 = 0.19$, $p = 0.39$) and fit poorly with the established calibration. These surfaces may weather particularly slowly or even non-linearly given the inappropriateness of quartz to SHED.

4.3.2 Sandstone

Sampling of sandstone surfaces has shown that there is no relationship between R values and exposure ages (Figure 3B). Weathering of Sandstone is evident (Turkington & Paradise, 2005) with time-dependent variation in edge-roundness (Kirkbride & Bell, 2010) but this clear relationship does not appear to have any appreciable effect on R values, despite a smooth surface texture and limited lichen colonisation. As there is evidently no link between R values and exposure ages, Sandstone is clearly not suitable for SHED over large spatial scales. Non-linear weathering of Sandstone surfaces, meso-micro scale variation in weathering processes, or case-hardening due to carbonate precipitation must account for some of the R value variability, as it not explained by any factor considered in this study.

4.3.3. Quartz

Surface texture and mineralogical content are key factors in determining the rate and magnitude of rock surface disintegration. In particular, quartz exhibits limited R value variation (Owen *et al.*, 2007) as it is thermodynamically stable and therefore is particularly resistant to chemical weathering (Figure 3C; Pope, 1995). Additionally, its strong crystalline silicate structure (SiO_2) ensures that physical weathering occurs at a much slower rate than other minerals, as it is less sensitive to granular disintegration. While it is affected by freeze-thaw cycles (Schwamborn *et al.*, 2012) and does undergo weathering, as the differential lowering of Quartz veins indicates (Owen *et al.*, 2007), it does so slowly that the high magnitude ‘noise’ generated by the Schmidt Hammer, due to differential surface texture, moisture content, discontinuities or lichen cover, obscures the low magnitude signal from weathering (Figure 3Cc). Moreover, quartzite and quartz veins break down along fractures that can be several centimetres in thickness, which would lead to a stochastic signal in TCND ages and make R Value calibration difficult.

4.3.4. Gneiss

The unsuitability of gneiss to SHED (Figure 3D) can be attributed to surface texture. Gneiss is predominantly coarse grained (Feuten *et al.*, 1991) and as the Schmidt Hammer is most effective when applied to flat, planar surfaces (Williams & Robinson, 1983), it is inferred that the significant R value variation evident in this study ($\text{range} = 15.1$) is representative of the highly variable surface texture and not a function of weathering through time. Gneiss is also highly

variable in its mineralogical content (Marshak, 2009) and therefore, this lithology is particularly unsuitable, as there is considerable variability both between gneissic surfaces, due to mineralogical content, and on individual surfaces, due to its coarse grained texture.

4.3.5. Ordovician felsic ash-flow tuffs

Unfortunately, despite relatively consistent R values and TCN ages, the data are not internally consistent, indicating that this lithology is unsuitable for SHED numerical dating (Figure 3E). A simple explanation is that the ^{10}Be ages were derived from quartz veins (see Glasser *et al.*, 2012) while R values were generated from the surrounding bedrock although theoretically, this should have little impact, as the bedrock and quartz veins should have undergone deglaciation concurrently. Another possibility is wide variability in the physical properties of the felsic ash-flow tuffs, which also occur in association with intrusive and extrusive rhyolites (Glasser *et al.*, 2012). The sample size for felsic ash-flow tuffs was relatively small ($n = 5$) and recent TCN ages from this and related lithologies has recently become available (Hughes *et al.*, 2016). As new TCN data becomes available, the efficacy of SHED numerical dating on different lithologies can be explored further in the coming years.

5. Implementing SHED

5.1. Case Study

As a first step to apply the SHED method presented in this study, 31 undated Shap granite erratics were sampled from Shap Fell, NW England (Figure 4, Table-4). This site is useful for a number of reasons. Firstly, there are numerous granite erratics derived from the nearby Shap granite pluton (Wilson *et al.*, 2013c). Secondly, it is in a region of similar climate but a significant distance from the majority of other tested granite samples in Scotland, thus providing an important testbed for SHED. Thirdly, the region is reasonably well dated with TCN dated samples on Shap Fell (Wilson *et al.*, 2013c) providing direct dates for glacial retreat. OSL dating at New Close and Warton Crag (Telfer *et al.*, 2009) can be used to indicate the commencement of ice free conditions further south. Dating using ^{14}C from Wray Bay (Coope and Pennington, 1977), Ravenglass (Lloyd *et al.*, 2013), Hallsenna Moor (Walker, 2004) and St Bees (Coope and Joachim, 1980) and ^{10}Be TCN dating from Wasdale (McCarroll *et al.*, 2010) are also useful and provide a timeframe for Lake District deglaciation to the west. Finally, and perhaps most importantly, accurate dating of Shap Fell deglaciation is key to the BIS as it retreated northwards from Shap Fell through the Vale of Eden (Clark *et al.*, 2012). This in turn determines the timing of retreat into the Solway Lowlands (Livingstone *et al.*, 2012) and provides context for the development of an independent ice centre in the Galloway Hills (Ballantyne *et al.*, 2013).

5.2. Study site

Shap Fell exhibits numerous north-south oriented drumlins (Clark *et al.*, 2004), formed due to deformation of the basal till layer (Delaney, 2003) and representative of rapid ice flow (Alley *et*

al., 1986), as well as abundant meltwater channels (Arthurton & Wadge, 1981). Although glacial deposits indicate that ice flow alternated during periods of internal ice sheet reorganisation (Livingstone *et al.*, 2012), the existence of oriented drumlin fields suggests the presence of an ice stream retreating northwards (Livingstone *et al.*, 2008).

A total of 32 samples were obtained from previously unmapped hummocks and drumlins, close to the erratic limit as identified by Clark *et al.*, (2004). These erratics are ~10km from the bedrock source and are sufficiently large (1-3m in length) to render post-glacial fluvial transport unlikely. The restricted distribution of these erratics could indicate post-depositional disturbance. However, any disturbance would be unlikely to affect rock hardness, unless significant rock breakup occurred. After excluding one outlier (CR-1), the R value consistency of these samples (37.6 ± 1.1 , average and mean absolute deviation) indicates that disturbance is not an important factor.

These erratics could have been deposited due to the breakup of an ice dammed lake. However, Clark *et al.*, (2004) report no major ice dammed lakes in the Vale of Eden, in contrast to initial research (Jowett & Charlesworth, 1929) although early assessments are not verified by glacial deposits (Evans *et al.*, 2005). The only major glacial lake in the area was proglacial, and developed to the north before Scottish ice readvance at 16.8 ka (Livingstone *et al.*, 2010a). The minimum extent of this lake and ice front positions before 16.8 ka are shown in Figure 4. While this was sufficient to flood the Solway Lowlands ($< 1350\text{km}^2$), a floodwater origin for the CR erratics is unlikely, as they are located over ~50km upslope. Flooding was more likely to the north and west as glacial retreat of the Scottish ice front could trigger rapid meltwater drainage. Therefore, we conclude these erratics are representative of glacial retreat.

5.3. Deglaciation

Based on SHED, Shap Fell underwent deglaciation at 16.5 ± 0.5 ka (Figure 5A; Table 4; uncertainty = mean absolute deviation). Modelling of BIS retreat indicates that Shap Fell underwent deglaciation between 17-18 ka (Clark *et al.*, 2012), after 16 ka (Livingstone *et al.*, 2012), at 16 ka (Hubbard *et al.*, 2009) and after 15.5 ka (Evans *et al.*, 2009). The result generated by SHED is in direct agreement with the youngest TCN date from Wilson *et al.*, (2013c) of 16.5 ± 1.0 ka. If the OSL dates from Telfer *et al.*, (2009) are accepted, this result suggests that deglaciation to the south of the coastal and upland regions occurred by 19.3 ± 2.6 and 16.5 ± 1.7 ka respectively although the level of uncertainty for these samples makes direct comparison with SHED ages difficult. The ^{14}C age of 17.8 ± 0.9 ka from Wray Bay (Coope and Pennington, 1977) indicates that lowland deglaciation in the Lake District predated BIS retreat from Shap Fell, although ^{14}C dates from coastal sites (Coope and Joachim, 1980; Walker, 2004; Lloyd *et al.*, 2013) of 14.7-16.5 ka indicate that final deglaciation of the western Lake District may have post-dated Shap Fell retreat due to the readvance of marine ice during the Killard Point Stadial (McCabe *et al.*, 2007b). The close correlation of two datasets (SHED, Wilson *et al.*, 2013c)

which are in agreement with the timeframe for deglaciation provided by modelling, is sufficiently strong to narrow the possible age range of deglaciation at Shap Fell to 16.5 ± 0.5 ka (Figure 5B).

For granitic lithologies, individual calibrated SHED ages and uncertainties (mean uncertainty = 1.5 ka) are comparable to the 1.1 ka mean uncertainty from TCND (Wilson *et al.*, 2013c). Moreover, our SHED ages in aggregate provide the strongest estimate for deglaciation simply due to the number of samples ($n = 31$), which highlights the value of low-cost high-sample volume dating.

Despite the R value consistency of these erratics, sampling of a sufficiently large number of surfaces (>20) is still necessary, as anomalous results (e.g. CR1) can occur. SHED of Shap granite is of particular value as this lithology is an important marker erratic (Evans *et al.*, 2005) given its restricted source area, but widespread local and regional distribution. It is commonly found in the Vale of York (Letzer, 1978) and indicates that ice from Scotland and the Lake District breached the Pennines through the Stainmore Gap (Catt, 2007). Therefore, SHED could be applied to these erratics to constrain the timing of Yorkshire deglaciation and indicate the extent to which North Sea ice shelves extended inland (Davies *et al.*, 2009; Clark *et al.*, 2012) and delayed terrestrial deglaciation, or if they were confined to the periphery of the landmass (Everest *et al.*, 2005).

6. Conclusions

This study has established SHED as a viable and cost efficient technique for dating of glacial landforms in the UK. SHED works effectively for granite surfaces and may be applicable beyond 20 ka although further sampling of LLS and Holocene surfaces is recommended. Fieldwork is time consuming, as many samples are required to generate statistically significant results. However, the cost of TCND and the potential for significant errors means that the choice of sampling site is critical (Ballantyne, 2010). This is less important for SHED as multi-sample studies permit straightforward identification of anomalous results, while data can be quickly compiled and analysed. Importantly, it is capable of providing direct ages for landforms and is applicable to many glacial environments, and thus has clear advantages over other techniques which provide indirect or minimum ages (^{14}C , OSL). This study should provide the foundations for and encourage future applications of SHED.

However, accurate SHED is completely dependent on the correct calculation of calibration curves as minor errors can significantly alter the calculated exposure age. Previous studies have used a limited number of calibration points but given the potential for geological uncertainty, relying on sparse and isolated control points and assuming a linear relationship between them is not recommended and sampling of multiple dated surfaces is necessary. In this study, we show that this strategy is effective, as a robust granite calibration curve has been applied in a region of

similar climate and generates results that conform to our understanding of regional deglaciation. The retreat of the BIS from Shap Fell at 16.5 ± 0.5 ka has now been dated using both TCND and SHED, and fits well with a regional pattern of deglaciation established using OSL and ^{14}C dating. This study has shown that SHED can provide independent verification of existing results. However, of greater value will be its application to poorly dated regions where it may provide a cost efficient and reliable first estimate of deglaciation.

Acknowledgements

We would like to thank C. Ballantyne, T. Bradwell, J. Everest, D. Small and P. Wilson for providing sample photographs which were essential when conducting fieldwork. D. Fabel also kindly provided unpublished calibration data for the LL LPR which permitted exposure age recalibration. We would also like to thank D. Tomkins whose assistance during fieldwork was essential to this research. This study was partially funded by the University of Manchester Zochonis Travel Award 2013/2014 and an Undergraduate Dissertation Bursary provided by the Manchester Geographical Society. Hughes and Dortch would like to thank the University of Manchester Research Stimulation Fund.

References

- Alley, R.B., Blankenship, D.D., Bentley, C.R., Rooney, S.T., 1986. Deformation of till beneath ice stream B, West Antarctica. *Nature* 322, 57–59. doi:10.1038/322057a0
- Applegate, P.J., Urban, N.M., Keller, K., Lowell, T. V., Laabs, B.J.C., Kelly, M. A., Alley, R.B., 2012. Improved moraine age interpretations through explicit matching of geomorphic process models to cosmogenic nuclide measurements from single landforms. *Quat. Res.* 77, 293–304. doi:10.1016/j.yqres.2011.12.002
- Arthurton, R.S., Wadge, J.A., 1981. Geology of the country around Penrith. *Mem. Geol. Surv. Gt. Britian*.
- Aydin, A., 2015. The ISRM Suggested Methods for Rock Characterization, Testing and Monitoring: 2007–2014 2007–2014. doi:10.1007/978-3-319-07713-0
- Aydin, A., Basu, A., 2005. The Schmidt hammer in rock material characterization. *Eng. Geol.* 81, 1–14. doi:10.1016/j.enggeo.2005.06.006
- Balco, G., 2011. Contributions and unrealized potential contributions of cosmogenic-nuclide exposure dating to glacier chronology, 1990–2010. *Quat. Sci. Rev.* 30, 3–27. doi:10.1016/j.quascirev.2010.11.003
- Balco, G., Rovey, C.W., 2008. An isochron method for cosmogenic-nuclide dating of buried soils and sediments. *Am. J. Sci.* 308, 1083–1114. doi:10.2475/10.2008.02
- Balco, G., Stone, J.O., Lifton, N. A., Dunai, T.J., 2008. A complete and easily accessible means of calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements. *Quat. Geochronol.* 3, 174–195. doi:10.1016/j.quageo.2007.12.001

Ballantyne, C.K., 2010. Extent and deglacial chronology of the last British-Irish Ice Sheet: Implications of exposure dating using cosmogenic isotopes. *J. Quat. Sci.* 25, 515–534. doi:10.1002/jqs.1310

Ballantyne, C.K., 2007. The Loch Lomond Readvance on north Arran, Scotland: Glacier reconstruction and palaeoclimatic implications. *J. Quat. Sci.* 22, 343–359. doi:10.1002/jqs.1059

Ballantyne, C.K., 2002. The Loch Lomond Stadial on the Isle of Mull, Scotland glacial reconstruction and palaeoclimatic implications. *J. Quat. Sci.* 17, 759–771. doi:10.1002/jqs.729

Ballantyne, C.K., 1989. The Loch Lomond Readvance on the Isle of Skye, Scotland: glacier reconstruction and palaeoclimatic implications. *J. Quat. Sci.* 4, 95–108.

Ballantyne, C.K., 1983. Precipitation gradients in Wester Ross, North-West Scotland,. *Weather* 38, 95–108.

Ballantyne, C.K., Black, N.M., Finlay, D.P., 1990. Use of the Schmidt test hammer to detect enhanced boulder weathering under late-lying snowpatches. *Earth Surf. Process. Landforms* 15, 471–474.

Ballantyne, C.K., Hall, A.M., 2008. The altitude of the last ice sheet in Caithness and east Sutherland, northern Scotland. *Scottish J. Geol.* 44, 169–181. doi:10.1144/sjg44020169

Ballantyne, C.K., Hall, A.M., Phillips, W., Binnie, S., Kubik, P.W., 2007. Age and significance of former low-altitude corrie glaciers on Hoy, Orkney Islands. *Scottish J. Geol.* 43, 107–114. doi:10.1144/sjg43020107

Ballantyne, C.K., McCarroll, D., Stone, J.O., 2006. Vertical dimensions and age of the Wicklow Mountains ice dome, Eastern Ireland, and implications for the extent of the last Irish Ice Sheet. *Quat. Sci. Rev.* 25, 2048–2058. doi:10.1016/j.quascirev.2006.01.026

Ballantyne, C.K., Rinterknecht, V., Gheorghiu, D.M., 2013. Deglaciation chronology of the Galloway Hills Ice Centre, southwest Scotland. *J. Quat. Sci.* 28, 412–420. doi:10.1002/jqs.2635

Ballantyne, C.K., Sandeman, G.F., Stone, J.O., Wilson, P., 2014. Rock-slope failure following Late Pleistocene deglaciation on tectonically stable mountainous terrain. *Quat. Sci. Rev.* 86, 144–157. doi:10.1016/j.quascirev.2013.12.021

Ballantyne, C.K., Schnabel, C., Xu, S., 2009a. Readvance of the last British-Irish Ice Sheet during Greenland Interstade 1 (GI-1): the Wester Ross Readvance, NW Scotland. *Quat. Sci. Rev.* 28, 783–789. doi:10.1016/j.quascirev.2009.01.011

Ballantyne, C.K., Stone, J.O., 2012. Did large ice caps persist on low ground in north-west Scotland during the Lateglacial Interstade? *J. Quat. Sci.* 27, 297–306.

Ballantyne, C.K., Stone, J.O., Fifield, L.K., 2009b. Glaciation and deglaciation of the SW Lake District, England: implications of cosmogenic ³⁶Cl exposure dating. *Proc. Geol. Assoc.* 120, 139–144. doi:10.1016/j.pgeola.2009.08.003

Ballantyne, C.K., Stone, J.O., McCarroll, D., 2008. Dimensions and chronology of the last ice sheet in Western Ireland. *Quat. Sci. Rev.* 27, 185–200. doi:10.1016/j.quascirev.2007.10.019

Blaauw, M., Bakker, R., Christen, J.A., Hall, V. A, Plicht, J. Van Der, 2007. A Bayesian framework for gae modelin of radiocarbon-dates peat deposits: case studies from the Netherlands. *Radiocarbon* 49, 357–367.

Bowen, D., 1986. Correlation of Quaternary glaciations in England, Ireland, Scotland and Wales. *Quat. Sci. Rev.* 5, 299–340. doi:10.1016/S0277-3791(86)80024-5

Bowen, D.Q., Phillips, F.M., McCabe, A.M., Knutz, P.C., Sykes, G. A., 2002. New data for the Last Glacial Maximum in Great Britain and Ireland. *Quat. Sci. Rev.* 21, 89–101. doi:10.1016/S0277-3791(01)00102-0

Bradwell, T., Stoker, M.S., Golledge, N.R., Wilson, C.K., Merritt, J.W., Long, D., Everest, J.D., Hestvik, O.B., Stevenson, A.G., Hubbard, A.L., Finlayson, A.G., Mathers, H.E., 2008. The northern sector of the last British Ice Sheet: Maximum extent and demise. *Earth-Science Rev.* 88, 207–226. doi:10.1016/j.earscirev.2008.01.008

Briner, J.P., 2011. Dating glacial landforms, in: Singh, V.P., Singh, P., Haritashya, U.K. (Eds.), *Encyclopedia of Snow, Ice and Glaciers*. Springer Science, pp. 175–186. doi:10.1007/978-90-481-2642-2

Briner, J.P., Kaufman, D.S., Manley, W.F., Finkel, R.C., Caffee, M.W., 2005. Cosmogenic exposure dating of late Pleistocene moraine stabilization in Alaska. *Bull. Geol. Soc. Am.* 117, 1108–1120. doi:10.1130/B25649.1

Brook, M.S., Kirkbride, M.P., Brock, B.W., 2004. Rock strength and development of glacial valley morphology in the Scottish Highlands and northwest Iceland. *Geogr. Ann. Ser. A Phys. Geogr.* 86, 225–234. doi:10.1111/j.0435-3676.2004.00227.x

Brooks, S., Axford, Y., Heiri, O., Langdon, P., Larocque-Tobler, I., 2012. Chironomids can be reliable proxies for Holocene temperatures. A comment on Velle et al., 2010. doi:10.1177/0959683612449757

Brooks, S., Birks, H., 2000. Chironomid-inferred late-glacial air temperatures at Whitrig Bog, southeast Scotland 15, 759–764. doi:10.1002/1099-1417(200012)15:8<759::AID-JQS590>3.0.CO;2-V

Brunsdon, C., McClatchey, J., Unwin, D.J., 2001. Spatial variations in the average rainfall-altitude relationship in Great Britain: An approach using geographically weighted regression. *Int. J. Climatol.* 21, 455–466. doi:10.1002/joc.614

Bursik, M., 1991. Relative dating of moraines based on landform degradation, Lee Vining Canyon, California. *Quat. Res.* 35, 451–455. doi:10.1016/0033-5894(91)90057-C

Catt, J., 2007. The Pleistocene glaciations of eastern Yorkshire: a review 56, 177–207. doi:10.1144/pygs.56.3.177

- Chiverrell, R.C., Thomas, G.S.P., 2010. Extent and timing of the Last Glacial Maximum (LGM) in Britain and Ireland: A review. *J. Quat. Sci.* 25, 535–549. doi:10.1002/jqs.1404
- Clark, C.D., Evans, D.J., Khatwa, A., Bradwell, T., Jordan, C.J., Marsh, S., Mitchell, W.A., Bateman, M.D., 2004. Map and GIS database of glacial landforms and features related to the last British Ice Sheet. *Boreas* 33, 359–375.
- Clark, C.D., Hughes, A.L.C., Greenwood, S.L., Jordan, C., Sejrup, H.P., 2012. Pattern and timing of retreat of the last British-Irish Ice Sheet. *Quat. Sci. Rev.* 44, 112–146. doi:10.1016/j.quascirev.2010.07.019
- Cockburn, H. A. P., Summerfield, M. A., 2004. Geomorphological applications of cosmogenic isotope analysis. *Prog. Phys. Geogr.* 28, 1–42. doi:10.1191/0309133304pp395oa
- Colman, S.M., 1981. Rock-weathering rates as functions of time. *Quat. Res.* 15, 250–264. doi:10.1016/0033-5894(81)90029-6
- Coope, G.R., Joachim, M.J., 1980. Lateglacial environmental changes interpreted from fossil coleoptera from St. Bees, Cumbria, NW England, in: Lowe, J.J., Gray, J., Robinson, J. (Eds.), *Studies in the Lateglacial of North-West Europe*. Pergamon Journals Ltd., Oxford, pp. 55–68.
- Coope, G.R., Pennington, W., 1977. The Windermere Interstadial of the Late Devensian. *Philos. Trans. R. Soc. London B* 280, 337–340.
- Croll, J., 1875. *Climate and Time in Their Geological Relations. A theory of secular changes of the earth's climate*. New York.
- Dahl, S.O., Nesje, A., Lie, Ø., Fjorheim, K., Matthews, J. a., 2002. Timing, equilibrium-line altitudes and climatic implications of two early-Holocene glacier readvances during the Erdalen Event at Jostedalsgreen, western Norway. *The Holocene* 12, 17–25. doi:10.1191/0959683602hl516rp
- Darvill, C.M., 2013. Cosmogenic nuclide analysis. *Geomorphological Tech.* 10, 1–25.
- Davies, B.J., Bridgland, D.R., Roberts, D.H., Cofaigh, C.Ó., Pawley, S.M., Candy, I., Demarchi, B., Penkman, K.E.H., Austin, W.E.N., 2009. The age and stratigraphic context of the Easington Raised Beach, County Durham, UK. *Proc. Geol. Assoc.* 120, 183–198. doi:10.1016/j.pgeola.2009.04.001
- Day, M.J., Goudie, A.S., 1977. Field assessment of rock hardness using the Schmidt test hammer. *BGR Tech. Bull.* 18, 19–29.
- Deere, D.U., Miller, R.P., 1966. Engineering classifications and index properties of intact rock, Technical Report AFWL-TR 65-115.
- Dehnert, A., Schlüchter, C., 2008. Sediment burial dating using terrestrial cosmogenic nuclides. *Eiszeitalter und Gegenwart - Quat. Sci. J.* 57, 210–225. doi:http://dx.doi.org/10.3285/eg.57.1-2.8
- Delaney, C., 2003. The Last Glacial Stage (the Devensian) in Northwest England 3.

Demirdag, S., Yavuz, H., Altindag, R., 2009. The effect of sample size on Schmidt rebound hardness value of rocks. *Int. J. Rock Mech. Min. Sci.* 46, 725–730. doi:10.1016/j.ijrmms.2008.09.004

Dortch, J.M., Owen, L. A., Caffee, M.W., 2013. Timing and climatic drivers for glaciation across semi-arid western Himalayan–Tibetan orogen. *Quat. Sci. Rev.* 78, 188–208. doi:10.1016/j.quascirev.2013.07.025

Dortch, J.M., Owen, L. A., Caffee, M.W., Brease, P., 2009a. Late Quaternary glaciation and equilibrium line altitude variations of the McKinley River region, central Alaska Range. *Boreas* 39, 233–246. doi:10.1111/j.1502-3885.2009.00121.x

Dortch, J.M., Owen, L. A., Haneberg, W.C., Caffee, M.W., Dietsch, C., Kamp, U., 2009b. Nature and timing of large landslides in the Himalaya and Transhimalaya of northern India. *Quat. Sci. Rev.* 28, 1037–1054. doi:10.1016/j.quascirev.2008.05.002

Dortch, J.M., Owen, L.A., Caffee, M.W., 2010a. Quaternary glaciation in the Nubra and Shyok valley confluence, northernmost Ladakh, India. *Quat. Res.* 74, 132–144.

Dortch, J.M., Owen, L.A., Caffee, M.W., Li, D., Lowell, T. V., 2010b. Beryllium-10 surface exposure dating of glacial successions in the Central Alaska Range. *J. Quat. Sci.* 25, 1259–1269.

Dunne, J., Elmore, D., Muzikar, P., 1999. Scaling factors for the rates of production of cosmogenic nuclides for geometric shielding and attenuation at depth on sloped surfaces. *Geomorphology* 27, 3–11. doi:10.1016/S0169-555X(98)00086-5

Engel, Z., 2007. Measurement and age assignment of intact rock strength in the Krkonoše Mountains, Czech Republic. *Zeitschrift für Geomorphol.* 51, 69–80.

Engineering geology field manual: field index tests, 1998. . U.S. Bureau of Reclamation.

Ericson, K., 2004. Geomorphological surfaces of different age and origin in granite landscapes: An evaluation of the Schmidt hammer test. *Earth Surf. Process. Landforms* 29, 495–509. doi:10.1002/esp.1048

Evans, D.J. A, Livingstone, S.J., Vieli, A., Ó Cofaigh, C., 2009. The palaeoglaciology of the central sector of the British and Irish Ice Sheet: reconciling glacial geomorphology and preliminary ice sheet modelling. *Quat. Sci. Rev.* 28, 739–757. doi:10.1016/j.quascirev.2008.05.011

Evans, D.J. A., Archer, S., Wilson, D.J.H., 1999. A comparison of the lichenometric and Schmidt hammer dating techniques based on data from the proglacial areas of some Icelandic glaciers. *Quat. Sci. Rev.* 18, 13–41. doi:10.1016/S0277-3791(98)00098-5

Evans, D.J.A., Clark, C., Mitchell, W.A., 2005. The last British Ice Sheet: A review of the evidence utilised in the compilation of the Glacial Map of Britain. *Earth Sci. Rev.* 70, 253–312.

Everest, J., Bradwell, T., Golledge, N., 2005. Subglacial landforms of the tweed palaeo-ice stream. *Scottish Geogr. J.* 121, 163–173. doi:10.1080/00369220518737229

- Everest, J., Kubik, P., 2006. The deglaciation of eastern Scotland: Cosmogenic ^{10}Be evidence for a Lateglacial stillstand. *J. Quat. Sci.* 21, 95–104. doi:10.1002/jqs.961
- Everest, J.D., Bradwell, T., Fogwill, C.J., Kubik, P.W., 2006. Cosmogenic ^{10}Be age constraints for the Wester Ross Readvance moraine: Insights into British Ice-Sheet behaviour. *Geogr. Ann. Ser. A Phys. Geogr.* 88, 9–17. doi:10.1111/j.0435-3676.2006.00279.x
- Everest, J.D., Bradwell, T., Stoker, M., Dewey, S., 2013. New age constraints for the maximum extent of the last British-Irish Ice Sheet (NW sector). *J. Quat. Sci.* 28, 2–7. doi:10.1002/jqs.2603
- Fabel, D., Ballantyne, C.K., Xu, S., 2012. Trimlines, blockfields, mountain-top erratics and the vertical dimensions of the last British-Irish Ice Sheet in NW Scotland. *Quat. Sci. Rev.* 55, 91–102. doi:10.1016/j.quascirev.2012.09.002
- Ffoulkes, C., Harrison, S., 2014. Evaluating the Schmidt Hammer as a method for distinguishing the relative age of late Holocene moraines: Svellnosbreen, Jotunheimen, Norway. *Geogr. Ann. Ser. A, Phys. Geogr.* 96, 393–402.
- Fowell, R.J., McFeat Smith, I., 1976. Factors influencing the cutting performance of a selective tunnelling machine. *Tunn. proc int symp IMM.*
- Geikie, J., 1984. *The Great Ice Age and Its Relation to the Antiquity of Man.* Edward Stanford, London.
- Glasser, N.F., Hughes, P.D., Fenton, C., Schnabel, C., Rother, H., 2012. ^{10}Be and ^{26}Al exposure-age dating of bedrock surfaces on the Aran ridge, Wales: Evidence for a thick Welsh Ice Cap at the Last Glacial Maximum. *J. Quat. Sci.* 27, 97–104. doi:10.1002/jqs.1519
- Goktan, R., Ayday, C., 1993. A suggested improvement to the Schmidt rebound hardness ISRM suggested method with particular reference to rock machineability. *Int. J. Rock Mech. Min. Sci.* 30, 321–322.
- Goktan, R.M., Gunes, N., 2005. A comparative study of Schmidt hammer testing procedures with reference to rock cutting machine performance prediction. *Int. J. Rock Mech. Min. Sci.* 42, 466–472. doi:10.1016/j.ijrmms.2004.12.002
- Golledge, N., Fabel, D., Everest, J., Freeman, S., Binn, 2007. First cosmogenic ^{10}Be age constrain on the timing of Younger Dryas glaciaton and ice cap thickness, western Scottish Highlands. *J. Quat. Sci.* 22, 785–791. doi:10.1002/jqs
- Golledge, N.R., 2010. Glaciation of Scotland during the Younger Dryas stadial: A review. *J. Quat. Sci.* 25, 550–566. doi:10.1002/jqs.1319
- Gosse, J.C., Phillips, F.M., 2001. Terrestrial in situ cosmogenic nuclides:theory and application. *Quat. Sci. Rev.* 20, 1475–1560.

- Goudie, A., Mignon, P., Allison, R., Rosser, N., 2002. Sandstone geomorphology of the Al Quwayra area of south Jordan. *Zeitschrift für Geomorphol.* 46, 365–390.
- Goudie, A.S., 2006. The Schmidt Hammer in geomorphological research. *Prog. Phys. Geogr.* 30, 703–718. doi:10.1016/B978-0-12-374739-6.00398-5
- Hall, K., 1993. Enhanced bedrock weathering in association with late-lying snowpatches: Evidence from Livingston Island, Antarctica. *Earth Surf. Process. Landforms* 18, 121–129.
- Hallet, B., Putkonen, J., 1994. Surface dating of dynamic landforms: Young boulders on aging moraines. *Science* 265, 937–940. doi:10.1126/science.265.5174.937
- Harrison, S., Rowan, A. V., Glasser, N.F., Knight, J., Plummer, M. A., Mills, S.C., 2014. Little Ice Age glaciers in Britain: Glacier-climate modelling in the Cairngorm Mountains. *The Holocene* 24, 135–140. doi:10.1177/0959683613516170
- Heyman, J., Stroeve, A.P., Harbor, J.M., Caffee, M.W., 2011. Too young or too old: Evaluating cosmogenic exposure dating based on an analysis of compiled boulder exposure ages. *Earth Planet. Sci. Lett.* 302, 71–80. doi:10.1016/j.epsl.2010.11.040
- Holmes, R., 1997. Reflections on the Ice Age in Scotland: An Update on Quaternary Studies, in: Gordon, J.E. (Ed.), . The Scottish Association of Geography Teachers and Scottish Natural Heritage, Glasgow, pp. 72–94.
- Hubbard, A., Bradwell, T., Gollledge, N., Hall, A., Patton, H., Sugden, D., Cooper, R., Stoker, M., 2009. Dynamic cycles, ice streams and their impact on the extent, chronology and deglaciation of the British-Irish ice sheet. *Quat. Sci. Rev.* 28, 758–776. doi:10.1016/j.quascirev.2008.12.026
- Hucka, V., 1965. A rapid method of determining the strength of rocks in situ. *Int. J. Rock Mech. Min. Sci.* 2, 127–134. doi:10.1016/0148-9062(65)90009-4
- Hughes, A.L.C., Greenwood, S.L., Clark, C.D., 2011. Dating constraints on the last British-Irish Ice Sheet: a map and database. *J. Maps* 7, 156–184. doi:10.4113/jom.2011.1145
- Hughes, P.D., Braithwaite, R.J., Fenton, C.R., Schnabel, C., 2012. Two Younger Dryas glacier phases in the English Lake District: geomorphological evidence and preliminary ^{10}Be exposure ages. *North West Geogr.* 12, 10–19.
- Hughes, P.D., Glasser, N.F., Fink, D., 2016. Rapid thinning of the Welsh Ice Cap at 20-19 ka based on ^{10}Be ages. *Quaternary Research*. DOI: 10.1016/j.yqres.2015.11.003
- International Society for Rock Mechanics, 1978. Suggested Methods for Determining Hardness and Abrasiveness of Rocks. *Int. J. Rock Mech. Min. Sci.* 15, 89–97.
- Ivy-Ochs, S., Kerschner, H., Reuther, A., Preusser, F., Heine, K., Maisch, M., Kubik, P., Schlüchter, C., 2008. Chronology of the last glacial cycle in the European Alps. *J. Quat. Sci.* 23, 559–575. doi:10.1002/jqs

Jowett, A., Charlesworth, J., 1929. The glacial geology of the Derbyshire Dome and western slopes of the southern Pennines. *J. Geol. Soc. London* 85, 307–334.

Katz, O., Reches, Z., Roegiers, J.C., 2000. Evaluation of mechanical rock properties using a Schmidt Hammer. *Int. J. Rock Mech. Min. Sci.* 37, 723–728. doi:10.1016/S1365-1609(00)00004-6

Kazi, A., Al-Mansour, R., 1980. Empirical relationship between Los Angeles abrasion and Schmidt hammer strength tests with application to aggregates around Jeddah. *Q. J. Eng. Geol. Hydrogeol.* 13, 45–52.

Kirkbride, M., Everest, J., Benn, D., Gheorghiu, D., Dawson, A., 2014. Late-Holocene and Younger Dryas glaciers in the northern Cairngorm Mountains, Scotland. *The Holocene* 24, 141–148. doi:10.1177/0959683613516171

Kirkbride, M.P., Bell, C.M., 2010. Edge-roundness of boulders of Torridonian Sandstone (northwest Scotland): Applications for relative dating and implications for warm and cold climate weathering rates. *Boreas* 39, 187–198. doi:10.1111/j.1502-3885.2009.00131.x

Kłapyta, P., 2013. Application of Schmidt hammer relative age dating to Late Pleistocene moraines and rock glaciers in the Western Tatra Mountains, Slovakia. *Catena* 111, 104–121. doi:10.1016/j.catena.2013.07.004

Lal, D., 1991. Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models. *Earth Planet. Sci. Lett.* 104, 424–439. doi:10.1016/0012-821X(91)90220-C

Letzer, J., 1978. The glacial geomorphology of the region bounded by Shap Fells, Stainmore and the Howgill Fells in East Cumbria. University of London.

Li, Y.K., 2013. Determining topographic shielding from digital elevation models for cosmogenic nuclide analysis: A GIS approach and field validation. *J. Mt. Sci.* 10, 355–362. doi:10.1007/s11629-013-2564-1

Licciardi, J.M., Pierce, K.L., 2008. Cosmogenic exposure-age chronologies of Pinedale and Bull Lake glaciations in greater Yellowstone and the Teton Range, USA. *Quat. Sci. Rev.* 27, 814–831. doi:10.1016/j.quascirev.2007.12.005

Livingstone, S.J., Evans, D.J. a, Ó Cofaigh, C., Davies, B.J., Merritt, J.W., Huddart, D., Mitchell, W. A., Roberts, D.H., Yorke, L., 2012. Glaciodynamics of the central sector of the last British-Irish Ice Sheet in Northern England. *Earth-Science Rev.* 111, 25–55. doi:10.1016/j.earscirev.2011.12.006

Livingstone, S.J., O’Cofaigh, C., Evans, D.J. A., 2008. Glacial geomorphology of the central sector of the last British-Irish ice sheet. 37–41. doi:10.4113/jom.2008.1032

Livingstone, S.J., O’Cofaigh, C., Evans, D.J.A., Palmer, A., 2010. Glaciolacustrine sedimentation in the Solway Lowlands (Cumbria,UK): evidence for a major glacial oscillation during Late Devensian deglaciation. *Boreas* 39, 505–527.

Lloyd, J.M., Zong, Y., Fish, P., Innes, J.B., 2013. Holocene and Lateglacial relative sea-level change in north-west England: Implications for glacial isostatic adjustment models. *J. Quat. Sci.* 28, 59–70. doi:10.1002/jqs.2587

Lowe, J.J., Rasmussen, S.O., Björck, S., Hoek, W.Z., Steffensen, J.P., Walker, M.J.C., Yu, Z.C., 2008. Synchronisation of palaeoenvironmental events in the North Atlantic region during the Last Termination: a revised protocol recommended by the INTIMATE group. *Quat. Sci. Rev.* 27, 6–17. doi:10.1016/j.quascirev.2007.09.016

Mann, M.E., Zhang, Z., Hughes, M.K., Bradley, R.S., Miller, S.K., Rutherford, S., Ni, F., 2008. Proxy-based reconstructions of hemispheric and global surface temperature variations over the past two millennia. *Proc. Natl. Acad. Sci. U. S. A.* 105, 13252–13257. doi:10.1073/pnas.0805721105

Marrero, S., Phillips, F., Borchers, B., Lifton, N., 2016. Cosmogenic Nuclide Systematics and the CRONUScalc Program. *Quat. Geochronol.* 31, 1–72. doi:10.1016/j.quageo.2015.09.005

Marshak, S., 2009. *Essentials of Geology*, 3rd ed. W.W. Norton.

Matsukura, Y., Matsuoka, N., 1996. Effect of rock properties on rates of tafoni growth in coastal environments. *Zeitschrift für Geomorphol.* 106, 57–72.

Matsukura, Y., Tanaka, Y., 2000. Effect of Rock Hardness and Moisture Content on Tafoni Weathering in the Granite of Mount Doeg-Sung, Korea. *Geogr. Ann. Ser. A, Phys. Geogr.* 82, 59–67. doi:10.2307/521442

Matthews, J. A., 2005. Little Ice Age' glacier variations in Jotunheimen, southern Norway: a study in regionally controlled lichenometric dating of recessional moraines with implications for climate and lichen growth rates 1, 1–19. doi:10.1191/0959683605hl779rp

Matthews, J. A., Dresser, P.Q., 2008. Holocene glacier variation chronology of the Smorstabbtindan massif, Jotunheimen, southern Norway, and the recognition of century- to millennial-scale European Neoglacial Events. *The Holocene* 18, 181–201. doi:10.1177/0959683607085608

Matthews, J. A., Owen, G., 2010. Schmidt hammer exposure-age dating: Developing linear age-calibration curves using Holocene bedrock surfaces from the Jotunheimen-Jostedalsbreen regions of southern Norway. *Boreas* 39, 105–115. doi:10.1111/j.1502-3885.2009.00107.x

Matthews, J. A., Owen, G., 2008. Endolithic lichens, rapid biological weathering and schmidt hammer r-values on recently exposed rock surfaces: Storbreen glacier foreland, jotunheimen, Norway. *Geogr. Ann. Ser. A Phys. Geogr.* 90, 287–297. doi:10.1111/j.1468-0459.2008.00346.x

Matthews, J. A., Winkler, S., 2011. Schmidt-hammer exposure-age dating (SHD): Application to early Holocene moraines and a reappraisal of the reliability of terrestrial cosmogenic-nuclide dating (TCND) at Austanbotnbreen, Jotunheimen, Norway. *Boreas* 40, 256–270. doi:10.1111/j.1502-3885.2010.00178.x

Matthews, J. A., Shakesby, R., 1984. The status of the “Little Ice Age” in southern Norway: relative-age dating of Neoglacial moraines with Schmidt hammer and lichenometry. *Boreas* 13, 333–346.

McCabe, A. M., Clark, P.U., Clark, J., 2007. Radiocarbon constraints on the history of the western Irish ice sheet prior to the Last Glacial Maximum. *Geology* 35, 147–150. doi:10.1130/G23167A.1

McCarroll, D., 1989. Schmidt hammer relative-age evaluation of a possible pre-“Little Ice Age” Neoglacial moraine, Leirbreen, southern Norway. *Nor. Geol. Tidsskr.* 69, 125–130.

McCarroll, D., 1987. The Schmidt hammer in Geomorphology: five sources of instrument error. *Br. Geomorphol. Res. Gr. Tech. Bull.* 36, 16–27.

McCarroll, D., Stone, J.O., Ballantyne, C.K., Scourse, J.D., Fifield, L.K., Evans, D.J. A, Hiemstra, J.F., 2010. Exposure-age constraints on the extent, timing and rate of retreat of the last Irish Sea ice stream. *Quat. Sci. Rev.* 29, 1844–1852. doi:10.1016/j.quascirev.2010.04.002

Mellor, A, Short, J., Kirkby, S.J., 1997. Tafoni in the El Chorro Area, Andalucia, Southern Spain. *Earth Surf. Process. Landforms* 22, 817–833. doi:10.1002/(SICI)1096-9837(199709)22:9<817::AID-ESP768>3.0.CO;2-T

Mignon, P., Placek, A., 2007. Rock control and geomorphology of a rocky sandstone scarp, Middle Sudetes Mountains, SW Poland. *Zeitschrift für Geomorphol.* 51, 41–55.

Owen, G., Owen, G., Matthews, J. A., Albert, P.G., 2007. Rates of Holocene chemical weathering, “Little Ice Age” glacial erosion, and implications for Schmidt-hammer dating at a glacier-foreland boundary, Fåbergstølsbreen, southern Norway 6, 829–834. doi:10.1177/0959683607081419

Owen, L. A., Dortch, J.M., 2014. Nature and timing of Quaternary glaciation in the Himalayan–Tibetan orogen. *Quat. Sci. Rev.* 88, 14–54. doi:10.1016/j.quascirev.2013.11.016

Phillips, F.M., 1994. Surface Exposure Dating of Glacial Features in Great Britain Using Cosmogenic Chlorine-36: Preliminary Results. *Mineral. Mag.* 58A, 722–723. doi:10.1180/minmag.1994.58A.2.113

Phillips, F.M., Argento, D.C., Balco, G., Caffee, M.W., Clem, J., Dunai, T.J., Finkel, R., Goehring, B., Gosse, J.C., Hudson, A.M., Jull, A. J.T., Kelly, M. A., Kurz, M., Lal, D., Lifton, N., Marrero, S.M., Nishiizumi, K., Reedy, R.C., Schaefer, J., Stone, J.O.H., Swanson, T., Zreda, M.G., 2015. The CRONUS-Earth Project: A Synthesis. *Quat. Geochronol.* 31, 119–154. doi:10.1016/j.quageo.2015.09.006

Phillips, W., Hall, A.M., Ballantyne, C.K., Binnie, S.A., Kubik, P.W., Freeman, S.P.H.T., 2008. Extent of the last ice sheet in northern Scotland tested with cosmogenic ¹⁰Be exposure ages. *J. Quat. Sci.* 23, 101–107. doi:10.1002/jqs

Poole, R., Farmer, I., 1980. Consistency and repeatability of Schmidt hammer rebound data during field testing. *Int. J. Rock Mech. Min. Sci.* 17, 167–171.

- Pope, G., 1995. Newly Discovered Submicron-Scale Weathering in Quartz: Geographical Implications. *Prof. Geogr.* 47, 375–387.
- Putkonen, J., Swanson, T., 2003. Accuracy of cosmogenic ages for moraines. *Quat. Res.* 59, 255–261. doi:10.1016/S0033-5894(03)00006-1
- Rode, M., Kellerer-Pirklbauer, A., 2012. Schmidt-hammer exposure-age dating (SHD) of rock glaciers in the Schoderkogel-Eisenhut area, Schladminger Tauern Range, Austria. *The Holocene* 22, 761–771. doi:10.1177/0959683611430410
- Rose, J., 2009. Early and Middle Pleistocene landscapes of eastern England. *Proc. Geol. Assoc.* 120, 3–33. doi:10.1016/j.pgeola.2009.05.003
- Rose, J., Moorlock, B.S.P., Hamblin, R.J.O., 2001. Pre-Anglian fluvial and coastal deposits in Eastern England: Lithostratigraphy and palaeoenvironments. *Quat. Int.* 79, 5–22. doi:10.1016/S1040-6182(00)00119-1
- Sánchez, J.S., Fernández Mosquera, D., Vidal Romaní, J., 2009. Assessing the age-weathering correspondence of cosmogenic ^{21}Ne dated Pleistocene surfaces by the Schmidt Hammer. *Earth Surf. Process. Landforms* 34, 1121–1125. doi:10.1002/esp
- Schimmelpfennig, I., Benedetti, L., Finkel, R., Pik, R., Blard, P.H., Bourlès, D., Burnard, P., Williams, A., 2009. Sources of in-situ ^{36}Cl in basaltic rocks. Implications for calibration of production rates. *Quat. Geochronol.* 4, 441–461. doi:10.1016/j.quageo.2009.06.003
- Schwamborn, G., Schirrmeister, L., Frütsch, F., Diekmann, B., 2012. Quartz weathering in freeze-thaw cycles: Experiment and application to the el'gygytgyn crater lake record for tracing Siberian permafrost history. *Geogr. Ann. Ser. A Phys. Geogr.* 94, 481–499. doi:10.1111/j.1468-0459.2012.00472.x
- Scourse, J.D., Haapaniemi, A.I., Colmenero-Hidalgo, E., Peck, V.L., Hall, I.R., Austin, W.E.N., Knutz, P.C., Zahn, R., 2009. Growth, dynamics and deglaciation of the last British-Irish ice sheet: the deep-sea ice-rafted detritus record. *Quat. Sci. Rev.* 28, 3066–3084. doi:10.1016/j.quascirev.2009.08.009
- Sejrup, H.P., Hjelstuen, B.O., Dahlgren, K.I.T., Hafliðason, H., Kuijpers, A., Nygård, A., Praeg, D., Stoker, M.S., Vorren, T.O., 2005. Pleistocene glacial history of the NW European continental margin. *Mar. Pet. Geol.* 22, 1111–1129. doi:10.1016/j.marpetgeo.2004.09.007
- Selby, M., 1993. Hillslope materials and processes. Oxford University Press, Oxford.
- Selby, M., 1980. A rock-mass strength classification for geomorphic purposes: with tests from Antarctica and New Zealand. *Zeitschrift für Geomorphol.* N.F 24, 31–51.
- Shakesby, R. A., Matthews, J. A., Owen, G., 2006. The Schmidt hammer as a relative-age dating tool and its potential for calibrated-age dating in Holocene glaciated environments. *Quat. Sci. Rev.* 25, 2846–2867. doi:10.1016/j.quascirev.2006.07.011

- Shakesby, R. A., Matthews, J. A., Karlén, W., Los, S., 2011. The Schmidt hammer as a Holocene calibrated-age dating technique: Testing the form of the R-value–age relationship and defining the predicted-age errors. doi:10.1177/0959683610391322
- Sissons, J.B., 1979. The Loch Lomond Stadial in the British Isles. *Nature* 280, 199–203. doi:10.1038/280199a0
- Sjöberg, R., Broadbent, N., 1991. Measurement and calibration of weathering, using the Schmidt hammer, on wave washed moraines on the upper Norrland Coast, Sweden. *Earth Surf. Process. Landforms* 16, 57–64. doi:10.1016/0148-9062(91)90609-P
- Small, D., Rinterknecht, V., Austin, W., Fabel, D., Miguens-Rodriguez, M., Xu, S., 2012. In situ cosmogenic exposure ages from the Isle of Skye, northwest Scotland: Implications for the timing of deglaciation and readvance from 15 to 11ka. *J. Quat. Sci.* 27, 150–158. doi:10.1002/jqs.1522
- Soiltest, 1976. Operating instructions-concrete test hammer.
- Stahl, T., Winkler, S., Quigley, M., Bebbington, M., Duffy, B., Duke, D., 2013. Schmidt hammer exposure-age dating (SHD) of late Quaternary fluvial terraces in New Zealand. *Earth Surf. Process. Landforms* 38, 1838–1850. doi:10.1002/esp.3427
- Stone, J.O., 2000. isotope production T , T , 105, 753–759.
- Stone, J.O., Ballantyne, C.K., Fifield, L.K., 1998. Exposure dating and validation of periglacial weathering limits, northwest Scotland. *Geology* 26, 587–590. doi:10.1130/0091-7613(1998)026<0587:EDAVOP>2.3.CO
- Sumner, P., Nel, W., 2002. The effect of rock moisture on Schmidt hammer rebound: Tests on rock samples from Marion Island and South Africa. *Earth Surf. Process. Landforms* 27, 1137–1142. doi:10.1002/esp.402
- Sumner, P., Nel, W., Holness, S., Boelhouwers, J., 2002. Rock Weathering Characteristics As Relative-Age Indicators for Glacial and Post-Glacial Landforms on Marion Island. *South African Geogr. J.* 84, 153–157. doi:10.1080/03736245.2002.9713766
- Swanson, T.W., Caffee, M.L., 2001. Determination of ³⁶Cl Production Rates Derived from the Well-Dated Deglaciation Surfaces of Whidbey and Fidalgo Islands, Washington. *Quat. Res.* 56, 366–382. doi:10.1006/qres.2001.2278
- Telfer, M.W., Wilson, P., Lord, T.C., Vincent, P.J., 2009. New constraints on the age of the last ice sheet glaciation in NW England using optically stimulated luminescence dating. *J. Quat. Sci.* 24, 906–915. doi:10.1002/jqs
- Turkington, A. V., Paradise, T.R., 2005. Sandstone weathering: A century of research and innovation. *Geomorphology* 67, 229–253. doi:10.1016/j.geomorph.2004.09.028

- Viles, H., Goudie, A., Grab, S., Lalley, J., 2011. The use of the Schmidt Hammer and Equotip for rock hardness assessment in geomorphology and heritage science: A comparative analysis. *Earth Surf. Process. Landforms* 36, 320–333. doi:10.1002/esp.2040
- Vincent, P.J., Wilson, P., Lord, T.C., Schnabel, C., Wilcken, K.M., 2010. Cosmogenic isotope (^{36}Cl) surface exposure dating of the Norber erratics, Yorkshire Dales: Further constraints on the timing of the LGM deglaciation in Britain. *Proc. Geol. Assoc.* 121, 24–31. doi:10.1016/j.pgeola.2009.12.009
- Walker, M.J.C., 2004. A Lateglacial pollen record from Hallsenna Moor, near Seascale, Cumbria, NW England, with evidence for arid conditions during the Loch Lomond (Younger Dryas) Stadial and early Holocene. *Proc. Yorksh. Geol. Soc.* 55, 33–42.
- White, A.F., Brantley, S.L., 2003. The effect of time on the weathering of silicate minerals: Why do weathering rates differ in the laboratory and field? *Chem. Geol.* 202, 479–506. doi:10.1016/j.chemgeo.2003.03.001
- White, K., Bryant, R., Drake, N., 1998. Techniques for measuring rock weathering: Application to a dated fan segment sequence in southern Tunisia. *Earth Surf. Process. Landforms* 23, 1031–1043. doi:10.1002/(SICI)1096-9837(1998110)23:11<1031::AID-ESP919>3.0.CO;2-G
- Williams, R.B.G., Robinson, D. A., 1983. The effect of surface texture on the determination of the surface hardness of rock using the Schmidt hammer. *Earth Surf. Process. Landforms* 8, 289–292.
- Wilson, P., Lord, T., Rodés, Á., 2013a. Deglaciation of the eastern Cumbria glaciokarst, northwest England, as determined by cosmogenic nuclide (^{10}Be) surface exposure dating, and the pattern and significance of subsequent environmental changes. *Cave Karst Sci.* 40, 22–27.
- Wilson, P., Schnabel, C., Wilcken, K.M., Vincent, P.J., 2013b. Surface exposure dating (^{36}Cl and ^{10}Be) of post-Last Glacial Maximum valley moraines, Lake District, northwest England: Some issues and implications. *J. Quat. Sci.* 28, 379–390. doi:10.1002/jqs.2628
- Winkler, S., 2009. First attempt to combine terrestrial cosmogenic nuclide (^{10}Be) and Schmidt hammer relative-age dating: Strauchon Glacier, Southern Alps, New Zealand. *Cent. Eur. J. Geosci.* 1, 274–290. doi:10.2478/v10085-009-0026-3
- Winkler, S., 2005. The Schmidt hammer as a relative-age dating technique: Potential and limitations of its application on Holocene moraines in Mt Cook National Park, Southern Alps, New Zealand. *New Zeal. J. Geol. Geophys.* 48, 105–116. doi:10.1080/00288306.2005.9515102
- Young, R., Fowell, R.J., 1978. Assessing rock discontinuities. *Tunnels & Tunnelling* 45–48.

Figure 1: Digital elevation model (DEM) of the UK showing the location of TCND ages and SHED sampling sites used in this study. Traditional extent of BIS redrawn from Bowen et al. (1986) and updated extent from Sejrup et al. (2005).

Figure 2: Selected photos of previously dated boulders utilized for Schmidt hammer calibration showing rock removal for cosmogenic analysis (outlined). A) Gneiss boulder in the Orkneys, B) Granite boulder in Coire an Lochan, Cairngorms, C) Sandstone boulder on the Redpoint moraine, Wester Ross, and D) Quartz boulder on Maol Chean Dearg, Torridon

Figure 3: R Value vs. ^{10}Be age calibration curves for lithologies presented in this study. A) Granite calibration curve, B) Sandstone calibration curve, C) Quartz calibration curve, D) Gneiss calibration curve, and E) Tuff calibration curve. Regression lines are demarked by solid red lines. Blue and grey dashed lines mark one and two sigma boundaries. Data used is shown by red dots with black error bars and outliers excluded from the calibration are shown by yellow dots with green error bars.

Figure 4: DEM of NW England showing location of SHED, TCND, OSL and ^{14}C samples used to date BIS retreat. Data taken from Coope and Pennington (1977), Coope and Joachim (1980), Walker (2004), Telfer et al. (2009), McCarroll (2010), Wilson et al. (2013c), Lloyd et al. (2013). Erratic source areas, erratic limits and drumlins after Clark et al. (2004) and ice front and Blackhall Wood lake extent from Livingstone et al. (2010).

Figure 5: A) Plot of 32 SHED ages after calibration from R Value to numerical age (black dots with error bars). The mean and absolute mean deviation is shown by red dot and error bars with limits marked by a horizontal grey bar. B) Comparison of TCND (green), OSL (blue) and ^{14}C (black) ages with the SHED mean age (red) and associated error bars for NW England. Green horizontal bar denotes preferred TCND age range from Wilson et al. (2013c). Horizontal grey bar marks the age range of the SHED mean age. Note the overlap between methods and the reduction of uncertainty provided by the SHED method.

Figure

Figure

A

R value = 60.0 ± 0.8

Ork 2, Phillips et al. 2008

B

R value = 47.5 ± 1.2

LCO #3, Kirkbride et al. 2014

C

R value = 56.1 ± 0.6

RM-01, Ballantyne et al. 2009

D

R value = 63.4 ± 0.6

DF-0808, Fabel et al. 2012

Figure

SHED ages

Age comparison

Table-1: Schmidt hammer sample methods arranged by year. This list is compiled from Goudie, (2006); Niedzielski et al. (2009) and Özkan & Bilim (2008).

Authors	Test Procedure
Hucka (1965)	Select the peak rebound value from 10 continuous impacts at a point. Average the peaks of the three sets of tests conducted at three separate points
Deere & Miller (1966)	Record three readings along the length of an Nxsize core for each 451 rotation. Average a total of 24 readings, disregarding the erroneous readings
Soiltest (1976)	Record 15 rebound values from single impacts and average the highest 10. The maximum deviations should be less than 2.5
Fowell & McFeat Smith (1976)	Mean of the last 5 values from 10 continuous impacts at a point
Day & Goudie (1977)	10 or 15 impacts, a greater sample size may be desirable if the material is highly variable
ISRM (1978); Young & Fowell (1978)	20 rebound values from single impacts separated by at least a plunger diameter and average the upper 10 values
Selby (1980)	25 impacts at each site
Kazi & Al-Mansour (1980)	Record at least 35 rebound readings, drop the 10 lowest readings and average the remaining 25
Poole & Farmer (1980)	Select the peak rebound value from 5 continuous impacts at a point. Average the peaks of the three sets of tests conducted at three separate points
Williams & Robinson (1983)	5 readings on each of 20 blocks
Matthews & Shakesby (1984)	15 rebound values for each sample, with 5 values that deviate most from the mean being discarded
Matthews et al. (1986); Winkler (2005); Shakesby et al. (2006)	50 boulders; one blow per boulder
Ballantyne et al. (1990)	5 readings at 5 points on 20 boulders at a site
Hall (1993)	A minimum of 100 values for each side of a dyke
Goktan & Ayday (1993)	Record 20 rebound values from single impacts separated by at least a plunger diameter. Reject outlier values using Chauvenet's criterion, and average the remaining readings
Matsukura & Matsuoka (1996)	30 rebounds
Mellor et al., (1997)	9 readings: 3 sample points with 3 readings from the corner of a 5 cm triangle centred on the point
White et al. (1998)	20 boulders sampled 8 times (i.e. on 8 different faces)
USBR (1998)	Ten readings at various locations on each surface. Discount the 5 lowest readings and average the highest 5
Katz et al., (2000)	32-40 individual impacts and averaged the upper 50%
Matsukura & Tanaka (2000)	15 at a site, with the initial 2-3 omitted
Goudie et al. (2002)	10 readings at each site
Sumner & Nel (2002)	Take 15 readings at different points and discard the 5 worst outliers to obtain a mean value from the remaining 10 values
Ericson (2004)	10 readings randomly across 2-3 dm ²
Engel (2007)	25 impacts at each site
Migoń & Placek (2007)	40 readings at each site
Owen et al., (2007)	50 impacts on quartz veins

Table-2: Details of samples used for SHED calibration with R Values and TCND ages.

Sample Name	Publication	Location (N/W)	Bedrock/ Erratic	Altitude (m)	Lithology	R Value	Uncertainty	Age (ka)	Uncertainty	Accepted?
RM-03	Ballantyne <i>et al.</i> , 2009a	57.63158, -5.66000	Erratic	329	Sandstone	50.69	0.70	15.9	0.9	✓
RM-02		57.63157, -5.65985	Erratic	333		50.85	0.55	15.0	0.9	✓
RM-01		57.63088, -5.65043	Erratic	345		56.11	0.64	15.1	0.9	✓
GM-06		57.78719, -5.70969	Erratic	223		50.10	0.48	15.0	0.9	✓
GM-03		57.76403, -5.72199	Erratic	130		47.97	0.67	16.3	1.0	✓
GM-02		57.76213, -5.73453	Erratic	101		44.78	0.71	14.2	0.8	✓
BR-05		57.85526, -5.70468	Bedrock	106		54.02	0.46	16.3	1.0	✓
BR-04		57.85479, -5.70511	Bedrock	96		43.19	0.68	17.3	1.0	✓
BR-03		57.85481, -5.70519	Bedrock	93		45.57	0.77	18.1	1.1	✓
AM-05		57.50681, -5.74867	Erratic	255		49.55	0.85	16.3	1.0	✓
AM-04		57.50669, -5.74964	Erratic	255		45.83	0.62	14.8	0.9	✓
AM-03		57.50697, -5.74968	Erratic	255		45.57	0.77	14.4	0.8	✓
AM-01		57.50642, -5.74929	Erratic	259		50.54	1.09	14.3	0.8	✓
AC-03		57.98310, -5.26063	Erratic	325		49.40	0.54	14.7	0.9	✓
AC-02		57.98536, -5.26284	Erratic	335		50.88	0.55	14.6	0.9	✓
AC-01		57.98542, -5.26306	Erratic	333		51.95	0.59	15.1	0.9	✓
LV0603	Bradwell <i>et al.</i> , 2008	58.12391, -4.98157	Erratic	101	Quartz	62.28	0.55	15.3	1.0	✓
LV0602		58.12379, -4.98118	Erratic	105		65.62	0.58	14.6	1.0	✓
LV0601		58.12404, -4.98179	Erratic	102		66.68	0.49	16.0	1.1	✓
CE9	Everest & Kubik, 2006	57.09855, -3.80015	Erratic	618	Granite	34.47	0.63	16.0	1.2	✗ ^d
CE8		57.09830, -3.80012	Erratic	617		35.13	0.70	14.3	1.2	✗ ^d
CE4		57.13184, -3.77971	Erratic	421		35.16	0.63	16.7	1.6	✗ ^d
CE3		57.12981, -3.78208	Erratic	457		39.57	0.81	16.9	1.1	✗ ^d
CE2		57.12924, -3.78420	Erratic	479		36.31	0.82	15.2	1.3	✗ ^d
CE1		57.13185, -3.78143	Erratic	422		36.00	0.60	16.1	1.2	✗ ^d
WRR/8	Everest <i>et al.</i> , 2006	57.76021, -5.73915	Erratic	84	Gneiss	47.68	1.10	14.0	2.0	✓
WRR/7		57.75983, -5.73964	Erratic	80		43.23	1.25	19.9	3.4	✓
WRR/6		57.75990, -5.73977	Erratic	79		38.86	0.86	17.8	3.1	✓
WRR/4		57.75019, -5.74571	Erratic	80		48.37	1.77	16.9	2.6	✓
DF-0815	Fabel <i>et al.</i> , 2012	57.51209, -5.40122	Erratic	915	Sandstone	56.15	0.63	15.6	0.9	✓
DF-0814		57.51197, -5.40145	Erratic	917		51.44	0.68	16.6	1.0	✓
DF-0809		57.48858, -5.46339	Erratic	835	Quartz	64.18	0.49	13.9	0.8	✓
DF-0808		57.48865, -5.46426	Erratic	838		63.39	0.61	39.0	2.1	✗ ^a

^a Inherited, ^b Transient shielding or site disturbance, ^c Overturning, ^d Quartz phenocrysts, ^e Glacially polished, ^f Anomolously soft

Table-2 continued:

Sample Name	Publication	Location (N/W)	Bedrock/ Erratic	Altitude (m)	Lithology	R Value	Uncertainty	Age (ka)	Uncertainty	Accepted?
DF-0807	Fabel et al., 2012	57.48851, -5.46355	Bedrock	837	Sandstone	49.47	0.57	14.8	1.5	✓
DF-0806		57.48870, -5.46476	Bedrock	844		52.04	0.56	13.2	0.8	✓
DF-0805		57.49104, -5.46532	Erratic	929	Quartz	59.69	0.65	50.7	2.7	✗ ^a
DF-0804		57.49150, -5.46400	Erratic	918		62.74	0.45	15.8	0.9	✓
DF-0803		57.49209, -5.46437	Erratic	918		62.94	0.56	15.9	0.9	✓
DF-0801		57.49171, -5.46310	Erratic	906		62.74	0.54	63.1	3.4	✗ ^a
AF04	Glasser <i>et al.</i> , 2012	52.77397, -3.69943	Bedrock	639	Ordovician felsic ash-flow tuffs	38.22	0.91	18.5	1.2	✓
AF03		52.78718, -3.68809	Bedrock	903		38.82	0.75	22.1	1.3	✓
AF02		52.78302, -3.69214	Bedrock	880		38.63	0.79	22.4	1.3	✓
AF01		52.78154, -3.69035	Bedrock	816		40.21	0.77	19.7	1.2	✓
AB01		52.80182, -3.68454	Bedrock	878		37.96	0.78	19.6	1.1	✓
LCO #6	Kirkbride <i>et al.</i> , 2014	57.10913, -3.68461	Erratic	895	Granite	39.25	0.76	15.3	0.8	✓
LCO #5		57.10909, -3.68562	Erratic	898		36.73	1.05	13.8	0.8	✓
LCO #4		57.10875, -3.68435	Erratic	906		39.57	1.33	14.3	1.1	✓
LCO #3		57.10859, -3.68351	Erratic	915		47.52	1.16	13.6	0.7	✓
LCO #2		57.10815, -3.68317	Erratic	928		49.01	0.94	13.2	0.7	✓
LCO #1		57.10920, -3.68513	Erratic	898		39.41	0.90	16.1	0.8	✓
LCI #6		57.10858, -3.68500	Erratic	905		42.55	1.21	14.3	0.8	✓
LCI #5		57.10789, -3.68443	Erratic	911		39.86	1.09	4.9	0.3	✗ ^c
LCI #4		57.10867, -3.68498	Erratic	902		42.49	0.79	12.7	0.7	✓
LCI #3		57.10748, -3.68256	Erratic	928		41.79	1.26	13.4	0.7	✓
LCI #2		57.10830, -3.68414	Erratic	909		44.68	1.30	5.7	0.3	✗ ^c
LCI #1		57.10834, -3.68425	Erratic	910		43.65	0.89	3.0	0.2	✗ ^c
WH2	Phillips <i>et al.</i> , 2008	58.89154, -3.36132	Erratic	342	Sandstone	38.11	0.47	35.6	2.2	✗ ^a
WH1		58.90318, -3.33989	Erratic	446		27.64	0.48	19.7	1.2	✗ ^f
PF5		57.24349, -2.55792	Bedrock	384	Granite	27.88	1.17	20.3	2.1	✓
PF4		57.24343, -2.55800	Erratic	384		30.67	1.62	19.9	2.5	✓
PF3		57.24328, -2.55800	Bedrock	385		31.00	1.13	22.3	2.7	✓
PF1		57.24342, -2.55806	Erratic	383		28.17	1.10	16.5	3.7	✗ ^b
ORK4		58.96765, -3.30502	Bedrock	94	Gneiss	45.70	1.05	14.7	2.0	✓
ORK3		58.96768, -3.30511	Bedrock	92		49.28	0.88	17.0	2.4	✓
ORK2		58.96772, -3.30520	Erratic	94		51.52	1.05	20.6	2.3	✓
ORK1		58.96772, -3.30570	Erratic	88		53.98	0.82	15.9	2.0	✓

^a Inherited, ^b Transient shielding or site disturbance, ^c Overturning, ^d Quartz phenocrysts, ^e Glacially polished, ^f Anomolously soft

Table-2 continued:

Sample Name	Publication	Location (N/W)	Bedrock/ Erratic	Altitude (m)	Lithology	R Value	Uncertainty	Age (ka)	Uncertainty	Accepted?
ORK14	Phillips et al., 2008	59.00910, -3.33618	Erratic	97	Gneiss	52.07	0.93	16.3	2.2	✓
ORK13		59.00840, -3.33670	Erratic	102		49.09	1.10	15.8	2.2	✓
ORK12		59.01373, -3.33193	Erratic	64	Sandstone	46.87	0.71	16.2	2.2	✓
ORK11		59.01318, -3.33066	Erratic	61		43.53	0.43	13.5	1.8	✗ ^b
CAITH9		58.65741, -3.38211	Bedrock	89	Sandstone	42.47	0.73	17.5	1.6	✓
CAITH8		58.32618, -3.19616	Erratic	84		40.87	0.84	17.0	1.1	✓
CAITH7		58.32618, -3.19687	Bedrock	84		55.84	0.73	15.5	1.1	✓
CAITH6		58.32627, -3.19702	Erratic	79		39.56	1.26	13.3	1.1	✗ ^b
CAITH5		58.32618, -3.19620	Bedrock	79		50.86	0.76	12.1	1.2	✗ ^b
CAITH13		58.35656, -3.19412	Erratic	131		46.74	0.90	18.9	1.3	✓
CAITH12		58.36465, -3.18856	Bedrock	164		48.69	1.36	19.6	3.0	✓
CAITH11		58.66029, -3.38128	Erratic	108		44.10	0.58	16.6	1.1	✓
CAITH10		58.66076, -3.38053	Erratic	110		44.16	0.38	61.3	3.8	✗ ^a
BEN4		57.29250, -2.56062	Bedrock	531	Granite	26.53	0.86	22.6	1.3	✓
BEN3		57.29244, -2.56153	Bedrock	524		29.99	0.61	33.6	2.0	✗ ^a
STR05-08	Small <i>et al.</i> , 2012	57.27057, -6.00919	Erratic	147	Granite	41.08	0.96	15.2	1.9	✓
STR04-08		57.27053, -6.00871	Erratic	147		41.67	0.61	14.9	0.7	✓
STR03-08		57.26892, -6.00708	Erratic	201		47.27	0.88	16.0	1.2	✗ ^e
STR02-08		57.26899, -6.00717	Erratic	198		40.39	0.70	15.6	1.1	✓
STR01-08		57.26840, -6.00664	Erratic	207		54.38	1.36	15.7	0.9	✗ ^e
SLAP-5		57.19839, -6.00131	Erratic	23		45.70	0.81	11.8	0.9	✓
SLAP-4		57.19812, -6.00149	Erratic	21		41.39	0.58	12.6	0.8	✓
SLAP-3		57.19868, -6.00110	Erratic	26		41.69	1.14	12.1	0.8	✓
SLAP-2		57.19865, -6.00110	Erratic	26		45.40	0.85	12.7	0.9	✓
SLAP-1		57.19865, -6.00111	Erratic	26		43.30	0.74	13.4	0.9	✓
AT-3	Stone <i>et al.</i> , 1998	57.82861, -5.23541	Bedrock	611	Sandstone	48.12	0.58	14.4	1.2	✓
AT-2		57.82940, -5.23432	Bedrock	609		44.81	0.51	14.8	1.1	✓
AT-1		57.83052, -5.23368	Bedrock	607		45.46	0.40	15.0	1.2	✓
SHAP-08	Wilson <i>et al.</i> , 2013c	54.47993, -2.61753	Erratic	330	Granite	28.85	0.52	21.0	1.2	✓
SHAP-07		54.49098, -2.61322	Erratic	325		38.87	0.68	16.5	1.0	✓
SHAP-03		54.48643, -2.63406	Erratic	220		35.84	0.54	18.8	1.1	✓
SHAP-02		54.48320, -2.63761	Erratic	280		39.67	0.80	17.1	1.0	✓

^a Inherited, ^b Transient shielding or site disturbance, ^c Overturning, ^d Quartz phenocrysts, ^e Glacially polished, ^f Anomolously soft

Table-3: Linear regression and significance of R Value calibration for each lithology

Lithology	<i>Number of samples</i>	R^2	Regression	<i>p value</i>
Granite	25	0.81	-0.4882x + 34.8394	< 0.01
Sandstone	30	0.08	-0.1053x + 20.9012	0.12
Quartz	6	0.02	-0.0592x + 19.0378	0.81
Gneiss	10	0.02	-0.0731x + 20.3917	0.67
Ordovician felsic ash-flow tuffs	5	0.01	0.1584x + 14.3161	0.90

Table-4: Sample location, R Values, and calibrated SHED ages for SHAP area samples.

Sample Name	Location (N/W)	Sample type	Altitude (m)	R Value	Uncer tainty	Sample type	Age (ka)	Uncer tainty
^a CR-1	54.51377, -2.58928	Erratic	221	50.50	0.94	SHED	10.4	1.6
CR-2	54.51384, -2.58917	Erratic	219	33.17	0.88	SHED	18.6	1.5
CR-3	54.51392, -2.58910	Erratic	217	36.76	0.86	SHED	16.9	1.5
CR-4	54.51355, -2.58822	Erratic	213	36.44	0.84	SHED	17.0	1.5
CR-5	54.51358, -2.58792	Erratic	214	37.42	0.78	SHED	16.6	1.5
CR-6	54.51454, -2.58748	Erratic	210	38.46	0.88	SHED	16.1	1.5
CR-7	54.51458, -2.58746	Erratic	208	39.01	0.86	SHED	15.8	1.5
CR-8	54.51448, -2.58776	Erratic	209	38.05	0.74	SHED	16.3	1.5
CR-9	54.51443, -2.58789	Erratic	208	37.88	0.79	SHED	16.3	1.5
CR-10	54.51454, -2.58821	Erratic	206	36.46	0.84	SHED	17.0	1.5
CR-11	54.51461, -2.58833	Erratic	207	38.93	0.82	SHED	15.8	1.5
CR-12	54.51461, -2.58856	Erratic	206	36.47	0.82	SHED	17.0	1.5
CR-13	54.51469, -2.58862	Erratic	207	38.13	0.74	SHED	16.2	1.5
CR-14	54.51483, -2.58849	Erratic	208	38.96	0.89	SHED	15.8	1.5
CR-15	54.51484, -2.58848	Erratic	208	37.89	0.89	SHED	16.3	1.5
CR-16	54.51488, -2.58839	Erratic	210	36.72	0.77	SHED	16.9	1.5
CR-17	54.51493, -2.58833	Erratic	208	37.62	0.70	SHED	16.5	1.5
CR-18	54.51493, -2.58818	Erratic	208	36.27	0.84	SHED	17.1	1.5
CR-19	54.51473, -2.58806	Erratic	208	38.07	0.94	SHED	16.3	1.5
CR-20	54.51476, -2.58823	Erratic	209	37.04	0.77	SHED	16.8	1.5
CR-21	54.51468, -2.58835	Erratic	210	34.68	0.64	SHED	17.9	1.5
CR-22	54.51467, -2.58826	Erratic	210	36.26	0.57	SHED	17.1	1.5
CR-23	54.51464, -2.58820	Erratic	209	37.13	0.72	SHED	16.7	1.5
CR-24	54.51463, -2.58807	Erratic	210	37.86	0.86	SHED	16.4	1.5
CR-25	54.51453, -2.58794	Erratic	211	38.66	0.88	SHED	16.0	1.5
CR-26	54.51463, -2.58764	Erratic	211	38.16	0.84	SHED	16.2	1.5
CR-27	54.51470, -2.58752	Erratic	212	37.92	0.92	SHED	16.3	1.5
CR-28	54.51481, -2.58777	Erratic	213	38.79	0.85	SHED	15.9	1.5
CR-29	54.51491, -2.58754	Erratic	215	37.22	0.83	SHED	16.7	1.5
CR-30	54.51490, -2.58748	Erratic	215	38.91	0.92	SHED	15.8	1.5
CR-31	54.51509, -2.58789	Erratic	214	39.93	0.90	SHED	15.3	1.5
CR-32	54.53400, -2.64785	Erratic	295	39.65	0.79	SHED	15.5	1.5
OxL-1802	54.077, -2.136	Sediment	365	N/A	N/A	OSL ^b	16.5	1.7
OxL-1806	54.147, -2.775	Sediment	103	N/A	N/A	OSL ^b	19.3	2.6
OxL-1790	54.117, -2.071	Sediment	365	N/A	N/A	OSL ^b	27.8	2.6
OxL-1789	54.117, -2.071	Sediment	365	N/A	N/A	OSL ^b	27.2	2.6
SHAP-08	54.47993, -2.61753	Erratic	330	N/A	N/A	TCND ^c	21.01	1.22
SHAP-03	54.48643, -2.63406	Erratic	220	N/A	N/A	TCND ^c	18.78	1.08
SHAP-02	54.48320, -2.63761	Erratic	280	N/A	N/A	TCND ^c	17.09	1.03
SHAP-07	54.49098, -2.61322	Erratic	325	N/A	N/A	TCND ^c	16.45	1.00

^aGlacially polished; ^bdata from Telfer et al., 2009; ^cdata from Wilson et al., 2013c