Nature and timing of Quaternary glaciation in the Himalayan-Tibetan orogen

DOI:

[10.1016/j.quascirev.2013.11.016](https://doi.org/10.1016/j.quascirev.2013.11.016)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Owen, L. A., & Dortch, J. M. (2014). Nature and timing of Quaternary glaciation in the Himalayan-Tibetan orogen. *Quaternary Science Reviews*, 88, 14-54. <https://doi.org/10.1016/j.quascirev.2013.11.016>

Published in:

Quaternary Science Reviews

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Highlights

- Evidence for Himalayan-Tibetan Quaternary glaciation is reviewed and evaluated
- Numerical dating methods used to define timing of glaciation are evaluated
- Strong regional contrasts in timing and extent of glaciation occur across the orogen
- Glaciation influenced by Asian monsoon, mid-latitude westerlies and topography
- Recommendations made to help advance future Himalayan-Tibetan glacial research

1 Nature and timing of Quaternary glaciation in the Himalayan-Tibetan 2 orogen

3
4 Lewis A. Owen^{a*} and Jason M. Dortch^b

5
6 ^a*Department of Geology, University of Cincinnati, Cincinnati, OH 45221, USA*

7 ^b*School of Environment and Development, The University of Manchester, M0 1QD, UK*

8 9 ABSTRACT

10 Much effort has been made in recent years to define the timing and extent of Quaternary
11 glaciation throughout the Himalayan-Tibetan orogen. These studies are challenging
12 because of the logistical and political inaccessibility of the region, and the inherent
13 problems associated with the application of numerical dating techniques. Nevertheless,
14 the studies are providing abundant evidence for significant glacial advances throughout
15 the last several glacial cycles and are beginning to accurately define the extent and timing
16 of glaciation in selected regions. Studies are showing that Himalayan-Tibetan glaciers in
17 arid regions during the last glacial cycle reached their maximum extent early in the cycle
18 and that global Last Glacial Maximum glacier advances were significantly less extensive.
19 However, along the more monsoonal-influenced Greater Himalaya, there is increasing
20 evidence to suggest that glaciation was more extensive later in the last glacial cycle, but
21 this has yet to be fully assessed. In addition, the new studies are showing that throughout

* Corresponding author.

E-mail address: Lewis.Owen@uc.edu (L.A. Owen)

most Himalayan-Tibetan regions, significant glacier advances occurred during the Late Glacial and early Holocene, with minor advances in some regions during the mid-Holocene. The still relatively poor chronological control in the Himalayan-Tibetan orogen, however, makes it difficult to construct correlations across the region, and with regions elsewhere in the world. This in turn makes it hard to assess the relative importance of the different climatic mechanisms that force glaciation across the Himalayan-Tibetan orogen, and to quantify paleoclimate change in this high altitude subtropical region. The Late Glacial and Holocene glacial records, however, are particularly well preserved in several Himalayan-Tibetan regions. Glacial successions such as these have the greatest potential to be examined in detail using newly developing numerical dating, and geomorphic and sedimentologic methods to derive high-resolution terrestrial records of glaciation that will help in paleoclimatic reconstruction for high altitude subtropical regions.

Keywords: Himalaya; Tibet; diamicts; glaciers; radiocarbon dates; cosmogenic dating; OSL dating; Asian monsoon; mid-latitude westerlies

1. Introduction

The mountains of the Himalayan-Tibetan orogen are the highest on Earth and the orogen is the most glaciated region outside of the polar realms (with ~ 126 200 km² glacier cover; Haeberli et al., 1989). Glaciers in Himalayan and Tibetan mountains are the source for numerous rivers that flow across the Indo-Gangetic Plain and central and eastern China, which provide freshwater to billions of people. Furthermore, the Himalayan-

Tibetan orogen has a profound influence on regional and global atmospheric circulation and it is therefore important for understanding the dynamics of global environmental change (Ruddiman & Kutzbach 1989; Molnar & England 1990; Prell & Kutzbach 1992; Owen et al. 2002d). The study of glaciation in the Himalayan-Tibetan orogen is also important for providing a framework to examine glacial controls on the development of topography, conditioning landscapes in mountain systems and focusing erosion in them, and governing tectonic feedbacks to the evolution of mountains (Brozović *et al.*, 1997, Zietler *et al.*, 2001; Norton *et al.*, 2010, Willett, 2010). Contention exists over the rate at which glaciers are currently receding to the extent that erroneous estimates have suggested that they are receding faster than in any other part of the world and are likely to disappear by at least the AD 2035 (Intergovernmental Panel on Climate Change, 2007; Bagla, 2009). Impressive glacial landforms and successions of Quaternary glacial sediments are abundant throughout the orogen, recording the nature and style of glacial oscillations.

Yet despite the importance of the region and its potentially useful geologic archive, the nature and dynamics of Himalayan-Tibetan glaciers throughout the Quaternary are not well understood. In particular, the timing and extent of past glacier fluctuations are poorly defined. This is partially because the logistical and political inaccessibility of the region has made research there particularly challenging. Nevertheless, the region has had a long history of glacial research, but the older studies were quite restricted in extent because of the logistical and political difficulties. Notably significant historical studies include those of Drew (1873), Dainelli (1922, 1934, 1935), Norin (1925), Klute (1930),

Trinkler (1930) and De Terra and Paterson (1939). In recent years, numerous researchers have undertaken detailed studies of the region's Quaternary glacial geology aided by better accessibility, newly developing remote sensing technologies and numerical dating, and a renewed interest in the region and its glaciers. This has resulted in a plethora of publications that are beginning to allow us to develop a dynamic picture for the Quaternary glaciation of the Himalayan-Tibetan orogen.

The aim of this manuscript is to present a review of the Quaternary glacial geology of the Himalayan-Tibetan orogen and to speculate on the role of climate and other forcing factors on glaciation in the region. Our manuscript builds on reviews by Shi et al. (1986), Derbyshire et al. (1991), Benn and Owen (1998), Lehmkuhl and Owen (2005), Owen et al. (2008a), Owen (2009, 2013), and regional syntheses in Elhers et al. (2004, 2011), Dortch et al. (2013), Murari et al (2013), and notably, Kamp and Owen (2011), Owen (2011) and Zhou et al. (2011).

2. The Physical Environment

The Himalayan-Tibetan orogen is a consequence of the collision of island arcs with the Eurasian continental plate, and the collision of the Indian and Eurasian continental plates at ~ 50 Ma and the subsequent continued northward movement of the Indian plate at ~ 50 mm/a (Yin and Harrison, 2000; Searle and Richard, 2007). This has resulted in a mountain mass made up of several approximately east-west trending ranges, with a combined average elevation of ~ 5000 m asl (Fielding, 1996), composed of a complex assemblage of rocks of different ages. The Himalayan-Tibetan orogen now stretches ~

2000 km and ~ 1500 km in an east–west and north–south direction, respectively, the main ranges comprising, from south to north, the Siwaliks, Lesser Himalaya, Greater Himalaya, Transhimalaya, Nyaingentanglha Shan, Tanggula Shan, Bayan Har Shan, Kunlun Shan, Altun Shan and Qilian Shan. In addition, the active mountains of the Pamir, Tian Shan and Altai Mountains are also part of the orogen (Fig. 1). However, we do not discuss the Altai Mountains in this paper as we do not consider them contiguous with the main mountain mass, and so these regions are beyond the scope of this paper.

The mid-latitude westerlies and the south Asian monsoon are the two dominant climatic systems influencing the region (Benn and Owen, 1998; Fig. 2). As Lehmkuhl and Owen (2005), Owen and Benn (2005), Owen et al. (2008a) and Owen (2009) highlight, numerous studies have attempted to link long-term (thousands of years) variations in these climatic systems, influenced by changes in Northern Hemisphere insolation, and shorter-term variations (years to decades), explained by changes within the climate system, to mountain glacier fluctuations throughout the Quaternary.

The two climatic systems vary spatially with the majority of the southern and eastern regions experiencing a pronounced summer precipitation maximum, reflecting moisture advected northwards from the Indian Ocean by the southwest monsoon. Summer precipitation declines sharply northward across the main Himalayan chain and is very low over western Tibet (Benn and Owen, 1998; Owen, 2009). The mid-latitude westerlies produce a winter precipitation maximum on the western end of the Himalaya, Transhimalaya and western Tibet as a consequence of moisture being advected from the

Mediterranean, Black, and Caspian Seas (Benn and Owen, 1998; Owen, 2009). These climate systems produce strong north-south and west-east precipitation gradients, as illustrated in Figure 3, ranging, for example, from one of the wettest places on Earth, the East Khasi Hills (with average annual precipitation >12 m) to one of the driest, the hyperarid Takla Makan Desert; these gradients strongly influence the rate of a variety of surface processes including mass movements, fluvial incision, alluvial fan deposition, and glaciation (Bookhagen et al., 2005; Bookhagen and Burbank, 2006; Dortch et al., 2011a). There are also strong microclimatic variations within individual mountain ranges and valleys, potentially exerting strong controls over glaciation (Böhner, 2006; Dortch et al., 2011b). Benn and Owen (1998) hypothesized that the relative role of the two dominant climate systems, the south Asian monsoon and mid-latitude westerlies, varied significantly throughout the Quaternary, resulting in asynchronous glaciation across the orogen. Trying to test this hypothesis has resulted in numerous studies in recent years, which will be discussed in more detail below.

Vegetation varies as a consequence of climate and altitude from thick subtropical forests in the foothills of the Himalaya and SE Tibet to deciduous forests at higher altitude and eventually alpine meadow and mountain tundra at the highest elevations with xerophytic vegetation in the arid interiors of the orogen. The equilibrium-line altitudes (ELAs) for glaciers across the orogen vary considerably as a consequence of the different roles of the two main climatic systems and the precipitation gradients, for example, ranging from 4300 m asl in southeast Tibet to over 6000 m asl in western Tibet (Owen and Benn, 2005). Benn and Lehmkuhl (2000) and Owen and Benn (2005) describe the controls on

mass balance and variation of present and past ELAs through the Himalaya and Tibet. Owen and Benn (2005) highlight that the present regional ELA varies in altitude across ranges; for example, it is 4600–5700 to 6000–6200 m asl south to north across Mount Everest, 5200–5700 m asl south to north across the Garhwal Himalaya, 4500–4700 m asl around Nanga Parbat, and 4600–5300 to 5200–5700 m asl south to north across the Karakoram Mountains.

Climate and topography have a strong influence on the nature of glacial systems throughout the Himalayan-Tibetan orogen. In essence, glacial systems are highly variable. Derbyshire (1982) divided the glaciers into three main types: 1) continental interior types in the central and western parts of the Tibet Plateau; 2) maritime monsoonal types in the Himalaya and in southeastern Tibet; and 3) continental monsoonal types in eastern and northeastern Tibet (Fig. 4). As Owen and Benn (2005) point out, most glaciers within the region occupy steep, high relief catchments, and have cold, high altitude (> 5000 m asl) accumulation areas. Where mass turnover is high, or where debris cover allows glaciers to descend to lower altitudes, ablation zones may be temperate whereas glaciers located in dry high altitude environments are cold throughout. Annual temperature cycles vary with the degree of continentality, altitude and insolation.

The maritime and continental monsoonal glaciers of the Himalaya and southeastern and eastern Tibet are warm-based and have summer accumulation and ablation; they are also highly diverse, including avalanche- and snowfall-fed cirque and valley glaciers, and very steep hanging glaciers (Derbyshire, 1981; Benn and Lehmkuhl, 2000; Benn and

Owen, 2002). Benn and Owen (2002) emphasized that snow avalanching from precipitous slopes forms an important component of accumulation on many of these glaciers, and glacier ablation areas can be separated from source areas by steep icefalls or avalanche tracks. These are usually high activity glaciers that may have velocities up to several hundred meters per year (e.g. Seong et al., 2009d). Basin topography, therefore, plays an important role in determining the geometry and mass balance of these monsoonal glaciers.

Alpine continental glaciers in the western part of the orogen (Transhimalaya, Pamir, and Tian Shan) are significantly smaller than their monsoonal counterparts and are generally low flux due to an annual precipitation typically < 1000 mm combined with high and cold accumulation areas. In spite of this, they produce landforms (roche moutonnées and large moraines) indicating mixed based if not warm based conditions at times (Koppes et al., 2008; Dortch et al. 2010, 2011b,c; Röhringer et al., 2012). In contrast, the continental ice caps of central and western Tibet have basal ice temperature $\ll 0^{\circ}\text{C}$ and they have an annual precipitation $\ll 1000\text{mm}$ (at altitudes > 5000 m asl). The Guilya icecap, for example, has an ice temperature of -18°C and has an annual precipitation of 300–400 mm (Shi, 2002). Shi (2002) noted that these continental glaciers are similar to sub-polar types, with low surface velocities, usually between 2 and 10 m/a, and may extend outside the permafrost zone. In addition, these glaciers are usually less topographically constrained and generally have limited debris cover compared to those in the Himalaya (Owen and Benn, 2005).

Benn and Owen (2002) provided a review of Himalayan glacial sedimentary systems, describing the complexity of associated landform and lithofacies, and the importance of erosional, depositional and deformational processes. They showed that many Himalayan glaciers have extensive mantles of supraglacial debris on their ablation zones, whereas others have little or no supraglacial debris; a spectrum of glacier types exists between these end members. Benn and Owen (2002) reasoned that the amount of debris cover on glacier surfaces is controlled by a number of factors, the most important of which is the distribution of steep slopes in the glacial catchment from which avalanches can deliver rock debris, either from bedrock or pre-existing glacial and paraglacial sediments. Other factors influencing debris cover include: (1) precipitation, which governs the amount of snowfall relative to rock inputs, and hence the debris concentration on the ice; (2) glacier size (long valley glaciers are most likely to have extensive debris mantles); and (3) bedrock erodibility (resistant, massive rocks such as granite will yield much less debris than highly fractured schists and sedimentary rocks). The abundance of supraglacial debris helps produce complex latero-frontal moraines that may record one or more glacial oscillations (Figs. 4D and E). Other important studies of the glacial sedimentary system include those of Derbyshire (1981, 1982, 1984, 1986), Li et al. (1984) and Owen and Derbyshire (1989).

Supraglacial debris may be so dominant in some glaciated catchments that morainic rock glaciers develop (Owen and England, 1998; Fig. 4F). Protalus rock glaciers may also form in permafrost regions (Owen and England, 1998; Fort, 2003; Iwata et al., 2003). Owen and England (1998) showed that, in the Karakoram and Himalaya of Northern

Pakistan and India, morainic and protalus rock glaciers are restricted altitudinally and climatically to sites above ~ 4000 m asl and in regions where annual precipitation is < 1000 mm. Generally in these regions, morainic rock glaciers are large, usually > 1 km long, > 100 m wide and > 15 m thick. Owen and England (1998) suggested that morainic rock glaciers record the advance of ice-cored moraines following retreat of glaciers, likely since the Little Ice Age.

3. Reconstructing past glaciations in the orogen

Many researchers throughout the twentieth century and increasingly over the last few decades have reconstructed the extent of Late Quaternary glaciation throughout the Himalayan–Tibetan orogen. Important compilations and reconstructions of glaciation throughout the orogen include those of Klute (1930), Frenzel (1960), Li et al. (1991) and Shi (1992), which show expanded ice caps and extensive valley glaciers (Fig. 5). The studies of Klute (1930) and Frenzel (1960) are remarkable given the limited studies that were available to these researchers. The most recent reconstructions by Li et al. (1991) and Shi (1992) represent an impressive compilation of studies and are in general agreement with most recent research showing alpine-style glaciation throughout the region during the last few glacial cycles. Numerous field studies provide regional maps of reconstruction (e.g. Porter, 1970; Derbyshire et al., 1984; Cronin et al., 1989; Holmes and Street-Perrott, 1989; Watanabe et al., 1989; Shiraiwa and Watanabe, 1991; Lehmkuhl and Lui, 1994; Osmaston, 1984; Owen et al., 1995, 1996, 1997, 2000; Taylor and Mitchell, 2000; Lehmkuhl et al., 2004; Heyman et al., 2010, 2011). Most recently, remote sensing has enabled very impressive glacial geologic maps to be reconstructed, notably by

Duncan et al. (1998), Heyman et al. (2008) and Moren et al. (2011).

In contrast to most studies, Kuhle (1985, 1986, 1987, 1988a,b, 1990a,b, 1991, 1993, 1995) argued for an extensive ice sheet covering most of the Tibetan Plateau during the Last Glacial. However, much evidence has been presented against the possibility that an extensive ice sheet could have existed across Tibet during the last 500 ka (Derbyshire, 1987; Burbank and Kang, 1991; Derbyshire et al., 1991; Shi et al., 1992; Hövermann et al., 1993a,b; Lehmkuhl, 1995, 1997, 1998; Rutter, 1995; Lehmkuhl et al., 1998; Zheng and Rutter, 1998; Schäfer et al., 2002; Owen et al., 2003a, 2008a; Seong et al., 2008; Dortch et al. 2013). In particular, Lehmkuhl et al. (1998), Owen et al. (2008), Seong et al. (2008), and Owen (2010) argue that Kuhle's (1985, 1986, 1987, 1988a,b, 1990a,b, 1991, 1993, 1995) reconstructions likely differ from other researchers because of his misinterpretation of landforms and sediments, the misuse of ELAs for defining former ice extent, and poor chronological control. Specifically, numerical dating that has been undertaken in Tibet shows that moraines for the majority of the most extensive glacial advances formed early in the Last Glacial, the penultimate glacial cycle or much earlier, proving that an ice sheet could not have existed during the Last Glacial, otherwise the moraines would have been destroyed. In an attempt to try to reconcile this evidence with his model for an ice sheet over Tibet, in contrast to all the other research being undertaken with radiocarbon, OSL and TCN dating in the Himalayan-Tibetan orogen and other high mountain regions, Kuhle (2011) dismisses numerical dating, arguing that numerical dating cannot be applied at high altitudes as it is at lower elevations. He argues that all radiometric ages need calibrating to make ages $1/6^{\text{th}}$ or 83% younger than

published ages to fit his ice sheet model. However, his conversion factor arguments are spurious, there being no validity to his suggestions. Ironically, this now resolved controversy with Kuhle (1985, 1986, 1987, 1988a,b, 1990a,b, 1991, 1993, 1995) has left the idea for an ice sheet over Tibet firmly out of favor but it may have been possible, although highly unlikely, for an ice sheet to form over Tibet during the early- to mid-Quaternary. Surficial glacial landforms of this period have not been identified, but further insight might come from marine and lake core stratigraphy.

Clearly, accurate reconstructions of former glacial extent rely on the sound interpretation of landforms and sediments as well as geochronology. In particular, researchers such as Derbyshire (1983, 1996), Owen and Derbyshire (1988), Fort (1986), Fort and Derbyshire (1988), Derbyshire and Owen (1990, 1997), Hewitt (1998, 1999), Benn and Owen (2002) and Hewitt et al. (2011), have warned of the dangers of misinterpreting glacial and mass movement landforms in the Himalaya and Tibet. Large landslides are common throughout the Himalayan-Tibetan orogen and their forms can often resemble moraines (Fig. 6). In addition, a continuum of landforms including moraines, rock glaciers, talus slopes and landslides exists that exacerbates the problems of distinguishing landform types (Owen, 1991; Fort, 2003). Moreover, intense fluvial and glacial erosion often destroys diagnostic morphologies of glacial and mass movement landforms, making their identification difficult (Owen, 1991; Benn and Owen, 2002). Furthermore, the diamictos that comprise mass movement and glacial landforms look very similar and have similar particle size distributions and particle shapes. All this makes it very challenging to identify landforms and sediments correctly unless sediments are examined with due care.

275
276 Hewitt (1998, 1999) and Hewitt et al. (2011) provide important examples from the
277 Karakoram of Northern Pakistan where landslides have been misidentified as moraines
278 (Fig. 7). Unfortunately, few papers provide detailed descriptions of the landforms and
279 sediments that are used to reconstruct former glacier extents. This makes it difficult to
280 fully assess the validity of many of the reconstructions. Benn and Owen (2002) provide a
281 useful overview to help researchers recognize the problems associated with identifying
282 landforms and sediments in Himalayan environments and suggest approaches to
283 accurately interpret the glacial geomorphic and sedimentological evidence.
284
285 Most detailed reconstructions of former glacial extents within individual study areas
286 apply standard morphostratigraphic methods similar to those outlined in Rose and
287 Menzies (1995). In essence, glacial and associated landforms are mapped, based on their
288 morphostratigraphic positions and the relative weathering of their forms and surfaces in
289 each detailed study area. Often this is made easy because large relief within a landscape
290 helps to separate moraines of different glacial advances at different altitudinal positions.
291 Relative dating methods have included the use of soil development (Guggenberger et al.,
292 1998; Bäumler et al., 1999; Zech et al., 2000), and weathering characteristics and rock
293 varnish development (Owen et al., 1996, 1997; Waragai, 1998, 2005). Commonly,
294 researchers will assign a glacial stage name to a particular suite of glacial and associated
295 landforms that they recognize to have formed during a distinct glacial advance (e.g.
296 Derbyshire et al., 1984; Burbank and Fort, 1985; Seong et al., 2007; 2009b, Owen et al.,
297 2003, 2012). Some studies provide type locations for a particular glacial stage, which

may be used by future researchers as a basis for comparing and extending the glacial chronologies (e.g. Seong et al., 2009b; Owen et al., 2012). Other studies simply number moraines of different stages (e.g. Zech et al., 2005; Abramowski et al., 2006; Koopes et al. 2008; Hendrick et al., 2011) or provide no number or name at all (e.g. Kong et al., 2009b). Applying glacial stage names or a numbering scheme allows landforms to be tentatively correlated across a study region and between adjacent regions before a quantitative chronostratigraphy has been established; however, correlation based on the number of stages in a sequence without chronology should be undertaken with extreme caution. Most recent studies are applying the approach described by Hughes et al. (2005) to develop formal stratigraphies. Figure 8 shows an example of the morphostratigraphic approach that has been applied to glacial successions within one valley of the Himalayan-Tibetan orogen.

Once a morphostratigraphy has been developed, it can form the framework for numerical dating to help define the timing of each glacial advance/glacial stage. The application of numerical dating will be discussed in detail in the next section. Detailed mapping is also commonly used to reconstruct the size of former glaciers as a basis for determining former ELAs (Benn and Lehmkuhl, 2000; Benn et al., 2005; Owen and Benn, 2005).

4. Dating methods: challenges and advances

Much effort has been undertaken in recent years to define the timing of glaciation throughout the Himalaya and Tibet but, until the turn of the millennium, few studies presented numerical ages for glacial successions in the Himalayan-Tibetan orogen.

Gillespie and Molnar's (1995) review of the timing of mountain glaciation throughout the world, and its asynchronicity with continental glaciers, highlighted the paucity of data for the Himalayan-Tibetan region. Similarly, Benn and Owen's (1998) review and speculation on the relative roles of the summer monsoon and mid-latitude westerlies in driving Himalayan-Tibetan glaciation emphasizes the dearth of numerical ages.

The former lack of numerical ages mainly resulted from a scarcity of organic material needed for radiocarbon dating in most regions. Where organic material was present, ~95% (142 of 149) of ages were limited to the Holocene. This is illustrated by the compilation of radiocarbon dating we present in Table 1 and Figure 9.

Röthlisberger and Geyh (1985a,b, 1986) undertook an extensive study using radiocarbon dating (including 68 radiocarbon ages). They examined eight different regions throughout the Himalaya and Karakoram of Pakistan, India and Nepal to show that glaciers had undergone at least 11 advances since the global Last Glacial Maximum (LGM) (Fig. 9A). However, only relatively modest studies have been undertaken using radiocarbon dating since the work of Röthlisberger and Geyh (1985a,b, 1986). Most studies have dated charcoal or wood (e.g. Benedict, 1976; Fushimi, 1977, 1978; Derbyshire et al., 1984; Geyh et al., 1985; Röthlisberger and Geyh, 1985a,b; Zheng and Li, 1986; Wang and Fan, 1987; Zhang, 1988; Zheng and Ma, 1994; Chinese Academy of Sciences, 1996; Narama, 2002; Su et al., 2002; Meyer et al., 2007; Lee et al. 2013), but many have used humic matter (Li and Jiao, 1990; Li and Li, 1992; Jiao and Iwata, 1993; Zheng et al., 1995; Iwata and Jiao, 1993; Owen et al., 1997; Iwata et al., 2002; Jiao et al., 2005; Zheng,

2006; Jiao and Zheng, 2006). Carbonate precipitates have also been dated but these are subject to large uncertainties because of hardwater effects; also the carbonates may have developed significantly later than the formation of the landform that is being dated yielding very minimum ages (e.g. Mann et al., 1996; Yi et al., 2004).

Yi et al. (2008) reviewed much of the radiocarbon dating Holocene glacial chronologies for Tibet and the surrounding mountains. They compiled 58 ages, but notably omitted the studies of Röthlisberger and Geyh (1985a,b, 1986). In addition, Yang et al. (2008) reviewed the radiocarbon dating undertaken to establish the Late Holocene monsoon temperate glacier fluctuations on the Tibetan Plateau. Building on these compilations, we compile the radiocarbon ages for studies throughout the Himalayan-Tibetan orogen and present these in Table 1 and Figure 9. We plot minimum radiocarbon ages, i.e., ages that are younger than the age of the moraine that is being dated, which are then used to calculate a probability distribution density graph. This allows us to determine whether these age data define the timing of glacial oscillations throughout the Late Quaternary, since the peaks of the probability distributions likely represent revegetation of the glacial foreland, which is assumed to approximate the time of deglaciation. The frequency of peaks is similar to those of the Holocene rapid climate change as suggested by Mayewski et al. (2004). However, this similarity relies on the assumption that glaciations dated by radiocarbon methods are synchronous throughout the orogen. This assumption needs to be tested in future radiocarbon-based studies by targeting Holocene landforms in several climatic regions of the orogen.

Following the significant developments in OSL and TCN surface exposure dating during the 1990s and early 2000s, numerous researchers began to apply these methods throughout the Himalayan-Tibetan orogen. OSL methods were applied before TCN methods, initially using thermoluminescence (TL; e.g., Derbyshire et al., 1984). The first extensive program of OSL dating in the Himalaya was undertaken by Richards (1999) resulting in detailed studies in the Khumbu Himal, Nepal (Richards et al. 2000b), and the Swat and Nanga Parbat regions in Northern Pakistan (Richards et al. 2000a). Other significant studies using OSL include those of Tsukamoto et al. (2002) in the Kanchenjunga Himalaya, Derbyshire et al. (1984) and Spencer and Owen (2004) in the Hunza valley of northern Pakistan, Narama et al. (2007) in the Terskey Alatau Range (Kyrgyz Republic), and Narama et al. (2009) and Zhao et al. (2012) in the Tien Shan. Richards (2000) discusses the problems associated with the application of OSL methods in the Himalaya, which include those associated with inadequate bleaching of sediment and low luminescence signals (see Fuchs and Owen (2008) for more details about OSL dating of glaciogenic sediments). Table 2 summarizes the OSL dating studies that have been undertaken throughout the Himalayan-Tibetan orogen.

Another potentially important technique, electron spin resonance (ESR) dating has been undertaken in a few areas (e.g. Wu et al., 2001; Zhao et al., 2009, 2012). Researchers using ESR contend that the ESR signal is reset in till by pressure during glacial grinding. However, there is much contention over whether this mechanism can fully reset the ESR signal and, as such, ESR ages should be considered very conservatively.

Arguably, TCN dating has been applied more to the Himalayan-Tibetan glacial successions than any other glaciated mountain region in the world. Currently, for example, there are > 1700 ^{10}Be ages on moraine boulders in addition to numerous glacially eroded bedrock surfaces, plus boulder ages and depth profiles on associated landforms. Table 3 and Appendix 1 list the ^{10}Be ages for chronologies throughout the Himalayan-Tibetan orogen. Most TCN studies use ^{10}Be for dating, but ^{26}Al has also been applied, usually on the same sample that is being dated with ^{10}Be (e.g. Owen et al., 2001, 2002a, 2003b,c; Schäfer et al., 2002; Tschudi et al., 2003). In addition, ^3He and ^{21}Ne have been applied in several studies (Tschudi et al., 2003; Gayer et al., 2006; Graf et al., 2008; Schäfer et al., 2008; Strasky et al., 2009).

Problems associated with the application of TCN methods to date moraines in the Himalaya, Tibet and elsewhere, have been discussed in depth in numerous studies (e.g. Hallet and Putkonen, 1994; Benn and Owen, 2002; Owen et al., 2002a, 2003a,c, 2005, 2006a,b, 2008a, 2009, 2010, 2012; Putkonen and Swanson, 2003; Putkonen and O'Neal, 2006; Seong et al., 2007, 2008, 2009a; Putkonen et al., 2008; Dortch et al. (2010b); Chevalier et al., 2011; Heyman et al., 2011b). In particular, Owen et al (2008) highlighted the uncertainty introduced when calculating TCN ages for Himalayan and Tibetan glacial successions using different scaling models and that this may result in ages that differ by as much as 30% using 777 ages (see Figs. 5 and 6 in Owen et al., 2008). Here, updated plots including the now > 1700 published ages from the orogen show that ages from various researchers differ by up to ~ 40% from recalculated Lal (1991) – Stone (2002) time independent ages (CRONUS Calculator Version 2.2; Fig. 10A). We highlight this

to illustrate the problems with using published data at face value. Moreover, as Balco et al. (2008) points out, the greatest difference between ages calculated by the different scaling models are those for low latitudes and high altitudes, which encompasses the Himalaya-Tibetan orogen. The resulting differences can be seen in Figure 10B, where percent difference after using Lifton et al. (2005), Desilets et al. (2003, 2006), Dunai (2001), and Lal (1991) – Stone (2000) time dependent scaling schemes have been plotted with respect to Lal (1991) – Stone (2002) time independent ages. These large differences present challenges when comparing TCN ages with radiocarbon, OSL, and other numerical age methods, as well as climatic records. The accuracy of the reference sea level, high-latitude (SLHL) ^{10}Be productions rate has been challenged by the regional rates of Balco et al. (2009), Putnam et al. (2010), Fenton et al. (2011), Kaplan et al. (2011), Briner et al. (2012), and Goehring et al. (2012). The new regional ^{10}Be production rates range between 5-15% lower than SLHL and result in older ages. Figure 10C shows the difference in percent between Balco et al. (2009) and Putnam et al. (2010) regional production rates with respect to Lal (1991) – Stone (2000) SLHL of $4.5 \text{ at } \text{g}^{-1} \text{ yr}^{-1}$ time-independent ages. Although these studies show broad agreement, they only cover a limited temporal ($< 15 \text{ ka}$) and elevation ($< 1500 \text{ m asl}$) scale and a regional calibration site comparing radiocarbon and TCN methods has not been defined in the Himalayan-Tibetan orogen. Moreover, differences in isostatic rebound, air-pressure, and differing paleomagnetic field effects could create significant offsets when extrapolating to dissimilar regions (e.g. Balco et al. 2009). Thus, extrapolation of production rates from calibration sites to high elevation ($>4000 \text{ m}$) and old landforms (50-700 ka) should only be undertaken with extreme caution. Moreover, the TCN data used in calibration papers

do not have perfect gaussian distributions; thus, when using the subsequent productions rates, researchers should use the same statistical methods to examine clustering of ages. Unfortunately, there is no scaling scheme or production rate that has been rigorously proven to be the most accurate, leaving researchers free to choose the scheme that they deem best, or best fits their interpretations. Ironically, if carefully chosen, using the combination of new production rates and scaling scheme, ~ 78% (n=1331) of all ages yield results within 10% of time-independent Lal (1991) – Stone (2000) ages when between 11-700 ka; whereas only ~ 10% (n=169) of all ages are within 15% between 5-11 ka (Fig. 10D). These correction factors do not appear to be quite as significant when taken in the context of normal individual age errors of 5-10%.

Irrespective of these various correction schemes, the geologic error associated with TCN dating and the statistical treatment of ages to identify outliers has a far more significant effect on dating moraines and other landforms. As Hallet and Putkonen (1994), Owen et al. (2008) and numerous other researchers point out, geological factors present the largest uncertainty in applying TCN surface exposure dating methods. These include early stabilization and denudation of the landform, as well as weathering, exhumation, prior exposure and shielding of the surface that is being dated by snow and/or sediment. Generally these processes reduce the concentration of TCNs, resulting in an underestimate of the true age of the landform, with the exception of prior exposure that results in an overestimation of the landform's age. The net result of these processes can be a large spread in apparent exposure ages of individual boulders on a single landform, such as a moraine, or in different positions on an extensive rock surface such as glacially

eroded bedrock. For example, Dortch et al. (2013) show that 40% of all ages in the western semi-arid part of the orogen are outliers with age underestimation occurring four times more often than inheritance. They show that inherited outliers are 60% higher than landform age on average, with numerous samples having up to 300% inheritance, while underestimation diverges from moraine age by an average of ~ 55% younger with numerous samples diverging by up to 90%. Due to possible post-processes, researchers typically interpret TCN ages as representing the timing of deglaciation. Researchers usually collect multiple samples on a surface in an attempt to assess these effects. The combined results of Dortch et al. (2013; n=95) and Murari et al. (2013; n=208) that included the examination of 303 local glacial stages shows that when ≤ 4 samples are collected, the probability of obtaining three ages that fit a normal distribution is ~27%, but dramatically shifts to at 5 samples to is ~77%, and increases with ≥ 6 samples to ~98% (Fig. 11). The basic rationale is that the presence of multiple boulders or surface samples having similar apparent ages is taken as evidence that the boulders were not derived from a range of older surfaces and/or were not weathered, exhumed or covered. Additionally, sequences of landforms are often dated to assess the relative behavior of clustered TCN ages. Hence the ages are then considered to be representative of the true age of the surface. However, it is not common to have strong clustering of ages, particularly for Pleistocene moraines, and the spread of ages is generally greater with antiquity.

Figure 12 provides an example of a large, reasonably well-clustered data set for a study area within the Himalayan-Tibetan orogen. Each rectangle encloses individual ^{10}Be ages

for a particular glacial stage. Clearly, the clustering of ^{10}Be ages is tightest for the younger glacial stages. The greatest challenge is interpreting the broad spread of ages. Many researchers, including Hallet and Putkonen (1994), Zreda et al. (1994), Putkonen and Swanson (2003), Briner et al. (2005), Zech et al. (2008), Applegate et al. (2010) and Heyman et al. (2011), favor assigning the glacial age based on the age of the oldest boulder (if not an “obvious” outlier) determined for a surface. The oldest boulder method implies that the boulder has been least weathered, shielded or exhumed. However, the oldest age might be a consequence of inherited TCNs. Moreover, the method of determining if a boulder is an “obvious” outlier varies from a simple judgment call to using a non-stringent statistical test (e.g. beyond 2σ of the mean of the population). In contrast, the youngest boulder might be considered most representative since it is least likely to have inherited TCNs from former exposure. Putkonen and Swanson (2003) and Heyman et al. (2011) argue that only a small percent of dated moraine boulders have had prior exposure, which lends favor to the oldest boulder method. To reiterate the point, Dortch et al. (2013) caution that 8% of all glacial boulders dated in the semi-arid western portion of the orogen show inheritance typically ranging from 5-300% with rarer examples with up to 1200% of the landform age (for ages $< 1\text{--}350\text{ ka}$) and that outliers are not always obvious. Similar analysis by Murari et al. (2013) supports this view. They show that 10% of all glacial boulders dated in the monsoon influenced regions of Himalaya and Tibet show inheritance typically ranging from 6-300% with rare examples exceeding 1000%. To add complication, both studies show that young outliers occur 3–4 times more often than inherited outliers and argue that statistical analysis is required to identify the most correct ages.

505
506 Numerous researchers suggest statistical analysis of ages, such as applying the mean
507 square of weighted deviates method (MSWD); iterative reduced chi-squared, is more
508 appropriate because it is a much stricter test and easily identifies outliers (e.g. Gosse et
509 al., 1995; Finkel et al., 2003; Meriaux et al., 2004; Chevalier et al., 2005; Owen et al.,
510 2005; Dortch et al., 2010, 2011; Hedrick et al., 2011). However, Ivy-Ochs et al. (2007)
511 argue that using MSWD and the resulting weighted mean unjustly skews ages because
512 younger ages typically have smaller uncertainties. Dortch et al. (2013) separated
513 gaussians from cumulative probability frequency distributions to obtain best fits on TCN
514 data for semi-arid regions at the western end of the Himalayan-Tibetan orogen, which
515 they then used to delimit local glacial stages. Student's t-test was used to correlate TCN
516 age clusters across the region, the results of which will be discussed in more detail below.
517 Clearly, the issue of understanding age data needs to be resolved and care must be taken
518 when considering the ages of glacial events in the Himalayan-Tibetan orogen.
519 Fortunately, most researchers present all their primary data so that future researchers can
520 manipulate published data and make their own interpretations. However, we suggest that
521 studies focused on the timing of glaciation and the links to climatic records should use
522 the Lal (1991) – Stone (2000) time-independent scaling scheme for comparability until
523 geomagnetic correction and SLHL productions rate uncertainties are better constrained
524 and that significant attention needs to be given to developing a standard for the statistical
525 treatment of ages.

526

527 A strategy taken by some researchers to try to overcome problems with dating is to apply

both OSL and TCN methods within a study area. This has the advantage of providing depositional ages (OSL dating) and deglaciation ages (minimum ages with TCN dating). Benn and Owen (2002) set out possible sampling strategies that might be applied using both OSL and TCN methods, as illustrated in Figure 13. Unfortunately, one of the main limitations of using this strategy is the availability of good exposures of sediment for OSL dating. This strategy has been successfully applied in several regions, most notably in the Khumbu Himal and Rongbuk valley on the south and north sides of Mount Everest, respectively. Figure 14 shows an example of a laterofrontal moraine complex that was dated using both OSL and TCN methods at the Lhotse Nup glacier terminus at Chhukung in the Khumbu Himal (Richards et al. 2000b; Finkel et al., 2003). ^{10}Be surface exposure ages on this moraine are between 2.6-3.5 ka whereas an OSL age on the upper sediment unit is 1.1 +/- 0.2 ka, supporting the view that the moraine represents a Late Holocene glacial advance. In this case, the older TCN ages might reflect a thousand or years of prior exposure. Unlike TCN methods, OSL dating also allowed an earlier advance to be defined to the early Holocene (~10 ka) by dating the lower sediment unit at this site. Figure 15 shows the chronologies for the Rongbuk valley, which are well defined by both ^{10}Be TCN and OSL dating methods.

5. Evidence of extremely old glaciations

There is abundant geomorphic and sedimentological evidence for glaciation predating the penultimate glacial cycle in the Himalayan-Tibet orogen. In some cases moraines are well preserved, especially in the semi-arid regions of Tibet and the Transhimalaya, but in most regions old moraines are denuded. Amongst the oldest moraines are those of the

Indus glacial stage in Ladakh. Owen et al. (2006a) showed these to have been formed when extensive valley glaciers advanced to an elevation below 3250 m asl; they are dated to > 430 ka (Fig. 16A). Other extremely old moraines are present on the west side of Gurla Mandata in southernmost central Tibet (Fig. 16B). Based on ^{10}Be dating of boulders, Owen et al. (2010) argued that these moraines are considerably older than 300 ka and thus likely formed during MIS 10 or during an earlier glacial cycle when ice caps expanded during the Naimona'nyi glaciation. In addition, in the Rongbuk valley on the northern side of Mt Everest, Owen et al. (2009) dated erratics to > 330 ka, which they argued formed by extensive valley glaciers advancing northwards from Mount Everest. In the Central Karakoram, Seong et al. (2007) recognized an extensive valley glacier system, which they dated to MIS 6 or older. Other moraines of great antiquity (penultimate or earlier glacial cycle) have been dated using TCN methods by Owen et al. (2006b), Chevalier et al. (2005, 2011), Schäfer et al. (2008) and Hedrick et al. (2011) in the Kunlun Shan, Ayilari Range, Nyalam, Xainza graben in south central Tibet, and the Zaskar, respectively. On Muztag Ata, in the Chinese Pamir, Seong et al. (2009a) showed that ice caps expanded during the Karasu glacial stage, which they dated to the penultimate glacial cycle or earlier using ^{10}Be dating. Just south of Muztag Ata in the Tashkurgan valley, Owen et al. (2012) dated moraines of the Dabudaer glacial stage using ^{10}Be to the penultimate glacial cycle and/or earlier (Fig. 16C).

Using ESR and TL methods, Wu et al. (2001) dated tills of the Wangkun Glaciation on the Kunlun Pass to between 0.7-0.5 Ma. In addition, using ESR methods, Zhao et al. (2001) dated till of the Gaowangfeng Glacial Stage in the headwaters of the Urumqi

River in the Eastern Tien Shan to ~ 400-470 ka, while Zhao et al. (2009) dated till of the Qingshantou Glacial Stage in the Ateaoyinake River Valley of the central Tien Shan to ~440 ka. Till of the Zhonglianggan glaciation in the Qilian Mountains was also dated using ESR methods by Zhou et al. (2001, 2006) yielding an age of ~ 463 ka.

The large uncertainty associated with dating these old moraines and sediments precludes regional correlation. Moreover, it is difficult to correlate convincingly the old moraines within a region. Owen et al. (2012) for example, stated that moraines assigned to their Dabudaer glacial stage in the Tashkurgan valley might represent several glaciations, since the strong degree of weathering of Dabudaer glacial stage landforms does not allow subtle morphostratigraphic distinctions to be made.

Old moraines (penultimate glacial cycle or older) and sediments in high-precipitation monsoon-influenced Himalayan regions are not as common as for the semi-arid regions, however. This might simply be because the preservation potential for old moraines is low due to intense denudation, mainly by fluvial and mass movement processes, which characterize these wetter environments. It is also possible that Late Glacial and Holocene glacial advances may have been more extensive than early glaciations and hence may have destroyed any landform or sedimentary evidence of earlier glaciations in the monsoon-influenced Himalayan regions. Due to the great relief in these settings, some evidence of older glaciations is expected because of the large altitudinal separation of landforms. On the other hand, the high relief yields more gravitational potential, which would drive quicker reworking of landforms through various surface processes.

597

598 Owen et al. (2005, 2006a, 2008, 2010, 2012) and Seong et al. (2009) noted, that in
599 regions where there is evidence for very old glaciation there is a progressive change in
600 style of glaciation with a significant reduction in glacial extent over successive glacial
601 cycles. Owen (2005, 2006a, 2008, 2012) and Seong et al. (2009) hypothesized that this
602 could reflect a change in regional climatic forcing that might be the result of the
603 progressive surface uplift of adjacent mountain ranges that restricted moist input and
604 hence precipitation in their regions. They cautioned that drawing links between mountain
605 uplift and changes in regional climate is very tentative because it is not possible to define
606 the timing and magnitude of surface uplift for most mountain ranges. However, since this
607 pattern of glaciation occurs in disparate regions across the Transhimalaya and Tibet,
608 Owen et al. (2005, 2006a, 2008, 2010, 2012) and Seong et al. (2009) argued that it might
609 reflect regional climate change rather than localized tectonics. In contrast, Dortch et al.
610 (2013) argue that most major glacial events are driven by monsoonal precipitation and
611 that change in regional climate is not likely responsible for decreased glacial extent
612 during the late Quaternary. An alternative mechanism is the adjustment of topographic
613 hypsometry to glacial erosion and the lowering of accumulation area driving the
614 reduction of glacial extent during late Quaternary cycles (e.g. Kaplan et al. 2009;
615 Pedersen and Egholm, 2013). The reason for this apparent changing pattern of glaciation
616 through time, however, has yet to be resolved.

617

618 **6. The Last Glacial**

619 There is abundant evidence in the Himalayan-Tibetan orogen for multiple glacial

advances throughout the last glacial cycle. In some regions up to five distinct glacial advances have been recognized, for example, in the Khumbu Himal (Finkel et al. 2003), Muztag Ata-Kongus Shan (Seong et al. 2009a), and Xainza Range (Chevalier et al., 2011). Care must be taken, however, when interpreting and trying to correlate glacial successions because the preservation of the glacial geologic record may well be biased by likely preservation potential (Gibbons et al., 1984).

Given the large number of ^{10}Be ages that have been generated in recent years, Owen et al. (2008a) and Chevalier et al. (2011) examined the timing of glaciation by generating probability distribution frequency plots for the ^{10}Be ages for selected time slices and regions. Figure 17 builds on the analysis of Owen et al. (2008) by increasing their data set from 777 to > 1700 ^{10}Be ages. The new plots in Figure 17 use only moraine boulder ages. As in Owen et al. (2008), we recognize that pooling all the ages runs the risk of including ages that may be too old or too young to represent the true age of a glacial advance because of the geological problems highlighted above. Given the great number of ages, however, clear patterns emerge. As in the study by Owen et al. (2008a), the probability plots shown in Figure 17 demonstrate a decrease in sample frequency with increasing age. This is essentially a function of preservation, with younger landforms being better preserved, more common and less difficult to date. Multiple peaks are apparent for most of the probability plots during MIS 3 and the Late Glacial, and several peaks occur within the Holocene. The new plots support the view of Owen et al. (2008a) that there is not a dominant peak for the global LGM centered on 21 ka but, instead, a dominant peak occurs in the Late Glacial and is centered around 12–14 ka. In addition, the multiple

peaks during the Holocene suggest that Himalayan-Tibetan glaciers may be sensitive to climate oscillations, reflecting times of Holocene rapid climate change emphasized by Mayewski et al. (2004).

The probability distribution frequency plots presented in Figure 17, however, may be spurious since they combine data from climatically different regions where glaciers may be responding differently to different climate forcing. To test for regional patterns, as in Owen et al. (2008a), the ^{10}Be data have been sorted into four climatic-topographic regions (Fig. 18): 1) Pamir and Tien Shan dominated by the mid-latitude westerlies; 2) semi-arid Transhimalaya and western Tibet dominated by the mid-latitude westerlies; 3) monsoon-influenced Himalaya including northern Pakistan, India and Nepal; 4) monsoon-influenced Tibet, including southern central, central, SE and NE Tibet. Three time slices are considered: 1) all ^{10}Be boulder ages back to 200 ka; 2) all ^{10}Be boulder ages younger than 74 (MIS 5a); and 3) all ^{10}Be boulder ages younger than 15 ka (to include Late Glacial and Holocene ages). The study areas are listed in Table 3 and Appendix 1.

The caveats in Figure 17 apply to the plots in Figure 18 when interpreting the data. Clearly the probability plots still reflect bias in occurrence and sampling of younger moraines and the distributions are broadly similar. It is notable that there is no peak centered on ~ 30-50 ka (MIS 3) for the Pamir and Tien Shan data, but rather there is a peak at 60-75 ka (MIS 4); however, there is no peak at this time for the other three regions (Fig. 18A). However, due to significantly fewer samples collected from

landforms >30 ka in age, erosional processes could easily diminish any > 30 ka peak into the background so that the sampled regions of Pamir and Tien Shan may only reflect a particular peak due to very dry conditions promoting preservation. The Pamir and Tien Shan data have apparent bimodal peaks between the Late Glacial through to the global LGM, whereas no other data set has a peak at the global LGM (Fig. 18B). As pointed out by Owen et al. (2008a), this suggests that glaciation in the Pamir and Tien Shan has responded differently compared to the other three regions. In contrast, Dortch et al. (2013) shows a strong regional glacial signal at 20 ± 2 ka in the Pamir and Western Transhimalaya by analyzing individual datasets, highlighting the weakness of combining datasets without first eliminating outliers. It is likely that glaciation in the Pamir and Tien Shan is forced by northern hemisphere climate change and is broadly synchronous with oscillations in the northern hemisphere ice sheets. In contrast, as suggested by numerous researchers, in other areas of the Himalayan-Tibetan orogen, glaciers respond to changes in the south Asian monsoon. In particular, the peak centered around MIS 3 supports the view that glaciers advanced during a time of increased insolation, which helped drive the monsoon influence farther into the orogen resulting in increased precipitation and cloudiness, further resulting in positive glacier mass balances (e.g. Finkel et al., 2003; Owen et al., 2003a, 2005, 2006b, 2009; Rupper et al., 2008). Moreover, the ages and glacial geologic evidence shows that, in many regions of the Himalayan-Tibetan orogen, glaciers advanced to their largest extents earlier in the last glacial cycle than the northern hemisphere ice sheets, thus supporting the views of Gillespie and Molnar (1995) and Thackray et al. (2008).

The probability distribution frequency plots also show a significant Late Glacial advance (Fig. 18C). The peaks are broad and it is not possible to assign the advance to a particular stadial such as the Younger Dryas.

Figure 19 examines regional variability in more detail. The picture that emerges is complex, it being evident that many of the glacial advances span long durations, in some cases many tens of thousands of years. This largely reflects the uncertainty in the chronological control. As a consequence, it is clear that a lot of age data are dispersed within probability distribution frequency plots but, nevertheless, the plots are useful for examining possible broad trends. The bars representing glacial advances shown in Figure 19 are color coded to help recognize correlations. Particularly notable is the lack of glacial advances prior to the Late Glacial in many of the monsoon-influenced regions of the Himalaya and Tibet. Possible reasons for this were discussed above. Notable glacial advances occurred during the Late Glacial, early Holocene, mid-Holocene, Neoglacial and during the last millennia in nearly all areas. The data suggest broad patterns with glaciers responding at Milankovitch and sub-Milankovitch frequencies; it is likely that, for most regions both the monsoon and mid-latitude westlies force glaciation, albeit to different extents. This makes it difficult to correlate glacial stages across the orogen.

When considered in more detail, the patterns are even more complex. Figure 20, for example, highlights the main study areas at the western end of the Himalayan-Tibetan orogen. The timing of the local last glacial maximum is indicated on the left of the figure, with approximate maximum extents of glaciation. ELA data are not presented because of

the complexity of determining them for each region (see Benn and Lehmkuhl, 2000, Benn et al., 2005 and Owen and Benn, 2005 for more discussion). The pattern that emerges is very complex with the local last glacial maximum occurring at very different times across the western end of the orogen. Figure 20 also compares these data with the climate modeling presented in Bishop et al. (2010) that covers the same aerial extent. The variation in snowfall across the region and between time slices is striking. Although the modeling focuses on the global LGM and the Holocene, it illustrates the complex and differing patterns across the region at different times. It is therefore not surprising that the style and timing of glaciation across the orogen is complex.

Despite this complexity, Dortch et al. (2013) examined the glacial successions and ^{10}Be ages across the semi-arid regions at the western end of the Himalayan-Tibetan orogen to develop a regional framework for glaciation across these dryland regions. Using Gaussian separation, Dortch et al. (2013) recognized nineteen regional glacial stages that they named the semi-arid western Himalayan-Tibetan stages (SWHTS). These are illustrated in Figure 21. The data suggest that regional glacial stages older than 21 ka are broadly correlated with a greater monsoonal influence. SWHTS that are 21 ka or younger have smaller uncertainties and broadly correlate with global ice volume as defined by marine oxygen isotope stages and northern hemisphere climatic events (Oldest Dryas, Older Dryas, Younger Dryas, Roman Humid Period, and Little Ice Age). The pattern of glaciation that emerges shows no significant spatial clustering within the study area even though several correlative advances throughout the region were identified. The lack of spatial grouping and correlation between glacial chronologies and climatic records led

Dortch et al. (2013) to speculate that the arid regions respond to small fluctuations in precipitation brought via the mid-latitude westerlies, sediments being subsequently wiped out by larger, less frequent monsoon driven advances. This view is in agreement with Seong et al. (2009) regarding westerly control of post-LGM advances outlined below.

Using the methods of Dortch et al. (2013), Murari et al. (2013), together with new ^{10}Be -ages to define the timing of glaciation in SW Garhwal, examined the glacial chronologies throughout the monsoon-influenced regions of the Himalaya. They recognized 29 regional glacial stages, which they called monsoon-influenced Himalayan-Tibet stages (MOHITS). Murari et al. (2013) suggest that there are strong correlations with both periods of strong monsoons and northern hemisphere events throughout the entire chronologic range. They identify 16 stages linked to the monsoon and 11 stages linked to the mid-latitude westerlies (two unassigned) with a complex pattern of glaciation influenced by two climatic systems since the mid/late Quaternary. The larger number of regional glacial stages compared to the semi-arid regions suggests more frequent moraine deposition during the last glacial cycle and the Holocene. However, this may be an artifact of the larger number of dated glacial sediments and landforms in the monsoon influenced regions.

Comparison between the semi-arid and monsoon influenced regions are shown in Figure 21 and visually compared with the SWHITS, and statically compared using the student t-test in Table 4. We recognize that much care must be taken when making such correlations within between regions as large as the Himalayan-Tibetan orogen because

some detailed complexity may be lost with grouping of data statistically. However, the correlations provide a framework for future studies. Moreover, it is very striking the large dataset is now developing to define the timing of glacier oscillations throughout the region over the last half million years, and the high frequency of those oscillations.

7. Holocene Glaciation

Zhou et al. (1991), Yi et al. (2008) and Owen (2011) provide useful summaries of the nature of Holocene glacier fluctuations in the Himalaya and Tibet. In particular, Owen (2011) drew attention to the paucity of comprehensive studies of Holocene glaciation in the region, but emphasized the potential to derive a high-resolution (sub-millennial scale) record of glaciation for paleoclimatic change using the Holocene glacial record. This is because the Holocene glacial record is better preserved than older moraine successions and numerical dating can be more readily and successfully applied to younger moraines. The Yi et al. (2008) review included a compilation of 53 radiocarbon ages for Holocene glacial advances in Tibet, which they argued that glaciers advanced at 9.4–8.8 ka, 3.5–1.4 ka, and 1.0–0.13 ka, suggesting that glaciation in Tibet is synchronous with cooling periods identified in the $\delta^{18}\text{O}$ record of ice core. Zhou et al. (1991) showed that Holocene continental glaciers in northwest China advanced at ~ 9.3 ka, 6.4 ka, 4.5 ka, and 0.5 ka (~ 8300, 5700, 4000 and 400 ^{14}C years BP), while maritime-influenced glaciers in southeastern Tibet advanced at 3.1 ka, 1.9 ka, 0.9 ka, and 0.3 ka (~ 3000, 2000, 1000, and 200 ^{14}C years BP), arguing for a 2500-year climate cycle at ~ 9.3 ka, 6.4 ka, 3.2 ka and 0.3 ka (8300, 5700, 3000 and 200 ^{14}C years BP), and a ~ 1000-year climate cycle at 3.2 ka, 1.9 ka, 0.9 ka, and 0.3 ka (~3000, 2000, 1000, and 200 ^{14}C years BP).

781
 782 Röthlisberger and Geyh (1985a,b) and Seong et al. (2009a) are arguably the two most
 783 significant quantitative studies of Holocene moraines in the Himalayan-Tibetan orogen
 784 that present primary data. Both studies also included an examination of global LGM and
 785 Late Glacial moraines. Röthlisberger and Geyh (1985a,b), presented 68 radiocarbon ages
 786 from around 16 different glaciers in Pakistan, India and Nepal to show that glaciers
 787 advanced at ~ 8.3 ka, 5.4–5.1 ka, 4.2–3.3 ka, and 2.7–2.2 ka (7400, 4900–4600, 3700–
 788 3100, 2700–2100 ^{14}C years BP) with relatively small extensions at 2.6–2.4 ka (2500–
 789 2300 ^{14}C years BP), 1.7–1.4 ka, 1.3–0.9 ka, 0.8–0.55 ka and 0.5–0.1 ka (1700–1500,
 790 1200–950, 800, 550 and 400–100 ^{14}C years BP; Fig. 9).

791
 792 The study of Seong et al. (2009a) utilized ^{10}Be TCN dating of moraines around Mustah
 793 Ata and Kongur Shan in the Chinese Pamir to show that glaciers advanced at ~ 11.2 ka,
 794 10.2 ka, 8.4 ka, 6.7 ka, 4.2 ka, 3.3 ka, 1.4 ka, and at a few hundred years before the
 795 present (Fig. 22). They argued that, since the global LGM, glaciers in western Tibet
 796 likely responded to Northern Hemisphere climate oscillations, with minor influences
 797 from the south Asian monsoon, responding on millennial timescales to Holocene rapid
 798 climate changes that were highlighted by Bond et al. (1997, 2001) and Mayewski et al.
 799 (2004) (Fig. 22).

800
 801 The probability distribution plots in Figure 18C, and introduced in the previous section,
 802 show significant early Holocene peaks. The early Holocene advance likely reflects
 803 similar monsoonal forcing to the MIS 3 advance, since this was a time of increased

insolation and enhanced monsoonal influence. Climate modeling supports the view that increased precipitation and snowfall throughout much of the Himalayan-Tibetan region took place during the early Holocene (Bush, 2004, Rupper et al., 2008; Li and Harrison, 2008; Bishop et al., 2010). Rupper et al. (2008) argued that an increase in cloudiness reduces incoming shortwave radiation that results in greater cooling at the surface of the glacier and accounts for more of the ELA lowering than an increase in monsoon precipitation. Multiple peaks are evident for the mid- and late Holocene, representing millennial-scale glacier oscillations. These glacial advances may reflect periods of rapid climate change influenced by teleconnections to Northern Hemisphere climate systems as suggested by Seong et al. (2009a) and Owen (2009).

As Owen (2009) underscored, these studies support the view that Himalayan and Tibetan glaciers oscillated at a comparable frequency to that recognized in ice cores from the Greenland Ice Sheet and in deep-sea sediment cores from the North Atlantic, where a quasi-periodicity of ~ 1470 years is apparent throughout the Late Glacial and Holocene (Bond et al., 1997, 2001). Seong et al. (2009a) and Owen (2009) pointed out that the duration of quasi-periodicity of glacial oscillations throughout the Holocene is similar to climatic oscillations recognized in the North Atlantic by Mayewski et al. (2004), and suggested that regional climate change in Tibet may well be forced by teleconnections via mid-latitude westerlies and Atlantic climate change. Moreover, Owen (2009) suggested that it is likely that as many as eight significant glacial advances would be expected to have occurred throughout the Holocene in Himalayan-Tibetan regions. He also stressed that rapid climate change and accompanied glacial oscillations is also

supported by the ice core data from Tibetan ice sheets, which show short (millennial-centennial) abrupt oscillations in climate throughout the Late Quaternary (Thompson et al., 1989, 1997; Thompson, 2000; Yao et al., 1996, 1997). In addition, Dortch et al. (2013) recognized late-Pleistocene and early Holocene glacial advances that correlate with Heinrich events. They documented three regional glacial advances during the latter part of the Holocene in the semi-arid western end of the Himalayan-Tibetan orogen at 3.7 ± 0.6 ka (SWHT 1C), 1.6 ± 0.3 ka (SWHTS 1B) and 0.4 ± 0.1 ka (SWHTS 1A) that they argue broadly correlate with Northern Hemisphere climatic events (Fig. 21).

Unlike many other mountain regions throughout the world, the most extensive glacial advance during the Holocene in the Himalayan-Tibetan orogen occurred during the early Holocene (cf. Grove, 2004; Clapperton, 1993; Porter, 2000). This advance is represented by impressive suites of sharp-crested moraines which have been dated by numerous researchers using radiocarbon, OSL and TCN methods to between ~ 11.5 and 8.0 ka (Shiraiwa, 1993; Sharma and Owen, 1996; Phillips et al., 2000; Richards et al., 2000a; Owen et al., 2001, 2002b, 2003a,c, 2005, 2006a,b; Finkel et al., 2003; Zech et al., 2003, 2005; Barnard et al., 2004a,b, 2006a,b; Spencer and Owen, 2004; Abramowski et al., 2006; Jiao and Shen, 2006; Seong et al., 2007, 2009a; Meyer et al., 2009; Chevalier et al., 2011). The extent of glacial advances and ELA depressions during the early Holocene varies considerably between regions; for example, glaciers only advanced a few kilometers in the Muztag Ata-Konga Shan and Hunza regions with an ELA depression of < 200 m (Benn and Owen, 2005; Seong et al., 2009c) as compared to the Central Karakoram where glaciers advanced several tens of kilometers with an ELA depression

of > 500 m (Seong et al., 2009b). The disparity of ELA depression may be due to relief control of precipitation as outlined by Bookhagen and Burbank (2006), with the high-relief Karakoram receiving more precipitation due to orographic effects.

Figure 19 highlights this early Holocene regional signal, but the resolution of the dating in most of the studies, which generally utilize TCN methods, is too coarse to determine whether this apparent correlation is synchronous on a millennial scale. Moreover, most of the dating does not allow us to distinguish whether the early Holocene advance was synchronous with the early Holocene isolation maximum, and hence is a monsoon signal, or the 8.5 ka cold event, or both. Reducing the dating uncertainty on early Holocene moraines is clearly an area that needs attention in order to advance understanding of the nature of early Holocene glaciation and climate change.

Owen (2011) pointed out that mid-Holocene (~ 8.0–3.0 ka, Hypsithermal) moraines have been dated in several regions of the Himalaya and Tibet. These moraines are generally well preserved and often comprise latero-frontal moraine complexes. As Owen (2011) points out, the Hypsithermal has yet to be adequately defined by glacial geologic proxies for the Himalaya and Tibet and the use of the term in this region should therefore be used very conservatively. Moreover, it may be diachronous in monsoonal Asia (An et al., 2000). Significant glacial advances (a few kilometers beyond present ice margins) have been recognized, however, for selected glaciers in Tibet (Zhou et al., 1991), in the Khumbu Himal (Finkel et al., 2003), the Diancang Mountain in the Hengduan Range (Yang et al., 2006), Qilian Shan (Wu, 1984a), and in Muztag Ata and Konga Shan (Seong

et al., 2009a). Climatic simulations support the view that increased monsoon precipitation and cloudiness, and summer cooling over the Himalaya and Tibet would have helped force glaciation during the mid- Holocene, albeit not as great as during the early Holocene (Bush, 2002, 2004; Li and Harrison, 2008; Rupper et al., 2008).

Many studies have attributed sharp-crested, well-preserved moraines within a kilometer of the present glacier margin to the Neoglaciation, which is younger than ~ 3-2.7 ka (e.g. Benedict, 1976; Fushimi, 1978; Derbyshire et al., 1984; Wu, 1984a; Zheng, 1997; Richards et al., 2000a; Finkel et al., 2003; Zheng, 1997; Owen et al., 2005; Jiao and Shen, 2006; Seong et al., 2009a, Meyer et al., 2009). Yet many of these studies do not have adequate numerical dating to show moraines are, indeed, Neoglacial and the extensive early Holocene advance can be easily misinterpreted as Neoglacial. Moreover, as Owen (2011) points out, the onset of the Neoglaciation in the Himalayan-Tibetan orogen may be asynchronous.

Pu (1991), for example, defines four Neoglacial advances (Neoglaciation I–IV) in western China, including Tibet, which date to ~ 9.2–7.8 ka, 6.7–5.6 ka, 3.6–2.4 ka, and 0.5–0.03 ka (~ 7000–8200, 4900–5800, 2400–3320 and 450–30 ¹⁴C years BP). Clearly, by the definition of Neoglaciation (see Denton and Karlén, 1973, Porter, 2000; Porter and Denton, 1967), the first two of these advances did not occur in the Neoglacial. Therefore, as Owen (2011) emphasizes, confused nomenclature as well as the challenges of dating moraines adds to the complexity of understanding Holocene glaciation in the Himalayan-Tibetan orogen. Yang et al. (2008) showed three main periods of glacial advance at

around 1.8–1.4 ka, 1.2–1.15 ka, and 0.6–0.08 ka in the Hengduan Mountains, the central and eastern parts of the Himalaya and the eastern Nyainquentanghlha Range of SE Tibet. This complex array of ages suggests that glaciers likely would have been responding to climate change throughout the Himalayan-Tibetan orogen in a very complicated fashion.

Glacial advances during the Little Ice Age (LIA) in the Himalayan-Tibetan orogen are not well defined. There are few definitive studies providing ages on moraines that can be attributed to the LIA. The most notable ones include studies in the Khumbu Himal that have been dated by both radiocarbon and TCN methods (Benedict, 1976; Fushimi, 1978; Finkel et al., 2003). Yi et al. (2008) collates all the published radiocarbon ages for LIA moraines from Nyainqentanglha, Karakoram, Langtang, the western slopes of Namjagbarwa Peak in the eastern Himalaya and Gonga in SE Tibet to show that the LIA in Tibet had three substages between 1.0 to 0.13 ka. Dortch et al. (2013) showed that a significant regional glacial stage is recognized across the semi-arid western end of the Himalayan-Tibetan orogen at 0.4 ± 0.1 ka (Fig. 21).

8. Historical and predicted future changes

In most areas of the Himalayan-Tibetan orogen, glaciers began to retreat after the LIA at the beginning of the 20th Century and have continued to retreat since (Mayewski and Jeschke, 1979; Mayewski et al., 1980; Ono, 1984, 1985, 1986; Ono et al., 1997; Shi and Lui, 2000; Su and Shi, 2002; Fujita et al., 1997; Ren et al., 2004; Bolch et al., 2008). Cogley (2011) provides a comprehensive glacier inventory for the Himalaya and Karakoram containing 20,812 glaciers covering an area of 43,178 km². Undertaking a

glacier-by-glacier analysis, Cogley (2011) suggests that up to about one-fifth of the glaciers present in 1985 may have disappeared. Furthermore, Cogley (2011) argues that, if mass loss were to remain constant at the average rate for 1975-2008, from 3000 to 13,000 more glaciers might disappear by 2035. Moreover, he suggests that if mass loss were to continue to accelerate as inferred for 1985-2008, only a few thousand to a few hundred glaciers might remain in 2035. Cogley (2011) acknowledges that these projections are uncertain and neglect some possibly important mitigating controls, but they demonstrate the need for more complete analyses to help define the reduction and assess its potential effects.

Su and Shi (2002) argue that, since the height of the LIA during 17th century, the mean temperature of temperate monsoonal glaciers in China has increased by 0.8°C and their area has decreased by $\sim 3900 \text{ km}^2$, an amount equivalent to 30% of the area of modern glaciers. Moreover, Su and Shi (2002) argue that if temperature rises by 2.1°C as may be likely by the year 2100, then temperate monsoonal glaciers in China are likely to decrease by 75%, an $\sim 9900 \text{ km}^2$ loss of area. Furthermore, if precipitation decreases as well, then the retreat of the glaciers will be even faster. However, Su and Shi (2002) argue that the area reduction percentage of the glaciers is unlikely to exceed 80%. Some regions have experienced accelerated glacier retreat in recent decades often resulting in the formation of impressive supraglacial lakes (Benn et al., 2001).

The rate of retreat over the past 30 years in some regions, however, has been less than during the earlier part of the 20th century (Copland et al., 2011). In some regions, certain

glaciers have begun to stabilize and/or advance during the past few years. Some glaciers have even been remarkably stable, such as Baltoro Glacier in northern Pakistan that has varied by no more than ± 200 m since the 1850s (Mayer et al., 2006). In some regions, such as the Karakoram, individual glaciers have advanced and retreated during the last century, sometimes surging (Hewitt, 1969, 2005; Barrand and Murray, 2006; Copland et al., 2011).

Predicting increasing mean annual temperatures due to human activity will cause glaciers to melt, retreat and eventually disappear is challenging. Other factors including, for example, increased deposition of black carbon on glaciers, will enhance melting. However, the large altitudinal range, and the complex topographic setting, and microclimatic variations make it difficult to quantify future temperature change and its net effect on glacier melting. In monsoon-influenced regions, predicted increased snowfall and increased cloudiness may lead to positive glacier mass balances. Consequently, some glaciers may thicken and advance. These glaciers may eventually retreat as the increase of mean annual temperature out-competes the increased glacier growth due to enhanced snowfall and cloudiness; this change may be rapid if summer snows become summer rain.

9. Future progress

Understanding of the nature, timing, and extent of Late Quaternary glaciation in the Himalayan-Tibetan orogen has advanced rapidly and significantly in the past decade or so, specifically aided by improvements in remote sensing and dating methods, but the

vastness and the political and logistical inaccessibility of much of the region still present significant challenges. Numerical dating studies suggest that, on millennial timescales, glaciation is broadly synchronous throughout the orogen, but the exact timing and extent of glaciation varies considerably within and between different mountain ranges within Highland Asia. The complex patterns of glaciation that are emerging imply that glaciation throughout the Himalayan-Tibetan orogen reflects temporal and spatial variability of forcing by the different climatic systems that dominate the region. As Owen (2009; 2013) indicated the style and timing of glaciation appears to be strongly controlled by climatic gradients, but topography may also have an important control. Correlation between regions is hindered because of large uncertainties related to the dating methods, mainly TCN and OSL methods. Reducing these uncertainties and developing stronger correlations can be achieved by improving the dating methods, and by using two or more dating techniques (for example Owen et al. 2009, 2001). A productive area of research might include numerical modeling of landform degradation to help explain the distribution of TCN ages on landforms since a post-depositional modification of landforms results can result in significant underestimates of TCN ages. In spite of these challenges, geologic error outweighs uncertainties in current dating methodologies. In addition, the lack of a standard way in which to interpret TCN results can lead to erroneous correlations or the missing of valid correlations. Thorough numerical analysis of dating results using methods such as those described by Dortch et al. (2013) is beginning to allow regional glacial stages to be recognized, forming a stronger foundation for the study of Quaternary glaciation in the Himalayan-Tibetan realm.

Owen et al. (2005, 2009, 2010, 2012) and Owen (2013) pointed out that the extent of glaciation has become increasingly restricted throughout the Late Quaternary in the more arid regions of Highland Asia, which has helped preserve very old (pre-last glacial cycle) glacial landforms. In regions very strongly influenced by the monsoon, the preservation potential of pre-Late Glacial moraines is poor. Owen et al. (2005) suggest that this may be because Late Glacial and Holocene glacial advances were the most extensive, destroying the geomorphic or sedimentary evidence of earlier glaciations, together with the intense fluvial and mass movement denudation that is characteristic of the wetter regions.

Mid-Holocene Neoglaciation, LIA, and historical glacial records testify to a progressive retreat of glaciers during the twentieth century and warn of a continued retreat during the coming century, but the pattern is complex and the glaciers may be asynchronous during these climatostratigraphic times, issues that need to be considered in more detail. Fortunately, because of the better preservation of the Holocene glacial record and because numerical dating can more readily be applied to Holocene landforms and sediment, we should target interesting times of climatic instability: the early Holocene, Neoglaciation, and specifically, the LIA. The use of other geologic proxies such as lake, aeolian and ice core records should be further utilized. Climatic and glacial modeling also has great potential to provide valuable information regarding the patterns and possible complexity of change.

In sum, the Himalayan-Tibetan orogen provides an exciting natural laboratory for the

examination of the links between climatic and glacial systems which, in turn has important implications for the geomorphology and hydrology of the high mountains and their forelands in Asia. The numerous studies that have been undertaken in this region have established a broad framework, which can be also be used to develop and identify intriguing, more specific questions about climatic-glacial-erosional linkages.

Acknowledgements

Sincere thanks to Neil Glasser and Tim Horscroft for encouraging us to write this paper and being patient with our submission time. Many thanks to three anonymous referees who provided very constructive and useful comments to help us improve our paper and to Craig Dietsch for his advice of our manuscript. This paper was presented in an early form by LAO as an invited lecture at the American Geophysical Union Annual Meeting in San Francisco in December 2012.

References

- Abramowski, U., 2004. The use of ^{10}Be surface exposure dating of erratic boulders in the reconstruction of the late Pleistocene glaciation history of mountainous regions, with examples from Nepal and Central Asia. Unpublished PhD., Universität Bayreuth, Germany, 167p.
- Abramowski, U., Bergau, A., Seebach, D., Zech, R., Glaser, B., Sosin, P., Kubik, P.W., Zech, W., 2006. Pleistocene glaciations of Central Asia: results from ^{10}Be surface exposure ages of erratic boulders from the Pamir (Tajikistan) and the Alay-Turkestan range (Kyrgyzstan). Quaternary Science Reviews 25, 1080-1096.

- 1035 Ali, S.N., Biswas, R.H., Shukla, A.D., Juyal, N., 2013. Chronology and climatic implications of Late
 1036 Quaternary glaciations in the Goriganga valley, central Himalaya, India. *Quaternary Science Reviews*
 1037 73, 59-76.
 1038
- 1039 An, Z., Porter, S.C., Kutzbach, J.E., Wu, X., Wang, S., Liu, X., Li, X., Zhou, W., 2000.
 1040 Asynchronous Holocene optimum of the East Asian monsoon. *Quaternary Science Reviews*, 19,
 1041 743-762.
 1042
- 1043 Aoki, T., Imamura, M., 1999. Reconstructing the glacial chronology based on the ^{10}Be exposure
 1044 age in the Khumbu Glacier, Eastern Nepal Himalaya. *Proceedings of the Korea-Japan/Japan-*
 1045 *Korea Geomorphological Conference*, 134–135.
 1046
- 1047 Applegate, P.J., Urban, N.M., Laabs, B.J.C., Keller, K., Alley, R.B., 2010. Modeling the
 1048 statistical distributions of cosmogenic exposure dates from moraines. *Geoscientific Model*
 1049 *Development* 3, 293-307.
 1050
- 1051 Bagla, P., 2009. No sign yet of Himalayan meltdown, Indian report finds. *Science* 326, 924-925.
 1052
- 1053 Balco, G., Stone, J.O., Lifton, N.A., Dunai T.J., 2008. A complete and easily accessible means of
 1054 calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements.
 1055 *Quaternary Geochronology* 8, 174-195.
 1056
- 1057 Balco, G., Briner, J., Finkel, R.C., Rayburn, J.A., Ridge, J.C., Schaefer, J.M., 2009, Regional
 1058 beryllium-10 production rate calibration for northeastern North America. *Quaternary*
 1059 *Geochronology* 4,. 93–107.
 1060

- 1061 Barnard, P.L., Owen, L.A., Finkel, R.C., 2004a. Style and timing of glacial and paraglacial
 1062 sedimentation in a monsoonal influenced high Himalayan environment, the upper Bhagirathi
 1063 Valley, Garhwal Himalaya. *Sedimentary Geology* 165, 199-221.
 1064
- 1065 Barnard, P.L., Owen, L.A., Finkel, R.C., Asahi, K., 2006. Landscape response to deglaciation in a
 1066 high relief, monsoon-influenced alpine environment, Langtang Himal, Nepal. *Quaternary Science*
 1067 *Reviews* 25, 2162-2176.
 1068
- 1069 Barnard, P.L., Owen, L.A., Sharma, M.C., Finkel, R.C., 2004b. Late Quaternary (Holocene)
 1070 landscape evolution of a monsoon-influenced high Himalayan valley, Gori Ganga, Nanda Devi,
 1071 NE Garhwal. *Geomorphology* 61, 91-110.
 1072
- 1073 Barrand, N. E., Murray, T., 2006. Multivariate controls on the incidence of glacier surging in the
 1074 Karakoram Himalaya. *Arctic, Antarctic, and Alpine Research* 38, 489–498.
 1075
- 1076 Bäumler R, Zech, W., Ni, A.A., Savoskul, O., 1999. Soil geographical and pedogeochemical
 1077 studies on Pleistocene and Holocen glaciation in the Northwestern Tien Shan (Uzbekistan)
 1078 *Zeitschrift fur Gletscherkunde und Glazialgeologie* 35, 147-173.
 1079
- 1080 Benedict, J.B., 1976. Khumbu glacier series, Nepal. In: Buckley, J., *Isotopes' radiocarbon*
 1081 *measurements XI Radiocarbon* 18, 177–178.
 1082
- 1083 Benn DI, Lehmkuhl F., 2000. Mass balance and equilibrium-line altitudes of glaciers in high
 1084 mountain environments. *Quaternary International* 65/66, 15–29.
 1085

- 1086 Benn, D.I., Owen, L.A., Osmaston, H.A., Seltzer G.O., Porter C.S., Mark B., 2005.
 1087 Reconstruction of equilibrium-line altitudes for tropical and sub-tropical glaciers. *Quaternary*
 1088 *International* 138/139, 8-21.
 1089
- 1090 Benn, D.I., Owen L.A., 1998. The role of the Indian summer monsoon and the mid-latitude
 1091 westerlies in Himalayan glaciation: review and speculative discussion. *Journal of the Geological*
 1092 *Society* 155, 353-363.
 1093
- 1094 Benn, D.I., Owen, L.A., 2002. Himalayan glacial sedimentary environments: a framework for
 1095 reconstructing and dating former glacial extents in high mountain regions. *Quaternary*
 1096 *International* 97/98, 3-26.
 1097
- 1098 Benn, D.I., Wiseman, S., Hands, K.A., 2001. Growth and drainage of supraglacial lakes on
 1099 debris-mantled Ngozumpa Glacier, Khumbu Himal, Nepal. *Journal of Glaciology* 47, 626-638.
 1100
- 1101 Bishop, M.P., Bush, A., Copland, L., Kamp, U., Owen, L.A., Seong, Y.B., Shroder, J.F., 2010.
 1102 Climate Change and Mountain Topographic Evolution in the Central Karakoram, Pakistan.
 1103 *Annals of Geography* 100, 1-22.
 1104
- 1105 Böhner, J. 2006. General climatic controls and topoclimatic variations in Central and High Asia.
 1106 *Boreas* 35, 279-295.
 1107
- 1108 Bolch, T., Buchroithner, M. F., Pieczonka, T., Kunert, A., 2008. Planimetric and Volumetric
 1109 Glacier Changes in the Khumbu Himalaya since 1962 Using Corona, Landsat TM and ASTER
 1110 Data. *Journal of Glaciology*, 54, 592-600.
 1111

- 1112 Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., deMenocal, P., Priore, P., Cullen, H.,
 1113 Hajdas I., Bonani, G., 1997. A pervasive millennial-scale cycle in North Atlantic Holocene and
 1114 glacial climates. *Science* 278, 1257-1266.
 1115
- 1116 Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M., Showers, W., Hoffmann, S., Lotti-
 1117 Bond, R., Hajdas, I., Bonani, G., 2001. Persistent solar influence on North Atlantic climate during
 1118 the Holocene. *Science* 294, 2130-2136.
 1119
- 1120 Bookhagen, B. & Burbank, D.W. 2006. Topography, relief, and TRMM-derived rainfall
 1121 variations along the Himalaya. *Geophysical Research Letters*, 33, L08405.
 1122
- 1123 Bookhagen, B., Thiede, R.C. & Strecker, M.R. 2005. Late Quaternary intensified monsoon
 1124 phases control landscape evolution in the northwest Himalaya. *Geology*, 33, 149–152.
 1125
- 1126 Bookhagen, B. (in review): High resolution spatiotemporal distribution of rainfall
 1127 seasonality and extreme events based on a 12-year TRMM time series, in review.
 1128
- 1129 Briner, J.P., Young, N.E., Goehring, B.M., Schaefer, J.M., 2012. Constraining Holocene ^{10}Be
 1130 production rates in Greenland. *Journal of Quaternary Science* 27, 2–6.
 1131
- 1132 Brown, E.T., Bendick, R., Bourles, D.L., Gaur, V., Molnar, P., Raisbeck, G.M., Yiou, F., 2002.
 1133 Slip rates of the Karakoram fault, Ladakh, India, determined determined using cosmic ray
 1134 exposure dating of debris flows and moraines. *Journal of Geophysical Research* 107, B9, 2192, 7-
 1135 1–7-8.
 1136

- 1137 Brozović, N., Burbank, D.W., Meigs, A.J., 1997. Climatic limits on landscape development in the
 1138 northwestern Himalaya. *Science* 276, 571-574.
 1139
- 1140 Burbank, D.W., Fort, M.B., 1985. Bedrock control on glacial limits: Examples from the Ladakh
 1141 and Zaskar Ranges, north-western Himalaya, India. *Journal of Glaciology* 31, 143–149.
 1142
- 1143 Burbank, D.W., Kang, J.C., 1991. Relative dating of Quaternary moraines, Rongbuk Valley,
 1144 Mount Everest, Tibet: Implications for an Ice Sheet on the Tibetan Plateau. *Quaternary Research*
 1145 36, 1-18.
 1146
- 1147 Bush, A.B.G., 2000. A positive climatic feedback mechanism for Himalayan glaciation.
 1148 *Quaternary International* 65/66, 3-13.
 1149
- 1150 Bush, A.B.G. 2002. A comparison of simulated monsoon circulations and snow accumulation in
 1151 Asia during the mid-Holocene and at the Last Glacial Maximum. *Global and Planetary Change*
 1152 32, 331-347.
 1153
- 1154 Bush, A.B.G., 2001. Pacific sea surface temperature forcing dominates orbital forcing of the
 1155 Early Holocene monsoon. *Quaternary Research* 55, 25-32.
 1156
- 1157 Bush, A.B.G., 2004. Modeling of late Quaternary climate over Asia; a synthesis. *Boreas* 33, 155-
 1158 163.
 1159
- 1160 Chevalier, M.-L., Ryerson, F.J., Tapponnier, P., Finkel, R.C., Van Der Woerd, J., Haibing, L.,
 1161 Qing, L., 2005. Slip-rate measurements on the Karakoram Fault may imply secular variations in
 1162 fault motion. *Science* 307, 411-414.

1163

1164 Chevalier, M.-L., Hilley, G., Tapponnier, P., Van DerWoerd, J., Liu-Zeng, J., Finkel, R.C.,
 1165 Ryerson, F.J., Li Haibing, L., Liu, X., 2011. Constraints on the late Quaternary glaciations in
 1166 Tibet from cosmogenic exposure ages of moraine surfaces. *Quaternary Science Reviews* 30, 528-
 1167 554.

1168

1169 Chinese Academy of Sciences, 1996. *Physical Geography and Natural Resources in the Mount*
 1170 *Nanjiabawa Region*. Science Press, Beijing, pp. 1–210 (in Chinese).

1171

1172 Cogley, J.G. 2011. Present and future states of Himalaya and Karakoram glaciers. *Annals of*
 1173 *Glaciology* 52, 69-73.

1174

1175 Clapperton, C., 1993. *Quaternary Geology and Geomorphology of South America*. Elsevier,
 1176 Amsterdam.

1177

1178 Colgan, P.M., Munroe, J.S., Zhou, S., 2006. Cosmogenic radionuclide evidence for the limited
 1179 extent of last glacial maximum glaciers in the Tanggula Shan of the central Tibetan Plateau.
 1180 *Quaternary Research* 65, 336-339

1181

1182 Copland, L., Sylvestre, T., Bishop, M.P., Shroder, J.F., Seong, Y.B., Owen, L.A., Bush, A., Kamp,
 1183 U., 2011. Expanded and recently increased glacier surging the Karakoram. *Arctic, Alpine and*
 1184 *Antarctic Research* 43, 503-516.

1185

1186 Cronin, V.S., Johnson, W.P., Johnson, N.M., Johnson, G.D., 1989. Chronostratigraphy of the
 1187 upper Cenozoic Bunthang sequence and possible mechanisms controlling base level in Skardu

- 1188 intermontane basin, Karakoram Himalaya, Pakistan. Geological Society of America Special
 1189 Paper 232, 295-309.
 1190
- 1191 Dainelli, G., 1922. Studi sul glaciale Spedizone Italian de Filippi nell' Himalaia, Caracorum e
 1192 Turchestan/Cinese (1913-1914). Ser II, 3: Zanichelli, Bologna, 658 pp.
 1193
- 1194 Dainelli, G. 1934. Studi sul Glaciale: Spedizone Italian de Filippi nel'Himalaia, Caracorum e
 1195 Turchestan Cinese (1913-1914). Ser. II., 3: Zanichelli, Bologna, 658 pp.
 1196
- 1197 Dainelli, G., 1935. La Series dei Terreni: Spedizione Ilaliana De Fillippi nell'Himalaia,
 1198 Caracorum e Turchestan Cinese (1913-1914). Ser. II, 2: Zanichelli, Bologna, 230 pp.
 1199
- 1200 De Terra, H., Paterson, T.T., 1939. Studies on the Ice Age in India and associated human
 1201 cultures. Carnegie Institute of Washington Publications 493, 354 pp.
 1202
- 1203 Denton, G.H., Karlén, 1973. Holocene climatic variations – their pattern and possible cause.
 1204 Quaternary Research 3, 155–205.
 1205
- 1206 Derbyshire, E., Li, J., Perrott, F.A., Xu, S., Waters, R.S., 1984. Quaternary glacial history of the
 1207 Hunza valley Karakoram Mountains, Pakistan. In International Karakoram Project, Miller K.
 1208 (ed), CambridgeUniversity Press: Cambridge; 456–495.
 1209
- 1210 Derbyshire, E., 1981. Glacier regime and glacial sediment facies: a hypothetical framework for
 1211 the Qinghai-Xizang Plateau. In: Proceedings of Symposium on Qinghai-Xizang (Tibet) Plateau,
 1212 Beijing, China. Geological and Ecological studies of Qinghai-Xizang Plateau. Vol. 2, Science
 1213 Press, Beijing, 1981, pp. 1649-1656.

1214

1215 Derbyshire, E., 1983. The Lushan Dilemma: Pleistocene glaciation south of the ChangJiang

1216 (Yangste River). *Zeitschrift für Geomorphologie* 27, 445-471.

1217

1218 Derbyshire, E., 1984. Sedimentological analysis of glacial and proglacial debris: a framework for

1219 the study of Karakoram glaciers. In: Miller, K.J. (Ed.), *The International Karakoram Project*, 1.

1220 Cambridge University Press, Cambridge, pp. 347-364.

1221

1222 Derbyshire, E., 1987. A history of the glacial stratigraphy in China. *Quaternary Science Reviews*

1223 6, 301-314.

1224

1225 Derbyshire, E., 1996. Quaternary glacial sediments, glaciation style, climate and uplift in the

1226 Karakoram and northwest Himalaya: review and speculations. *Palaeogeography,*1227 *Palaeoclimatology, Palaeoecology* 120, 147-157.

1228

1229 Derbyshire, E., Owen, L.A., 1990. Quaternary Alluvial Fans in the Karakoram Mountains. In:

1230 Rachocki, A.H., Church, M. (Eds.), *Alluvial Fans: A field approach*. John Wiley & Sons Ltd.,

1231 pp. 27-53.

1232

1233 Derbyshire, E., Owen, L.A., 1997. Quaternary glacial history of the Karakoram Mountains and

1234 Northwest Himalayas: a review. *Quaternary International* 38/39, 85-102.

1235

1236 Derbyshire, E., Shi, Y., Li, J., Zheng, B., Li, S., Wang, J., 1991. Quaternary glaciation of Tibet:

1237 the geological evidence. *Quaternary Science Reviews* 10, 485–510.

1238

- 1239 Desilets, D., Zreda, M., 2003. Spatial and temporal distribution of secondary cosmic-ray nucleon
 1240 intensities and applications to in situ cosmogenic dating. *Earth and Planetary Science Letters* 206,
 1241 21-42.
 1242
- 1243 Desilets, D., Zreda, M., 2006. Elevation dependence of cosmogenic ^{36}Cl production in Hawaiian
 1244 lava flows. *Earth and Planetary Science Letters* 246, 277–287.
 1245
- 1246 Dortch, J.M., Owen, L.A., Haneberg, W.C., Caffee, M.W., Dietsch, C., Kamp, U., 2009. Nature
 1247 and timing of mega-landslides in northern India. *Quaternary Science Reviews* 28, 1037-1056.
 1248
- 1249 Dortch, J.M., Owen, L.A., Caffee, M.W., 2010a. Quaternary glaciation in the Nubra and Shyok
 1250 valley confluence, northernmost Ladakh, India. *Quaternary Research* 74, 132-144.
 1251
- 1252 Dortch, J. M., Owen, L. A., Caffee, M. W. & Brease, P. 2010b. Late Quaternary glaciation and
 1253 equilibrium line altitude variations of the McKinley River region, central Alaska Range.
 1254 *Boreas* 39, 233–246.
 1255
- 1256 Dortch, J.M., Owen, L.A., Dietsch, C., Caffee, M.W., Bovard, K., 2011a. Episodic fluvial
 1257 incision of rivers and rock uplift in the Himalaya and Transhimalaya. *Journal of the Geological*
 1258 *Society, London* 168, 783-804.
 1259
- 1260 Dortch, J.M., Owen, L.A., Schoenbohm, L.M., Caffee, M.W., 2011b. Asymmetrical erosion and
 1261 morphological development of the central Ladakh Range, northern India. *Geomorphology* 135,
 1262 167–180.
 1263

- 1264 Dortch, J.M.*, Owen, L.A., Caffee, M.W., Kamp, U., 2011. Catastrophic partial drainage of
 1265 Pangong Tso, northern India and Tibet. *Geomorphology* 125, 109-121.
- 1266 Dortch, J.M., Owen, L.A., Caffee, M.W., 2013. Timing and climatic drivers for glaciation across
 1267 semi-arid western Himalayan-Tibetan orogen. *Quaternary Science Reviews*, in revision.
- 1268
- 1269 Drew, F., 1873. Alluvial and lacustrine deposits and glacial records of the upper Indus basin; Part
 1270 1, Alluvial deposits. *Geological Society of London Quarterly Journal* 29, 449-471.
- 1271
- 1272 Dunai, T.J., 2000. Scaling factors for production rates of in situ produced cosmogenic nuclides: a
 1273 critical reevaluation. *Earth and Planetary Science Letters* 176, 157-169.
- 1274
- 1275 Duncan, C.C., Klein, A.J., Masek, J.G, Isacks, B.L., 1998. Late Pleistocene and modern
 1276 glaciations in Central Nepal from digital elevation data and satellite imagery. *Quaternary*
 1277 *Research* 49, 241-254.
- 1278
- 1279 Ehlers, J., Gibbard, P., (eds.) 2004. Quaternary glaciations – extent and chronologies. Part III:
 1280 South America, Asia, Africa, Australia, Antarctica. *Developments in Quaternary Science* 2, 380
 1281 pp.
- 1282
- 1283 Ehlers, J., Gibbard, P., Hughes, P.D., (eds.) 2011, Quaternary Glaciations - Extent and
 1284 Chronology: a closer look. *Developments in Quaternary Science*, volume 15, Elsevier,
 1285 Amsterdam, 2nd Edition, 929-942.
- 1286
- 1287 Eyles, N., Eyles, C.H., Miall, A.D., 1983. Lithofacies types and vertical profile models, An
 1288 alternative approach to the description and environmental interpretation of glacial diamict and

- 1289 diamictite sequences. *Sedimentology* 30, 393–410.
- 1290
- 1291 Fenton, C.R., Hermanns, R.L., Blikra, L.H., Kubik, P.W., Bryant, C., Niedermann, S., Meixner,
- 1292 Anette, Goethals, M.M., 2011. Regional ^{10}Be production rate calibration for the past 12 ka
- 1293 deduced from the radiocarbon-dated Grøtlandsura and Russenes rock avalanches at 69° N,
- 1294 Norway. *Quaternary Geochronology* 6, 437–452.
- 1295
- 1296 Fielding, E., Isacks, B., Barazangi, M., Duncan, C., 1994. How flat is Tibet? *Geology* 22, 163-
- 1297 167.
- 1298
- 1299 Finkel, R.C., Owen, L.A., Barnard, P.L., Caffee, M.W., 2003. Beryllium-10 dating of Mount
- 1300 Everest moraines indicates a strong monsoonal influence and glacial synchronicity throughout the
- 1301 Himalaya. *Geology* 31, 561-564.
- 1302
- 1303 Finkel, R.C., Owen, L.A., Barnard, P.L., Caffee, M.W., 2003. Beryllium-10 dating of Mount
- 1304 Everest moraines indicates a strong monsoonal influence and glacial synchronicity throughout the
- 1305 Himalaya. *Geology* 31, 561-564.
- 1306
- 1307 Fort, M., 1986. Glacial extension and catastrophic dynamics along the Annapurna front, Nepal
- 1308 Himalaya. In: Kuhle, M. (Ed.) *Internationales über Tibet und Hochasien vom 8.-11. Oktober*
- 1309 *1985 im Geographischen Institut der Universität Göttingen*, Verlag Erich Goltze GmbH & Co.
- 1310 KG, Göttingen, 105-125.
- 1311
- 1312 Fort, M., Derbyshire, E., 1988. Some characteristics of tills in the Annapurna Range, Nepal. In:
- 1313 Chen, E. (Ed.), *Proceedings of the Second Conference on the palaeoenvironment of East Asia*

- 1314 from the Mid-Tertiary. Vol. 1 Geology, Sea level changes, palaeoclimatology and palaeobotany.
 1315 University of Hong Kong, Hong Kong, 195-214.
 1316
- 1317 Fort, M., 2003. Are high altitude, lava stream-like, debris mixtures all rock glaciers? A
 1318 perspective from the Western Himalaya. *Zeitschrift für Geomorphology* 130, 11-29.
 1319
- 1320 Frenzel, B., 1960. Die Vegetations- und Landschaftszonen Nordeurasiens während der letzten
 1321 Eiszeit und während der Postglazialen Warmezeit. Akademie der Wissenschaften und der
 1322 Literatur in Mainz, Abhandlungen der Mathematisch-Naturwissenschaftlichen Klasse 13, 937-
 1323 1099.
 1324
- 1325 Fu, P, Stroeven, A.P., Harbor, J.M., Hättestrand, C., Heyman, J., Caffee, M.W., Zhou, L. 2013.
 1326 Paleoglaciation of Shaluli Shan, southeastern Tibetan Plateau. *Quaternary Science Reviews* 64, 121-
 1327 135.
 1328
- 1329 Fuchs, M., Owen, L.A., 2008. Luminescence dating of glacial and associated sediments: review,
 1330 recommendations and future directions. *Boreas* 37, 636-659.
 1331
- 1332 Fujita, K., Nakawo, M., Fujii, Y., Paudyal, P., 1997. Changes in glaciers in Hidden Valley,
 1333 Mukut Himal, Nepal Himalayas, from 1974 to 1994. *Journal of Glaciology* 43, 583–588.
 1334
- 1335 Fushimi, H., 1977. Glaciations in the Khumbu Himal (1). *Seppyo* 39, 60–67.
 1336
- 1337 Fushimi, H., 1978. Glaciations in the Khumbu Himal (2). *Seppyo* 40, 71–77.
 1338

- 1339 Gayer, E., Lavé, J., Pik, R., France-Lanord, C., 2006. Monsoonal forcing of Holocene glacier
1340 fluctuations in Ganesh Himal (central Nepal) constrained by cosmogenic ^3He exposure ages of
1341 garnets. *Earth Planetary Science Letters* 252, 275–288.
- 1342
- 1343 Geyh, M.A., Rothlisberger, F. and Gellantly, A.F., 1985. Reliability tests of C14 dates from
1344 paleosol in glacier environments. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 21, 275-
1345 281
- 1346
- 1347 Gibbons, A.B., Megeath, J.D., Pierce, K.L., 1984. Probability of moraine survival in a succession
1348 of glacial advances. *Geology* 12, 327-330.
- 1349
- 1350 Gillespie, A., Molnar, P., 1995. Asynchronous maximum advances of mountain and continental
1351 glaciers. *Reviews of Geophysics* 33, 311-364.
- 1352 Goehring, B.M., Lohne, Ø.S., Mangerud, J., Svendsen, J.I., Gyllencreutz, R., Schaefer, J., Finkel,
1353 R., 2012. Late glacial and Holocene ^{10}Be production rates for western Norway. *Journal of*
1354 *Quaternary Science* 27, 89–96.
- 1355
- 1356 Gosse, J., Klein, J., Evenson, E.B., Lawn, B., Middleton, R., 1995. Beryllium-10 dating of the
1357 duration and retreat of the last Pinedale glacial Sequence. *Science* 228, 1329-1333.
- 1358
- 1359 Graf, A.A., Strasky, S., Zhao, Z.Z., Akçar, N., Ivy-Ochs, S., Kubik, P.W., Christl, M., Kasper,
1360 H.U., Wieler, R., Schlüchter, C., 2008. Glacier extension on the eastern Tibetan Plateau in
1361 response to MIS 2 cooling, with a contribution to ^{10}Be and ^{21}Ne methodology. In: Strasky,
1362 S., *Glacial response to global climate changes: cosmogenic nuclide chronologies from high and*
1363 *low latitudes. PhD Thesis, ETH Zürich.*

1364

1365 Grove, J.M., 2004. The Little Ice Age: ancient and modern. Volume 1, 2nd edition. Routledge,
1366 London, 432 p.

1367

1368 Guggenberger, G., Bäumler, R., Zech, W., 1998. Weathering of soils developed in eolian material
1369 overlying glacial deposits in Eastern Nepal. Soil Science 163, 325-337.

1370

1371 Haeberli, W., Bosch, H., Scherler, K., Ostrem, G., Wallen, C.C., 1989. World Glacier Inventory:
1372 Status 1988. Compiled by the World Glacier Monitoring Service, IAHS-UNEP-UNESCO:
1373 Wallingford, UK.

1374

1375 Hallet, B., Putkonen, J., 1994. Surface dating of dynamic landforms: young boulders on aging
1376 moraines. Science 265, 937–940.

1377

1378 Hedrick, K.A., Seong, Y.B., Owen, L.A., Caffee, M.C., Dietsch, C., 2011. Towards defining the
1379 transition in style and timing of Quaternary glaciation between the monsoon-influenced Greater
1380 Himalaya and the semi-arid Transhimalaya of Northern India. Quaternary International 236, 21-
1381 33.

1382

1383 Heimsath, A.M., McGlynn, R., 2008. Quantifying periglacial erosion in the Nepal high Himalaya.
1384 Geomorphology 97, 5–23.

1385

1386 Hewitt, K., 1969. Glacier surges in the Karakoram Himalaya (Central Asia). Canadian Journal of
1387 Earth Sciences 6, 1009-1018.

1388

- 1389 Hewitt, K., 1998. Catastrophic landslides and their effects on the Upper Indus streams,
 1390 Karakoram Himalaya, northern Pakistan. *Geomorphology* 26, 47-80.
 1391
- 1392 Hewitt, K., 1999. Quaternary moraines vs catastrophic avalanches in the Karakoram Himalaya,
 1393 northern Pakistan. *Quaternary Research* 51, 220-237.
 1394
- 1395 Hewitt, K., 2005. The Karakoram anomaly? Glacier Expansion and the 'Elevation Effect,'
 1396 Karakoram Himalaya. *Mountain Research and Development* 25, 332-340.
 1397
- 1398 Hewitt, K., Gosse, J., Clague, J.J., 2011. Rock avalanches and the pace of late Quaternary
 1399 development of river valleys in the Karakoram Himalaya. *Geological Society of America Bulletin*
 1400 123, 1836-1850.
 1401
- 1402 Heyman, J., Hattestrand, C., Stroeven, A.P., 2008. Glacial geomorphology of the Bayan Har
 1403 sector of the NE Tibetan plateau. *Journal of Maps*, 2008, 42-62.
 1404
- 1405 Heyman, J., Stroeven, A.P., Alexanderson, H., Hättstrand, C., Harbor, J., Li, Y.K., Caffee,
 1406 M.W., Zhou, L.P., Veres, D., Liu, F., Machiedo, M., 2009. Palaeoglaciation of Bayan Har Shan,
 1407 northeastern Tibetan Plateau: Glacial geology indicates maximum extents limited to ice cap and
 1408 ice-field scales. *Journal of Quaternary Science* 24, 710-727.
 1409
- 1410 Heyman, J., Stroeven, A.P., Caffee, M.W., Hättstrand, C., Harbor, J., Li, Y.K., Alexanderson, H.,
 1411 Zhou, L.P., Hubbard, A., 2010. Palaeoglaciology of Bayan Har Shan, NE Tibetan Plateau: the
 1412 case of a missing LGM expansion, In: Heyman, J., Palaeoglaciology of the northeastern Tibetan
 1413 Plateau. PhD thesis, Stockholm University.
 1414

- 1415 Heyman, J., Stroeven, A.P., Caffee, M.W., Hättestrand, C., Harbor, J.M., Li, Y.K., Alexanderson,
 1416 H., Zhou, L.P., Hubbard, A., 2011a. Palaeoglaciology of Bayan Har Shan, NE Tibetan Plateau:
 1417 exposure ages reveal a missing LGM expansion. *Quaternary Science Reviews* 30, 1988-2001.
 1418
- 1419 Heyman, J., Stroeven, A., Harbor, J., Caffee, M.W. 2011b. Too young or too old: evaluating 884
 1420 cosmogenic exposure dating based on an analysis of compiled boulder exposure 885 ages. *Earth
 1421 and Planetary Science Letters* 302, 71-80.
 1422
- 1423 Holmes, J.A., Street-Perrott, F.A. 1989: The Quaternary Glacial History of Kashmir, North-West
 1424 Himalaya: A Revision of de Terra and Paterson's Sequence. *Zeitschrift für Geomorphologie* 76,
 1425 195-212.
 1426
- 1427 Hövermann J, Lehmkuhl F, Pörtge K-H., 1993a. Pleistocene glaciations in Eastern and Central
 1428 Tibet- Preliminary results of the Chinese-German joint expeditions. *Zeitschrift für
 1429 Geomorphologie* 92, 85-96.
 1430
- 1431 Hövermann, J., Lehmkuhl, F., Süssenberger, H., 1993b. Neue Befunde zur Paläoklimatologie
 1432 Nordafrikas und Zentralasiens. *Abhandlungen der Braunschweigischen Wissenschaftlichen
 1433 Gesellschaft* 43, 127-150.
 1434
- 1435 Hughes, P.D., Gibbard, P.L., Woodward, J.C., 2005. Quaternary glacial records in mountain
 1436 regions: a formal stratigraphical approach. *Episodes* 28, 85-92.
 1437
- 1438 Intergovernmental Panel on Climate Change, 2007. *Climate Change 2007: Impacts, Adaptations
 1439 and Vulnerability*. Parry, M., Canziani, O., Palutikof, J., Van der Linden, P., Hanson, C. (Eds.),

- 1440 Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental
 1441 Panel on Climate Change. Cambridge University Press, Cambridge, UK, 976 pp.
- 1442 Ivy-Ochs, S., Kerschner, H., Schlüchter, C. 2007. Cosmogenic nuclides and the dating of
 1443 lateglacial and EarlyHolocene glacier variations: The Alpine perspective. *Quaternary*
 1444 *international* 164–165, 53–63.
- 1445
- 1446 Iwata, S.J., Jiao, K.Q.. 1993. Fluctuations of the Zepu Glacier in late Holocene Epoch, the eastern
 1447 Nyainqentanglha Mountains. In: Yao T.D., Ageda, Y. (Eds.) *Glaciation Climate and Environment*
 1448 *in Qinghai–Tibet Plateau*, Science Press, Beijing, 130–138.
- 1449
- 1450 Iwata S.J., Naramaa, C.K., Karma, 2002. Three Holocene and late Pleistocene glacial stages
 1451 inferred from moraines in the Lingshi and Thanza village areas, Bhutan. *Quaternary International*
 1452 *97/98*, 69–78.
- 1453
- 1454 Iwata, S., Naito, N., Narama, C., 2003. Rock glaciers and the lower limit of permafrost in the
 1455 Bhutan Himalayas. *Zeitschrift für Geomorphologie* 130, 129-143.
- 1456
- 1457 Jiao, K.Q., Iwata, S.J., 1993. The glacier variation in Kunlun pass area and southeast Tibet area
 1458 since last glaciation. In: Yao, T.D., Ageda, Y. (Eds.), *Glaciation Climate and Environment in*
 1459 *Qinghai–Tibet Plateau*, Science Press, Beijing, 120–129.
- 1460
- 1461 Jiao, K.Q., Shen, Y.P., 2006. Quaternary glaciations in Tanggulha Mountains. In: Shi, Y.F., Su,
 1462 Z., Cui, Z.J. (Eds.) *The Quaternary glaciations and environmental variations in China*. Science
 1463 *and Technology Press of Hebei Province*, Shijiazhuang, 326-356.
- 1464

- 1465 Jiao, K.Q., Zheng, B.X., 2006. Quaternary glaciations in Kunlun Mountains. In: Shi, Y.F., Su, Z.,
 1466 Cui, Z.J., (Eds.), The Quaternary Glaciations and Environmental variations in China, Science and
 1467 Technology Press
 1468 of Hebei Province, Shijiazhuang, 326–356.
 1469
- 1470 Jiao, K., Iwata, S., Yao, T., Jing, Z., Li, Z., 2005. Variation of Zepu Glacier and environmental
 1471 change in the eastern Nyainqentanglha Range since 3.2 ka BP. Journal of Glaciology and
 1472 Geocryology 27, 74–79.
 1473
- 1474 Kamp, U., Owen, L.A., 2011. Late Quaternary Glaciation of Northern Pakistan. . In: Elhers, J.,
 1475 Gibbard, P. and Hughes, P.D. (Eds.), Quaternary Glaciations - Extent and Chronology: a closer
 1476 look. Developments in Quaternary Science, volume 15, Elsevier, Amsterdam, 909-927.
- 1477 Kaplan, M.R., Hein, A.S., Hubbard, A., Lax, S.M., 2009. Can glacial erosion limit the extent of
 1478 glaciation? Geomorphology 103, 172–179.
 1479
- 1480 Kaplan, M.R., Strelin, J.A., Schaefer, J.M., Denton, G.H., Finkel, R.C., Schwartz, R., Putnam,
 1481 A.E., Vandergoes, M.J., Goehring, B.M., Travis, S.G., 2011. In-Situ cosmogenic ^{10}Be
 1482 production rate at Lago Argentino, Patagonia: Implications for late-glacial climate chronology.
 1483 Earth and Planetary Science Letters 309, 21–32.
 1484
- 1485 Klute, F., 1930. Verschiebung der Klimagebiete der letzten Eiszeit. Petermanns Mitteilungen
 1486 Ergänzungsheft 209, 166-182.
 1487
- 1488 Kong, P., Fink, D., Na, C.G., Huang, F.X., 2009a. Late Quaternary glaciation of the Tianshan,
 1489 central Asia, using cosmogenic ^{10}Be surface exposure dating. Quaternary Research 72, 229–233.

- 1490
- 1491 Kong, P., Na, C.G., Fink, D., Zhao, X.T., Xiao, W., 2009b. Moraine dam related to late
1492 Quaternary glaciation in the Yulong Mountains, southwest China, and impacts on the Jinsha
1493 River. *Quaternary Science Reviews* 28, 3224–3235
- 1494
- 1495 Koppes, M., Gillespie, A.R., Burke, R.M., Thompson, S.C., Stone, J., 2008. Late Quaternary
1496 glaciation in the Kyrgyz Tien Shan. *Quaternary Science Reviews* 27, 846-866.
- 1497
- 1498 Kuhle, M., 1985. Ein subtropisches Inlandeis als Eiszeitauslöser, Südtibet und Mt. Everest
1499 expedition 1984. *Georgia Augusta, Nachrichten aus der Universität Göttingen*, May, 1-17.
- 1500
- 1501 Kuhle, M., 1986. The upper limit of glaciation in the Himalayas. *GeoJournal* 13, 331-346.
- 1502
- 1503 Kuhle, M., 1987. The Problem of a Pleistocene Inland Glaciation of the Northeastern Qinghai-
1504 Xizang Plateau. In *Reports of the Qinghai- Xizang (Tibet) Plateau*, Hövermann J, Wang W. (eds).
1505 Beijing; 250-315.
- 1506
- 1507 Kuhle, M., 1988a. Geomorphological findings on the built-up of Pleistocene glaciation in
1508 Southern Tibet and on the problem of inland ice. *GeoJournal* 17, 457-512.
- 1509
- 1510 Kuhle, M., 1988b. Topography as a fundamental element of glacial systems. *GeoJournal* 17, 545-
1511 568.
- 1512
- 1513 Kuhle, M., 1990a. The cold deserts of high Asia (Tibet and contiguous mountains). *GeoJournal*
1514 20, 319-323.
- 1515

- 1516 Kuhle, M., 1990b. Ice marginal ramps and alluvial fans in semiarid mountains: convergence and
 1517 difference. In: Rachocki AH, Church M. (Eds.), Alluvial Fans: a field approach, John Wiley and
 1518 Sons: Chichester; 55-68.
 1519
- 1520 Kuhle, M., 1991. Observations supporting the Pleistocene inland glaciation of High Asia.
 1521 GeoJournal 25, 131-231.
 1522
- 1523 Kuhle, M., 1993. A short report of the Tibet excursion 14-A, Part of the XIII INQUA Congress
 1524 1991 in Beijing. GeoJournal 29, 426-427.
 1525
- 1526 Kuhle, M., 1995. Glacial isostatic uplift of Tibet as a consequence of a former ice sheet.
 1527 GeoJournal 37, 431-449.
 1528
- 1529 Kuhle, M., 2011. The High Glacial (Last Ice Age and Last Glacial Maximum) ice cover of high
 1530 and Central Asia, with critical review of some recent OSL and TCN dates. In: Elhers, J., Gibbard,
 1531 P. and Hughes, P.D. (Eds.), Quaternary Glaciations - Extent and Chronology: a closer look.
 1532 Developments in Quaternary Science, volume 15, Elsevier, Amsterdam, 2nd Edition, 943-965.
 1533
- 1534 Lal, D., 1991. Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and
 1535 erosion models. Earth and Planetary Science Letters 104, 429-439.
 1536
- 1537 Lasserre, C., Gaudemer, Y., Tapponnier, P., Mériaux, A.-S., Van der Woerd, J., Yuan, D.,
 1538 Ryerson, F.J., Finkel, R.C., Caffee, M.W., 2002. Fast late Pleistocene slip rate on the Leng Long
 1539 Ling segment of the Haiyuan fault, Qinghai, China. Journal of Geophysical Research 107, B11,
 1540 2276.
 1541

- 1542 Lee, S.Y., Seong, Y.B., Owen, L.A., Murari, M.K., Lim, H.S., Yoon, H.I., You, K.-C, 2013. Late
 1543 Quaternary glaciation in the Nun-Kun massif, northwestern India. *Boreas*, in press.
 1544
- 1545 Lehmkuhl, F., 1995. Geomorphologische Untersuchungen zum Klima des Holozäns und
 1546 Jungpleistozäns Osttibets. *Göttinger Geographische Abhandlungen* 102, 1–184.
 1547
- 1548 Lehmkuhl, F., 1997. Late Pleistocene, Late-glacial and Holocene glacier advances on the Tibetan
 1549 Plateau. *Quaternary International* 38/39, 77-83.
 1550
- 1551 Lehmkuhl, F., 1998. Extent and spatial distribution of Pleistocene glaciations in Eastern Tibet.
 1552 *Quaternary International* 45/46, 123-134.
 1553
- 1554 Lehmkuhl, F., Lui S., 1994. An outline of physical geography including Pleistocene glacial
 1555 landforms of Eastern Tibet (Provinces Sichuan and Qinghai). *GeoJournal* 34, 7–30.
 1556
- 1557 Lehmkuhl, F., Owen L.A., 2005. Late Quaternary glaciation of Tibet and the bordering
 1558 mountains: a review. *Boreas* 34, 87-100.
 1559
- 1560 Lehmkuhl, F., Owen, L.A., Derbyshire, E., 1998. Late Quaternary glacial history of northeastern
 1561 Tibet. *Quaternary Proceedings* 6, 121-142.
 1562
- 1563 Lehmkuhl, F., Klinge, M., Stauch, G. 2004. The extent of Late Pleistocene Glaciations in the
 1564 Altai and Khangai Mountains. – In: Ehlers, J. & P.L. Gibbard (Eds.): *Quaternary glaciations –*
 1565 *extent and chronologies, Part III: South America, Asia, Africa, Australia, Antarctica*, p.243-254.
 1566 Elsevier, Oxford .
 1567

- 1568 Leuschner, D.C., Sirocko, F., 2003. Orbital insolation forcing of the Indian monsoon – A motor
 1569 for global climate change. *Paleogeography, Paleoclimatology and Paleoecology* 197, 83-95.
 1570
- 1571 Li, Y., Harrison, S.P., 2008. Simulations of the impact of orbital forcing and ocean on the Asian
 1572 summer monsoon during the Holocene. *Global and Planetary Change* 60, 505–522.
 1573
- 1574 Li, S.J., Jiao, K.Q., 1990. Glacier variation on the south slope of west Kunlun Mountains since
 1575 30000 ^{14}C yr BP. (Chinese with English summary). *Journal of Glaciology and Geocryology* 12,
 1576 311–318.
 1577
- 1578 Li, S.J., Li, S.D., 1992. Quaternary glacial and environmental changes in the region of Hoh Xil,
 1579 Qinghai Province. (Chinese with English summary). *Journal of Glaciology and Geocryology* 14,
 1580 316–324.
 1581
- 1582 Li, J.J., Derbyshire, E., Street-Perrott, F.A., Xu, S.Y., Waters, R.S., 1984. Glacial and paraglacial
 1583 sediments of the Hunza valley, north-west Pakistan: a preliminary analysis. In: Miller, K. (Ed.),
 1584 The International Karakoram Project, Cambridge University Press: Cambridge, 496-535.
 1585
- 1586 Li, J.J., Zheng, B.X., Yang, X.J., 1986. Glacier in Tibet. Science Press, Beijing, 1-275.
 1587
- 1588 Li, B., Li, J., Cui, Z., (eds.) 1991. Quaternary glacial distribution map of Qinghai-Xizang (Tibet)
 1589 Plateau 1:3,000,000. Shi Y. (Scientific Advisor), Quaternary Glacier, and Environment Research
 1590 Center, Lanzhou University.
 1591
- 1592 Lifton, N.A., Bieber, J.W., Clem, J.M., Duldig, M.L., Evenson, P., Humble, J.E., Pyle, R., 2005.
 1593 Addressing solar modulation and long-term uncertainties in scaling secondary cosmic rays for in

- 1594 situ cosmogenic nuclide applications. *Earth and Planetary Science Letters* 239, 140–161.
- 1595
- 1596 Lisiecki, L.E., Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed
- 1597 benthic $\delta^{18}\text{O}$ records. *Paleoceanography* 20, PA1003, DOI: 10.1029/2004PA001071
- 1598
- 1599 Mann, D.H, Sletten, R.S, Reanier, R.E. 1996. Quaternary glaciation of the Rongbuk Valley,
- 1600 Tibet. *Journal of Quaternary Science* 11, 267-280.
- 1601
- 1602 Martinson, D.G., Pisias, N.G., Hays, J.D., Imbrie, J., Moore, T.C., Shackleton, N.J., 1987. Age
- 1603 dating and the orbital theory of ice ages: Development of a high resolution 0 to 300,000-year
- 1604 chronology. *Quaternary Research* 27, 1–29.
- 1605
- 1606 Mayer, C., Lambrecht, A., Belò, M., Smiraglia, C., and Diolaiuti, G., 2006: Glaciological
- 1607 characteristics of the ablation zone of Baltoro glacier, Karakoram, Pakistan. *Annals of Glaciology*
- 1608 43: 123–131.
- 1609
- 1610 Mayewski, P.A., Jeschke, P.A., 1979. Himalayan and Trans-Himalayan glacier fluctuations since
- 1611 AD 1812. *Arctic and Alpine Research* 11, 267–287.
- 1612
- 1613 Mayewski, P.A., Pregent, G.P., Jeschke, P.A., Ahmad, N., 1980. Himalayan and Trans
- 1614 Himalayan glacier fluctuations and the South Asian Monsoon record. *Arctic and Alpine Research*
- 1615 12, 171–182.
- 1616
- 1617 Mayewski, P.A., Rohling, E.E., Stager, C.J., Karlén, W., Maasch, A., Meeker, L.D., Meyerson,
- 1618 E.A., Gasse, F., Van Kreveld, S., Holmgren, K., Lee-Thorp, J., Rosqvist, G., Rack, F.,

- 1619 Staubwasser, M., Schneider, R.R., Steig, E.J., 2004. Holocene climate variability. *Quaternary*
1620 *Research* 62, 243-255.
- 1621
- 1622 Mériaux, A.-S., Ryerson, F.J., Tapponnier, P., Van der Woerd, J., Finkel, R.C., Xu, X., Xu, Z.,
1623 Caffee, M.W. 2004. Rapid slip along the central Altyn Tagh Fault: morphochronologic evidence
1624 from Cherchen He and Sulamu Tagh. *Journal of Geophysical Research* 109, B06401.
- 1625
- 1626 Meyer, M.C., Hofmann, Ch.-Ch., Gemmell, A.M.D., Haslinger, E., Häusler, H., Wangda, D.,
1627 2009. Holocene glacier fluctuations and migration of Neolithic yak pastoralists into the high
1628 valleys of northwest Bhutan. *Quaternary Science Reviews* 28, 1217-1237.
- 1629
- 1630 Morén, B., Heyman, J., Stroeve, A.P., 2011. Glacial geomorphology of the central Tibetan
1631 Plateau. *Journal of Maps* 2011, 115-125.
- 1632
- 1633 Muller, F., 1980. Present and late Pleistocene equilibrium line altitudes in the Mt. Everest
1634 region—An application of the glacier inventory. *World Glacier Inventory* 126, 75–94.
- 1635
- 1636 Murari, M.K., Owen, L.A., Dortch, J.M., Caffee, M.W., Dietsch, C., Fuchs, M., Haneberg, W.,
1637 Sharma, M.C., Townsend-Small, A. 2013. Timing and climatic drivers for glaciation across
1638 monsoon-influenced regions of the Himalayan-Tibetan orogen. *Quaternary Science Reviews*,
1639 in review.
- 1640
- 1641 Narama, C., 2002. Late Holocene variation of the Raigorodskogo Glacier and climate change in
1642 the Pamir-Alai, central Asia. *Catena* 48, 21-37.
- 1643

- 1644 Narama, C., Kondo, R., Tsukamoto, S., Kajiura, T., Ormukov, C., Abdrakhmatov, K., 2007. OSL
1645 dating of glacial deposits during the Last Glacial in the Terskey-Alatoo Range, Kyrgyz Republic.
1646 Quaternary Geochronology 2, 249–254.
1647
- 1648 Narama, C., Kondo, R., Tsukamoto, S., Kajiura, T., Duishonakunov, M., Abdrakhmatov, K.,
1649 2009. Timing of glacier expansion during the Last Glacial in the inner Tien Shan, Yrgyz Republic
1650 by OSL dating. Quaternary International 199, 147–156.
1651
- 1652 NGRIP members, 2004. High-resolution record of Northern Hemisphere climate extending into
1653 the last interglacial period. Nature 431, 147–151.
1654
- 1655 Norin, E., 1925. Preliminary notes on the Late Quaternary glaciation of the North Western
1656 Himalaya. Geografiska Annaler 7, 165-194.
1657
- 1658 Norton, K.P., Abbühl, L.M., Schlunegger, F., 2010. Glacial conditioning as an erosional driving
1659 force in the Central Alps. Geology 38, 655-658.
1660
- 1661 Ono, Y., 1984. Annual moraine ridges and recent fluctuation of Yala (Dakpatsen) Glacier,
1662 Langtang Himal. In: Higuchi, K. (Ed.) Glacier studies in Langtang Valley. Data Center for
1663 Glacier Research, Japan Society for Snow and Ice 2, 73-83.
1664
- 1665 Ono, Y., 1985. Recent fluctuations of the Yala (Dakpatsen) Glacier, Langtang Himal,
1666 reconstructed from Annual moraine ridges. Zeitschrift für Gletscherkunde und Glaziaklgeologie
1667 21, 251-258.
1668

- 1669 Ono, Y., 1986. Glacial fluctuations in the Langtang valley Nepal Himalaya. In: Kuhle, M. (Ed.)
 1670 Internationales über Tibet und Hochasien vom 8.-11. Oktober 1985 im Geographischen Institut
 1671 der Universität Göttingen, Verlag Erich Goltze GmbH & Co. KG, Göttingen, 31-38.
 1672
- 1673 Ono, Y., Liu, D., Zhao, Y., 1997. Paleoenvironments of Tibetan Plateau from glacial fluctuations
 1674 in the northern foot of the West Kunlun Mountains. *Journal of Geography* 106, 184-198 (In
 1675 Japanese).
- 1676 Osmaston, H., 1994. The geology, geomorphology and Quaternary history of Zangskar. In:
 1677 Crook, J., Osmaston, H. (Eds.) *Himalayan Buddhist Villages*. University of Bristol Press, UK, 1-
 1678 36.
 1679
- 1680 Owen, L.A., 1988. Wet-sediment deformation of Quaternary and Recent sediments in the Skardu
 1681 Basin, Karakoram Mountains, Pakistan. In: Croots, D. (Ed.) *Glaciotectonics*, A. A. Balkema,
 1682 Rotterdam, pp. 123-147.
 1683
- 1684 Owen, L.A., 1991. Mass movement deposits in the Karakoram Mountains: their sedimentary
 1685 characteristics, recognition and role in Karakoram landform evolution. *Zeitschrift für*
 1686 *Geomorphologie* 35, 4, 401-424.
 1687
- 1688 Owen, L.A., 2009. Latest Pleistocene and Holocene glacier fluctuations in the Himalaya and
 1689 Tibet. *Quaternary Science Reviews* 28, 2150-2164.
 1690
- 1691 Owen, L.A., 2010. Landscape development of the Himalayan-Tibetan orogen: a review. *Special*
 1692 *Publication of the Geological Society of London* 338, 389-407.
 1693

- 1694 Owen, L.A., 2011. Quaternary glaciation of Northern India. In: Elhers, J., Gibbard, P. and
 1695 Hughes, P.D. (Eds.), Quaternary Glaciations - Extent and Chronology: a closer look.
 1696 Developments in Quaternary Science, volume 15, Elsevier, Amsterdam, 2nd Edition, 929-942.
 1697
- 1698 Owen, L.A., 2013. Late Quaternary glaciations in Highland Asia. In: Scott E.A (Ed.)
 1699 Encyclopedia of Quaternary Science, Vol. 2, pp. 236-244. Elsevier, Amsterdam.
 1700
- 1701 Owen, L.A., Benn, D.I., 2005. Equilibrium-line altitudes of the Last Glacial Maximum for the
 1702 Himalaya and Tibet: an assessment and evaluation of results. Quaternary International 138/139,
 1703 55-78.
 1704
- 1705 Owen, L.A., Derbyshire, E., 1988. Glacially deformed diamictos in the Karakoram Mountains,
 1706 Northern Pakistan. In: Croots, D. (Ed.) Glaciotectonics, A. A. Balkema, Rotterdam, pp. 149-176.
 1707
- 1708 Owen, L.A., Derbyshire, E., 1989. The Karakoram glacial depositional system. Zeitschrift für
 1709 Geomorphologie 76, 33-74.
 1710
- 1711 Owen, L.A., England, J., 1998. Observations on rock glaciers in the Himalayas and Karakoram
 1712 Mountains of northern Pakistan and India. Geomorphology, 26, 199-213.
 1713
- 1714 Owen, L.A., Benn, D.I., Derbyshire, E., Evans, D.J.A., Mitchell, W.A., Thompson, D.,
 1715 Richardson, S., Lloyd, M., Holden, C., 1995. The geomorphology and landscape evolution of the
 1716 Lahul Himalaya, Northern India. Zeitschrift für Geomorphologie 39, 145-174.
 1717

- 1718 Owen, L.A., Derbyshire, E., Richardson, S., Benn, D.I., Evans, D.J.A., Mitchell, W.A., 1996. The
 1719 Quaternary glacial history of the Lahul Himalaya, Northern India. *Journal of Quaternary Science*
 1720 11, 25-42.
 1721
- 1722 Owen, L.A., Mitchell, W., Bailey, R.M., Coxon, P., Rhodes, E., 1997. Style and timing of
 1723 Glaciation in the Lahul Himalaya, northern India: a framework for reconstructing late Quaternary
 1724 palaeoclimatic change in the western Himalayas. *Journal of Quaternary Science* 12, 83-109.
 1725
- 1726 Owen, L.A., Scott, C.H., Derbyshire, E., 2000. The Quaternary glacial history of Nanga Parbat.
 1727 *Quaternary International* 65/66, 63-79.
 1728
- 1729 Owen, L.A., White, B., Rendell, H., Derbyshire, E., 1992. Loessic silts in the western Himalayas:
 1730 their sedimentology, genesis and age. *Catena* 19, 493-509.
 1731
- 1732 Owen, L.A., Gualtieri, L., Finkel, R.C., Caffee, M.W., Benn, D.I., Sharma, M.C., 2001.
 1733 Cosmogenic radionuclide dating of glacial landforms in the Lahul Himalaya, Northern India:
 1734 defining the timing of Late Quaternary glaciation. *Journal of Quaternary Science* 16, 555-563.
 1735
- 1736 Owen, L.A., Finkel, R.C., Caffee, M.W., Gualtieri, L., 2002a. Timing of multiple glaciations
 1737 during the Late Quaternary in the Hunza Valley, Karakoram Mountains, Northern Pakistan:
 1738 defined by cosmogenic radionuclide dating of moraines. *Geological Society of America Bulletin*
 1739 114, 593-604.
 1740
- 1741 Owen, L.A., Kamp, U., Spencer, J.Q., Haserodt, K., 2002b. Timing and style of Late Quaternary
 1742 glaciation in the eastern Hindu Kush, Chitral, northern Pakistan: a review and revision of the

1743 glacial chronology based on new optically stimulated luminescence dating. *Quaternary*
 1744 *International* 97-98, 41-56.
 1745
 1746 Owen, L.A., Finkel, R.C., Caffee, M.W., 2002d. A note on the extent of glaciation throughout the
 1747 Himalaya during the global Last Glacial Maximum. *Quaternary Science Reviews* 21, 147-157.
 1748
 1749 Owen, L.A., Finkel, R.C., Ma, H., Spencer, J.Q., Derbyshire, E., Barnard, P.L., Caffee, M.W.,
 1750 2003a. Timing and style of Late Quaternary glaciations in NE Tibet. *Geological Society of*
 1751 *America Bulletin*, 11, 1356-1364.
 1752
 1753 Owen, L.A., Ma Haizhou, Derbyshire, E., Spencer, J.Q., Barnard, P.L., Zeng Yong Nian, Finkel,
 1754 R.C., Caffee, M.W., 2003b. The timing and style of Late Quaternary glaciation in the La Ji
 1755 Mountains, NE Tibet: evidence for restricted glaciation during the latter part of the Last Glacial.
 1756 *Zeitschrift für Geomorphologie*, 130, 263-276.
 1757
 1758 Owen, L.A., Spencer, J.Q., Ma, H., Barnard, P.L., Derbyshire, E., Finkel, R.C., Caffee, M.W.,
 1759 Zeng Yong Nian, 2003c. Timing of Late Quaternary glaciation along the southwestern slopes of
 1760 the Qilian Shan. *Boreas* 32, 281-291.
 1761
 1762 Owen, L.A., Finkel, R.C., Barnard, P.L., Ma, H., Asahi, K., Caffee, M.W., Derbyshire, E., 2005.
 1763 Climatic and topographic controls on the style and timing of Late Quaternary glaciation
 1764 throughout Tibet and the Himalaya defined by ^{10}Be cosmogenic radionuclide surface exposure
 1765 dating. *Quaternary Science Reviews* 24, 1391-1411.
 1766

- 1767 Owen, L.A., Caffee, M., Bovard, K., Finkel, R.C., Sharma, M., 2006a. Terrestrial cosmogenic
 1768 surface exposure dating of the oldest glacial successions in the Himalayan orogen. Geological
 1769 Society of America Bulletin 118, 383-392.
 1770
- 1771 Owen, L.A., Finkel, R. C., Ma Haizhou, Barnard, P.L., 2006b. Late Quaternary landscape
 1772 evolution in the Kunlun Mountains and Qaidam Basin, Northern Tibet: a framework for
 1773 examining the links between glaciation, lake level changes and alluvial fan formation. Quaternary
 1774 International 154-155, 73-86.
 1775
- 1776 Owen, L.A., Caffee, M.W., Finkel, R.C., Seong, B.S., 2008a. Quaternary glaciation of the
 1777 Himalayan–Tibetan orogen. Journal of Quaternary Science 23, 513–532.
 1778
- 1779 Owen, L.A., Kamp, U., Khattak, G.A., Harp, E.L., Keefer, D.K., Bauer, M.A., 2008b. Landslides
 1780 triggered by the October 8, 2005, Kashmir Earthquake. Geomorphology 94, 1-9.
 1781
- 1782 Owen, L.A., Robinson, R., Benn, D.I., Finkel, R.C., Davis, N.K., Yi, C., Putkonen, J., Li, D.,
 1783 Murray, A.S., 2009. Quaternary glaciation of Mount Everest. Quaternary Science Reviews 28,
 1784 1412-1433.
 1785
- 1786 Owen, L.A., Yi, C., Finkel, R.C., Davis, N., 2010. Quaternary glaciation of Gurla Mandata
 1787 (Naimon'anyi). Quaternary Science Reviews 29, 1817-1830.
 1788
- 1789 Owen, L.A., Chen, J., Hedrick, K.A., Caffee, M.W., Robinson, A., Schoenbohm, L.M., Zhaode,
 1790 Y., Li, W., Imrecke, D., Liu, J., 2012. Quaternary glaciation of the Tashkurgan Valley, Southeast
 1791 Pamir. Quaternary Science Reviews 47, 56-72.
 1792

- 1793 Pendersen, V.K., Egholm, D.L., 2013. Glaciations in response to climate variations
 1794 preconditioned by evolving topography. *Nature* 493, 206–210.
 1795
- 1796 Phillips, W.M., Sloan, V.F., Shroder, J.F. Jr, Sharma, P., Clarke, M.L., Rendell, H.M., 2000.
 1797 Asynchronous glaciation at Nanga Parbat, northwestern Himalaya Mountains, Pakistan. *Geology*
 1798 28, 431–434.
 1799
- 1800 Porter, S.C., 1970. Quaternary glacial record in the Swat Kohistan, West Pakistan. *Geological*
 1801 *Society of America Bulletin* 81, 1421–1446.
 1802
- 1803 Porter, S.C., 2000. Onset of neoglaciation in the Southern Hemisphere. *Journal of Quaternary*
 1804 *Science* 15, 395–408.
 1805
- 1806 Porter, S.C., Denton, G.H., 1967. Chronology of neoglaciation in the North American cordillera.
 1807 *American Journal of Science* 265, 177–210.
 1808
- 1809 Pratt-Sitaula, B., 2005. Glaciers, climate, and topography in the Nepalese Himalaya. PhD Thesis,
 1810 University of California, Santa Barbara.
 1811
- 1812 Pratt-Sitaula, B., Burbank, D.W., Heimsath, A.M., Humphrey, N.F., Oskin, M., Putkonen, J.,
 1813 2011. Topographic control of asynchronous glacial advances: a case study from Annapurna,
 1814 Nepal. *Geophysical Research Letters*, 38, L245092, doi:10.1029/2011GL049940.
 1815
- 1816 Prell, W. L. Kutzbach, J. F. 1992. Sensitivity of the Indian monsoon to forcing parameters and
 1817 implications for its evolution. *Nature* 360, 647–652.
 1818

- 1819 Pu, Q., 1991. Quaternary glaciers in China. In: Zhang, Z., Shao, S., Tong, G., Cao, J. (Eds.), The
 1820 Quaternary of China. China Ocean Press, Beijing, 240-273 pp.
 1821
- 1822 Putkonen, J., Swanson, T., 2003. Accuracy of cosmogenic ages for moraines. Quaternary
 1823 Research 59, 255-261.
 1824
- 1825 Putkonen, J., O'Neal, M.A., 2006. Degradation of unconsolidated Quaternary landforms in the
 1826 western North America. Geomorphology 75, 408-419.
 1827
- 1828 Putkonen, J., Connolly, J., Orloff, T., 2008. Landscape evolution degrades the geologic signature
 1829 of past glaciations. Geomorphology 97, 208-217.
 1830
- 1831 Putnam, A., Schaefer, J.M., Barrell, D.J.A., vandergoes, M., denton, G.H., Kaplan, M.R., Finkel,
 1832 R.C., Schwartz, R., Goehring, B.M., Kelley, S.E., 2010. In situ cosmogenic ^{10}Be production-rate
 1833 calibration from the Southern Alps, New Zealand. Quaternary Geochronology 5, 392–409.
 1834
- 1835 Reimer, P.J., Baillie, M.G.L., Bard, E., Bayliss, A., Beck, J.W., Bertrand, C.J.H., Blackwell, P.G.,
 1836 Buck, C.E., Burr, G.S., Cutler, K.B., Damon, P.E., Edwards, R.L., Fairbanks, R.G., Friedrich, M.,
 1837 Guilderson, T.P., Hogg, A.G., Hughen, K.A., Kromer, B., McCormac, G., Manning, S., Ramsey,
 1838 C.B. Reimer, R.W., Remmele, S., Southon, J.R. Stuiver, M., Talamo, S., Taylor, F.W., van der
 1839 Plicht, J., Weyhenmeyer, C.E. 2004. INTCAL04 terrestrial radiocarbon age calibration, 0–26 Cal
 1840 Kyr BP. Radiocarbon 46, 1029-1058.
 1841
- 1842 Ren, J., Qin, D., Kang, S., Hou, S., Pu, J., Jing, Z., 2004. Glacier variations and climate warming
 1843 and drying in the central Himalayas. Chinese Science Bulletin, 49, 65–69.
 1844

- 1845 Richards, B.W.M., 1999. Paleoclimate of South Asia over the last 80 ka: luminescence ages of
 1846 sediment from former glaciations in Nepal and Pakistan. Unpublished PhD thesis, Royal
 1847 Holloway, University of London, UK.
 1848
- 1849 Richards, B.W.M., 2000. Luminescence dating of Quaternary sediments in the Himalaya and
 1850 High Asia: a practical guide to its use and limitations for constraining the timing of glaciation.
 1851 Quaternary International 65/66, 49–61.
 1852
- 1853 Richards, B.W.M., Owen, L.A., Rhodes, E.J., 2000a. Timing of Late Quaternary glaciations in
 1854 the Himalayas of northern Pakistan. Journal of Quaternary Science 15, 283-297.
 1855
- 1856 Richards, B.W.M., Benn, D., Owen, L.A., Rhodes, E.J., Spencer, J.Q., 2000b. Timing of Late
 1857 Quaternary glaciations south of Mount Everest in the Khumbu Himal, Nepal. Geological Society
 1858 of America Bulletin 112, 1621-1632.
 1859
- 1860 Röhringer, I., Zech, R., Abramowski, U., Sosin, P., Aldahan, A., Kubik, P.W., Zoller, L., Zech,
 1861 W., 2012. The late Pleistocene glaciation in the Bogchigir Valleys (Pamir, Tajikistan) based on
 1862 ¹⁰Be surface exposure dating. Quaternary Research 78, 590-597.
 1863
- 1864 Röthlisberger, F., Geyh, M.A., 1985a. Gletscherschwankungen der letzten 10.000 Jahre – Ein
 1865 Vergleich zwischen Nord- und Südhemisphäre (Alpen, Himalaya, Alaska, Südamerika,
 1866 Neuseeland). Verlag Sauerländer, Aarau.
 1867
- 1868 Röthlisberger, F., Geyh, M., 1985b. Glacier Variations in Himalayas and Karakoram. Zeitschrift
 1869 für Gletscherkunde und Glazialgeologie 21, 237-249.
 1870

- 1871 Rose, J., Menzies, J., 1995. Glacial stratigraphy, in Menzies, J., ed., Past glacial environments:
 1872 Sediments, forms and techniques: Oxford, Butterworth Heinemann, p. 253–284.
 1873
- 1874 Ruddiman, W. F., Kutzbach, J. E., 1991. Plateau uplift and climate change. *Scientific American*
 1875 264, 42-51.
 1876
- 1877 Rupper, S., Roe, G., Gillespie, A., 2008. Spatial patterns of Holocene glacier advance and retreat
 1878 in central Asia. *Quaternary Research*, in press.
 1879
- 1880 Rutter, N.W., 1995. Problematic Ice Sheets. *Quaternary International* 28, 19-37.
 1881
- 1882 Schäfer, J.M., Tschudi, S., Zhao, Z., Wu, X., Ivy-Ochs, S., Wieler, R., Baur, H., Kubik, P.W.,
 1883 Schluchter, C., 2002. The limited influence of glaciations in Tibet on global climate over the past
 1884 170000 yr. *Earth and Planetary Science Letters* 194, 287-297.
 1885
- 1886 Schäfer, J.M., 2000. Reconstruction of landscape evolution and continental paleoglaciations using
 1887 in-situ cosmogenic nuclides: Examples from Antarctica and the Tibetan Plateau. Diss. ETH
 1888 Zürich, Der Andere Verlag, Osnabrück.
 1889
- 1890 Schaefer, J.M., Oberholzer, P., Zhao, Z.Z., Ivy-Ochs, S., Wieler, R., Baur, H., Kubik, P.W.,
 1891 Schluchter, C., 2008. Cosmogenic beryllium-10 and neon-21 dating of late Pleistocene
 1892 glaciations in Nyalam, monsoonal Himalayas. *Quaternary Science Reviews* 27, 295–311.
 1893
- 1894 Scherler, D., Bookhagen, B., Strecker, M.R., von Blanckenburg, F., Rood, D., 2010. Timing and
 1895 extent of late Quaternary glaciation in the western Himalaya constrained by ^{10}Be moraine dating
 1896 in Garhwal, India. *Quaternary Science Reviews* 29, 815–831.

1897

1898 Searle, M.P., Richard, J.P., 2007. Relationships between right-lateral shear along the Karakoram
1899 fault and metamorphism, magmatism, exhumation and uplift: Evidence from the K2–
1900 Gasherbrum, Pangong Ranges, north Pakistan and Ladakh. *Journal of the Geological Society*,
1901 London, 164, 439–450.

1902

1903 Seong, Y.B., Owen, L.A., Bishop, M.P., Bush, A., Clendon, P., Copland, P., Finkel, R.C., Kamp,
1904 U., Shroder, J.F., 2007. Quaternary glacial history of the Central Karakoram. *Quaternary Science*
1905 *Reviews* 26, 3384–3405.

1906

1907 Seong, Y.B., Owen, L.A., Bishop, M.P., Bush, A., Clendon, P., Copland, P., Finkel, R.C., Kamp,
1908 U., Shroder, J.F., 2008. Reply to comments by Matthias Kuhle on Seong, Y.B., Owen, L.A.,
1909 Bishop, M.P., Bush, A., Clendon, P., Copland, P., Finkel, R.C., Kamp, U., Shroder, J.F., 2007.
1910 Quaternary Glacial History of the Central Karakoram. *Quaternary Science Reviews* 27, 1656–
1911 1658.

1912

1913 Seong, Y.B., Owen, L.A., Yi, C., Finkel, R.C., 2009a. Quaternary glaciation of Muztag Ata and
1914 Kongur Shan: evidence for glacier response to rapid climate changes throughout the Late Glacial
1915 and Holocene in westernmost Tibet. *Geological Society of America, Bulletin* 121, 348–365.

1916

1917 Seong, Y.B., Bishop, M.P., Bush, A., Clendon, P., Copland, P., Finkel, R., Kamp, U., Owen,
1918 L.A., Shroder, J.F., 2009b. Landforms and landscape evolution in the Skardu, Shigar and Braldu
1919 Valleys, Central Karakoram. *Geomorphology* 103, 251–267.

1920

- 1921 Seong, Y.B., Owen, L.A., Yi, C., Finkel, R.C., Schoenbohm, L., 2009c. Geomorphology of
 1922 anomalously high glaciated mountains at the northwestern end of Tibet: Muztag Ata and Kongur
 1923 Shan. *Geomorphology* 103, 227-250.
 1924
- 1925 Seong, Y.B., Owen, L.A., Caffee, M.W., Kamp, U., Bishop, M.P., Bush, A., Copland, L. Shroder,
 1926 J.F. 2009d Rates of basin-wide rockwall retreat in the K2 region of the Central Karakoram
 1927 defined by terrestrial cosmogenic nuclide ^{10}Be . *Geomorphology* 107, 254-262.
 1928
- 1929 Sharma, M.C., Owen, L.A., 1996. Quaternary glacial history of NW Garhwal Himalayas.
 1930 *Quaternary Science Reviews* 15, 335-365.
 1931
- 1932 Shi, Y., Ren, B., Wang, J., Derbyshire, E., 1986. Quaternary Glaciation in China. *Quaternary*
 1933 *Science Reviews* 5, 503-510.
 1934
- 1935 Shi, Y., Zheng, B., Li, S., 1992. Last glaciation and maximum glaciation in the Qinghai-Xizang
 1936 (Tibet) Plateau: A controversy to M.Kuhle's ice sheet hypothesis. *Zeitschrift für Geomorphologie*
 1937 84, 19-35.
 1938
- 1939 Shi, Y. Lui, S., 2000. Estimation on the response of glaciers in China to global warming in the
 1940 21st century. *Chinese Science Bulletin*, 45, 668-672.
 1941
- 1942 Shi, Y., 1992. Glaciers and glacial geomorphology in China. *Zeitschrift für Geomorphologie* 86,
 1943 19-35.
 1944

- 1945 Shiraiwa, T., Watanabe, T., 1991. Late Quaternary glacial fluctuations in the Langtang Valley,
 1946 Nepal Himalaya, reconstructed by relative dating methods. *Arctic and Alpine Research* 23, 404-
 1947 416.
 1948
- 1949 Shroder, J., Owen, L.A., Seong, Y.B.*, Bishop, M.P., Bush, B., Caffee, M.W., Finkel, R.C.,
 1950 Kamp, U., 2011. The role of mass movement on landscape evolution in the Central Karakoram:
 1951 Discussion and speculation. *Quaternary International* 236, 34-47.
 1952
- 1953 Spencer, J.Q., Owen, L.A., 2004. Optically stimulated luminescence dating of Late Quaternary
 1954 glaciogenic sediments in the upper Hunza valley: validating the timing of glaciation and assessing
 1955 dating methods. *Quaternary Sciences Reviews* 23, 175-191.
 1956
- 1957 Strasky, S., Graf, A.A., Zhao, Z.Z., Kubik, P.W., Baur, H., Schluchter, C., Wieler, R., 2009. Late
 1958 glacial ice advances in southeast Tibet. *Journal of Asian Earth Science* 34, 458–465.
 1959
- 1960 Stone, J.O., 2000. Air pressure and cosmogenic isotope production. *Journal of Geophysical*
 1961 *Research* 105, 23753–23759.
 1962
- 1963 Su, Z., Shi, Y., 2002. Response of monsoonal temperate glaciers to global warming since the
 1964 Little Ice Age. *Quaternary International* 97-98, 123-132.
 1965
- 1966 Su, Z., Shi, Y., Zheng, B., 2002. Quaternary glacial remains on the Gongga Mountain and the
 1967 division of glacial period. *Advance in Earth Sciences* 17, 647–2002.
 1968
- 1969 Taylor, P.J., Mitchell, W.A., 2000. Late Quaternary glacial history of the Zaskar Range, North-
 1970 west Indian Himalaya. *Quaternary International* 65/66, 81-100.

- 1971
- 1972 Thompson, L.G., 2000. Ice core evidence for climate change in the Tropics: Implications for our
- 1973 future. *Quaternary Science Reviews* 19, 19-35.
- 1974
- 1975 Thompson, L.G., Mosley-Thompson, E., Davis, M.E., Bolzan, J.F., Dai, J., Yao, T., Gundestrup,
- 1976 N., Wu, X., Klein, L., Xie, Z., 1989. Holocene–Late Pleistocene climatic ice core records from
- 1977 Qinghai-Tibetan Plateau. *Science* 246, 474-477.
- 1978
- 1979 Thompson, L.G., Yao, T., Davis, M.E., Henderson, K.A., Mosley-Thompson, E., Lin, P.-N.,
- 1980 Beer, J., Synal, H.A., Cole-Dai, J., Bolzan, J.F., 1997. Tropical climate instability: the Last
- 1981 Glacial Cycle from a Qinghai–Tibetan ice core. *Science* 276, 1821–1825.
- 1982
- 1983 Thompson, L.G., Yao, T., Mosley-Thompson, E., Davis, M.E., Henderson, K.A., Lin, P.-N., 2000.
- 1984 A high-resolution millennial record of the South Asian monsoon from Himalayan ice cores.
- 1985 *Science* 289, 1916-1919.
- 1986
- 1987 Trinkler, E., 1930. The Ice-Age on the Tibetan Plateau and in the adjacent region. *Geography*
- 1988 *Journal* 75, 225-232.
- 1989
- 1990 Tschudi, S., Schäfer, J.M., Zhao, Z.Z., Wu, X.H., Ivy-Ochs, S., Kubik, P.W., Schlüchter, C.,
- 1991 2003. Glacial advances in Tibet during the Younger Dryas? Evidence from cosmogenic ^{10}Be ,
- 1992 ^{26}Al , and ^{21}Ne . *Journal of Asian Earth Science* 22, 301–306.
- 1993
- 1994 Tsukamoto, S., Asahi, K., Watanabe, T., Kondo, R., Rink, W.J., 2002. Timing of past glaciation
- 1995 in Kanchenjunga Himal, Nepal by optically stimulated luminescence dating of tills. *Quaternary*
- 1996 *International* 97/98, 57-68.

- 1997
- 1998 Wang, F., Fan, C.Y., 1987. Climatic changes in the Qinghai-Xizang (Tibetan) region of China
- 1999 during the Holocene. *Quaternary Research* 28, 50–60.
- 2000
- 2001 Wang, J., Raisbeck, G., Xu, X.B., Yiou, F., Bai, S.B., 2006. In situ cosmogenic ^{10}Be dating of the
- 2002 Quaternary glaciations in the southern Shaluli Mountain on the southeastern Tibetan Plateau.
- 2003 *Science in China Series D: Earth Sciences* 49, 1291–1298.
- 2004
- 2005 Wang, J., Kassab, C., Harbor, J.M., Caffee, M.W., Cui, H., Zhang, G., 2013. Cosmogenic nuclide constraints
- 2006 on late Quaternary glacial chronology on the Dalijia Shan, northeastern Tibetan Plateau. *Quaternary*
- 2007 *Research* 79, 439-451.
- 2008
- 2009 Waragai, T. 2005. Holocene calcrete crust deposits on the moraine of Batura Glacier, northern
- 2010 Pakistan. *The Island Arc* 14, 368-377.
- 2011
- 2012 Waragai, T., 1998. Effects of rock surface temperature on exfoliation, rock varnish, and lichens
- 2013 on a boulder in the Hunza Valley, Karakoram Mountains, Pakistan. *Arctic and Alpine Research*
- 2014 30, 184-192.
- 2015
- 2016 Watanabe, T., Shiraiwa, T., Ono, Y. 1989. Distribution of periglacial landforms in the Langtang
- 2017 Valley, Nepal Himalaya. *Bulletin of Glacier Research* 7, 209-220.
- 2018
- 2019 Willett, S.D., 2010. Erosion on a line. *Tectonophysics* 484, 168-180.
- 2020
- 2021 Williams, Van S., 1983, Present and former equilibrium-line altitudes near Mount Everest, Nepal

- 2022 and Tibet: Arctic and Alpine Research, v. 15, p. 201–211.
- 2023
- 2024 Wu, G.H., 1984a. The preliminary observation of Neoglacial in the Qilian Mountains. (In
- 2025 Chinese with English summary). Journal of Glaciology and Geocryology 6, 53-60.
- 2026
- 2027 Wu, G.H., 1984b. The Quaternary glaciation questions in Qilian mountains. In: Memoirs of
- 2028 Lanzhou Institute of Glaciology and Geocryology, Chinese Academy of Science (No. 5): Glacier
- 2029 Changes and Utilizations in Qilian Mountains. Lanzhou Institute of Glaciology and Geocryology,
- 2030 Chinese Academy of Science. Science Press, Beijing, 116–123.
- 2031
- 2032 Wu, Y., Cui, Z., Liu, G., Ge, D., Yin, J., Xu, Q., Pang, Q. 2001. Quaternary geomorphological
- 2033 evolution of the Kunlun Pass area and uplift of the Qinghai-Xizang (Tibet) Plateau.
- 2034 Geomorphology 36, 203-216.
- 2035
- 2036 Von Wissmann H., 1959. Die heutige Vergletscherung und Schneegrenze in Hochasien mit
- 2037 Hinweisen auf die Vergletscherung der letzten Eiszeit. Akademie der Wissenschaften und der
- 2038 Literatur in Mainz, Abhandlungen der Mathematisch-Naturwissenschaftlichen Klasse 14, 121-
- 2039 123.
- 2040
- 2041 Yao, T., Shi, Y., Thompson, L.G., 1997. High resolution record of paleoclimate since the Little
- 2042 Ice Age from the Tibetan ice cores. Quaternary International 37, 19-23.
- 2043
- 2044 Yao, T., Thompson, L.G., Mosley-Thompson, E., Yang, Z., Zhang, X., Lin, P., 1996.
- 2045 Climatological significance of $\delta^{18}\text{O}$ in north Tibetan ice cores. Journal of Geophysical Research
- 2046 101, D23, 29,531-29,537.
- 2047

- 2048 Yang, B., Brauning, A., Dong, Z., Zhang, Z., Keqing, J. 2008. Late Holocene monsoonal
2049 temperate glacier fluctuations on the Tibetan Plateau. *Global and Planetary Change*, 60, 126-140.
2050
- 2051 Yang, J.Q., Zhang, W., Cui, Z.J., Yi, C.L., Liu, K.X., Ju, Y.J., 2006. Late Pleistocene glaciation
2052 of Diancang Mountains and Gongwang Mountains, southeastern margin of the Tibetan Plateau.
2053 *Quaternary International* 154-155, 52-62.
2054
- 2055 Yi, C., Owen, L.A., 2006. Quaternary palaeoenvironmental change in Tibet and the bordering
2056 mountains. *Quaternary International* 154/155, 157 pp.
2057
- 2058 Yi, C., Li, X., Qu, J., 2002. Quaternary glaciation of Puruogangri-the largest modern ice field in
2059 Tibet. *Quaternary International* 97/98, 111-123.
2060
- 2061 Yi, C., Liu, K.X., Cui, Z.J., Jiao, K.Q., Yao, T.D., He, Y.Q., 2004. AMS radiocarbon dating of
2062 late Quaternary glacial landforms, the source area of the Urumqi River, Tien Shan: a pilot study
2063 of ^{14}C dating on inorganic carbon. *Quaternary International* 121, 99-107.
2064
- 2065 Yi, C., Chen, H., Yang, J., Liu, B., Fu, P., Liu, K., Li, S., 2008. Review of Holocene glacial
2066 chronologies based on radiocarbon dating in Tibet and its surrounding mountains. *Journal of*
2067 *Quaternary Science* 23, 533-558.
2068
- 2069 Yin, A., Harrison, T.M., 2000. Geologic evolution of the Himalayan-Tibetan orogen. *Annual*
2070 *Reviews of Earth and Planetary Sciences* 28, 211-80.
2071
- 2072 Zech, R., 2012a. A Late Pleistocene glacial chronology from the Kitschi-Kurumdu Valley, Tien
2073 Shan (Kyrgyzstan), based on ^{10}Be surface exposure dating. *Quaternary Research* 77, 281-288.

2074

2075 Zech, R., Abramowski, U., Glaser, B., Sosin, P., Kubik, P.W., Zech, W., 2005. Late Quaternary
 2076 glacier and climate history of the Pamir Mountains derived from cosmogenic ^{10}Be exposure ages.
 2077 Quaternary Research 64, 212-220.

2078

2079 Zech, W., Glaser, B., Abramowski, U., Dittmar, C., Kubik, P.W., 2003. Reconstruction of the
 2080 Late Quaternary Glaciation of the Macha Khola valley (Gorkha Himal, Nepal) using relative and
 2081 absolute (^{14}C , ^{10}Be , dendrochronology) dating techniques. Quaternary Science Reviews 22, 2253-
 2082 2265.

2083

2084 Zech, W., Glaser, B., Ni, A., Petrov, M., Lemzin, I., 2000. Soil as indicators of the Pleistocene
 2085 and Holocene landscape history: Alay Range (Khyrgstan). Quaternary International 65/66: 161-
 2086 170.

2087

2088 Zech, W., Glaser, B., Sosin, P., Kubik, P.W., Zech, W., 2003. Evidence for long-lasting landform
 2089 surface instability on hummocky moraines in the Pamir Mountains (Tajikistan) from ^{10}Be surface
 2090 exposure dating. Earth and Planetary Science Letters 237, 453-461.

2091

2092 Zech, R., Zech, M., Kubik, P.W., Kharki, K., Zech, W., 2009. Deglaciation and landscape history
 2093 around Annapurna, Nepal, based on ^{10}Be surface exposure dating. Quaternary Science Reviews
 2094 28, 1106–1118.

2095

2096 Zech, R., Abramowski, U., Glaser, B., Sosin, P., Kubik, P.W., Zech, W., 2005a. Late Quaternary
 2097 glacial and climate history of the Pamir Mountains derived from cosmogenic ^{10}Be exposure ages.
 2098 Quat. Res. 64, 212–220.

2099

- 2100 Zech R., Röhringer, I., Sosin, P., Kabgov, H., Merchel, S., Akhmadaliev, S., Zech, W., 2013.
 2101 Late Pleistocene glaciations in the Gissar Range, Tajikistan, based on ^{10}Be surface exposure
 2102 dating. *Palaeogeography, Palaeoclimatology, Palaeoecology* 369, 253–261.
 2103
- 2104 Zeitler, P.K., Meltzer, A.S., Koons, P.O., Craw, D., Hallet, B., Chamberlain, C.P., Kidd, W.S.F.,
 2105 Park, S.K., Seeber, L., Bishop, M., Shroder, J.F., 2001. Erosion, Himalayan geodynamics and the
 2106 geomorphology of metamorphism. *GSA Today* 11, 4-8.
 2107
- 2108 Zhang, Z.S., 1988. Fluctuation of glaciers on the northwest slope of Mt. Nanjagbarwa since the
 2109 last glaciation. (Chinese with English summary). *Journal of Glaciology and Geocryology* 10,
 2110 180–188.
 2111
- 2112 Zheng, B., 1997. Glacier variation in the monsoon maritime glacial region since the last
 2113 glaciation on the Qinghai-Xizang (Tibetan) plateau. In: *The Changing Face of East Asia During*
 2114 *the Tertiary and Quaternary*. Center of Asian Studies, The University of Hong Kong, 103–112.
 2115
- 2116 Zheng, B.X., 2006. Quaternary glaciations in Nyainqentanglha Mountains. In: Shi, Y.F., Su, Z.,
 2117 Cui, Z.J. (Eds.), *The Quaternary Glaciations and Environmental Variations in China*. Science and
 2118 Technology Press of Hebei Province, Shijiazhuang, 374–400.
 2119
- 2120 Zheng, B.X., Ma, Q.H., 1994. The glacier variation, climatic change and the river valley
 2121 development in the Holocene on the Gongga Mountains. (Chinese with English summary). *Acta*
 2122 *Geographica Sinica* 49, 500–508.
 2123
- 2124 Zheng, B.X., Li, J.J., 1986. Regional divisions of existing glaciers in Xizang region. In: Li, J.J.,
 2125 Zheng, B.X., Yang, X.J. (Eds.), *Glaciers of Xizang (Tibet)*. Science Press, Beijing, 130–148.

- 2126
- 2127 Zheng, B., Rutter, N., 1998. On the problem of Quaternary glaciations, and the extent and
 2128 patterns of Pleistocene ice cover in the Qinghai-Xizang (Tibet) plateau. *Quaternary International*
 2129 45/46, 109-122.
- 2130
- 2131 Zheng, B.X., Li, S.J., Wang, S.M., 1995. The Quaternary glacier evolution history in the
 2132 surrounding high mountains, Zoige Basin. In: Expert Committee on Qingzang Program (Eds.),
 2133 The Formation Evolution, Environmental Variance, and Ecosystem Research in Qinghai–Tibet
 2134 Plateau, Science Press, Beijing, 218–226.
- 2135
- 2136 Zhao, J., Liu, S., He, Y., Song, Y., 2009. Quaternary glacial chronology of the Ateayinake River
 2137 Valley, Tianshan Mountains, China. *Geomorphology* 103, 276–284.
- 2138
- 2139 Zhao, J., Lai, Z., Liu, S., Song, Y., Li, Z., Yin, X., 2012. OSL and ESR dating of glacial deposits
 2140 and its implications for glacial landform evolution in the Bogeda Peak area, Tianshan range,
 2141 China. *Quaternary Geochronology*, 10, 237-243.
- 2142 Zhou, S., Li, J., Zhao, J., Wang, J., Zheng, J. 2011. Quaternary glaciations: extent and chronology
 2143 in China. In: Elhers, J., Gibbard, P. and Hughes, P.D. (Eds.), *Quaternary Glaciations - Extent and*
 2144 *Chronology: a closer look. Developments in Quaternary Science*, volume 15, Elsevier,
 2145 Amsterdam, 2nd Edition, 981-1002.
- 2146
- 2147 Zhou, S.Z., Chen, F.H., Pan, B.T., Cao, J.J. Li, J., Derbyshire, E., 1991. Environmental change
 2148 during the Holocene in western China on a millennial timescale. *Holocene* 1, 151–156.
- 2149
- 2150 Zhou, S.Z., Xu, L.B., Colgan, P.M., Mickelson, D.M., Wang, X.L., Wang, J., Zhong, W., 2007.

- 2151 Cosmogenic ^{10}Be dating of Guxiang and Baiyu glaciations. Chinese Science Bulletin 52, 1387–
2152 1393.
2153
- 2154 Zhou, S.Z., Yi, C.L., Shi, Y.F., Ye, Y.G., 2001. Study on the ice age MIS 12 in Western China.
2155 Journal of Geomechanics 7 (4), 321–327 (in Chinese with English Abstract).
2156
- 2157 Zhou, S.Z., Wang, X.L., Wang, J., Xu, L.B., 2006. A preliminary study on timing of the oldest
2158 Pleistocene glaciation in Qinghai–Tibetan Plateau. Quaternary International 154/155, 44–51
2159
- 2160 Zhou, S.Z., Wang, J., Xu, L.B., Wang, X.L., Colgan, P.M., Mickelson, D.M., 2010. Glacial
2161 advances in southeastern Tibet during late Quaternary and their implications for climatic changes.
2162 Quaternary International 218, 58–66.
2163

FIGURES

Fig. 1. Digital elevation model produced using GeoMapApp

(<http://www.geomapapp.org>) showing the area discussed in this paper.

Fig. 2. Simplified views of characteristic summer (A) and winter (B) air circulation over southern and central Asia. The Tibetan Plateau and bordering mountains above 5000 m asl are shown within the areas shaded brown. The blue solid lines in (A) and (B) indicate airflow at about 6000 and 3000 m asl, respectively, and the dark blue dashed lines airflow at about 600 m asl (adapted from Owen et al. 1998 and Benn & Owen 1998).

Fig. 3. Precipitation maps with the TRMM data draped over the DEM with 36% opacity. Units are in mm/a and are averaged over 12 years (1998-2009). From data product 2B31, a combined Precipitation Radar (PR) / TRMM Microwave Imager (TMI) rain-rate product with path-integrated attenuation at 4 km horizontal and 250 m vertical resolutions (<http://www.geog.ucsb.edu/~bodo/TRMM/#ascii>). The inset map (from Owen et al., 2005) shows a larger area to compensate for the regions not covered by TRMM data.

Fig. 4. Glaciers of the Himalayan-Tibetan orogen. A and B) Small valley polar-type continental glaciers in the Zaskar Range of northern India. C) Rongbuk glacier in Tibet and its end moraine on the northern slopes of Mt. Everest. D) Ghulkin glacier in the Karakoram Mountains of Northern Pakistan showing an impressive laterofrontal moraine complex in the fore and middle ground and the accumulation area in the top right of the photograph. E) Maritime-type monsoonal influenced unnamed glacier in the Bhagarathi

valley in the Garhwal Himalaya of Northern India. The contemporary laterofrontal moraines and outlet glaciofluvial stream is inset into an older laterofrontal moraine complex. F) Morainic rock glacier in the Milang Valley in Lahul Himalaya of Northern India. The very last remnants of glacial ice can be seen in the top left part of the photograph.

Fig. 5. Selected reconstructions for the maximum extent of glaciation during the Last Glacial across Tibet and the bordering mountains (from Owen et al. 2008a, and Owen, 2010). Light grey, relief over 4000 m above sea level; dark grey, areas considered glaciated. A) Klute's (1930) reconstruction based on a temperature depression of $\sim 4^{\circ}\text{C}$, with a shift of climatic zones to the south and an intensification of atmospheric circulation such that precipitation increased towards the dry areas of central Asia. B) Frenzel's (1960) reconstruction based on the detailed work of von Wissmann (1959), who evaluated the observations of the earliest explorers. C) Kuhle's (1985) reconstruction based on field observations and extrapolation of large equilibrium-line altitude depressions (>1000 m) from the margins of Tibet into the interior regions. D) Reconstruction of Shi (1992) and Li et al. (1991) based on detailed field mapping of glacial and associated landforms and sediments.

Fig. 6. Examples of large runout landslides in the Himalaya and Karakoram. A) The Ghoru Cho rock avalanche in the Shigar valley, Karakoram, Northern Pakistan, dated by radiocarbon to ~ 7.1 ka by Hewitt (1999) and by ^{36}Cl to Late Glacial through early Holocene by Shroder et al. (2011). B) Katzarah landslides in the Indus valley of the

2210 Karakoram, Northern Pakistan mapped and studied by Hewitt (1999) and dated by Hewitt
 2211 et al. (2011) to ~7.8 ka using ^{10}Be . C) The Kelang Serai landslide in the Lahul Himalaya
 2212 dated by ^{10}Be to 6.6 ± 0.4 ka by Dortch et al. (2009). D) The Hattian Bala landslide
 2213 triggered by the 2005 Kashmir earthquake. The landslide buried four villages resulting in
 2214 ~450 fatalities (described in Dunning et al., 2007 and Owen et al., 2008b).

2215

2216 **Fig. 7.** Geomorphology of the Skardu Basin, Karakoram Mountains, Northern Pakistan
 2217 (after Owen et al., 1988). Hewitt (1998, 1999, 2011) argues that some of the mapped
 2218 moraines are long runout landslides; these are highlighted by the golden ellipses
 2219 numbered I-VIII (adapted from Kamp and Owen, 2011).

2220

2221 **Fig. 8.** Examples of geomorphic maps produced by Owen et al. (2012) for developing a
 2222 morphostratigraphy in the Tashkurgan valley of the Chinese Pamir. A) through D) are
 2223 individual study areas within the valley. The legends are separated into feature types in
 2224 different panels, but refer to all panels. The large white text on each moraine is the
 2225 relative assignment of glacial ages to the landforms. The letters K, J, HD and T refer to
 2226 the study areas (e.g. K = Kuzigun study area) and the numbers refer to the relative ages (1
 2227 being the oldest age glacial landform in the particular study area). Sampling points for
 2228 ^{10}Be TCN dating are shown to help illustrate a typical strategy for dating. The ^{10}Be ages
 2229 for these samples are shown in Figure 10.

2230

2231 **Fig. 9.** A) Röthlisberger and Geyh's (1985b) summary of glacier fluctuations in the
 2232 Himalaya and Karakoram defined by radiocarbon dating. The original curves of

Röthlisberger and Geyh (1985b) were plotted on a radiocarbon timescale, but this figure redrafted by Owen (2009) has the broad advances and retreats plotted for calibrated years using Reimer et al.'s timescale (2004). B) Probability distribution frequency plot and scatter graph for radiocarbon ages from the Himalaya-Tibetan orogen. Only minimum radiocarbon ages, that is, ones that are younger than the moraines they are dating are plotted on the probability distribution graph (see Table 2 for data and sources). Light gray bands indicate times of rapid climate changes after Mayewski et al. (2004).

Fig. 10. Log-normal plots of > 1700 TCN ages from the Tibetan-Himalayan orogen expressed as percentage difference from the Lal (1991) – Stone (2000) time-independent scaling scheme. A) Published ages, light gray shading shows ages with differences that would fit within geomagnetic corrections schemes. Dark gray area delineates ages that fit in the Lal (1991) – Stone (2000) time dependent scheme specifically, which covers the majority of geomagnetic corrected ages. B) scaling schemes including geomagnetic correction, C) time-independent ages using new regional production rates, and D) Balco et al. (2009) regional production rate combined with scaling schemes including geomagnetic corrections. Thick horizontal dashed lines enclose 10% deviation from time-independent ages. All primary data are presented in Appendix 1.

Fig. 11. Log-normal plot of 267 gaussians that represent regional glacial stages (combined local glacial stages) for the semi-arid western end and the monsoon-influenced areas of the Himalayan-Tibetan orogen of Dortch et al. (2013) and Murari et al. (2013), respectively. Y-axis position is determined by the total number of samples

from a landform(s). Black dots indicate gaussians that enclose 3 or more ages. Green dots represent gaussians that enclose two ages and are considered tentative. Red dots represent either the oldest gaussian from a dataset that was too scattered to fit a normal distribution or landforms with a single age. The 31 gaussians from landforms that do not follow morphostratigraphic order are excluded from this plot.

Fig. 12. An example of the spread of ^{10}Be ages for glacial boulders dated to define glacial stages from Owen et al. (2012) for the Tashkurgan valley in the Pamir. The glacial stages are plotted by relative age (oldest to the left). The horizontal black lines and the dark grey bands show the mean and 1s for each glacial stage, respectively. The light blue bands highlight interstadials and interglacials based on the Marine Oxygen Isotope Stages (MIS) of Martinson et al. (1987). Note how the spread of ages for each glacial stage becomes progressively larger with relative age, likely reflecting geologic effects such as weathering, exhumation and toppling of boulders resulting in anomalously young ages.

Fig. 13. Sampling strategies for OSL and TCN dating of moraines and associated deposits and landforms proposed by Benn and Owen (2002) showing the significance of each dating site.

Fig. 14. View of section through a composite latero-frontal moraine at Lhotse-Nup in the Khumbu Himal, Nepal. Measured drawing after Richards et al. (2000b). UC1, UC2, and UC3 represent three unconformities. OSL dating undertaken by Richards et al. (2000b) is shown to illustrate the long and complex history of this landform. The lithofacies codes

of Eyles et al. (1983) are used to describe the sedimentology (Dmm—massive matrix-supported diamict; Dms—stratified, matrix-supported diamict; Dcs—stratified, clast-supported diamict; Sh—low-angle, cross-stratified, very fine to very coarse sands; Gh—horizontally bedded gravels).

Fig. 15. Glacial chronology for Mount Everest after Finkel et al. (2003) and Owen et al. (2009). A) Schematic diagram showing the stratigraphic context and locations, and graphic sedimentary logs for samples for OSL (in red samples OSL-1 to OSL-22) and ^{10}Be (in black Ron 1 to 78) for the Rongbuk valley on the northern side of Mount Everest (after Owen et al., 2009). The ages for these samples are shown in part B. B) ^{10}Be and OSL ages for moraines for the Rongbuk valley. Ages for each dated moraine are enclosed by a different color rectangle. The proximal outwash terrace is also dated (after Owen et al., 2009). C) Probability distribution plots for ^{10}Be ages for glacial stages on either side of Mount Everest (upper plots = Rongbuk valley and lower plots = Khumbu Himal on the south side of Mt. Everest) are compared (after Owen et al., 2009).

Fig. 16. Examples of evidence for very old glaciation in the Himalayan-Tibetan orogen. A) View looking east along the Indus Valley from a Leh glacial stage moraine (penultimate or older glacial cycle) towards Indus glacial stage moraines in the middle distance that date to more 430 ka (Owen et al., 2006a). B) Naimona'nyi glacial stage moraine dating to MIS 10 or older in the foreland of the Ronggua Gorger on the west side of Gurla Mandata. This particular boulder was dated using ^{10}Be TCN to $\sim 482 \pm 48$ ka (Owen et al., 2010). C) Dabudaer glacial stage moraine near the mouth of the

Jialongquiete valley in the Tashkurgan valley, which dates to penultimate or an earlier glacial cycle. This particular boulder was dated to 291 ± 20 ka using ^{10}Be TCNs (Owen et al., 2012).

Fig. 17. Probability distribution plots for ^{10}Be TCN moraine boulder ages for the Himalayan-Tibetan orogen presented for different time slices. All ages were recalculated using the Lal (1991)-Stone (2000) time dependent model (see Appendix 1 for the data).

Fig. 18. Probability distribution frequency plots for ^{10}Be TCN moraine boulder ages for different time slices for four regional subdivisions of the Himalayan-Tibetan orogen building on the methods of Owen et al. (2008a). Compared to Owen et al. (2008a), the data set has increased from 777 to > 1700 ages (see Appendix 1 for data). All ages were recalculated using the Lal (1991)-Stone (2000) time dependent model.

Fig. 19. Simplified glacial chronologies for each region in the Himalayan-Tibetan orogen back to 100 ka (adapted from Owen et al., 2012). The color bars represent times of glacial advances and their duration includes the likely uncertainty in the ages. The color scheme is used to suggest tentative regional correlations. Marine Oxygen Isotope Stages are shown to the far right.

Fig. 20. Glacial geologic study areas at the western end of the Himalayan-Tibetan orogen showing the ages of local last glacial maxima and approximate extent of valley glaciation. Data from Owen et al. (1992, 2001, 2010, 2002b, 2007, 2002a, 2006a),

2325 Sharma and Owen (1996), Phillips et al. (2000), Richards et al. (2000a), Taylor and
 2326 Mitchell (2000), Barnard et al. (2004a, b), Zech et al. (2005a), Chevalier et al. (2005,
 2327 20011), Abramowski et al. (2006), Seong et al. (2009a), Dortch et al. (2010a,2011b,c),
 2328 Hedrick et al. (2011), Rohringer et al. (2012) and Lee et al. (in press). The area shown in
 2329 the right panel is same area that was modeled by Bishop et al. (2010) using a seventy-
 2330 year simulation with a fully coupled atmosphere-ocean general circulation model for the
 2331 global LGM, 16 ka, 9 ka, 6 ka and a doubling of CO₂.

2332

2333 **Fig. 21.** Age plot of regional glacial stages (semi-arid western Himalaya-Tibetan Stage:
 2334 SWHTS) for the semi-arid regions at the western end of the Himalayan-Tibetan orogen
 2335 defined by Dortch et al. (2013) and for the monsoon-influences Himalayan-Tibetan
 2336 orogen (monsoon Himalayan-Tibetan stages: MOHITS) defined by Murari et al. (2014).
 2337 Ages for regional glacial stages are distinct at the 95% confidence interval despite some
 2338 overlap. SWHTS calculations were based on 352 out of 695 ¹⁰Be ages that were
 2339 analyzed; 98 failed the cluster analysis, 192 and 50 we excluded for being too young and
 2340 too old, respectively. MOHITS calculations were based on 520 out of 1119 ¹⁰Be ages that
 2341 were analyzed; 185 failed the cluster analysis, 316 and 98 were excluded for being too
 2342 young and too old, respectively. Stacked marine $\delta^{18}\text{O}$ curves of Lisieki and Raymo
 2343 (2005), simulated monsoon index and 65°N insolation of Leuschner and Sirocko (2003)
 2344 and the NGRIP (2004) $\delta^{18}\text{O}$ curve are provided for comparison. The duration of specific
 2345 climatic events are marked by red and dark blue bars in the far right column. Light gray
 2346 horizontal bands in the far right column indicate times of rapid climate change after
 2347 Mayewski et al. (2004).

2348

2349 **Fig. 22.** Probability distribution plots for ^{10}Be TCN surface-exposure ages for glacial
2350 advances in Muztag Ata and Kongur Shan in the Chinese Pamir for the last 20 ka (after
2351 Seong et al., 2009a). Moraines of similar advances are grouped with each different color
2352 representing a distinct glacial advance. Light gray bands indicate times of rapid climate
2353 change after Mayewski et al. (2004). The number of boulders dated per each advance is
2354 shown at the bottom of the figure.

2355

2356

2357

2358 **Tables**

2359

2360 **Table 1**

2361 Summary of the radiocarbon dating that has been undertaken throughout the Himalayan-
2362 Tibetan orogen for glacial successions.

2363

2364 **Table 2**

2365 Major optically stimulated luminescence dating projects that has been undertaken for
2366 glacial geologic studies in the Himalayan-Tibetan orogen.

2367

2368 **Table 3**

2369 Summary of glacial chronologies dated by ^{10}Be methods throughout the Himalayan-
2370 Tibetan orogen listing the dating methods applied to define the ages of glacial advances.

2371

2372 **Table 4**

2373 Comparison of regional glaciers stages of Dortch et al. (2013) and Murari et al. (2013)
2374 for the semi-arid western end and the monsoonal-influenced areas of the Himalayan-
2375 Tibetan orogen, respectively. Correlations are assessed using student T-Test.

2376

2377 **Supplementary Item/Appendix**

2378

2379 Appendix 1. Table listing the Be-10 TCN exposure dating that has been undertaken in the
2380 Himalayan-Tibetan orogen with published ages and recalculated ages using different
2381 scaling models. All references are listed in the main text of the paper.

Table 1

Table 1. Summary of major radiocarbon dating studies for the Himalayan-Tibetan orogen.

Area	Location	Latitude	Longitude	Altitude	Dated material and geomorphic context	Age in relation to glacial advance	Radiocarbon age (years)		Calibrated age (cal years BP) ^a			Source
		(°N)	(°E)	(m asl)				uncertainty	From	To	%	
Pamir	Ayizhisu River, trib of Wuyitak River, northern slope of Kongur Peak	~38.7 ^b	75.3 ^b	2700 ^b	Wood root in terminal moraines	Younger	110	50	277	8	95.4	Ono et al. (1997)
						Younger	220	50	431	-4	95.4	
Tien Shan	Upper Urumqi River valley	43.12	86.82	~4200	Organic carbon coating on boulder	Younger	390	120	652	-3	95.3	Yi et al. (2004)
						Younger	420	150	676	-3	95.5	
					Organic coating on clast	Younger	1860	100	2035	1545	95.4	
						Younger	6560	150	7700	7165	95.4	
					Till matrix	Contemporaneous	23080	510	29201	26346	95.4	
						Contemporaneous	19010	450	23811	21557	95.4	
						Contemporaneous	19590	130	23856	22691	95.4	
Karakoram	Minapin Glacier, Hunza Valley	~36.25	~74.52	~2000	Wood in moraines	Younger	325	60	505	158	95.4	Derbyshire et al. (1984)
						Younger	685	45	694	553	95.4	
						Younger	2265	65	2453	2070	95.4	
	Gargo Glacier, Gilgit Agency	36.04	74.6	~3100	soil	Older	90	85	285	...	95.4	Röthlisberger and Geyh (1986)
						Younger	585	80	676	504	95.4	
						Older	3620	130	4381	3590	95.5	
Nun Kun	Rantac Glacier	34.05	75.94	3100	Humic matter in superposed fossil soil on end moraine	Contemporaneous	19490	1630	28473	19896	95.4	Röthlisberger and Geyh (1985)
						Younger	15570	770	20828	17054	95.4	

	Tarangoz Glacier	34.05	75.91	3800	Humic matter in fossil soil on lateral moraine	Younger	2230	95	2485	1952	95.4	
						Older	3280	90	3816	3336	95.4	
		34.05	75.91	3800	Humic matter in fossil soil buried by colluvium on lateral moraine	Younger	2395	170	2843	2005	95.4	
						Younger	3630	105	4247	3642	95.4	
		34.06	75.93	3150	Humic matter in organic horizon in loess on lateral moraine	Younger	4680	60	5583	5307	95.4	
						Younger	4860	60	5731	5469	95.4	
		34.05	75.91	3900	Humic matter in fossil soil within lateral moraine	Younger	7405	185	8590	7862	95.4	
						Younger	12750	190	16216	14189	95.4	
		34.06	75.93	3425	Buried soil in loess overlaying moraine	Younger	1620	40	1607	1408	95.4	Lee et al. (2013)
						Younger	3150	40	3455	3265	95.4	
						Younger	4480	40	5298	4975	95.4	
Kashmir	Taragoz Glacier	75.91	34.05	3800	soil	Older	2230	95	2485	1952	95.4	Röthlisberger and Geyh (1986)
					soil	Younger	3280	90	3816	3336	95.4	
					soil	Older	7505	185	8750	7940	95.4	
					soil	Younger	12750	1909	21950	10733	95.4	
	Risla Glacier	76.01	33.63	3700	soil	Older	3210	80	3634	3261	95.4	
					soil	Younger	3660	185	4520	3487	95.4	
	Kolohoi Glacier	75.32	34.27	3870	soil	Older	2190	90	2354	1950	95.4	
					soil	Younger	2440	60	2711	2352	95.4	
					wood	Contemporaneous	2575	80	2844	2364	95.3	
					soil	Older	2695	160	3210	2360	95.4	
					soil	Younger	3490	100	4074	3482	95.4	

Lahul	Chandra Tal	32.47	77.61	4300	Basal peat layer over moraine	Younger	9160	70	10509	10212	95.4	Owen et al. (1997)
						Younger						
Langtang Himal	Langtang Valley	28.2	85.5	~3700	Organic matter in soil on terminal moraines	Younger	550	70	665	500	95.4	Shiraiwa and Watanabe (1991)
						Younger	2800	110	3244	2741	95.4	
						Younger	2850	140	3356	2745	95.4	
						Younger	2980	110	3399	2865	95.4	
						Older	3650	320	4868	3214	95.4	
					Wood (not in situ)	Younger	3860	110	4572	3929	95.4	
Nepal	Gangapurna N. Glacier	84.01	28.64	3680	wood in situ	Older	135	50	283	0	95.4	Röthlisberger and Geyh (1986)
					soil	Older	1210	55	1274	986	95.4	
					soil	Younger	2330	45	2666	2160	95.5	
					soil	Older	2955	105	3373	2860	95.4	
					soil	Younger	4560	280	5904	4522	95.4	
Khumbu Himal	Khumbu Glacier	27.9	86.8	~4600	Charcoal in outwash terraces associated with moraines	Younger	550	85	680	335	95.4	Benedict (1976)
					Charcoal and soil humates in outwash terraces associated with moraines	Younger	530	165	897	-2	95.3	

				Charcoal and shrub fragments in outwash sands associated with moraines	Younger	1155	160	1348	742	95.4	
				Wood in moraine	Younger	410	110	654	-2	95.4	Fushimi (1978)
					Younger	2640	170	3207	2342	95.4	
				Lateral moraine	Younger	480	80	653	317	95.4	Muller (1980)
					Younger	1150	80	1262	931	95.4	
Khumbu 1 Glacier				soil	Older	960	70	1050	728	95.4	Röthlisberger and Geyh (1986)
Khumbu 1 Glacier				soil	Older	1310	115	1415	962	95.4	
				soil	Older	1340	75	1380	1076	95.4	
				soil	Older	1420	95	1538	1142	95.4	
				soil	Older	1635	115	1813	1316	95.4	
				soil	Younger	2215	165	2713	1873	95.4	
				soil	Younger	2250	190	2749	1834	95.4	
				soil	Younger	2540	145	2964	2208	95.4	
				soil	Older	2625	125	2995	2355	95.4	
				soil	Older	2665	160	3203	2352	95.4	
				soil	Younger	3115	370	4294	2364	95.4	
				soil	Younger	3290	550	5271	2314	95.4	
Khumbu 2a Glacier				plants	Younger	1150	80	1262	931	95.4	
Khumbu 2b Glacier				plants	Younger	480	80	653	317	95.4	
Kyuwo 3 Glacier	no data			wood	Contemporaneous	410	100	644	153	95.5	
				charcoal	Younger	1200	100	1292	936	95.4	
Llotse Nup 1 Glacie	86.895	27.9225	5050	soil	Older	1025	110	1226	725	95.4	
				soil	Older	1190	155	1384	789	95.4	
				soil	Older	1630	155	1893	1279	95.4	
				soil	Contemporaneous	3265	700	5576	1986	95.4	
				soil	Contemporaneous	3385	500	5213	2459	95.4	
Nupse 1 Glacier	86.87277778	27.92777778	5170	soil	Younger	565	55	655	515	95.4	
				soil		4745	730	7245	3700	95.4	

					soil	Older	5575	500	7555	5323	95.4	
	Lhotse Shar 1 Glacier	86.95	27.91	5170	organics	Older	610	40	660	541	95.4	
					soil	Older	775	110	922	553	95.4	
					organics	Younger	1135	70	1242	928	95.4	
					soil organic fragmer	Younger	1205	55	1269	984	95.4	
					soil	Older	1705	95	1860	1403	95.4	
					soil	Younger	2195	160	2704	1830	95.4	
Bhutan Himalaya	Raphstheng Glacier, upper reaches of Lingshi Chu (River) and Goyok Chu	90	28	~4600	Humid soil buried in a terminal moraine	Younger	1710	40	1709	1535	95.4	Iwata et al. (2002)
						Younger	2120	40	2302	1992	95.4	
Everest	Rongbuk valley				Carbonate precipitate on clast	Younger	3170	70	3563	3220	95.4	Mann et al. (1996)
Namche Barwa		29.63	95.63	2950	Wood in a terminal moraine in front of Zelongnong Glacier	Younger	287	93	514	-4	95.4	Zhang (1988), Chinese Academy of Sciences (1996)
						Younger	394	83	552	159	95.4	
						Younger	1481	134	1705	1094	95.4	
Qilian Shan	Baishui River valley	~38.4	~98.7	~4500	Humic matter in buried soil interlayer in lower part of an overlapped terminal moraines	Contemporaneous	2530	120	2856	2341	95.4	Wu (1984a, b)
						Contemporaneous	3110	120	3607	2972	95.4	

	Dunde ice cap	~38.1	94.4	~4800	Humic matter in a lacustrine interlayer in a terminal moraines	Older	8455	265	10194	8774	95.4	Li and Jiao (1990)
West Kunlun	Keliya and North glaciers, Chongce ice cap	~35.3	~81.5	~5600	Humic matter in a lacustrine interlayer in terminal moraines	Older	8134	176	9470	8605	95.4	Jiao and Zheng (2006)
						Older	8287	160	9549	8776	95.4	
East Kunlun	Guoluo Glacier, Bayanhar Mt.	~33	~101	4100	Humic matter 40-60 cm deep in two terminal moraines	Younger	2540	110	2845	2349	95.4	Zheng et al. (1995)
		33	~101	~4100		Younger	4090	95	4850	4305	95.4	
Tanggula	Gaerqu River	c	c	c	Humic matter at the bottom of soil covering terminal moraines	Earlier than	2690	110	3140	2467	95.3	Jiao and Shen (2006) cited in Ti et al. (2008)
	Page Glacier	32	95	4200-4300	Lichen ages on three end moraines	Minimum	5410, 1809, 1857-1888, 1920	225	6674	5662	95.4	Li et al. (1986)
Eastern Nyainqentangliha	Zepu Glacier	30.3	95.15	2960	Detrital wood, trees in moraines, charcoal and humic soil in moraines, Zepu Glacier	Minimum	190	80	430	-4	95.4	Jiao and lwata (1993); lwata and Jiao (1993); Jiao et al. (2005)
				3400		Younger	197	80	432	-4	95.4	
				3300		Younger	580	130	767	315	95.4	
				>3000		Younger	720	210	1120	306	95.4	
				>3000		Younger	950	120	1167	670	95.5	
				3200		Younger	1056	115	1256	737	95.4	

Qiangtang Plateau	Arza Glacier	29.77	95.7	>3000		Younger	1100	160	1305	725	95.4	Li et al. (1986)		
				>3000		Younger	1570	300	2302	923	95.4			
				>3000		Younger	3242	101	3716	3220	95.4			
				~3850	Wood in a lateral moraine, Arza Glacier	Younger	2980	150	3470	2785	95.4			
														Wang and Fan (1987)
	~3835	Tree branches buried in lateral moraine	Younger	1820	100	1987	1530	95.4						
	~3820	Extreme growth of trees on lateral moraines	Younger	1813-1852, 18884-1908, 1960							Wang and Fan (1987)			
	Ruoguo Glacier	30.3	94.73	3960	Wood in a moraine, Ruoguo Glacier	Younger	1920	110	2143	1573	95.4		Li et al. (1986), Zheng and Li (1986)	
				3880		Younger	1740	85	1870	1418	95.4		Li et al. (1986)	
				3820		Younger	1540	85	1610	1295	95.4		Li et al. (1986)	
				3640	Lichen dates on moraine	Younger	1822 AD							
	Bomi Glacier	29.77	95.7	4000	Age of trees	Contemporaneous	1580-1590, 1860-1890							Bräuning (2006)
	Laigu Glacier	29.33	96.8	~4000	Buried tree in lateral moraine	Contemporaneous/maximum	1150	80	1262	931	95.4		Wang and Fang (1987)	
	Luzi Glacier	not provided	not provided	5100	Humic matter in soil 1-2 m below the turf on the terminal moraine	Earlier than/minimum	3050	100	3461	2961	95.4		Zheng (2006)	
Qiangtang Plateau		32.61	89.77	~4900	Humic matter in an exposure of a terminal moraine, Gangzari Mt.	Contemporaneous	5760	170	7001	6212	95.4	Li and Li (1992)		
Hengduan Mountains	Hailuoguo Glacier, Gongga Shan	29.6	101.95	2850	Wood in lateral and terminal moraines	Younger	150	60	288	-2	95.4	Li et al. (1986); Zheng and Ma (1994)		

			2730		Younger	540	70	665	489	95.4	
			~3000		Younger	780	90	916	560	95.4	
			2930		Younger	940	50	933	740	95.4	
			~2900		Younger	1160	50	1234	960	95.4	
			3000		Younger	1550	80	1610	1300	95.4	
			2750		Younger	1580	60	1607	1345	95.4	
			~3000		Older	2170	110	2433	1877	95.4	
			~3000		Older	2350	65	2702	2159	95.3	
			2980		Younger	3080	80	3467	3065	95.4	
			~3050	Black clay at the bottom of lacustrine sediment between two moraines	Older	8010	150	9396	8482	95.5	Zheng and Ma (1995)
Diancangshan	25.67	100.09	3985	Organic silt over laying moraines	Younger	1310	60	1315	1075	95.4	Yang et al. (2006)
	25.67	100.09	4041		Younger	1600	110	1769	1296	95.4	
	25.72	100.07	4021		Younger	3240	60	3615	3358	95.4	
	25.72	100.07	4012		Younger	3960	60	4778	4184	95.4	
	25.67	100.09	3941		Younger	4830	60	5709	5330	95.4	
					Younger						
Lhamcoka Glacier	31.82	99.12	4150-4370	Maximum tree ages on moraines	Younger	1760-1780, 1807-1820, 1907-1920					Bräuning (2006)
Yanzigou Glacier	29.63	101.9	3850-3900	Willow trunk in paleosol between two tills	Contemporaneous	145	100	429	...	95.4	Smiraglia (1997)
					Contemporaneous	235	110	485	-3	95.4	
					Contemporaneous	1305	110	1405	975	95.4	
					Contemporaneous	1610	65	1691	1358	95.4	
					Contemporaneous	1785	65	1865	1558	95.4	

Nanmenguangou	29.67	101.98	3000	Buried rotten wood in till	Contemporaneous	780	90	916	560	95.4	Zheng and Ma (1994)
Gongba Glacier	30.5	101.87	3900	Buried rotten wood in till	Contemporaneous	440	50	545	319	95.4	Su et al. (2002)
			3700		Contemporaneous	620	40	662	546	95.4	

Notes

^a Radiocarbon ages calibrated using OxCal 4.2 applying IntCal 09

^b Exact location not described in paper, so values are very approximate

^c Exact locations not provided. Ages do not readily match with original publication (in Chinese).

Table 2

Major optically stimulated luminescence dating projects that has been undertaken for glacial geologic studies in the Himalayan-Tibetan orogen.

Area	Source	Material dated
Ateaoyinake River Valley, Tianshan ¹	Zhao et al. (2009)	Till and glaciofluvial outwash
Bogeda Peak area, Tianshan ¹	Zhao et al. (2012)	Tills and associated sediments
Terskey-Alatoo Range, Kyrgyz Republic	Narama et al. (2007)	Glacial and loess deposits
Tien Shan, Yrgyz Republic	Narama et al. (2009)	Glacial and loess deposits
Central Karakoram	Seong et al. (2008)	Glaciofluvial sediment
Hunza valley, Karakoram Mountains	Derbyshire et al. (1984), Spencer and Owen (2004)	Surpaglacial and proglacial sediments
Nun Kun	Lee et al. (2013)	Aeolian and colluvial deposits
Zaskar	Taylor and Mitchell (2000)	Glaciofluvial sediments
Chitral	Richards et al (2000a), Owen et al. (2002b)	Supraglacial and proglacial glaciofluvial sediments
Swat Himalaya	Owen et al. (1992), Richards et al. (2000a)	Colluviated loess overlaying moraines
Nanga Parbat	Richards et al. (2000a)	Proglacial glaciofluvial sediments
Lahul Himalaya	Owen et al. (1997)	Glaciodeltaic sediments
Garhwal Himal	Sharma and Owen (1996)	Englacial and supraglacial glaciofluvial sediments
Central Himalaya	Ali et al. (2013)	Outwash gravels and moraines
Khumbu Himal	Richards et al. (2000b)	Supraglacial and englacial glaciofluvial sediments
Kanchenjunga	Tsukamoto et al. (2000)	Glacial sediments
Rongbuk valley, Everest	Owen et al. (2009)	Supraglacial and proglacial glaciofluvial sediments
Kunlun Mountains	Owen et al. (2006b)	Fluvial and aeolian sediments on terraces
Anyemaquen Mountains	Owen et al. (2003a)	Surpaglaciofluvial sediment

Notes: 1. Also includes ESR dating

Table 3.

Summary of glacial chronologies dated by ^{10}Be methods throughout the Himalayan-Tibetan orogen listing the dating methods applied to define the ages of glacial advances.

Study region	Study	Glacial stage or moraine name
1) Pamir and Tien Shan		
Gissar Range, Western Tajikistan	Zech et al. (2013)	G1 G2 (not dated) G3 (not dated) G4 G5 (not dated) G6 G7 G8
Aksu Valley, Turkestan Range, Kyrgyzstan		
Aksu Valley, Turkestan Range, Kyrgyzstan	Abramowski et al. (2006)	AK4 moraine AK3 moraine AK2 moraine AK1 moraine
Koksu Valley, Alay Range Kyrgyzstan		
Koksu Valley, Alay Range Kyrgyzstan	Abramowski et al. (2006)	AV moraine KK moraine
Ailuitek Pass, north Central Pamir, Tajikistan		
Ailuitek Pass, north Central Pamir, Tajikistan	Abramowski et al. (2006)	AT moraine TK moraine
S-Alichur Range, Yashikul, Tajikistan		
S-Alichur Range, Yashikul, Tajikistan	Zech et al. (2005) Zech et al. (2005) Abramowski et al. (2006) Zech et al. (2005) Abramowski et al. (2006) Zech et al. (2005)	M4 moraine M3 moraine BO1 moraine (=M3/YK3 of Zech et al. 2005) M2 moraine BO2 moraine (=M2/YK2 of Zech et al. 2005) M1 moraine
S-Alichur Range, Kol-Uchkol, Tajikistan		
S-Alichur Range, Kol-Uchkol, Tajikistan	Abramowski et al. (2006)	UK6 moraine UK5 moraine UK4 moraine UK3 moraine UK2 moraine UK1 moraine
S-Alichur Range, Gurumdy, Tajikistan		
S-Alichur Range, Gurumdy, Tajikistan	Abramowski et al. (2006)	GU4 moraine GU3 moraine GU2 moraine GU1 moraine
Bogchigir Range, Pamir Tajikistan		
Bogchigir Range, Pamir Tajikistan	Röhringer et al. (2012)	BO8 BO7 BO6 BO5 BO4 BO3 BO1 M1
Mustagata and Kongur Shan, Chinese Pamir		
Mustagata and Kongur Shan, Chinese Pamir	Seong et al. (2009a)	Olimde glacial stage Subaxh glacial stage Karasu glacial stage
Tashkurgan valley, Chinese Pamir		
Tashkurgan valley, Chinese Pamir	Owen et al. (2012)	Kuzigun Hangdi Tashkurgan Dabudaer
W Tien Shan, Kyrgyzstan		
Ala Archa, Wn Tien Shan, Kyrgyzstan Aksai, Wn Tien Shan, Kyrgyzstan	Koppes et al. (2008)	V IV, V IV, V IV, V

Ala Bash, Wn Tien Shan, Kyrgyzstan		IV
Chor Kyrchak, Wn Tien Shan, Kyrgyzstan		III
Ala Bash, Wn Tien Shan, Kyrgyzstan		II
		I
Central Tien Shan, Kyrgyzstan		
Kitschi-Kurumdu Valley, Tien Shan, Kyrgyzstan	Zech (2012)	M1
		M2
		M3
		M4
		M5
		M6
		M7
Urumqi River valley, En Tian Shan, China		
Urumqi River valley, En Tian Shan, China	Kong et al. (2009a)	Lateral moraine, Glacier Observation Station
		Latero-terminal moraine, lower trough, Shangwanfeng
		Elevated platform, Gaowangfeng
2) Transhimalaya and Western Tibet		
Sulamu Tagh, Altyn Tagh Nn Tibet		
Sulamu Tagh, Altyn Tagh Nn Tibet	Mériaux et al. (2004)	M2 moraine
		M1 moraine
Central Karakoram (K2), Nn Pakistan		
Central Karakoram (K2), Nn Pakistan	Seong et al. (2007)	Askole glacial stage
		Mungo glacial stage
		Skardu glacial stage
Hunza valley, Karakoram Mts. Nn Pakistan		
Hunza valley, Karakoram Mts. Nn Pakistan	Owen et al. (2002b)	Batura glacial stage
		Ghulkin II glacial stage
		Ghulkin I glacial stage
		Borit Jheel glacial stage
		Yunz glacial stage
Ladakh Range, Nn India		
Ladakh Range, Nn India	Owen et al. (2006a)	Khalling glacial stage
		Bazgo glacial stage
		Kar glacial stage
		Leh glacial stage
		Indus glacial stage
Ladakh, Nn India		
Ladakh, Nn India	Dortch et al. (in press)	Pangong cirque
		Ladakh cirque
		Ladakh-2 stage
		Ladakh-4 stage
		Pangong-2 stage
Nubra and Shyok valley, N Ladakh		
Nubra valley, N Ladakh	Dortch et al. (2010a)	Deshkit 1
		Deshkit 2
Shyok valley, N Ladakh		Deshkit 3
Puga valley, Zaskar, Nn India		
Puga valley, Zaskar, Nn India	Hedrick et al. (2011)	PM-3
		PM-2
		PM-1
		PM-0
Karzok valley, Zaskar, Nn India		
Karzok valley, Zaskar, Nn India	Hedrick et al. (2011)	KM-4
		KM-3
		KM-2
		KM-1
		KM-0
Ayilari Range, Sn Tibet		
Ayilari Range, Sn Tibet	Chevalier et al. (2005)	M1 moraine
		M2W (oldest moraine, west)
		M2E (oldest moraine east)
Ayilari Range, SWn Tibet		
Ayilari Range, SWn Tibet	Chevalier et al. (2011)	17b: CK M2 inner
		17c: CK M2 outer
		17a: CK M3
Gurla Mandhata, Sn Tibet		
Muguru valley, Gurla Mandhata	Owen et al. (2010)	M10
		M9
		M8
		M7
		M6
		M4

Manarodi valley, Gurla Mandhata		M6 M5 M4b M4a M4c M4b M4b M4a M3 M2 M1c M1b M1a
Ronggua gorge foreland, Gurla Mandhata		
Gurla Mandhata, Sn Tibet Gurla Mandhata, Sn Tibet	Chevalier et al. (2011)	12a: Pulan M1W 12c: Pulan M1E 12B: Pulan M2
Kailas Range, SWn Tibet Kailas Range, SWn Tibet	Chevalier et al. (2011)	12: EXS 14: WXS 15: A Qu
Rongbuk valley, N. Mt. Everest Rongbuk valley, N. Mt. Everest	Owen et al. (2009)	T7 T6 T5c T5b T4 T4 T3' T3 T2 T1
3) Monsoon-influenced Himalaya Nanga Parbat, Nn Pakistan Nanga Parbat, Nn Pakistan	Phillips et al. (2000)	Mid Holocene advance Maximum Holocene extent MIS-3 advance
Nun Kun, Nn India Nun Kun, Nn India	Lee et al. (in press)	TG3 ST3 ST2 TG1 ST1 NKB1
Lahul Himalaya, Nn India Lahul Himalaya, Nn India	Owen et al. (2001)	Historical Kulti glacial stage Batal glacial stage Chandra glacial stage (not dated)
Gangotri, Garhwal Himalaya, Nn India Gangotri, Garhwal Himalaya, Nn India	Barnard et al. (2004a)	Bhujbas glacial stage Gangotri glacial stage Kedar glacial stage Bhagirathi glacial stage
Central Garhwal, Nn India Bhillangana and Dudhganag valleys, central Garhwal	Murari et al. (in review)	mbd4 mbd3 mbd2 mbd1
Mayali, central Garhwal	Murari et al. (in review)	mm1
Kedarnath, central Garhwal	Murari et al. (in review)	mk3 mk3 mk1
Western Garhwal Himalaya, Nn. India Tons Valley, Garhwal Himalaya, Nn. India	Scherler et al. (2010)	Tons Valley (location G) Tons Valley (location E) Tons Valley (location F) Tons Valley (location F') Tons Valley (location D) Tons Valley (location C) Tons Valley (location B): Gangar lateral moraine (no stage name) Tons Valley (location B): Gangar Kame deposit Thangi valley: Lateral moraine (no stage name) Thangi valley: Lateral moraine (no stage name) Pin Valley: Lateral moriane (no stage name)
Pin Valley, Wn Garhwal, Nn India		
Thangi Valley, Wn Garhwal, Nn India		

Nanda Devi, Garhwal, Nn India Nanda Devi, Garhwal, Nn India	Barnard et al. (2004b)	Moraine m4 Moraine m3 Moraine m2 Moraine m1
Annapurna Range, Nepal Annapurna Range	Pratt-Sitaula (2004) Pratt-Sitaula (2011) Pratt-Sitaula (2011) Pratt-Sitaula (2004) Pratt-Sitaula (2004) Pratt-Sitaula (2011) Pratt-Sitaula (2011) Pratt-Sitaula (2004) Pratt-Sitaula (2004) Pratt-Sitaula (2004) Pratt-Sitaula (2004) Pratt-Sitaula (2004)	Yak upper (early-mid Holocene) Danfe Glacier (early-mid Holocene) Syaktan Glacier (early-mid Holocene) Lyapche Glacier (early Holocene) Junam Glacier (Younger Dryas) Kicho Glacier (Younger Dryas) Midim Glacier (Younger Dryas) Khudi Glacier (Younger Dryas) Yak Glacier lower (Younger Dryas) Nar main (Late Glacial) Nar Glacier end (LGM) Pangri Glacier (LGM)
Annapurna, Nepal Dudh Khola Valley, Annapurna, Nepal Marsyandi Valley, Annapurna, Nepal	Zech et al. (2009)	Little Ice Age moraine Neoglacial diamicton
Macha Lhola Valley, Gorkha Himal, Nepal Macha Lhola Valley, Gorkha Himal, Nepal	Abramowski (2004)	MK4: LIA MK7: Neoglacial MK5: Late Glacial MK2: LGM
Milarepa's Glacier, Annapurna Range, Nepal Milarepa's Glacier, Annapurna Range, Nepal	Heimsath and McGlynn (2008)	E moraine crest W moraine crest
Mailun Khola, Ganesh Himal, Nepal Mailun Khola, Ganesh Himal, Nepal	Gayer et al. (2006)	M6 M3 post-M2
Langtang Himal, Nepal Langtang Valley, Langtang Himal, Nepal	Abramowski (2004) Barnard et al. (2006)	LT6 LT3: Langtang Stage LT2 Yala I glacial stage Langtang glacial stage
Nyalam County, Sn Xixabangma, Sn Tibet Nyalam County, Sn Xixabangma, Sn Tibet	Schaefer et al. (2008)	Naisa valley moraines: older ridge Naisa valley moraines: younger ridge Fu Qu valley, Puluo moraine sequence: Puluo 2 moraine Fu Qu valley, Puluo moraine sequence: Puluo 1 moraine Nyalam moraine sequence: Moraine #3 Nyalam moraine sequence: Moraine #2 Nyalam moraine sequence: Moraine #1
Ama Drime Range, Sn Tibet Ama Drime Range, Sn Tibet	Chevalier et al. (2011)	8: Dingye N 9b: Dingye S main #1 9c: Dingye S main #2 9d: Dingye S main #3 9a: Dingye S frontal
Khumbu Himal, Nepal Khumbu Himal, Nepal	Finkel et al. (2003)	Lobuche glacial stage and historical Thukliha glacial stage Chhukung glacial stage Periche II glacial stage Periche I glacial stage Thyangboche II glacial stage Thyangboche I glacial stage
4) Monsoon-influenced Tibet Xainza Range, Sn Tibet Xainza Range, Sn Tibet	Chevalier et al. (2011)	3a: M4 #1 3b: M4 #2 3c: M4 #3 4c: M3old 4a: M3 main 4b: M3 inner 5: M2 6: M1 7:M

Kunlun Shan Kunlun Shan (southern slopes)	Owen et al. (2006b)	M2 moraines M1 moraines M2 moraines M2' moraines M1 moraines
La Ji Shan, NE Tibet La Ji Shan, NE Tibet	Owen et al. (2003b)	t3 moraine t2 moraine t1 moraine
Qilian Shan, NE Tibet NW Menyuan, Qilian Shan, NE Tibet	Owen et al. (2003c)	Holocene moraine (Gangshiga valley) LGM moraine (Gangshiga valley) LGM moraine (Laolongwan valley) Unnamed moraine (Laotugou valley)
Qilian Shan, NE Tibet Xiyang He valley, Qilian Shan, NE Tibet	Laserre et al. (2002)	unnamed moraine
Nianbaoyeze Mountains Nianbaoyeze Mountains	Owen et al. (2003a)	Ximencuo glacial stage Jiukehe glacial stage
Anyemaqen Mountains Anyemaqen Mountains	Owen et al. (2003a)	Halong glacial stage Qiemuqu glacial stage Anyemaqen glacial stage
Bayan Har, NE Tibet Bayan Har, NE Tibet	Heyman et al. (2011)	BH-1 BH-2 BH-3 BH-4
Dalijia Shan, NE Tibet Qital Valley, Dalijia Shan, NE Tibet	Wang et al. (2013)	Group D moraines Group C moraines Group B moraines Group A moraines Dalijia Pass moraines
Cental Tanggula Shan Tanggula Pass, Central Tanggula Shan	Colgan et al. (2006) Owen et al. (2005), Colgan et al. (2006) Schäfer et al. (2002), Owen et al. (2005) Owen et al. (2005), Colgan et al. (2006)	Longxiazai glacial stage (youngest) Basicuo glacial stage Zhajiazangbo glacial stage Tanggula glacial stage
Nyainqentangliha Shan, Central Tibet SE Samdainkangsang, Nyainqentangliha Shan	Owen et al. (2005)	Un-named (youngest) moraine Un-named (intermediate age) moraine Un-named (oldest) moraine
Nyainqentanggula, South Central Tibet Nyainqentanggula, South Central Tibet	Chevalier et al. (2011)	2b: ybj inner 2a: ybj outer W 2c: ybj outer E 2d: ybj outer N 1a: Gulu W 1b: Gulu E
KungCo, Sn Tibet KungCo, Sn Tibet	Chevalier et al. (2011)	11: KungCo
Cho Oyu Range, S Central Tibet Cho Oyu Range, S Central Tibet	Chevalier et al. (2011)	10: Cho Oyo
Shaluli Shan, Chuanxi Plateau, En Tibet Shaluli Shan, Chuanxi Plateau, En Tibet	Graf et al. (2008) includes Ne-21 too	Cuo Ji Gang Wa and Cuo Naleng paleoglaciers: lateral moraine Cuo Ji Gang Wa and Cuo Naleng paleoglaciers: outer end moraine Cuo Ji Gang Wa and Cuo Naleng paleoglaciers: middle end moraine Cuo Ji Gang Wa and Cuo Naleng paleoglaciers: inner end moraine Chuanxi and northern outlet glacier: resting on HAI-2 Chuanxi and northern outlet glacier: on smaller moraine ridge Chuanxi and northern outlet glacier: embedded in basal till
Shaluli Shan, Haizishan Plateau, En Tibet Shaluli Shan, Haizishan Plateau, En Tibet	Fu et al. (2013)	innermost moraine sequence intermediate moraine sequence outermost moraine sequence
Shaluli Shan, Xinlong Plateau, En Tibet		

Shaluli Shan, Xinlong Plateau, En Tibet	Fu et al. (2013)	outermost moraine sequence
Shaluli Shan, Nata, En Tibet		
Shaluli Shan, Nata, En Tibet	Fu et al. (2013)	outermost moraine sequence
Shaluli Shan, Heranseba, En Tibet		
Shaluli Shan, Heranseba, En Tibet	Fu et al. (2013)	outermost moraine sequence
Litang County, En Tibet		
Litang County, En Tibet	Schäfer et al. (2002)	Un-named moraine (youngest) Un-named moraine (oldest)
Hengduan Mountains, SE Tibet		
Kangding, Daxue Shan, Hengduan Mts., SE Tibet	Tschudi et al. (2003), Strasky et al. (2003)	Zhheduo valley moraine ridge
Mt. Kaluxung, Sn Tibet		
Karola Pass, Mt. Kaluxung, Sn Tibet	Owen et al. (2005)	Un-named (youngest) moraine Un-named (intermediate age) moraine Un-named (oldest) moraine
Gangrigabu Shan, SE Tibet		
Bodui Zangbo River, Gangrigabu SE Tibet	Zhou et al. (2007, 2010)	East moraines: Baiyu Glaciation West younger moraine: Baiyu Glaciation West older moraine: Baiyu Glaciation Area C: Guxiang Glaciation
Gonga Shan, SE Tibet		
Hailuogou Valley, Gonga Shan, SE Tibet	Monsoon Tibet Owen et al. (2005)	Little Ice Age moraines Neoglacial moraines "Recessional moraine" Local LGM moraines
Shaluli Mountain, SE Tibet		
Haizi Shan, Shaluli Mountain, SE Tibet	Wang et al. (2006)	Kuzhaori end moraine
Yulong Mountains, SE Tibet		
Renhe valley, Yulong Mountains, SE Tibet	Kong et al. (2009b)	No stage name given
Xinlian valley, Yulong Mountains, SE Tibet	Kong et al. (2009b)	No stage name given
Baishui valley, Yulong Mountains, SE Tibet	Kong et al. (2009b)	No stage name given
Ganheba, Ganheba Mountains, SE Tibet	Kong et al. (2009b)	No stage name given

Table 4

Regional glaciers stages of Dortch et al. (2013) and Murari et al. (2013) for the semi-arid western end and the monsoonal-influenced areas of the Himalayan-Tibetan orogen, respectively. Student's T-Test is utilized to compare the populations of TCN ages between glacial stages in the two regions. The resulting P-values are presented in the middle column where bold numbers indicate failure to reject the null hypothesis and the possibility of the two stages being correlative is good. Italicized P-values indicate the regional stage is distinct from all other regional stages at $\geq 95\%$ confidence interval; in these cases, the highest P-value is presented.

Semi-arid western end of the Himalayan-Tibetan orogen			Monsoon-influenced Himalayan-Tibetan orogen	
Regional glacial stage name	Age	P-value	Regional glacial stage name	Age
		<i>0.00</i>	MOHITS10	483 $\pm$ 38
SWHTS9	311 $\pm$ 32	0.92	MOHITS9	314 $\pm$ 48
SWHTS7	234 $\pm$ 44	0.18	MOHITS7	192 $\pm$ 10
		<i>0.04</i>	MOHITS6B	168 $\pm$ 12
SWHTS6	146 $\pm$ 18	0.64	MOHITS6A	143 $\pm$ 12
SWHTS5E	121 $\pm$ 11	0.97	MOHITS5E	121 $\pm$ 5
		<i>0.02</i>	MOHITS5B	91 $\pm$ 15
SWHTS5A	80 $\pm$ 5	<i>0.02</i>		
SWHTS5A-	72 $\pm$ 8	<i>0.00</i>		
SWHTS4	61 $\pm$ 5	0.12	MOHITS4	68 $\pm$ 10
		<i>0.00</i>	MOHITS3C	55 $\pm$ 5
SWHTS3	46 $\pm$ 4	0.75	MOHITS3B	46 $\pm$ 5
		<i>0.00</i>	MOHITS3A	42 $\pm$ 2
SWHTS2F	30 $\pm$ 3	0.25	MOHITS2G	31 $\pm$ 3
		<i>0.01</i>	MOHITS2F	26 $\pm$ 2

		<i>0.00</i>	MOHITS2E	22±3
SWHTS2E	20±2	<i>0.03</i>		
		<i>0.03</i>	MOHITS2D	18.7±1.8
SWHTS2D	16.9±0.7	0.21	MOHITS2C	17.3±0.3
SWHTS2C	14.9±0.8	0.23	MOHITS2B	15.5±1.3
SWHTS2B	13.9±0.5	<i>0.00</i>		
SWHTS2A	12.2±0.8	<i>0.01</i>	MOHITS2A	12.9±0.9
		<i>0.00</i>	MOHITS1K	11.4±0.7
		<i>0.00</i>	MOHITS1J	10.1±0.5
		<i>0.00</i>	MOHITS1I	9.1±0.3
1E	8.7±0.2	0.16	MOHITS1H	8.1±0.8
1D	6.9±0.1	<i>0.00</i>	MOHITS1G	7.7±0.6
Askole 1	5.5±0.3	0.42	MOHITS1F	5.4±0.6
1C	3.8±0.6	0.37	MOHITS1E	3.5±0.4
PM-2	2.8±1.2	0.54	MOHITS1D	2.3±0.1
1B	1.7±0.2	0.15	MOHITS1C	1.5±0.2
Askole 3	1.1±0.1	<i>0.03</i>	MOHITS1B	0.7±0.1
1A	0.4±0.1	0.25	MOHITS1C	0.4±0.1

Figure 1

Figure 1

Figure 2

Figure 2

Figure 3

Figure 3

Figure 4

Figure 4

Figure 5

Figure 5

Figure 6

Figure 6

Figure 7

Topographic features

- Streams
- Terrace riser/edge of moraine
- Regolith/colluvium covered slopes

Mouth of Kuzigun valley

Glacial features

- Moraine ridge
- Kettle hole

Glacial stages

- Kuzigun
- Hangdi
- Tashkurgan
- Dabudaer

Map Labels: K1, K2, K3, XJ61, XJ62, XJ63, XJ64, XJ65, XJ66, XJ67, XJ70, XJ71, XJ72, Pamir9, Pamir10, Pamir11, Pamir12, Pamir13, Pamir14, N, 1 km.

Figure 8

Figure 9

Figure 9

Figure 10

A scatter plot showing the relationship between 'Gaussian age (ka)' on the x-axis and 'Number of samples' on the y-axis. The x-axis is logarithmic, ranging from 0.01 to 1000. The y-axis is linear, ranging from 1 to 50. Data points are categorized into three groups: black, red, and green. The black group shows a wide distribution of sample sizes across all ages, with a peak around 10-100 ka. The red group is concentrated at low sample sizes (1-5) across all ages. The green group shows a more structured distribution, with sample sizes increasing from 2 to 5 as age increases from 0.5 to 100 ka, then slightly decreasing at higher ages.

Figure 11

Figure 12

Figure 12

Figure 13

Figure 13

Figure 14

Figure 14

Figure 15

Figure 15

Figure 16

Figure 16

Figure 17

Figure 17

Figure 18

Figure 18

Figure 19

Figure 20

Figure 20

Figure 21

Figure 21

Figure 22

Figure 22

Supplementary Data - Appendix 1

[Click here to download Supplementary Data: Owen and Dortch_Appendix 1 Appendix 1.pdf](#)