Submarine channel evolution, terrace development, and preservation of intra-channel thin-bedded turbidites

DOI:

[10.1016/j.margeo.2016.11.011](https://doi.org/10.1016/j.margeo.2016.11.011)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Hansen, L., Janocko, M., Kane, I., & Kneller, B. C. (2017). Submarine channel evolution, terrace development, and preservation of intra-channel thin-bedded turbidites: Mahin and Avon channels, offshore Nigeria. *Marine Geology*. <https://doi.org/10.1016/j.margeo.2016.11.011>

Published in:

Marine Geology

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Submarine channel evolution, terrace development, and preservation of intra-channel thin-bedded turbidites: Mahin and Avon channels, offshore Nigeria

Author list:

Larissa Hansen¹, Michal Janocko², Ian Kane³, Ben Kneller¹

Email of corresponding author: Larissa.Hansen@abdn.ac.uk

¹ University of Aberdeen, Department of Geology and Petroleum Geology, Aberdeen, AB24 3UE, UK

² Statoil ASA, Research Centre Bergen, NO-5020 Bergen, Norway

³ University of Manchester, School of Earth and Environmental Sciences, Williamson Building, Oxford Road, Manchester, M13 9PL, UK

Abstract

Terraces on the modern seafloor are defined as topographically flat areas above the active submarine channel thalweg but within the confines of the channel-belt. They have been described from many modern submarine channels, but the controls on terrace distribution, evolution and stacking patterns are not well understood. In this study, we describe the architecture of the Mahin and Avon channel-belts and their associated terraces, located offshore Nigeria towards the northwest of the Niger Delta. The studied channel sections are less than 10 km apart up-dip and converge downslope. They are on slopes with similar gradients, yet they have significantly different morphologies indicating that the interplay between sedimentary processes and channel evolution must be different.

26 The surfaces defining the bases of the terrace bodies have been mapped along both channels
27 using high-resolution 3D seismic data. Spectral decomposition of the data reveals subtle
28 variations in seismic character that highlight sedimentological detail that can otherwise not be
29 recognized, allowing us to suggest the processes responsible for terrace formation and terrace
30 body composition.

31

32 The contrasting evolution of the two channels is reflected in the morphology and architecture
33 of their terraces. While the Mahin terrace bodies show a predictable pattern, typically
34 consisting of stacked channel-fill and overbank deposits in a circular planform shape (shape
35 controlled by cut-off channel bends), the deposits of the Avon terrace bodies up-dip of the
36 confluence with the Mahin are dominated by overbank deposits, with the planform terrace
37 shape heavily controlled by the topography of the underlying channel-belt deposits.

38

39 This study shows how spatially and temporally associated channels can have markedly
40 different architectures. The evolution of the channel and the abundance and stacking patterns
41 of compositional elements within terrace bodies are shown to be linked.

42

43 **Keywords:** Nigeria, channel-levee, internal levee, depositional terraces, thin-bedded
44 turbidites, 3D seismic, channel sinuosity

45

46 **1. Introduction**

47 Submarine canyons and channels form some of the most striking features on the continental
48 slope. They act as conduits for sediment transport from the shelf to the basin floor (e.g. Mutti
49 & Normark, 1991) by the mechanism of sediment gravity flows (Kuenen and Migliorini,
50 1950; Bouma and Ravenne, 2004; Meiburg and Kneller, 2010). Submarine channel-fills are
51 commonly associated with thin-bedded turbidites, which are deposited in channel overbank

52 areas, building levees and terraces. While levees have been described from numerous modern
53 seafloor, outcrop and subsurface datasets (Piper and Normark, 1983; Hiscott et al., 1997;
54 Peakall et al., 2000; Deptuck et al., 2003; Kane et al., 2007; Meiburg and Kneller, 2010; Kane
55 and Hodgson, 2011; Nakajima and Kneller, 2013; Morris et al., 2014) terraces have mainly
56 been identified in modern seafloor systems based on bathymetric and side-scan sonar data,
57 with little information available on the deposits below the terrace surfaces (O’Connell et al.,
58 1995; Hübscher et al., 1997; Torres et al., 1997; Von Rad and Tahir, 1997; Piper and Hiscott,
59 1999; Deptuck et al., 2003; Babonneau et al., 2004; Babonneau et al., 2010; Gamberi and
60 Rovere, 2011; Paull et al., 2013).

61

62 Terraces are topographically flat areas adjacent to the channel but within the channel-belt
63 (Babonneau et al., 2004) and are elevated above the adjacent channel base. The foundations of
64 these terraces can be formed by a number of different processes such as: punctuated lateral
65 channel migration or entrenchment within the channel-belt; continuous channel migration
66 within the channel-belt forming point bars; channel bend cut-offs; or by collapse of external
67 levee deposits into the channel-belt due to over-steepening of the inner face of the external
68 levee (Hansen et al., 2015). Once these terrace foundations are established, the process that
69 created them and which determine their composition may no longer be active. These
70 foundations can subsequently be subject to overspill processes from the adjacent channel,
71 resulting in deposition of thin-bedded turbidites. If the overspilling sediment creates a wedge
72 shaped deposit on top of these surfaces it is called an *internal levee* (Buffington, 1952;
73 Walker, 1975; Hiscott et al., 1997; Normark and Damuth, 1997; Skene et al., 2002; Deptuck
74 et al., 2003; Pirmez and Imran, 2003; Kane et al., 2007; Meiburg and Kneller, 2010; Kane and
75 Hodgson, 2011; Nakajima and Kneller, 2013; Morris et al., 2014) while a flat deposit is called
76 a *depositional terrace* (Hansen et al., 2015). However, these deposits most likely form a
77 continuum across a variety of different scales and end-members are rarely preserved (Hansen
78 et al., 2015). The *terrace body*, is defined as the sediment that comprises the thickness from

79 the interpreted base of the terrace (below the terrace foundation) to the seafloor, and it is
80 areally constrained by its planform geometry governed by the terrace forming process and the
81 processes occurring after terrace formation. Both internal levee and depositional terrace
82 deposits can include laterally extensive sand layers and can form significant hydrocarbon
83 reservoirs (e.g. Clemenceau et al, 2000).

84

85 In this study, we use 3D seismic data to describe the architecture of mid-slope sections of the
86 Mahin and Avon channels and their associated terraces on the seafloor, which are located on
87 the western Nigerian continental slope (Fig. 1). The main aims of this paper are to (1) identify
88 controls on variation in morphology between the Avon and Mahin channels, (2) identify
89 controls on terrace distribution along the Mahin and Avon channels, and (3) understand the
90 evolution and stacking patterns of deposits within the terrace bodies and how they are linked
91 to the geomorphic evolution of the channels.

92

93

94 **2. Geological setting**

95 **2.1. Western slope, offshore Nigeria**

96 The study area is located on the continental slope offshore and northwest of the prominent
97 Niger Delta in the Gulf of Guinea along the western margin of Africa (Fig. 1). The western
98 offshore area of Nigeria is flanked by a 50-70 km wide shelf, with a shelf break at 150-200 m
99 water depth (Fig. 1). The continental slope extends from the shelf break to depths of 2000-
100 3000 m with an average slope gradient of 1.7° (Damuth, 1994).

101

102 Sediment is supplied to the offshore areas by numerous tidal channels which are present along
103 the edge of the Niger delta and probably overlie a distributive network of incised valleys that
104 transported sediment across the shelf to shelf-edge deltas and ultimately into deep water
105 canyon heads during periods of low Pleistocene sea level (Deptuck et al., 2007).

106

107 Strong longshore drift currents are active along the edge of the delta, which transport
108 sediment north up the western face of the Niger delta due to prevailing south-westerly winds
109 (Burke, 1972) (Fig. 1). Longshore drift also transports sediment from the Volta Delta
110 eastwards towards Mahin and along the coast of Lagos (Fig. 1) (Allen, 1965; Burke, 1972;
111 Anthony, 2015; Dada et al., 2016). A sandy beach is present at Lagos but abruptly terminates
112 100 km to the east, giving way to the Mahin mud beach (Fig. 1). Longshore drift northwards
113 of the Niger delta picks up increasing amounts of sand from successive distributaries but sand
114 only extends as far as the Benin River, north of which the shore is muddy. Though
115 unconfirmed, it may be that some of the sediment is directed down the numerous submarine
116 canyons along the Niger delta even during the current sea level high-stand. However, during
117 the late Pleistocene to early Holocene the sediment supply to these canyons was different and
118 ample sand, from large barrier complexes, was available on the shelf edge to be transported
119 down the canyons along the Niger delta front (Allen, 1970).

120

121 The continental margin off the Niger Delta is undergoing outward radial gravity deformation
122 caused by rapid seaward progradation (Damuth, 1994; Wu et al., 2015). This has created an
123 upper extensional zone of listric faults, a translational zone of mud diapirs and ridges beneath
124 the upper slope (Hovland et al., 1997), and a lower compressional zone of imbricated thrust
125 structures beneath the lower slope and rise (Damuth, 1994; Morley and Guerin, 1996;
126 Corredor et al., 2005) (Fig. 1). This dataset shows that these structures have heavily
127 influenced the deepwater depositional systems in this area.

128

129

130 **2.2. Study area**

131 The study area is located at water depths between 1350 and 2500 m in the mid-slope
132 translational to compressional zone and is about 30 km downslope from the shelf-edge (Fig.

133 1). The seafloor in this area, particularly in the up-dip parts, exhibits a number of structural
134 highs elevated up to 100 m above the surrounding seafloor (Fig. 2). These structural highs are
135 underpinned by mud diapirs in the subsurface, which resulted from rapid sedimentation rates
136 causing the mobilization of unstable mud (Burke, 1972; Doust and Omatsola, 1989; Damuth,
137 1994; Morley and Guerin, 1996). The topographic lows between the mud diapirs have
138 controlled the pathway of the Mahin and Avon channels (Fig. 2). Average slope gradients
139 vary between 0.5° and 2.5°, but are locally as high as 7° in areas adjacent to the structural
140 highs.

141

142 The Mahin and Avon channels converge in the downslope part of the study area (Fig. 3) and
143 the merged channel continues downslope and likely terminates in a submarine fan (Damuth
144 1994). Above the confluence the sinuosity and geomorphology of the two channels is
145 markedly different, while below the confluence the Avon channel has a similar sinuosity to
146 the Mahin channel above the confluence (Fig. 3). Within the confines of both channel-belts
147 numerous terraces are present but their distribution and characteristics vary significantly (Fig.
148 3).

149

150 **3. Data and methods**

151 **3.1. Seismic data**

152 Conventional 3D seismic mapping techniques were used to interpret the 3D seismic
153 reflectivity volumes in the study area. The 3D seismic survey covers an area of 4500 km² (45
154 x 100 km) with a bin size spacing of 12.5 m by 12.5 m (in-line and cross-line spacing) and a
155 sampling interval of 4 ms. The average frequency of the full-stack data is 50 Hz, resulting in a
156 vertical resolution of approximately 7.5 m (assuming the vertical resolution is a quarter of the
157 wavelength (Brown, 1999) and using the velocity of seawater (1500 m s⁻¹) (Jones et al.,
158 2010)) for the shallow section under consideration in the seismic profiles.

159

160 The data is SEG negative standard polarity where a positive reflection coefficient indicates an
161 increase in acoustic impedance and is represented by a trough, where the data is assumed to
162 be approximately zero phase. On the sections shown a peak is represented by red/yellow and a
163 trough is represented by blue. These assumptions of polarity and phase are supported by the
164 prominent seafloor event, which is represented by an approximately symmetrical trough.

165

166 The seafloor horizon was contoured in ESRI ArcGIS and the slope spatial analyst tool was
167 used to calculate the dip of each grid cell based on the elevation of its nearest neighbours to
168 generate a slope gradient map. Using the seafloor morphological expression each terrace
169 within the Mahin and Avon channel-belts was constrained by a polygon. The ESRI ArcGIS
170 aspect spatial analyst tool was used to create a map showing the spatial orientation and dip
171 direction of each terrace surface on the seafloor. Additionally, a horizon was interpreted at the
172 base of each terrace body within the confines of the terrace polygon (Fig. 4). This was
173 performed manually and based on the interpretation of the base of the channel-levee system,
174 and an understanding of the evolution of each individual terrace based on the seafloor
175 morphological expression. The base of each terrace body is composed of the deposit that
176 formed the terrace foundation within the channel-belt, on top of which turbidity currents may
177 have deposited sediment due to overspill from the adjacent channel.

178

179

180 **3.1.1. Seismic facies**

181 Regularly spaced cross-sections through the terrace bodies allowed the identification and
182 interpretation of six seismic facies, which helped determine the compositional elements
183 present in each terrace body. They have been classified according to the continuity and
184 amplitude of the reflections (Fig. 5). Due to the absence of lithological calibration from
185 seafloor cores, the interpretation of their sedimentary environments is based on comparison

186 with previously published seismic and sub-bottom reflection studies (Abreu et al., 2003;
187 Deptuck et al., 2003; Posamentier and Kolla, 2003; Gamberi and Marani, 2006; Catterall et
188 al., 2010; Jobe et al., 2015). While not available from the present study area, seafloor cores
189 obtained from channel-fills, internal levees and depositional terraces about 100 km to the
190 south east have been used to calibrate seismic facies comparable to those in our study (Jobe et
191 al., 2015), and we use these as a basis for lithological interpretation.

192

193 Facies A is characterized by high amplitude, apparently chaotic reflectors and occurs in axial
194 channel environments or at the base of terraces (Abreu et al., 2003; Deptuck et al., 2003;
195 Posamentier and Kolla, 2003; Jobe et al., 2015); it is interpreted as coarse grained channel-
196 fill. Facies B is characterized by continuous straight reflectors that converge and dim away
197 from the adjacent channels and are interpreted as the external levees that confine both the
198 Avon and Mahin channel-belts (Hübscher et al., 1997; Deptuck et al., 2003; Nakajima and
199 Kneller, 2013). Facies C is composed of continuous straight and parallel reflectors that can
200 vary in amplitude and are interpreted to represent heterolithic strata that form either
201 depositional terraces (*sensu* Hansen et al., 2015) or internal levees (Deptuck et al., 2003;
202 Babonneau et al., 2004; Jobe et al., 2015). Facies D is characterized by straight continuous
203 reflectors which are inclined towards the channel and represent heterolithic slide blocks that
204 have slid into the channel (Deptuck et al., 2003; Babonneau et al., 2004). Facies E is
205 composed of chaotic reflectors with varying amplitude representing slumps that may have
206 originated from the side of the channels or from further up-dip (Deptuck et al., 2003;
207 Posamentier and Martinsen, 2011; Ortiz-Karpf et al., 2015). Facies F is characterized by flat
208 continuous reflectors with consistent amplitudes and represents the lobe deposit prior to
209 incision of the Avon and palaeo-Avon channel (Posamentier, 2003; Posamentier and Kolla,
210 2003; Gamberi and Rovere, 2011) (Fig. 5).

211 3.1.2. *Spectral decomposition*

212 Spectral decomposition combined with RGB (Red Green Blue) colour blending of selected
213 discrete frequency bands (McArdle and Ackers, 2012) is an imaging technique used to
214 highlight subtle features and geometries in seismic data. Spectral decomposition was used to
215 image subtle frequency variations at the base of each terrace to identify depositional elements
216 whose scale is below the average seismic resolution of the full-stack data. Since the peak
217 frequency response for a specific seismic interval varies with its temporal thickness (with
218 higher frequencies imaging thinner geological features), visually blending a number of
219 discrete frequency bands can reveal subtle changes in thickness. Therefore sedimentological
220 features can be identified that are otherwise difficult to detect on conventional amplitude
221 maps.

222

223 The seismic survey was decomposed into three amplitude-response volumes, each with a
224 different average frequency: 38.33 Hz, 47.63 Hz and 68.37 Hz. The average frequencies of
225 the decomposed volumes were selected to span the wide frequency distribution of the full-
226 stack data, which was determined from the terrace base horizons. The selected frequencies
227 were far spaced in order to separate and highlight geological features with a distinct frequency
228 content. The frequency volumes were then assigned separate RGB colours (38.33 Hz = red,
229 47.63 Hz = green, 68.37 Hz = blue) and merged together to produce a frequency colour blend.
230 The colour hue in the colour blend reflects the average frequency around the studied horizon
231 and the brightness of the colour represents the amplitude of the seismic signal. Furthermore,
232 the blended frequency volumes were calculated using different frequency filter lengths (i.e.
233 time intervals) and therefore the visualized colour blend is a composite of three time windows
234 centred around the chosen horizon. Typically the maps provide more detail than seismic
235 profiles because the observed geological features are much wider than they are thick while
236 both the vertical and horizontal resolution remains the same.

237

238 **3.2. Channel and terrace measurements**

239 In addition to seismic interpretation, the morphology of the channels and terraces at the
240 seafloor was parameterized by taking measurements from profiles located perpendicular to the
241 local orientation of the channels (spaced every 1 to 6 km along the channel thalweg). The zero
242 point for channel length measurements along the thalweg is located at the easternmost
243 (upstream) limit of the Mahin channel in the study area. The measurements in the Avon
244 channel are referenced to the measured Mahin channel distance at the confluence and hence
245 start at 42 km (Fig. 3b). The measurements permit comparisons to be made between two
246 adjacent channels that cross morphologically similar slope segments, but with substantially
247 different morphological characteristics. All measurements were taken in metres by depth
248 converting the seafloor map using an average seawater velocity of 1500 m s^{-1} (Jones et al.,
249 2010).

250

251 The Mahin and upper Avon channels were subdivided into three sections with the start and
252 end points of each section determined by significant changes in channel gradient and
253 sinuosity. The lower Avon channel was not further subdivided and is labelled as section 4.
254 The sections allowed quantitative comparison of channel and terrace properties along the
255 channels.

256

257 The measured parameters from these profiles include the channel-belt width (defined as the
258 perpendicular width between external levee crests) (Fig. 6), channel width (defined as the
259 perpendicular width between the channel-facing terrace edges, where terraces are present)
260 (Fig. 6), and channel-belt depth (defined as the depth between highest external levee crest and
261 the thalweg i.e. the deepest point in channel). The channel margin gradients were calculated
262 from the perpendicular channel profiles by calculating the gradient along the steepest section
263 of the channel margin. The down-dip gradient of both channels was calculated using the
264 thalweg depths and measuring the change in depth per kilometre along the thalweg (Fig. 7a,
265 b). A 4 km moving average of the gradient was also calculated to reduce local irregularities

(Fig. 7 b.1, b.2). Variations of the open slope gradient adjacent to the Mahin and Avon channels were also measured in order to determine whether differences in morphology of the two channels might be related to differences in the gradient of the adjacent slope. The slopes have an average of 0.85° . The sinuosity index (formula shown in Figure 6), in which a straight feature has an index of 1, was calculated for both channels over a 6 km sliding window (Fig. 6, 7 c.1, c.2).

For each terrace, the terrace area, perimeter, cross-stream width, downstream length, circularity, and height of terrace above the thalweg was measured at the seafloor (Fig. 6). The circularity was measured using the formula shown in Figure 6 (Friel, 2000).

4. Channel geomorphology

4.1. Mahin channel geomorphology

The present day expression of the 90 km long Mahin channel section that lies within the study area is a leveed channel-belt with external levees up to a maximum of 5 km wide. It is a highly sinuous channel that has some spatial variations in its cross-sectional (relief, symmetry, width) and planform geometry (more or less sinuous) (Fig. 7). It is characterized by the presence of terraces on almost every channel bend (34 terraces in total), which locally occur in stepped pairs representing multiple generations of terraces at different heights within the channel-belt (Fig. 4b). The channel mostly has a symmetrical V-shaped cross-section with a difference in conjugate levee crest height of less than 20 m (Fig. 8a, b). The channel is most asymmetric at bend apexes due to outer channel bend erosion and deposition focused on the inner bends (Fig. 8b). Channel margins have gradients mostly between 14° and 26° but can be as low as 10° on inner channel bends (Fig.8b).

292 The Mahin channel is divided into three sections. Up-dip in sections 1 and 2 the channel-levee
293 system is almost completely confined within the structural low created by the mud diapirs and
294 the levees are lapping onto these structural highs (Figs. 2b, 9a, b). Down-dip in section 3 the
295 topographic effect of the mud diapirs is less, and the system is consequently less confined
296 (Fig. 9c, d).

297

298 Section 1 is the steepest part of the Mahin channel with a channel thalweg gradient of 0.7° ,
299 and an average sinuosity of 1.57 (Table 1, Fig. 7 a.1 & c.1). In this section the channel-belt is
300 very wide (on average 3454 m wide, see Table 1), with multiple generations of terraces. The
301 channel thalweg gradient and channel-belt width decrease throughout section 2 and 3 to 0.36°
302 and 2214 m respectively (Table 1). Sinuosity is highest in section 2, at 2.37, and decreases to
303 1.75 in section 3 (Table 1, Fig. 7 c.1). At the start (between 72.5 and 73.6 km) and end
304 (between 88.6 and 90.7 km) of section 3 the channel gradient reverses with the channel floor
305 rising up by several metres (Fig. 7 a.1). At the start of this section this is due to sediments that
306 have collapsed into the channel from the inner external levee. At the end of section 3
307 (between 88.6 and 90.7 km) the channel thalweg rises just before it converges with the Avon
308 channel at which point there is a 30 m drop into the Avon channel (Fig. 8c).

309

310 **4.2. Upper Avon channel geomorphology**

311 The 49 km long Avon channel also lies within a leveed channel-belt (with levees up to 3 km
312 wide) but it is less sinuous (1.16 on average) compared to the Mahin channel (1.9 on average)
313 (Fig. 7 c.2) and has fewer terraces (18 compared to 34 in the Mahin channel-belt). The cross-
314 sectional form changes frequently from V-shaped to U-shaped and can be strongly
315 asymmetrical at channel bends (Fig. 8b). The steepness of the channel margins is mostly
316 between 18° and 26° but can be as low as 9° on inner channel bends (Fig. 8b).

317

318 The upper Avon channel is divided into three sections. Similarly to the Mahin channel, in
319 section 1 the channel-levee system is completely confined within the structural low created
320 between the mud diapirs with the levees lapping onto the structural highs (Fig. 9a, b). Down-
321 dip in sections 2 and 3 the upper Avon channel becomes less confined due to the reduced
322 topographic effects of the mud diapirs (Fig. 9c, d).

323

324 Throughout sections 1 to 3 the sinuosity is mostly consistent at around 1.1 to 1.2 (Table 1)
325 with the exception of a moderate sinuosity increase at the transition between sections 1 to 2
326 where the upper Avon channel makes a sharp bend (Figs. 3a and 7 c.2). This is likely
327 associated with a prominent topographic high obstructing the pre-bend channel direction (Fig.
328 3). The thalweg gradient (Table 1, Fig. 7 b.2) stays constant around 0.8° throughout section 1
329 and part of section 2 but increases sharply in the last part of section 2 to 1.6° , which is
330 interpreted as a knickpoint (Fig. 7 a.2 b.2). This is associated with an abrupt increase in the
331 channel-belt depth and width from 83 m to 184 m and 1469 m to 2484 m respectively (Table
332 1). The thalweg of the upper Avon channel appears to be smooth in section 1 to 3 where it is
333 straight. However, in the lower Avon channel, after the confluence with the Mahin channel,
334 the channel floor becomes more undulating, showing frequent bedforms, which are not seen
335 in the Mahin channel, with up to 10 m of relief, together with an increase in sinuosity (Fig. 7
336 a.2).

337

338 **4.3. Lower Avon channel geomorphology (after confluence)**

339 The 39 km long section of the lower Avon channel has a similar sinuosity to the Mahin
340 channel (1.76 in the lower Avon compared to 1.9 in the Mahin channel), which marks a
341 distinct change between the upper and lower parts of the Avon channel (1.16 in the lower
342 Avon vs. 1.76 in the upper Avon) (Table 1, Fig. 7c). The lower Avon channel-belt is confined
343 by levees and is characterized by the presence of terraces along the entire channel length (19
344 terraces in total). It shows an overall increase in channel-belt depth (201 m on average) and

channel-belt width (3281 m on average) compared to the upper Avon (on average 102 m deep and 1978 m wide) and Mahin channels (on average 171 m deep and 2740 m wide) (Table 1). Some straighter channel sections seem to be related to a decrease in thalweg gradient (Fig. 7 b, c). The cross-sectional relief is predominantly V-shaped, but again shows asymmetry at channel bends (Fig. 8b).

350

351

352

353 **5. Terrace morphometric characteristics**

354 **5.1. Mahin channel terraces**

Numerous terraces are present along almost the entire length of the Mahin channel-belt (34 in total). Terraces occur both on the inside and outside bends of the channel with their area often extending across two channel bends. Some terraces occur as stepped pairs, where two adjacent terraces are at different heights above the channel. They are generally topographically flat in parts but are dominated by local topographic irregularities with different terrace sections dipping in various direction (Fig. 10). Where terraces are directly next to the channel they can be inclined towards the channel. Some terraces, in section 1, are characterized by topographic highs in their centre (Figs. 3, 4, 10). Some terraces also show a distinct topographic elevation at the channel-facing side which tapers away from the channel (Fig. 10). All terraces have a greater downstream length than width (Table 2, Fig. 11). The terrace characteristics change frequently down channel, with an overall decrease in terrace area, length and width decreasing between sections 1 and 3 (Table 2, Fig. 11). In contrast, the terrace circularity and height stay comparably more constant between section 1 and 3 (Table 2). The number of terraces is greatest in section 3 (12 terraces present in section 3) where they also have the smallest area (Table 2, Fig. 11).

370

371 **5.2. Upper Avon channel terraces**

372 The abundance of terraces along the upper Avon channel-belt is less than the Mahin channel-
373 belt (18 compared to 34 terraces in the Mahin channel-belt) (Table 2). Terraces are more
374 abundant on the inside of channel bends and most commonly occur as stepped pairs at
375 different heights above the channel (Fig. 3). In sections 1 and 2, all terraces have topographic
376 irregularities and almost all of them dip downslope by up to 5° towards the SW (Fig. 10). In
377 section 3, which is below the knickpoint in the upper Avon channel (Figs. 3 and 7 a.2, b.2),
378 the terraces are smoother and do not dip downslope. They are almost 100 m higher above the
379 adjacent channel than the terraces in sections 1 and 2 of the upper Avon channel (Table 2,
380 Fig. 11). The majority of the terraces have a long downstream length and are bigger and less
381 circular than the Mahin terraces. Overall, the terrace area, width and length decrease
382 downslope (Table 2, Fig. 11).

383

384 **5.3. Lower Avon channel terraces (after confluence)**

385 Below the confluence terraces are again present both on the inside and outside of channel
386 bends and sometimes occur as stepped pairs (19 terraces in total). Some of the terraces are
387 topographically flat but the majority have local topographic irregularities (100 to 200 m in
388 scale) with different parts of the terrace dipping in various directions (Fig. 10). On average
389 the terraces are bigger than the Mahin terraces, have almost the same area as the upper Avon
390 terraces but are significantly wider than both the Mahin and upper Avon terraces (Table 2,
391 Fig. 11). The circularity is similar to the Mahin terraces and hence greater than the upper
392 Avon terraces. The terrace height above the thalweg is on average 26 m higher than the Mahin
393 terraces and 93 m higher than the upper Avon terraces in section 1 and 2. They are on average
394 almost equivalent in height to the upper Avon terraces in section 3 (Table 2). Within this
395 section the width and area of the terraces increases downslope (Table 2, Fig. 11).

396

397

398 **6. Terrace body compositional elements**

399 Interpretation of seismic facies (Fig. 5) within the terrace body, together with spectral
400 decomposition slices at the terrace base (Fig. 12) and throughout the terrace body, were used
401 to assist in the quantification of the compositional elements that make up each terrace body
402 along the Mahin and Avon channel-belts, and to reveal terrace forming processes.

403

404 **6.1. Spectral decomposition of terrace base horizons**

405 **6.1.1. Mahin terrace bases**

406 At the base of many of the terraces of the Mahin channel, several older channel-fills can be
407 recognized to make up the foundation of the terrace body. Some of them show the location of
408 older channel bends that have been cut-off. These are particularly obvious in Figure 12b
409 where the remnant channel bend fills can be seen at the base of four terraces (process
410 illustrated in Fig. 13c, d); on the seafloor these terraces have a distinct topographic high at
411 their centre which is the remnant topographic high in the centre of the older channel bend.
412 Several terrace bases show lateral accretion packages (LAPs) that may indicate progressive
413 channel migration as channel sinuosity changed (Fig. 12b). Some areas of the spectral
414 decomposition slices look incoherent and chaotic and are interpreted to be sediment that has
415 slumped towards the channel (Fig. 12c).

416

417 **6.1.2. Upper Avon terrace bases**

418 Regional seismic interpretation indicates that the current upper Avon channel is underlain by
419 older lobe and channel deposits (Fig. 13a, b). The terrace bases in channel section 1 are
420 dominated by older channel-fills making up the foundation of the terrace body (Fig. 12d),
421 which belong to the palaeo-Avon channel whose remnant is visible on the seafloor between
422 the Mahin and Avon channels (Figs. 3, 9, 13, 14). In channel section 2, the chute channel
423 remnants interpreted to have been present in the overbank areas at the initiation of the current

424 Avon channel are preserved at the terrace bases (Figs. 12d, 13b). In section 3, the terrace
425 bases are dominated by older channel-fill deposited in channel bend cut-offs similar to the
426 Mahin terraces into which the Avon channel is incising (Fig. 13c).

427

428 **6.1.3. Lower Avon terrace bases (after confluence)**

429 The base of the terraces in the lower Avon channel show an abundance of older channel-fills
430 imaged by the spectral decomposition slices (Fig 12e). LAPs and multiple generations of
431 older channel-fills can be seen, which indicates the lateral migration of this channel over time.
432 More complex channel migration patterns can be seen in the most basinward terrace of the
433 lower Avon channel, where several segments of channel-fill overprint one another (Fig. 12e).
434 Some more chaotic and less coherent areas of the spectral decomposition slices suggest
435 slumped material from the channel-belt margin that has moved towards the channel (Figs.
436 12e, 13d). It is unclear whether this slumped material is composed of coherent blocks or more
437 disaggregated strata.

438

439 **6.2. Volumetric assessment of terrace compositional elements**

440 By using both the seismic facies and the spectral decomposition slices at the base and within
441 the body of each terrace, it is possible to interpret the compositional elements that make up
442 the terrace bodies along the Mahin and Avon channel-belts. Three different compositional
443 elements were differentiated: Channel-fill, slumps and heterolithics (Fig. 14). Heterolithics
444 are a combination of seismic facies C and D (Fig. 5), which were grouped together for the
445 purpose of volume calculations for the terrace body because they likely underwent similar
446 sedimentological processes during initial deposition. Facies B represents the external levees
447 and is not present in the terrace bodies unless part of the external levee sediment slid into the
448 channel-belt (in which case it is classified as facies D). For each terrace body, the volume of
449 each compositional element was calculated (Fig. 14).

450

451 **6.2.1. Mahin channel terraces**

452 Almost all of the terrace bodies along the Mahin channel-belt are partially composed of
453 remnant channel-fill facies (up to 50% with an average of 29%) (Fig. 14), which can be seen
454 in the spectral decomposition slices from the base of each terrace (Fig. 12). The volume of
455 heterolithics in the terrace bodies varies down dip but is commonly the largest component of
456 each terrace body (between 36% and 83%) (Fig. 14). These heterolithics would have been
457 deposited by overbank flow after the initial elevated surface that created the terrace
458 foundation was established in the channel-belt. The proportion of slumps changes
459 significantly down dip and it only occurs in three terrace bodies in sections 1 and 2 whereas it
460 is dominant in section 3 of the Mahin channel (Fig. 14).

461

462 **6.2.2. Upper Avon channel terraces**

463 The terrace bodies of the upper Avon channel-belt consist mainly of heterolithic overbank
464 deposits (average 93%) related to the current Avon channel. They rest directly on top of
465 palaeo-Avon channel and lobe deposits belonging to older stratigraphic units of the Avon
466 system. Where overbank deposits overlie lobe deposits, the terrace bases show signs of wide
467 chute channel-like features, which are too subtle to be recognized in seismic profiles (Figs.
468 12c, 13b). Part of the first upper Avon terrace body is composed of slump deposits (Fig. 14).

469

470 **6.2.3. Lower Avon channel terraces (after confluence)**

471 In the lower Avon channel-belt, the percentage of channel-fill, for each terrace body is
472 slightly higher than in the Mahin channel (average 34% compared to 29% in the Mahin
473 channel) and slumps are still common at the start of this channel section. Three terrace bodies
474 consist of slumps and heterolithics only (Fig. 14).

475

476 **7. Discussion**

477 **7.1. Controls on morphological differences between the Mahin and Avon channels**

478 The main morphological difference between the Mahin and Avon channels above the
479 confluence is their sinuosity. The longitudinal evolution of the sinuosity along the Mahin
480 channel shows a high and non-linear variability, while there is less overall variability in the
481 upper Avon channel sinuosity as it is predominantly straight (Figs. 3, 7c). Sinuosity in
482 submarine channels is suggested to be controlled by a number of factors such as the slope
483 gradient, flow interaction with seafloor topography, the calibre of sediment the channel is
484 transporting and eroding into, and channel maturity (Clark et al., 1992; Nanson and Croke,
485 1992; Reading and Richards, 1994; Clark and Pickering, 1996; Pirmez et al., 2000; Schumm,
486 2005; Peakall et al., 2007; Kane et al., 2008; Kleinhans, 2010; Straub et al., 2011; Peakall et
487 al., 2012; Janocko et al., 2013b; Sylvester et al., 2013; Sylvester and Covault, 2016). Many
488 factors probably affect the sinuosity of the channels in the study area but it is likely that one
489 factor is dominant. These will be discussed below.

490

491 Variations in river channel sinuosity have been attributed to slope gradient in flume tank
492 models of open channels (Schumm and Khan, 1972; Schumm et al., 1972) and in modern
493 seafloor studies such as those of the Indus, Rhone, Mississippi, Monterey, and Amazon
494 submarine channels (Clark et al., 1992; Peakall et al., 2012). However, in the study area the
495 slope gradient measured on a line adjacent and beyond the external levees to both channels
496 shows that the gradient is similar and has an average of 0.85° . Given the substantially
497 different sinuosities of the two channels, slope gradient is therefore unlikely to be the
498 controlling factor.

499

500 Flow interaction with the seafloor topography can also impact channel sinuosity. Seafloor
501 topography can be either tectonically driven or created by the presence of other sedimentary

502 depositional bodies such as other channels, lobes or mass transport deposits (Reading and
503 Richards, 1994; Pirmez et al., 2000; Babonneau et al., 2002; Adeogba et al., 2005; Straub et
504 al., 2008; Catterall et al., 2010). In the study area the seafloor topography is largely shaped by
505 the underlying mud diapirs, which control the location of the Mahin and Avon channels. In
506 places diapirism has caused local increases in gradient, which has resulted in localized
507 straightening of the channel (Fig. 2a). While mud diapirism certainly affected the channel
508 sinuosity on a local scale, the topographic irregularities of the area occupied by the Mahin and
509 Avon channels are similar, making it unlikely that the differing sinuosity is driven by seafloor
510 topography.

511

512 Clark et al. (1992) interpret the grain size of sediment transported in channels to be an
513 important factor in determining the channel morphology, where channels with a coarser grain
514 size range tend towards lower sinuosities than those dominated by a finer grain size range.
515 Additionally, the cohesiveness of the substrate into which the channel is incising may impact
516 channel sinuosity, where erosion into a muddy substrate results in more sinuous channels than
517 erosion into more easily erodible sandy substrate, resulting in straighter channels (Nanson and
518 Croke, 1992; Schumm, 2005; Peakall et al., 2007; Kleinhans, 2010; McHargue et al., 2011;
519 Verhagen et al., 2013). Figure 15 shows a RMS amplitude map of the study area that
520 illustrates the striking difference in amplitudes along the Avon and Mahin channels above the
521 confluence. In this case the RMS amplitude has been attributed to the amount of coarse
522 grained sediment present in the interval of amplitude extraction. The RMS amplitudes suggest
523 that the Avon channel is fed by turbidity currents containing significantly coarser grained
524 material than the Mahin channel. Additionally, the lobate shape of the amplitudes shows that
525 the Avon channel is incising into older, probably sandy, palaeo-Avon lobe deposits. These
526 observations agree with the suggestion that channels carrying and eroding into coarser grained
527 sediments generally have lower sinuosities than those carrying and eroding into finer-grained
528 sediments (Clark et al., 1992; Peakall et al., 2007; Peakall et al., 2012).

529

530 Evidence for the difference in the sediment supply systems to the Avon and Mahin channels
531 can be traced back to the Late Pleistocene and earlier Holocene. In the Pleistocene (20,000
532 years ago) the Benin River extended across the shelf to the Mahin canyon (Burke, 1972),
533 which may have been responsible for the initiation of the Mahin channel. The sediment that
534 was transported in the channel then is unknown but at the maximum extend of the last
535 transgression, around 7000 years ago, old beach ridges, also referred to as the ‘older sands’ by
536 Allen (1964), were present at the head of the Avon canyon but these are more limited at the
537 head of the Mahin canyon (Allen, 1964). Additionally, at the present day the coastline at the
538 head of the Avon canyon is sandy while it is muddy at the head of the Mahin canyon, due to
539 the dominant longshore drift patterns (forming the Mahin mud beach, refer to Fig. 1). This
540 current apparent absence of a sand supply to the Mahin canyon may either be due to
541 reworking and transport of sediment away from the canyon head by waves and tides due to
542 the prevailing south-westerly wind, or due to the capture of the sand supplied by eastward
543 longshore drift by the Avon canyon (which cuts further landward than the Mahin canyon) or
544 by any of the numerous other canyons present along the Niger shelf edge (Allen, 1964; Burke,
545 1972) (Fig. 1). Hence, the Avon channel may be receiving coarse-grained sediment even
546 during the current sea level highstand and sediment supply and cohesiveness of the substrate
547 are interpreted to be contributing factors to the difference in sinuosity between the Avon and
548 Mahin channels.

549

550 Channel sinuosity has also been attributed to channel maturity where younger channels are
551 straight and become more sinuous over time (Peakall et al., 2000). This may be partly due to
552 flows becoming smaller and muddier through time as sediment supply is reduced, ultimately
553 causing an increase in sinuosity (Janocko et al., 2013b), or simply due to the time taken to
554 establish an equilibrium sinuosity. Though there is no direct evidence of the relative ages of
555 the Mahin and Avon channels, we interpret the Avon channel to be younger. Evidence for this

556 is the presence of the palaeo-Avon channel (Fig. 3, 9, 13, 15), which has eroded into the
557 palaeo-Avon lobe deposit and then avulsed to form the current Avon channel. Evidence for
558 the pre-existence of the Mahin channel is that below the confluence, the lower part of the
559 Avon channel adopts a similar sinuosity to the Mahin channel and we speculate that the
560 sinuosity of the lower Avon channel was created by the pre-existing Mahin channel. The 30 m
561 drop (Fig. 8c) at the mouth of the Mahin channel suggests that the Mahin channel is no longer
562 active, as otherwise this drop would have been eroded and smoothed by turbidity currents
563 active in the channel. The Mahin channel inactivity may also be the reason for the abrupt
564 brightening of the RMS amplitudes in and around the Mahin and Avon confluence. Channel
565 confluences have been documented mainly on seafloor datasets e.g. from the Espirito Santo
566 Basin, offshore Brazil (Gamboa et al., 2012) and the Tyrrhenian Sea, offshore Sicily
567 (Gamberi et al., 2013). They are commonly associated with a knickpoint, which can also be
568 seen between sections 2 and 3 of the upper Avon channel (Fig 7 a.2). This may be a result of
569 the straight Avon channel having to navigate the previously eroded sinuosity created by the
570 Mahin channel, which disrupted the equilibrium profile of the converging upper Avon
571 channel. To regain the equilibrium profile the knickpoint now is migrating up the Avon
572 channel causing increased channel erosion (and thus channel depth) below the knickpoint.
573 Knickpoint migration has commonly been documented from deepwater channel studies
574 (Adeogba et al., 2005; Antobreh and Krastel, 2006; Catteral et al., 2010; Gamberi and Rovere,
575 2011).

576

577 This suggests that, in the case of the Mahin and Avon channels, channel maturity is likely the
578 main reason for the difference in channel sinuosity. It is important to recognize the influence
579 of channel morphology when analysing terrace distribution and composition.

580

581

582

583 **7.2. How are the terraces formed?**

584 The spectral decomposition slices of the base of the terraces have shown sedimentological
585 detail that allows us to interpret how terraces were initially formed, which is reflected in the
586 composition of the terrace bodies. Four terrace forming processes can be observed in the
587 study area (detailed description of each process is summarized in Hansen et al. 2015) but in
588 some cases more than one of these processes has occurred. The dominant process for the
589 formation of each terrace is shown in Figure 16.

590

591 The majority of the terraces in the Mahin channel-belt and the upper Avon channel-belt are
592 formed by entrenchment. Entrenchment of a channel into older deposits can occur solely by
593 vertical incision, or with a component of lateral migration by the process referred to as
594 punctuated channel migration (Maier et al., 2012). The sediments below these elevated
595 surfaces will be composed of whatever the channel is incising into e.g. older channel-fills,
596 lobes, slope deposits, levee deposits. In the Mahin channel-belt, the formation of these
597 terraces is interpreted to be mainly due to the abrupt lateral migration of the channel as the
598 sinuosity of the channel changed over time, either as a response to local topography or due to
599 sediment collapse into the channel resulting in a shift of the channel position. Many of the
600 terrace bases are therefore composed of older channel-fill (Fig. 12, 13d, 14). In the upper
601 Avon channel-belt, the terraces have been incised into previously deposited channel and lobe
602 deposits by vertical entrenchment of the current Avon channel (Fig. 13a, b, c, 14). The
603 incision of channels into lobes has been documented from many turbidite systems such as the
604 Golo turbidite system, Corsica (Gervais et al., 2006) and the Karoo Basin in South Africa
605 (Brunt et al., 2013). This may be associated with some externally imposed forcing such as a
606 change in relative sea level and a resultant progradation of the whole deepwater depositional
607 system (Chen and Hiscott, 1999; Hodgson et al., 2016).

608

609 Point bar accretion is a result of continuous channel bend growth, and has been convincingly
610 demonstrated in only a few cases in turbidite systems (Abreu et al., 2003; Deptuck et al.,
611 2007; Dykstra and Kneller, 2009; Janocko et al., 2013b). True point bar accretion is
612 challenging to image on this seismic data and is not always distinguishable from punctuated
613 channel migration, but can be differentiated by progressive bend widening shown by scroll
614 bars. Figure 13d shows scenarios of how point bar accretion deposits may occur at terrace
615 bases. Point bar accretion can be interpreted at one place in section 1 of the Mahin channel
616 and in several places in the lower Avon channel (Fig. 16) due to the presence of scroll bars in
617 the spectral decomposition images (referred to as LAPs in Fig. 12).

618

619 Both the Mahin channel-belt and the lower Avon channel-belt have a number of terraces with
620 bases interpreted to have formed by channel bend cut-off (Fig. 16). Channel bend cut-offs are
621 a common phenomenon in highly sinuous, laterally migrating channels and result from the
622 progressive accentuation of bend curvature and eventual neck cut-off (McHargue, 1991;
623 Cronin, 1995; O'Connell et al., 1995; Mayall and Stewart, 2000; Kolla et al., 2001; Deptuck
624 et al., 2003; Samuel et al., 2003; Janocko et al., 2013b). Figure 13c and d show how channel
625 bend cut-offs can create elevated areas within the channel-belt. In some cases, like the most
626 basinward terrace in the lower Avon channel, multiple generations of channel bend cut-offs
627 can be seen at the base of the terrace body (Fig. 12e). This indicates a complicated history of
628 the lateral switching of the channel similar to what has been reported in Mchargue et al
629 (2011) and Hodgson et al (2011). The terrace bases created by channel bend cut-offs are the
630 largest in area (1.5km^2 on average) and have on average the most circular geometry.

631

632 Some of the smaller terraces along the Mahin and lower Avon channel-belts are interpreted to
633 have formed by margin failure. Figure 13d shows how margin failure deposits may occur at
634 terrace bases. They occur towards the end of the Mahin channel and at the start of the lower
635 Avon channel (Fig. 14), which may be a result of the increase in channel gradient at these

636 locations (Fig. 7b), causing increased channel erosion and creating steep margins prone to
637 collapse.

638

639 There is a direct link between the behaviour of the channel, which is in part controlled by the
640 channel maturity, and the terrace forming processes. It is noticeable how many more terraces
641 are present within the Mahin and lower Avon channel-belts than the upper Avon channel-belt.
642 Lateral channel movement and channel bend cut-offs are terrace forming processes that are
643 both also related to the evolution of the channel sinuosity, suggesting that terraces are more
644 abundant next to sinuous rather than straighter channels. In addition, terraces in the Mahin
645 channel-belt are on average 26 m lower above the thalweg than the terraces in the lower Avon
646 channel-belt (Table 2). Hence it seems that the knickpoint migration up the Avon channel as a
647 result of the upper Avon channel merging with the pre-existing Mahin channel has resulted in
648 roughly 30 m more channel incision in the lower Avon channel which agrees well with the 30
649 m drop observed at the Mahin channel mouth (Fig. 8c). The influence of the knickpoint is
650 most pronounced when comparing the height of the terraces above the thalweg between
651 upstream (sections 1 and 2) and downstream of the knickpoint (sections 3 and 4) within the
652 Avon channel-belt, which shows an almost 100 m height difference. In addition the increase
653 in occurrence of terraces originally formed by margin failure shows that the additional
654 channel incision due to the knickpoint migration can lead to channel margin instability. This
655 implies that terrace forming processes can be dependent on localized channel behaviour (even
656 as local as one channel bend), but can also be impacted by regional factors such as re-grading
657 of the channel equilibrium profile due to channel confluences or regional tectonics (Clark and
658 Cartwright, 2009).

659

660 **7.3. Terrace body compositional elements and stacking patterns**

661 By combining the spectral decomposition slices at the bases of each terrace and quantifying
662 the seismic facies within each terrace body a model of terrace evolution can be built. Stacking

663 patterns identified in terrace bodies can then be compared to stacking patterns commonly
664 observed in other thin-bedded depositional environments, which can assist in the
665 identification of these deposits in outcrop and the subsurface.
666
667 The most typical stacking pattern of deposits making up terrace bodies observed in the Mahin
668 and Avon terraces is channel-fill at the base with heterolithics deposited on top (Fig. 17
669 pseudo-core A). These heterolithics may at times occur as part of external levee slide blocks
670 that have slid into the channel-belt. The thickness of the channel-fill deposit at the base varies
671 but can be up to several 10s of metres thick, especially when multiple parts of older channel-
672 fill are preserved (Fig. 12e). The thickness of the heterolithics above the channel-fill deposits
673 depends on the amount of overspill and flow stripping the terrace has experienced, which is
674 related to the height of the terrace above the adjacent thalweg and the height and frequency of
675 the turbidity currents. In the Mahin and Avon terrace bodies the thickness of the heterolithics
676 varies between 20 – 100 m. The heterolithics within the terrace bodies predominantly form
677 depositional terraces (i.e. flat deposits, Hansen et al., 2015) but some internal levees are
678 recognized which are described in section 7.4. This stacking pattern and deposit thickness is
679 similar to what is described from the C/D ridge outcrop in the Karoo Basin, South Africa
680 (Hodgson et al., 2011). Another common stacking pattern is channel-fill at the base followed
681 by slump deposit and heterolithics above (Fig. 17 pseudo-core B). This sort of stacking
682 pattern would have developed initially by lateral channel migration or channel bend cut-offs,
683 followed by margin collapse due to oversteepening of the channel margin. On top of this
684 elevated area thin-bedded turbidites then deposit by overspill and flow stripping from the
685 adjacent channel (Fig. 13d). These stacking patterns can be a vital piece of information when
686 trying to identify heterolithic depositional environments in outcrop and the subsurface.

687

688

689

690 **7.4. Internal levees**

691 While a clear distinction exists between internal levees (*sensu* Kane and Hodgson, 2011) and
692 depositional terraces, it is likely that they form a continuum and that true internal levee and
693 depositional terrace examples rarely occur (Hansen et al., 2015). This hypothesis is further
694 strengthened by the varying dip directions of parts of each terrace surface seen in Figure 10,
695 which emphasizes the complexity of these deposits. By definition, internal levees can be
696 recognized by their wedge shape and form when the overspilling current has enough space to
697 decelerate and deposit the majority of its sediment before it hits a confining surface. The
698 orientation of the internal levee will depend on the dominant flow momentum of the
699 overspilling current, which most commonly is down channel. Hence, the wedge shape of the
700 internal levee may be orientated oblique to the channel direction. True internal levees were
701 identified on parts of seven terraces along the Mahin and Avon channel-belts (Fig. 18) but do
702 not occur in particularly unique locations adjacent to the channel that might imply a
703 relationship with the geometry of the channel. The dominant influence on internal levee
704 distribution seems to be the terrace height and width (Fig. 18). The height and width of the
705 terraces where internal levees occur do not stand out significantly from the background trend
706 but they are generally on the higher terraces with an average terrace width (Fig. 18). We
707 interpret that increased height of the terrace reduces the amount of sediment that is able to
708 overspill, and hence requires less space to decelerate and deposit most of its sediment before
709 hitting the confinement of the channel-belt.

710

711 While there does not seem to be a specific terrace width required to form an internal levee, the
712 terrace width impacts where on the terrace the internal levee develops. For example on terrace
713 number 30, which does not dip downslope, only the widest part of the terrace is covered by an
714 internal levee while on the narrower sections the deposits are either flat, dip downslope or
715 towards the channel (see Fig. 10 insert in Fig 18). This reinforces the hypothesis that there is
716 in fact a continuum between internal levee and depositional terrace deposits, and that the

717 amount of overspilling sediment and the space available adjacent to the channel determines
718 which type of deposit is able to form (Hansen et al., 2015).

719

720 **7.5. Terrace body preservation potential**

721 This study shows that terrace bodies are composed of a number of different types of deposits
722 including channel-fill or slumps at the base, and usually followed by heterolithics above.

723 Terraces form significant features next to submarine channels on the seafloor and can be over
724 1500 m wide and up to 7 km long in a downstream direction. They have been interpreted from
725 outcrop in the Cap Enragé Formation in Canada (Hein and Walker, 1982) as well as in
726 numerous shallow seismic examples (Deptuck et al., 2003; Babonneau et al., 2004),
727 suggesting that they can be preserved in the subsurface.

728

729 The channel systems studied here suggest that the preservation potential of terrace bodies is
730 directly related to channel sinuosity and the rate of channel incision. In the Mahin and lower
731 Avon channel-belts, the large number of terraces formed by the cut-off of sinuous channel
732 bends seem to have been preserved by the incision of the late Mahin and lower Avon
733 channels. Incision has been shown to inhibit lateral migration in both submarine (Janocko et
734 al., 2013a) and fluvial (Schumm et al., 1984) channels, and typically results in laterally-stable,
735 low aspect ratio channels, as can be seen in Mahin and lower Avon channels. Therefore,
736 unless re-incised during future rejuvenations, these terrace bodies are likely to be preserved
737 upon burial.

738

739 The upper Avon channel system shows a very different pattern. Its' highly aggradational
740 nature is responsible for frequent avulsions of the Avon channel, which reduces the terrace
741 development time and limits long term entrenchment and increases in sinuosity. The Avon
742 terrace bodies are therefore relatively thin and their planforms are to a large degree influenced
743 by the morphology of the underlying deposits. Their limited thickness, along with the

744 frequent channel avulsions within a narrow structural confinement, suggests a lower
745 preservation potential in comparison to the thick Mahin and lower Avon terrace body
746 deposits.

747

748

749 **8. Conclusions**

750 The mid-slope sections of the Mahin and Avon channels, located offshore western Nigeria,
751 have different architectures and are both associated with a number of terraces. The main
752 difference between the two channels is their sinuosity up-dip of their confluence. Substrate
753 cohesion is interpreted to be partially responsible for the geomorphological differences
754 between the two channels. However, the interpreted younger age and thus channel maturity
755 (linked to the channel avulsion frequency) of the upper Avon channel compared to the Mahin
756 and lower Avon channels is interpreted to be the primary control on the difference in
757 sinuosity.

758

759 The differing channel morphology and sinuosity greatly impact the distribution,
760 characteristics and stacking patterns of the deposits within terrace bodies that are adjacent to
761 these channels. Terrace surfaces can be formed by entrenchment, margin failure, point bar
762 accretion and channel bend cut-offs, which leave behind elevated areas within a channel-belt
763 on top of which heterolithics can be deposited by overbank flow. Terraces are much more
764 abundant in the channel-belts of the more sinuous Mahin and lower Avon channels compared
765 to the channel-belt of the upper Avon channel. Entrenchment and channel bend cut-offs are
766 linked to changes in channel sinuosity, resulting in greater abundance of terraces adjacent to
767 sinuous channels.

768

769 Many of the terrace bodies, particularly in the Mahin and lower Avon channel are composed
770 of channel-fill at the base and overlain by heterolithics, which predominantly form deposits
771 that lie on a continuum between flat depositional terraces and wedge-shaped internal levees.
772 True internal levees are only found when the terrace surface is high enough above the channel
773 that only the uppermost (more dilute) parts of the flow can overspill, and are only found in
774 parts of the terrace that is sufficiently wide to allow the overspilling current to decelerate and
775 deposit all its sediment before hitting the confinement.

776

777 The preservation potential of the terrace body deposits is dependent on the evolution of the
778 channel but given the abundance of terraces on the seafloor it is likely that they occur much
779 more commonly in outcrop and subsurface datasets than has been interpreted to date. The
780 abundance of sand prone heterolithics and channel-fill deposits makes these deposits
781 important to consider in hydrocarbon exploration.

782

783 **9. Acknowledgements**

784 The authors would like to thank PGS and Statoil ASA for permission to publish this dataset
785 and PGS also for collection and processing of the seismic data. Also thank you to Statoil ASA
786 for making the internship during which the bulk of this work was undertaken possible. We
787 acknowledge the support of the PRACSS Joint Industry Project at University of Aberdeen,
788 funded by BG Group, BP, DONG, Petrochina, RWE Dea, Statoil and Tullow Oil, which has
789 allowed us to undertake research in various areas where ideas for this paper were formulated.
790 We thank Dave Hodgson and Marzia Rovere for the insightful reviews that greatly improved
791 this paper.

792

793

794 **References**

- 795 Abreu, V., Sullivan, M., Pirmez, C., Mohrig, D., 2003. Lateral accretion packages (LAPs): An
796 important reservoir element in deep water sinuous channels. *Marine and Petroleum*
797 *Geology*, 20, 631–648. doi:10.1016/j.marpetgeo.2003.08.003
- 798 Adeogba, A., McHargue, T., & Graham, S., 2005. Transient fan architecture and depositional
799 controls from near-surface 3-D seismic data, Niger Delta continental slope. *AAPG*
800 *Bulletin*, 89(5), 627–643. doi:10.1306/11200404025
- 801 Allen, J.R., 1970. Sediments of the modern Niger Delta: A summary and review, in: Morgan,
802 J.P., Shaver, R.H. (Eds.), *Deltaic Sedimentation, Modern and Ancient*. SEPM Vol. 15,
803 pp. 138–151.
- 804 Allen, J.R., 1965. Late Quaternary Niger Delta, and adjacent areas: Sedimentary
805 environments and lithofacies. *AAPG Bulletin*, 49, 547–600.
- 806 Allen, J.R., 1964. The Nigerian continental margin: Bottom sediments, submarine
807 morphology and geological evolution. *Marine Geology*, 1, 289–332.
- 808 Anthony, E.J., 2015. Patterns of Sand Spit Development and Their Management Implications
809 on Deltaic, Drift-Aligned Coasts: The Cases of the Senegal and Volta River Delta Spits,
810 West Africa, in: Randazzo, G., Jackson, D., Cooper, J. (Eds.), *Sand and Gravel Spits*.
811 Springer International Publishing, pp. 21–36. doi:10.1007/978-3-319-13716-2
- 812 Antobreh, A., & Krastel, S., 2006. Morphology, seismic characteristics and development of
813 Cap Timiris Canyon, offshore Mauritania: A newly discovered canyon preserved-off a
814 major arid climatic region. *Marine and Petroleum Geology*, 23(1), 37–59.
815 doi:10.1016/j.marpetgeo.2005.06.003
- 816 Babonneau, N., Savoye, B., Cremer, M., & Klein, B., 2002. Morphology and architecture of
817 the present canyon and channel system of the Zaire deep-sea fan. *Marine and Petroleum*
818 *Geology*, 19(4), 445–467. doi:10.1016/S0264-8172(02)00009-0
- 819

820 Babonneau, N., Savoye, B., Cremer, M., Bez, M., 2004. Multiple terraces within the deep
 821 incised Zaire Valley (ZaiAngo Project): are they confined levees?, in: Lomas, S.A.,
 822 Joseph, P. (Eds.), *Confined Turbidite Systems*. Geological Society, London, Special
 823 Publications 222, pp. 91–114. doi:10.1144/GSL.SP.2004.222.01.06
 824 Babonneau, N., Savoye, B., Cremer, M., Bez, M., 2010. Sedimentary Architecture in
 825 Meanders of a Submarine Channel: Detailed Study of the Present Congo Turbidite
 826 Channel (Zaiango Project). *Journal of Sedimentary Research*, 80, 852–866.
 827 Beaubouef, R., 2004. Deep-water leveed-channel complexes of the Cerro Toro Formation,
 828 Upper Cretaceous, southern Chile. *AAPG Bulletin*, 88, 1471–1500.
 829 Bouma, A.H., Ravenne, C., 2004. The Bouma Sequence (1962) and the resurgence of
 830 geological interest in the French Maritime Alps (1980s): the influence of the Gres
 831 d’Annot in developing ideas of turbidite systems, in: Joseph, P., Lomas, S.A. (Eds.),
 832 *Deep-Water Sedimentation in the Alpine Basin of SE France: New Perspectives on the*
 833 *Gres d’Annot and Related Systems*. Geological Society of London, Special Publications,
 834 pp. 27–38. doi:10.1144/GSL.SP.2004.221.01.03
 835 Brown, A.R., 1999. *Interpretation of three-dimensional seismic data*, 5th ed. American
 836 Association of Petroleum Geologists, Tulsa, Oklahoma.
 837 Brunt, R.L., Hodgson, D., Flint, S.S., Pringle, J.K., Di Celma, C., Prlat, A., Grecula, M.,
 838 2013. Confined to unconfined: Anatomy of a base of slope succession, Karoo Basin,
 839 South Africa. *Marine and Petroleum Geology*, 41, 206–221.
 840 doi:10.1016/j.marpetgeo.2012.02.007
 841 Buffington, E.C., 1952. Submarine “Natural Levees.” *The Journal of Geology*, 60, 473–479.
 842 Burke, K., 1972. Longshore Drift, Submarine Canyons, and Submarine Fans in Development
 843 of Niger Delta. *AAPG Bulletin*, 56, 1975–1983. doi:10.1306/819A41A2-16C5-11D7-
 844 8645000102C1865D
 845
 846

- 847 Catterall, V., Redfern, J., Gawthorpe, R., Hansen, D., Thomas, M., 2010. Architectural Style
848 and Quantification of a Submarine Channel-Levee System Located in a Structurally
849 Complex Area: Offshore Nile Delta. *Journal of Sedimentary Research*, 80, 991–1017.
850 doi:10.2110/jsr.2010.084
- 851 Chen, C., Hiscott, R., 1999. Statistical analysis of turbidite cycles in submarine fan
852 successions; tests for short-term persistence. *Journal of Sedimentary Research*, 69, 486–
853 504. doi:10.2110/jsr.69.486
- 854 Clark, I. R., & Cartwright, J. A., 2009. Interactions between submarine channel systems and
855 deformation in deepwater fold belts: Examples from the Levant Basin, Eastern
856 Mediterranean sea. *Marine and Petroleum Geology*, 26(8), 1465–1482.
857 doi:10.1016/j.marpetgeo.2009.05.004
- 858 Clark, J.D., Kenyon, N.H., Pickering, K.T., 1992. Quantitative analysis of the geometry of
859 submarine channels: implications for the classification of submarine fans. *Geology*, 20,
860 633–636. doi:10.1130/0091-7613(1992)020<0633:QAOTGO>2.3.CO
- 861 Clark, J.D., Pickering, K.T., 1996. Architectural elements and growth patterns of submarine
862 channels; application to hydrocarbon exploration. *AAPG Bulletin*, 80, 194–221.
- 863 Clemenceau, G.R., Colbert, J., Edens, D., 2000. Production Results from Levee-Overbank
864 Turbidite Sands at Ram/Powell Field, Deepwater Gulf of Mexico, in: Weimer, P., Slatt,
865 M., Coleman, J., Rosen, N., Nelson, C., Bouma, A., Styzen, M., Lawrence, D. (Eds.),
866 Deep-Water Reservoirs of the World: GCSSEPM Foundation 20th Annual Research
867 Conference. pp. 241–251.
- 868 Corredor, F., Shaw, J.H. and Bilotti, F., 2005. Structural styles in the deep-water fold and
869 thrust belts of the Niger Delta. *AAPG Bulletin*, 89(6), pp.753-780
- 870 Cronin, B.T., 1995. Structurally-controlled deep-sea channel courses: examples from the
871 Miocene of south-east Spain and the Alboran Sea, south-west Mediterranean, in: Hartley,
872 A.J., Prosser, D.J. (Eds.), *Characterisation of Deep Marine Clastic Systems*. Geological
873 Society, London, Special Publication, Volume 94. pp. 113–133.

874 Dada, O., Qiao, L., Ding, D., Li, G., Ma, Y., Wang, L., 2015. Evolutionary trends of the Niger
875 Delta Shoreline during the last 100 years: Responses to rainfall and river discharge.
876 Marine Geology, 367, 202–211. doi:10.1016/j.margeo.2015.06.007

877 Dada, O.A., Li, G., Qiao, L., Ding, D., Ma, Y., 2016. Seasonal shoreline behaviours along the
878 arcuate Niger Delta coast: Complex interaction between fluvial and marine processes.
879 Continental Shelf Research, 122, 51–67. doi:10.1016/j.csr.2016.03.002

880 Damuth, J., 1994. Neogene gravity tectonics and depositional processes on the deep Niger
881 Delta continental margin. Marine and Petroleum Geology, 11, 320–346.
882 doi:10.1016/0264-8172(94)90053-1

883 Deptuck, M., Steffens, G.S., Barton, M., Pirmez, C., 2003. Architecture and evolution of
884 upper fan channel-belts on the Niger Delta slope and in the Arabian Sea. Marine and
885 Petroleum Geology, 20, 649–676. doi:10.1016/j.marpetgeo.2003.01.004

886 Deptuck, M., Sylvester, Z., Byrne, C., 2012. Pleistocene Seascape Evolution above a “simple”
887 stepped slope - Western Niger Delta, in: Prather, B.E., Deptuck, M.E., Mohrig, D., Van
888 Hoorn, B., Wynn, R.B. (Eds.), Application of the Principles of Seismic Geomorphology
889 to Continental-Slope and Base-of-Slope Systems: Case Studies from Seafloor and Near-
890 Seafloor Analogues. SEPM Special Publication No. 99, pp. 199–222.

891 Deptuck, M., Sylvester, Z., Pirmez, C., O’Byrne, C., 2007. Migration–aggradation history and
892 3-D seismic geomorphology of submarine channels in the Pleistocene Benin-major
893 Canyon, western Niger Delta slope. Marine and Petroleum Geology, 24, 406–433.
894 doi:10.1016/j.marpetgeo.2007.01.005

895 Doust, H., Omatsola, E., 1989. Niger Delta, in: Edwards, J.D., Santogrossi, P.A. (Eds.),
896 Divergent/Passive Margin Basins. American Association of Petroleum Geologists, Tulsa,
897 OK, pp. 201–238.

898

899

900 Dykstra, M., & Kneller, B., 2009. Lateral accretion in a deep-marine channel complex:
 901 implications for channellized flow processes in turbidity currents. *Sedimentology*, 56(5),
 902 1411–1432. <http://doi.org/10.1111/j.1365-3091.2008.01040.x>
 903 Friel, J., 2000. *Practical Guide to Image Analysis*. ASM International, 290 p.

904 Gamberi, F., Marani, M., 2006. Hinterland geology and continental margin growth: the case
 905 of the Gioia Basin (southeastern Tyrrhenian Sea), in: Moratti, G., Chalouan, A. (Eds.),
 906 *Tectonics of the Western Mediterranean and North Africa*. Geological Society of
 907 London, Special Publications, pp. 349–363. doi:10.1144/GSL.SP.2006.262.01.21

908 Gamberi, F., Rovere, M., 2011. Architecture of a modern transient slope fan (Villafranca fan,
 909 Gioia basin–Southeastern Tyrrhenian Sea). *Sedimentary Geology*, 236, 211–225.
 910 doi:10.1016/j.sedgeo.2011.01.007

911 Gamberi, F., Rovere, M., Dykstra, M., Kane, I., Kneller, B., 2013. Integrating modern
 912 seafloor and outcrop data in the analysis of slope channel architecture and fill. *Marine*
 913 *and Petroleum Geology*, 41, 83–103. doi:10.1016/j.marpetgeo.2012.04.002

914 Gamboa, D., Alves, T. M., & Cartwright, J., 2012. A submarine channel confluence
 915 classification for topographically confined slopes. *Marine and Petroleum Geology*, 35(1),
 916 176–189. doi:10.1016/j.marpetgeo.2012.02.011

917 Gervais, A., Savoye, B., Mulder, T., Gonthier, E., 2006. Sandy modern turbidite lobes: A new
 918 insight from high resolution seismic data. *Marine and Petroleum Geology*, 23, 485–502.
 919 doi:10.1016/j.marpetgeo.2005.10.006

920 Hansen, L.A.S., Callow, R.H.T., Kane, I., Gamberi, F., Rovere, M., Cronin, B.T., Kneller, B.,
 921 2015. Genesis and character of thin-bedded turbidites associated with submarine
 922 channels. *Marine and Petroleum Geology*, 67, 852–879.
 923 doi:10.1016/j.marpetgeo.2015.06.007

924 Hein, F., Walker, R., 1982. The Cambro-Ordovician Cap Enrage Formation, Quebec, Canada:
 925 conglomeratic deposits of a braided submarine channel with terraces. *Sedimentology*, 29,
 926 309–352.

927 Hiscott, R., Hall, F.R., Pirmez, C., 1997. 4. Turbidity-current overspill from the Amazon
 928 channel: texture of the silt/sand load, paleoflow from anisotropy of magnetic
 929 susceptibility and implications for flow processes, in: Flood, R., Piper, D., Klaus, A.,
 930 Peterson, L. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*, Vol.
 931 155. pp. 53–78.

932 Hodgson, D., Di Celma, C., Brunt, R., & Flint, S., 2011. Submarine slope degradation and
 933 aggradation and the stratigraphic evolution of channel-levee systems. *Journal of the*
 934 *Geological Society*, 168(3), 625–628. doi:10.1144/0016-76492010-177

935 Hodgson, D., Kane, I., Flint, S., Brunt, R., Ortiz-Karpf, A., 2016. Time transgressive
 936 confinement on the slope and the progradation of basin-floor fans: Implications for the
 937 sequence stratigraphy of deep-water deposits. *Journal of Sedimentary Research*, 86, 73–
 938 86. doi:10.1017/CBO9781107415324.004

939 Hovland, M., Gallagher, J.W., Clennell, M.B., Lekvam, K., 1997. Gas hydrate and free gas
 940 volumes in marine sediments: Example from the Niger Delta front. *Marine and*
 941 *Petroleum Geology*, 14, 245–255.

942 Hübscher, C., Spieß, V., Breitzke, M., Weber, M., 1997. The youngest channel-levee system
 943 of the Bengal Fan: results from digital sediment echosounder data. *Marine Geology*, 141,
 944 125–145.

945 Janocko, M., Cartigny, M.J.B., Nemec, W., Hansen, E.W.M., 2013a. Turbidity current
 946 hydraulics and sediment deposition in erodible sinuous channels: Laboratory
 947 experiments and numerical simulations. *Marine and Petroleum Geology*, 41, 222–249.
 948 doi:10.1016/j.marpetgeo.2012.08.012

949 Janocko, M., Nemec, W., Henriksen, S., Warchoń, M., 2013b. The diversity of deep-water
 950 sinuous channel belts and slope valley-fill complexes. *Marine and Petroleum Geology*,
 951 41, 7–34. doi:10.1016/j.marpetgeo.2012.06.012

952 Jobe, Z., Sylvester, N., Parker, A.O., Howes, N., Slowey, N., Pirmez, C., 2015. Rapid
 953 adjustment of submarine channel architecture to changes in sediment supply. *Journal of*
 954 *Sedimentary Research*, 85, 729–753.

955 Jones, S. M., Sutton, C., Hardy, R. J. J., and Hardy, D., 2010, Seismic imaging of variable
 956 water layer sound velocity in Rockall Trough, NE Atlantic, and implications for
 957 seismic surveying in deep water: *Petroleum Geology Conference Series*, 7, 549–558.

958 Kane, I., McCaffrey, W. D., & Peakall, J., 2008. Controls on sinuosity evolution within
 959 submarine channels. *Geology*, 36, 287–290. <http://doi.org/10.1130/G24588A.1>

960 Kane, I., Hodgson, D., 2011. Sedimentological criteria to differentiate submarine channel
 961 levee subenvironments: Exhumed examples from the Rosario Fm. (Upper Cretaceous) of
 962 Baja California, Mexico, and the Fort Brown Fm. (Permian), Karoo Basin, S. Africa.
 963 *Marine and Petroleum Geology*, 28, 807–823. doi:10.1016/j.marpetgeo.2010.05.009

964 Kane, I., Kneller, B., Dykstra, M., Kassem, A., McCaffrey, W.D., 2007. Anatomy of a
 965 submarine channel–levee: An example from Upper Cretaceous slope sediments, Rosario
 966 Formation, Baja California, Mexico. *Marine and Petroleum Geology*, 24, 540–563.
 967 doi:10.1016/j.marpetgeo.2007.01.003

968 Kendrick, J.W., 2000. Turbidite reservoir architecture in the Northern Gulf of Mexico
 969 deepwater: insights from the development of Auger, Tahoe, and Ram/Powell Fields, in:
 970 Weimer, P., Slatt, M., Coleman, J., Rosen, N., Nelson, C., Bouma, A., Styzen, M.,
 971 Lawrence, D. (Eds.), *Deep-Water Reservoirs of the World: GCSSEPM Foundation 20th*
 972 *Annual Research Conference*. pp. 450–468.

973 Kleinhans, M.G., 2010. Sorting out river channel patterns. *Progress in Physical Geography*,
 974 34, 287–326. doi:10.1177/0309133310365300

975 Kolla, V., Bourges, P., Urruty, J.-M., Safa, P., 2001. Evolution of deep-water Tertiary sinuous
 976 channels offshore Angola (west Africa) and implications for reservoir architecture.
 977 *AAPG Bulletin*, 85, 1373–1405.

978

979 Kuenen, P., Migliorini, C., 1950. Turbidity currents as a cause of graded bedding. The Journal
980 of Geology, 58, 91–127.

981 Maier, K., Fildani, a., McHargue, T.R., Paull, C., Graham, S. a., Caress, D.W., 2012.
982 Punctuated Deep-Water Channel Migration: High-Resolution Subsurface Data from the
983 Lucia Chica Channel System, Offshore California, U.S.A. Journal of Sedimentary
984 Research, 82, 1–8. doi:10.2110/jsr.2012.10

985 Manley, P., Pirmez, C., Busch, W., Cramp, A., 1997. 3. Grain-size characterization of
986 Amazon Fan deposits and comparison to seismic facies units, in: Flood, R., Piper, D.,
987 Klaus, A., Peterson, L. (Eds.), Proceedings of the Ocean Drilling Program, Scientific
988 Results, Vol. 155. pp. 35–52.

989 Mayall, M., Stewart, I., 2000. The Architecture of Turbidite Slope Channels, in: Weimer, P.,
990 Slatt, R.M., Coleman, J., Rosen, N.C., Nelson, H., Bouma, A.H., Styzen, M.J.,
991 Lawrence, D.T. (Eds.), Deep-Water Reservoirs of the World: GCSSEPM Foundation
992 20th Annual Research Conference. pp. 578–586.

993 McArdle, N., Ackers, M., 2012. Understanding seismic thin-bed responses using frequency
994 decomposition and RGB blending. First Break 30, 57–65.

995 McHargue, T., 1991. Seismic facies, processes, and evolution of Miocene inner fan channels,
996 Indus Submarine Fan, in: Weimer, P., Link, M. (Eds.), Seismic Facies and Sedimentary
997 Processes of Submarine Fans and Turbidite Systems. Springer-Verlag, New York, pp.
998 402–413.

999 McHargue, T., Pyrcz, M. J., Sullivan, M., Clark, J. D., Fildani, A., Romans, B. W.,
1000 Drinkwater, N. J., 2011. Architecture of turbidite channel systems on the continental
1001 slope: Patterns and predictions. Marine and Petroleum Geology, 28(3), 728–743.
1002 doi:10.1016/j.marpetgeo.2010.07.008

1003 Meiburg, E., Kneller, B., 2010. Turbidity Currents and Their Deposits. Annual Review of
1004 Fluid Mechanics, 42, 135–156. doi:10.1146/annurev-fluid-121108-145618
1005

- 1006 Morley, C.K., Guerin, G., 1996. Comparison of gravity-driven deformation styles and
1007 behavior associated with mobile shales and salt. *Tectonics*, 15, 1154.
1008 doi:10.1029/96TC01416
- 1009 Morris, E., Hodgson, D., Brunt, R., Flint, S., 2014. Origin, evolution and anatomy of silt-
1010 prone submarine external levées. *Sedimentology*, 61, 1734–1763. doi:10.1111/sed.12114
- 1011 Mutti, E., Normark, W., 1991. An integrated approach to the study of turbidite systems, in:
1012 Weimer, P., Link, M. (Eds.), *Seismic Facies and Sedimentary Processes of Submarine*
1013 *Fans and Turbidite Systems*. Springer-Verlag, New York, pp. 75–106.
- 1014 Nakajima, T., Kneller, B., 2013. Quantitative analysis of the geometry of submarine external
1015 levées. *Sedimentology*, 60, 877–910. doi:10.1111/j.1365-3091.2012.01366.x
- 1016 Nanson, G.C., Croke, J.C., 1992. A genetic classification of floodplains. *Geomorphology*, 4,
1017 459–486. doi:10.2113/gsecongeo.3.7.611
- 1018 NEDECO, 1961. *The waters of the Niger delta*. The Hague, North Holland Publishing Co., p.
1019 317.
- 1020 Normark, W., Damuth, J., 1997. 40. Sedimentary facies and associated depositional elements
1021 of the Amazon Fan, in: Flood, R.D., Piper, D.J.W., Klaus, A., Peterson, L.C. (Eds.),
1022 *Proceedings of the Ocean Drilling Program, Scientific Results*, Vol. 155. pp. 611–651.
- 1023 O’Connell, S., McHugh, C., Ryan, W., 1995. Unique fan morphology in an entrenched
1024 thalweg channel on the Rhone Fan, in: Pickering, K.T., Hiscott, N., Kenyon, N.H.,
1025 Lucchi, F.R., Smith, R. (Eds.), *Atlas of Deep Water Environments: Architectural Style in*
1026 *Turbidite Systems*. Chapman and Hall, London, pp. 80–83.
- 1027 Ortiz-Karpf, A., Hodgson, D.M., Mccaffrey, W.D., 2015. The role of mass-transport
1028 complexes in controlling channel avulsion and the subsequent sediment dispersal
1029 patterns on an active margin: The Magdalena Fan, offshore Colombia. *Marine and*
1030 *Petroleum Geology*, 64, 58–75. doi:10.1016/j.marpetgeo.2015.01.005

1031

1032

1033 Paull, C., Caress, D., Lundsten, E., Gwiazda, R., Anderson, K., McGann, M., Conrad, J.,
1034 Edwards, B., Sumner, E., 2013. Anatomy of the La Jolla Submarine Canyon system;
1035 offshore southern California. *Marine Geology*, 335, 16–34.
1036 doi:10.1016/j.margeo.2012.10.003

1037 Peakall, J., Ashworth, P.J., Best, J.L., 2007. Meander-Bend Evolution, Alluvial Architecture,
1038 and the Role of Cohesion in Sinuous River Channels: A Flume Study. *Journal of*
1039 *Sedimentary Research*, 77, 197–212. doi:10.2110/jsr.2007.017

1040 Peakall, J., Kane, I., Masson, D., Keevil, G., McCaffrey, W.D., Corney, R., 2012. Global
1041 (latitudinal) variation in submarine channel sinuosity. *Geology*, 40, 11–14.
1042 doi:10.1130/G32295.1

1043 Peakall, J., McCaffrey, W.D., Kneller, B., 2000. A Process Model for the Evolution,
1044 Morphology, and Architecture of Sinuous Submarine Channels. *Journal of Sedimentary*
1045 *Research*, 70, 434–448.

1046 Piper, D., Hiscott, R., 1999. Outcrop-scale acoustic facies analysis and latest Quaternary
1047 development of Hueneme and Dume submarine fans, offshore California.
1048 *Sedimentology*, 46, 47–78.

1049 Piper, D., Normark, W., 1983. Turbidite depositional patterns and flow characteristics, Navy
1050 submarine fan, California Borderland. *Sedimentology*, 30, 681–694.

1051 Pirmez, C., Beaubouef, R., Friedmann, S., Mohrig, D., 2000. Equilibrium profile and
1052 baselevel in submarine channels: examples from late Pleistocene systems and
1053 implications for the architecture of deepwater reservoirs, in: Weimer, P., Slatt, R.,
1054 Coleman, J., Rosen, N., Nelson, H., Bouma, A., Styzen, M., Lawrence, D. (Eds.), *Deep-*
1055 *Water Reservoirs of the World: GCSSEPM Foundation 20th Annual Research*
1056 *Conference*. pp. 782–805.

1057

1058

1059

- 1060 Pirmez, C., Hiscott, R., Kronen, J.D., 1997. 2. Sandy Turbidite successions at the base of
1061 channel-levee systems of the Amazon Fan revealed by FMS logs and cores: Unraveling
1062 the facies architecture of large submarine fans, in: Flood, R.D., Piper, D.J.W., Klaus, A.,
1063 Peterson, L.C. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*,
1064 Vol. 155. pp. 7–33.
- 1065 Pirmez, C., Imran, J., 2003. Reconstruction of turbidity currents in Amazon Channel. *Marine*
1066 *and Petroleum Geology*, 20, 823–849. doi:10.1016/j.marpetgeo.2003.03.005
- 1067 Posamentier, H., 2003. Depositional elements associated with a basin floor channel-levee
1068 system: case study from the Gulf of Mexico. *Marine and Petroleum Geology*, 20, 677–
1069 690. doi:10.1016/j.marpetgeo.2003.01.002
- 1070 Posamentier, H., Kolla, V., 2003. Seismic Geomorphology and Stratigraphy of Depositional
1071 Elements in Deep-Water Settings. *Journal of Sedimentary Research*, 73, 367–388.
1072 doi:10.1306/111302730367
- 1073 Posamentier, H., Martinsen, O., 2011. The character and genesis of submarine mass-transport
1074 deposits: insights from outcrop and 3D seismic data, in: Verwer, K., Playton, T., Harris,
1075 P. (Eds.), *Mass-Transport Deposits in Deepwater Settings*. SEPM Special Publication
1076 No. 96, pp. 7–38.
- 1077 Reading, H.G., Richards, M., 1994. Turbidite systems in Deep-Water basin margins classified
1078 by grain size and feeder system. *AAPG Bulletin*, 78, 792–822.
- 1079 Samuel, A., Kneller, B., Raslan, S., Sharp, A., Parson, C., 2003. Prolific deep-marine slope
1080 channels of the Nile Delta, Egypt. *AAPG Bulletin*, 87, 541–560.
- 1081 Schumm, S. A., & Khan, H. R., 1972. Experimental study of channel patterns. *Bulletin of the*
1082 *Geological Society of America*, 88, 1755–1770.
- 1083 Schumm, S. A., Khan, H. R., Winkley, B. R., & Robins, L. G., 1972. Variability of river
1084 patterns. *Nature; Physical Science*, 237, 75–76.
- 1085 Schumm, S., Harvey, M., Watson, C., 1984. *Incised channels: morphology, dynamics, and*
1086 *control*. Water Resources Publications.

1087 Schumm, S.A., 2005. River Variability and Complexity. Cambridge University Press, New
1088 York.

1089 Skene, K.I., Piper, D., Hill, P.S., 2002. Quantitative analysis of variations in depositional
1090 sequence thickness from submarine channel levees. *Sedimentology*, 49, 1411–1430.
1091 doi:10.1046/j.1365-3091.2002.00506.x

1092 Straub, K. M., Mohrig, D., McElroy, B., Buttles, J., & Pirmez, C., 2008. Interactions between
1093 turbidity currents and topography in aggrading sinuous submarine channels: A laboratory
1094 study. *Geological Society of America Bulletin*, 120 (3-4), 368–385.
1095 doi:10.1130/B25983.1

1096 Straub, K., Mohrig, D., Buttles, J., McElroy, B., Pirmez, C., 2011. Quantifying the influence
1097 of channel sinuosity on the depositional mechanics of channelized turbidity currents: A
1098 laboratory study. *Marine and Petroleum Geology*, 28, 744–760.
1099 doi:10.1016/j.marpetgeo.2010.05.014

1100 Sylvester, Z., Pirmez, C., Cantelli, A., & Jobe, Z. R., 2013. Global (latitudinal) variation in
1101 submarine channel sinuosity: COMMENT. *Geology*, 41(5), e287.
1102 <http://doi.org/10.1130/G33548C.1>

1103 Sylvester, Z., & Covault, J., 2016. Development of cutoff-related knickpoints during early
1104 evolution of submarine channels. *Geology*. <http://doi.org/10.1130/G38397.1>

1105 Torres, J., Droz, L., Savoye, B., Terentieva, E., Cochonat, P., Kenyon, N., Canals, M., 1997.
1106 Deep-sea avulsion and morphosedimentary evolution of the Rhone Fan valley and
1107 Neofan during the Late Quaternary (north-western Mediterranean Sea). *Sedimentology*,
1108 44, 457–477.

1109 Verhagen, I.T.E., Baas, J.H., Jacinto, R.S., McCaffrey, W.D., Davies, A.G., 2013. A first
1110 classification scheme of flow-bed interaction for clay-laden density currents and soft
1111 substrates. *Ocean Dynamics*, 63, 385–397. doi:10.1007/s10236-013-0602-8
1112
1113

- 1114 Von Rad, U., Tahir, M., 1997. Late Quaternary sedimentation on the outer Indus shelf and
1115 slope (Pakistan): evidence from high-resolution seismic data and coring. *Marine*
1116 *Geology*, 138, 193–236.
- 1117 Walker, R., 1975. Upper Cretaceous resedimented conglomerates at Wheeler Gorge,
1118 California: Description and Field guide. *Journal of Sedimentary Petrology*, 45, 105–112.
- 1119 Wu, J.E., McClay, K., Frankowicz, E., 2015. Niger Delta gravity-driven deformation above
1120 the relict Chain and Charcot oceanic fracture zones, Gulf of Guinea: Insights from
1121 analogue models. *Marine and Petroleum Geology*, 65, 43–62.
1122 doi:10.1016/j.marpetgeo.2015.03.008

Figure captions

Figure 1: Regional map of Western Niger slope with the study area outlined (adapted from Burke, 1972; Deptuck et al, 2007, 2012). Note that the area shown here is outlined in the map of Africa in the top right corner. Tidal channel names are from Allen (1965) and Burke (1972): Og (Ogun), On (Ona), Os (Oshun), Sh (Shasha), Be (Benin), Es (Escravos), Fo (Forcados), Ra (Ramos), Do (Dodo), Fi (Fishtown), Sa (Sangana), Nu (Nun), Br (Brass). Dashed arrows indicate dominant longshore drift directions and dashed lines to the north indicate the presence of beach ridges (from Burke, 1972).

Figure 2: Influence of seafloor expression of mud diapirs on the pathway of the Mahin and Avon channels (indicated by the black lines). a) Time depth map of interpreted regional top of mud diapirs in the study area. Interpreted horizon indicated in pink in the seismic line in the top left corner. Pathway of Mahin and Avon channels is guided through the structural lows between the mud diapirs. b) Thickness map between the interpreted regional base of the channel system and the seafloor indicated in yellow in seismic line in top left corner. White dashed lines highlight the main structural highs in the study area. Overbank areas of the Mahin and Avon channels are onlapping onto the structural highs. (Seismic dataset owner: PGS)

Figure 3: a) Depth map of the seafloor in the study area, derived from the seafloor reflection horizon in the 3D seismic dataset. Both channels were subdivided into sections, which are indicated. b) Line drawing showing the location of key geomorphic elements such as the channel-belts, channels, abandoned channels, terraces including terrace numbers, and structural highs on the seafloor. Distance measurements along the Mahin and Avon channels are also indicated. (Seismic dataset owner: PGS)

1150 Figure 4: Cross-sections through the Mahin channel showing the interpreted base of terrace
1151 horizons. Location of seismic lines is indicated in Fig. 3a. The high amplitude reflectors at the
1152 base of the terrace are interpreted as channel-fill deposits associated with the lateral migration
1153 of the channel over time. Depending on the evolution of the channel the base of the terrace
1154 may be stepped (in b), which indicates multiple generations of terraces. The base of the entire
1155 channel-belt is also indicated by the dashed lines. (Seismic dataset owner: PGS)

1156

1157 Figure 5: Seismic facies of the Mahin and Avon channel-belts

1158

1159 Figure 6: Descriptive geometrical parameters of a sinuous channel planform

1160

1161 Figure 7: Morphological characteristics of the Mahin and upper and lower Avon channels. a.1
1162 and a.2) Thalweg depth profile of the Mahin and Avon channels with gradients indicated for
1163 each section of the channel. Note that section 4 of the channel is repeated in the Mahin and
1164 Avon channel columns as this is the section of channel below the confluence. b.1 and b.2)
1165 The gradient of the Mahin and Avon channels measured along the thalweg over a 1 km and 4
1166 km window. The median gradient is also indicated. In b.1) the white arrow indicates increase
1167 in thalweg gradient where the Mahin channel crosses a mud diapir. c.1 and c.2) Sinuosity of
1168 the Mahin and Avon channels measured across a 6 km sliding window along the thalweg. The
1169 average sinuosity per channel section is also indicated.

1170

1171 Figure 8: a) Slope gradient map of the study area, with locations of cross-sections shown
1172 through both channels (indicated by the red and orange lines). b) Cross-sections through the
1173 channels indicated in a) with their orientation looking up-channel (SE to the right). Numbers
1174 indicate kilometres along channel from the zero point (which is 0 km for the Mahin channel
1175 and 42 km for the Avon channel). Note the steepness of the channel margins are indicated by
1176 the red numbers in the cross-sections. c) Confluence between the Mahin and Avon channels

1177 showing a 30 m drop between the Mahin channel mouth and the Avon channel. Location of
 1178 map shown in a). (Seismic dataset owner: PGS)

1179

1180 Figure 9: Interpretation of regional seismic lines (location indicated in Fig. 8) using seismic
 1181 facies described in Figure 5 and showing downslope changes in the channel morphology and
 1182 seismic facies. Note how the mud diapirs become less pronounced down-dip resulting in the
 1183 channels being less confined. (Seismic dataset owner: PGS)

1184

1185 Figure 10: Orientation and dip of terrace seafloor surface divided into N, NE, E, SE, S, SW,
 1186 W and NW. Note the irregular surfaces of all terraces. (Seismic dataset owner: PGS)

1187

1188 Figure 11: Characteristics of all terraces along the Mahin and Avon channels. Channel
 1189 sections are indicated for each parameter as well as the confluence between the Mahin and
 1190 Avon channels.

1191

1192 Figure 12: Spectral decomposition slices of the base of all terraces along the Mahin and Avon
 1193 channels. Locations of b), c), d) and e) are indicated in a). LAPs = lateral accretion packages.

1194

1195 Figure 13: Different scenarios of terrace forming processes. Different developmental stages of
 1196 the Mahin and Avon channel systems are indicated in the centre imitating different seismic
 1197 horizons. These include the proto-Avon and Mahin channels, palaeo-Avon lobe, and palaeo-
 1198 Avon channel. Four different terrace forming processes are shown. a) Avon section 1 is
 1199 dominated by the incision of the modern Avon channel into older channel-fill (CF) deposits,
 1200 leaving behind elevated areas on top of which heterolithics (HL) aggrade. b) Avon section 2 is
 1201 dominated by the incision of the modern Avon channel into older lobe deposits (LD) and
 1202 overbank chute channel (OCH) deposits that formed on top of terraces created by the incision

1203 of the modern Avon channel, leaving behind elevated areas on top of which heterolithics (HL)
1204 aggrade. c) Avon section 3 is dominated by the incision of the modern Avon channel into
1205 older channel bend channel-fill (CF) and abandonment deposits (AD), leaving behind
1206 elevated areas on top of which heterolithics (HL) aggrade. d) The Mahin and lower Avon
1207 channels are dominated by channel bend cut-offs, point bar accretion and the collapse of
1208 channel-margin sediment into the channel, forming elevated areas within the channel on top
1209 of which heterolithics (HL) aggrade.

1210

1211 Figure 14: Mahin and Avon channel depth profiles with histogram showing percentage of
1212 compositional elements within each terrace body. The histogram bars are located at the
1213 distance along the channel where the terraces occur and the vertical axis to the left of both
1214 graphs indicates the percentage of each compositional element. The knickpoint within the
1215 upper part of the Avon channel is also indicated.

1216

1217 Figure 15: RMS amplitude map extracted between 15 ms and 100 ms below the interpreted
1218 seafloor horizon. Higher RMS amplitude values suggest greater impedance contrasts and
1219 hence reflectivity within the measured interval. This may be the result either of increased
1220 impedance contrast between the lithologies present, or an increase in the frequency of
1221 reflection events due to increased interbedding of contrasting lithologies. A higher RMS value
1222 is therefore interpreted to indicate an increase in variability and interbedding of sands and
1223 muds, relative to the more homogeneous finer-grained background slope deposits. (Seismic
1224 dataset owner: PGS)

1225

1226

1227

1228 Figure 16: Quantification of terrace forming processes along the Mahin and Avon channels.
1229 Pie charts show the number of terraces formed by the processes indicated in the top left
1230 corner.

1231

1232 Figure 17: Schematic of a channel-levee system with terrace body deposits within the
1233 channel-belt. Stacking patterns of deposits in pseudo cores going through terrace body
1234 deposits. Stacking patterns can vary considerably. Channel is eroding into pre-existing lobe
1235 deposit as observed in the upper Avon channel. (modified from Hansen et al., 2015)

1236

1237 Figure 18: Slope gradient map of the study area highlighting the terraces in red where internal
1238 levee deposits are present. The graph in the top left corner shows a cross-plot of the terrace
1239 width vs terrace height with the terraces that have internal levees indicated in red and labelled
1240 with terrace numbers. The map in the bottom right corner is an insert from Figure 10 showing
1241 dip direction of the terrace surface on the seafloor. Note the irregularly dipping surface with
1242 part of it forming an internal levee as shown in cross-section A-A' (Seismic dataset owner:
1243 PGS)

1244

1245 **Table captions**

1246

1247 Table 1: Average geomorphological properties for the Mahin and Avon channels.

1248

1249 Table 2: Characteristics for all Mahin and Avon terraces.

Figure 1

Figure 5

Seismic facies	Description	Schematic	Architectural element	
<div>Facies A</div>	High amplitude apparently chaotic reflectors		Channel-fill	
<div>Facies B</div>	Continuous straight reflectors that converge and dim away from channel		Heterolithics	Levee heterolithics
<div>Facies C</div>	Continuous straight and parallel reflectors with varying amplitude			Channel- belt heterolithics
<div>Facies D</div>	Somewhat incoherent inclined but straight reflectors			Heterolithic slide block
<div>Facies E</div>	Chaotic reflectors with varying amplitude		Slump	
<div>Facies F</div>	Flat continuous reflectors with consistent amplitude		Lobe deposit	

Figure 6

L = Channel thalweg length

∂ = Average channel course (straight path)

Δ = Bend amplitude

λ = Bend length

$$\text{Terrace circularity} = \frac{4\pi \times A}{P^2}$$

$$\text{Channel sinuosity} = \frac{L}{\partial}$$

Figure 7

Mahin channel

Avon channel

Figure 8

- Facies A: Channel-fill
- Facies B: Heterolithics (external levee)
- Facies C: Heterolithics (within channel-belt)
- Facies D: Heterolithics (slide blocks)
- Facies E: Slumps
- Facies F: Lobe deposit

Figure 10

Figure 11 **Mahin channel**

Avon channel

Figure 13

b) Avon Section 2

- 1) Deposition of Palaeo-Avon lobe
- 2) Inception of modern Avon channel (an incised channel with wide, chute-dominated overbank areas)
- 3) Aggradation of heterolithics & maintenance of a bypass-dominated channel

c) Avon Section 3

- 1) Development of Proto-Avon (an incised meander-belt)
- 2) Abandonment & partial burial of Proto-Avon
- 3) Rejuvenation of Proto-Avon channel-belt by incision of modern Avon, aggradation of heterolithics in abandoned meander

a) Avon Section 1

- 1) Development of Palaeo-Avon channel (an incised channel-belt consisting of small, low-sinuosity channels)
- 2) Abandonment of Palaeo-Avon & inception of modern Avon channel
- 3) Incision of Avon thalweg & aggradation of heterolithics

d) Mahin & Lower Avon

- 1) Incision of Mahin channel meander-belt, point bar deposition on inner channel bend
- 2) Abandonment of meanders by neck & chute cut-off, collapse of channel-belt margins
- 3) Incision of thalweg, aggradation of heterolithics, restriction of lateral channel migration

Architectural elements

Channel fill
Lobe deposits
Overbank deposits
Channel margin failure deposits
Background deposits
Channel-belt deposits undifferentiated

Figure 14

Mahin channel

Avon channel

Figure 15

Figure 17

Core A:

Core B:

Figure 18

Table 1
Click here to download Table: Table 1_re-submission_reviews accepted.docx

Channel geomorphology:	Channel sections	Mahin channel (before confluence)		Upper Avon channel (before confluence)		Lower Avon channel (Section 4)
		Average/Total for sections 1-3		Average/Total for sections 1-3		
Average channel-belt depth (m)	Section 1	157	171	72	102	201
	Section 2	169		83		
	Section 3	180		184		
Average channel-belt width (m)	Section 1	3454	2740	2344	1978	3281
	Section 2	2851		1469		
	Section 3	2214		2484		
Average channel width (m)	Section 1	762	815	751	715	1068
	Section 2	887		566		
	Section 3	773		964		
Sinuosity (higher number = higher sinuosity)	Section 1	1.57	1.90	1.21	1.16	1.76
	Section 2	2.37		1.17		
	Section 3	1.75		1.1		
Thalweg length (km)	Section 1	25	90.7	20	49	38.7
	Section 2	45		21		
	Section 3	20.7		8		
Straight line distance (km)	Section 1	15.9	44.6	16.5	38	22
	Section 2	19		18		
	Section 3	11.8		7.3		
Average thalweg gradient (m/km)	Section 1	12.9/0.7°	10.4/0.59°	14.1/0.8°	15.1/0.86°	8.7/0.5°
	Section 2	9.7/0.44°		17.3/1°		
	Section 3	8.5/0.36°		14.0/0.8°		
Average regional slope gradient (m/km)	Section 1	18.6/1.07°	15.4/0.89°	20.2	15.6/0.87°	Left side = 12.8/0.7° Right side = 14.1/0.8°
	Section 2	12.1/0.7°		15.5		
	Section 3	15.6/0.9°		11.1		

Table 2
Click here to download Table 2 for submission.docx

here to download Table 2 re-submission.docx

Upper Avon channel (before confluence)					
Terrace number	Terrace area (km ²)	Terrace width (m)	Terrace length (m)	Terrace circularity	Terrace height above thalweg (m)
2	0.5	500.0	1781.0	0.4	70.0
3	4.8	1205.0	6995.0	0.2	56.3
4	0.6	482.0	1751.0	0.6	22.5
5	0.9	689.0	1982.0	0.6	60.0
6	0.6	910.0	964.0	0.7	18.8
7	1.0	485.0	4190.0	0.2	23.3
8	2.7	1115.0	3756.0	0.5	38.3
9	0.5	377.0	1725.0	0.4	20.3
10	0.4	450.0	1521.0	0.5	42.8
11	0.5	347.0	2054.0	0.3	61.5
12	0.6	796.0	1113.0	0.6	52.5
13	0.6	570.0	1495.0	0.5	35.0
14	0.2	506.0	320.0	0.7	20.0
15	0.4	488.0	1372.0	0.6	42.8
16	0.1	213.0	633.0	0.6	90.0
17	0.5	585.0	1188.0	0.7	138.0
18	2.2	648.0	3434.0	0.4	174.8
19	0.8	571.0	2187.0	0.4	147.8
Average	1.0	628.3	2180.3	0.5	61.9

Lower Avon channel (after confluence)					
35	1.1	939.8	1615.9	0.6	138.3
36	0.8	775.0	1149.7	0.6	86.1
37	1.0	796.6	1559.7	0.7	176.2
38	0.3	235.3	1292.0	0.4	136.6
39	1.0	656.0	1483.9	0.6	104.9
40	0.5	569.7	1053.6	0.6	18.3
41	0.4	953.8	970.0	0.7	152.1
42	1.4	1324.1	1360.9	0.7	66.6
43	1.2	1019.5	1487.3	0.8	120.9
44	0.9	594.6	1688.7	0.5	106.8
45	2.8	1674.5	2161.4	0.8	94.2
46	4.3	1453.1	3392.0	0.7	107.4
47	0.8	760.0	1431.7	0.7	146.4
48	0.6	959.9	1214.0	0.7	211.0
49	0.4	522.9	1423.8	0.5	183.5
50	1.2	713.9	2375.9	0.4	177.7
51	1.0	823.4	1366.7	0.7	176.2
52	0.2	373.8	831.8	0.7	178.2
53	1.0	1021.5	1463.0	0.6	97.5
54	0.6	556.0	1621.0	0.5	211.4
Average	1.1	836.2	1547.2	0.6	134.5

Mahin channel (before confluence)						
Terrace number	Terrace area (km ²)	Terrace width (m)	Terrace length (m)	Terrace circularity	Terrace height above thalweg (m)	
1	1.3	880.0	2432.0	0.4	109.0	Section 1
2	1.4	653.9	2537.0	0.3	135.0	
3	1.1	961.0	1706.0	0.5	89.4	
4	2.2	962.0	2323.0	0.7	111.5	
5	1.5	1629.0	1214.7	0.6	123.7	
6	1.1	1231.0	1082.9	0.8	113.3	
7	1.5	1465.2	1263.6	0.8	52.0	
8	1.3	1536.9	1852.7	0.5	98.9	Section 2
9	1.4	787.6	2168.7	0.5	118.0	
10	0.5	402.4	1281.8	0.5	146.4	
14	0.8	929.0	1147.5	0.9	93.6	
15	0.4	494.5	1003.0	0.7	107.0	
16	0.5	533.0	1016.0	0.8	108.3	
17	0.3	269.0	1396.0	0.4	113.3	
18	0.6	1508.0	1238.0	0.3	74.1	
19	0.5	813.0	914.0	0.6	126.0	
20	0.5	577.0	1359.0	0.5	164.6	
21	0.3	372.0	1122.0	0.6	110.0	
22	0.5	1263.5	1179.8	0.3	178.0	
23	0.2	399.8	518.3	0.8	119.1	Section 3
24	0.3	375.6	800.8	0.7	80.5	
25	0.3	416.0	808.0	0.6	140.9	
26	0.1	246.8	597.0	0.6	67.0	
27	0.2	352.7	917.0	0.5	153.0	
28	0.1	125.8	862.8	0.3	55.8	
29	0.1	189.0	633.5	0.4	69.6	
30	1.2	955.6	2276.6	0.5	168.5	
31	0.2	321.0	879.5	0.6	136.0	
32	0.1	333.8	522.0	0.6	113.6	
33	0.0	193.6	674.5	0.7	75.9	
34	0.9	883.3	1337.0	0.8	51.2	
Average	0.7	711.6	1260.2	0.6	109.8	