Serious fungal infections in Canada

DOI:

[10.1007/s10096-017-2922-y](https://doi.org/10.1007/s10096-017-2922-y)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Dufresne, S. F., Cole, D. C., Denning, D., & Sheppard, D. C. (2017). Serious fungal infections in Canada. *European Journal of Clinical Microbiology and Infectious Diseases*, 36(6), 987-992. <https://doi.org/10.1007/s10096-017-2922-y>

Published in:

European Journal of Clinical Microbiology and Infectious Diseases

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Title: Serious fungal infections in Canada

Authors: Simon F. Dufresne¹, Donald C. Cole², David W. Denning³ and Donald C. Sheppard⁴

Affiliation:

1- Department of Infectious Diseases and Medical Microbiology, Maisonneuve-Rosemont Hospital Research Center, Université de Montréal, Montréal, Canada;

2- Dalla Lana School of Public Health, University of Toronto, Toronto, Canada;

3- The National Aspergillosis Centre, University Hospital of South Manchester, The University of Manchester, Manchester Academic Health Science Centre, Manchester, UK;

4- Department of Microbiology and Immunology and Department of Medicine, Infectious Diseases and Immunity in Global Health Program, Research Institute of the McGill University Health Center, McGill University, Montréal, Canada.

Corresponding author:

Simon F. Dufresne

Department of Infectious Diseases and Medical Microbiology

Hôpital Maisonneuve-Rosemont

5415 boul. De l'Assomption

Montréal (Québec) Canada, H1T 1M4

sf.dufresne@umontreal.ca

Fax: 514-252-3898

Abstract

There are currently no nationwide epidemiological data on fungal infections in Canada. We estimated the burden of serious fungal diseases using literature review and modelling, as per a methodology previously described by the LIFE program (www.LIFE-worldwide.org). Among the population of Canada (35.5 million in 2014), it was estimated that approximately 1.8% are affected by a serious fungal infection. Recurrent vulvovaginal candidiasis, severe asthma with fungal sensitization and allergic broncho-pulmonary aspergillosis are the most frequent infections with a population prevalence of 498,688 (1,403/100,000), 73,344 (206/100,000) and 61,854 (174/100,000) cases, respectively. Over 3,000 invasive fungal infections are estimated to occur annually, with an incidence of 2,068 cases (5.8/100,000) of invasive candidiasis, 566 cases (1.6/100,000) of invasive aspergillosis, 252 cases (0.69/100,000) of *Pneumocystis* pneumonia, 99 cases (0.28/100,000) of endemic mycoses and 63 cases (0.18/100,000) of cryptococcosis. These estimates warrant validation through more formal epidemiological studies in Canada.

Key words: fungal diseases, burden, epidemiology, Canada

1   **Introduction**

2
3   Canada is a high-income country with a gross domestic product of 1,551 billion dollars (USD)
4   in 2015 (GDP per capita, 43,249). Several studies have examined the epidemiology of fungal
5   infections in Canada, but most were local (single-centered, regional or provincial) and focused
6   on individual infections in specific populations. A single study reported on the burden of
7   invasive fungal infections at the national level, however these data are over 20-years-old [1],
8   and relied largely on clinical microbiology laboratory records review. Since that time, the
9   Canadian population has expanded by 26%; use of immunosuppressive therapies that increase
10   the risk of fungal infection have risen in importance, e.g. solid organ and stem cell
11   transplantation and immunosuppressive biologic therapies; fungal diagnostics have improved;
12   and medical care practices have evolved greatly. Hence, we sought to provide an estimate of
13   the national burden of serious fungal infections.

15   **Material and methods**

16   The current study targeted serious fungal infections, which were defined as fungal-associated
17   syndromes (infection or hypersensitivity) causing significant morbidity or mortality. The
18   LIFE methodology (www.life-worldwide.org) was followed as described previously [2,3],
19   with some modifications.

20
21   In brief, an annual burden was estimated for each targeted serious fungal disease and
22   presented as: i) absolute number of cases per year in Canada (representing either incident or
23   prevalent cases depending on the nature of the infection), and ii) annual rates (incidence or
24   prevalence), using the annual number of cases as the numerator and the entire Canadian
25   population as the denominator. For simplicity, the 2014 Canadian population (n=35,540,419

[4]) was used regardless of the year from which the numerator data originated. To calculate the absolute number of cases per year in Canada, data derived from various sources were aggregated in a step-wise hierarchic fashion, starting with notifiable fungal diseases obtained directly from national and provincial public health agencies, followed by local data from individual centers or health authorities cited in peer review or grey literature and extrapolated to the total Canadian population (with the exception of geographically-confined infections, *Cryptococcus gattii* and endemic mycoses) and finally extrapolations from other countries' estimates.

Sources included: Statistics Canada (total and adult female population [4], asthma prevalence [5]), the Canadian Cancer Society (acute leukemia [6] and overall cancer incidence [7]), the Canadian Institute for Health Information (number of solid organ transplants) [8], the Canadian Bone Marrow Transplantation Group (number of hematopoietic stem cell transplants), the Organization for Economic Co-operation and Development (number of chronic obstructive pulmonary disease [COPD] admissions) [9], the Public Health Agency of Canada (HIV prevalence and AIDS incidence) [10], the World Health Organization (pulmonary tuberculosis incidence) [11] and Cystic Fibrosis Canada (cystic fibrosis prevalence) [12]. Of note, AIDS cases from the province of Québec are not currently reported to the Public Health Agency of Canada, so the number was calculated at pro rata of Québec population, assuming the same incidence as the rest of the country.

A map showing the geographic distribution of endemic mycoses was created using SmartDraw (SmartDraw Software, LLC, San Diego, CA, USA).

Results and Discussion

Estimated annual numbers of cases of serious fungal diseases and the corresponding population rates are summarized in Table 1. The distribution of geographically-confined mycoses is shown in Figure 1.

Opportunistic invasive fungal infections

The burden of invasive candidiasis (IC) was extrapolated from data reported from the Calgary health region and the Province of Québec [13,14]. In the first study, candidemia incidence was estimated at 2.8 cases per 100,000 person-year over a 5-year period, based on blood and cerebrospinal fluid culture (approximately 93% of cases were candidemia). Of note, incidence was significantly higher (3.7/100,000) during the latter 3 years of the study. In the second study, the reported annual incidence of candidemia was 3 cases per 100,000 population, with 50% of cases arising from intensive care units. These rates translate to at least 1,034 cases of candidemia each year in Canada. Deep-seated *Candida* infections such as peritonitis and endophthalmitis are accompanied by positive blood cultures in at most 50% of cases [15]. This observation is consistent with a recent nationwide German study which reported that the incidence of IC without candidemia was the same as candidemia [3]. Applying this ratio, a total of 2,068 IC cases are estimated to occur annually in Canada.

Invasive aspergillosis (IA) was largely reported in the main high-risk populations including patients with hematologic malignancies, hematopoietic stem cell transplant (HSCT) recipients and solid-organ transplant recipients (SOT), as well as patients with severe chronic obstructive pulmonary disease (COPD). A recent study from a Montreal center found an incidence of 8.9% among adult acute leukemia patients (both myelogeneous [AML] and

lymphocytic [ALL]) [16], consistent with other international reports [17]. There were 1,215 cases of AML in Canada in 2010 [6], corresponding to approximately 108 cases of IA that year. Since AML and ALL were reported to account for approximately 75% of IA cases occurring among patients with hematologic malignancies [18], an additional 36 IA cases are estimated to occur in patients with other hematologic malignancies. The incidence of IA amongst patients undergoing SOT was estimated using data from a prospective North American cohort study of these populations (12-month cumulative incidence: kidney, 0.2%; liver, 0.5%; lung, 3.8%; heart, 0.8%; pancreas, 0.2%) [19]. The incidence of IA after allogeneic HSCT was estimated at 7.5% based on recent observational and interventional studies [20-23]. In 2012, there were 1,358, 494, 194, 164 and 77 kidney, liver, lung, heart and pancreas transplants in Canada, respectively [8], corresponding to an estimated 14 cases of IA. A total of 1,200 allogeneic HSCT are performed annually in Canada (personal communication, Canadian Bone Marrow Transplantation Group), resulting in an additional estimated 90 cases of IA per year. Of note, large prospective cohort studies found that least 20% of IA cases occur more than a year after SOT and HSCT [24,19], representing an additional 26 cases. In addition, data from the North American Path Alliance registry showed that 22.3% of IA cases were associated with other underlying diseases, including solid tumors, HIV/AIDS and autologous HSCT, which translates to 78 cases. Hence, 352 IA cases are estimated to occur annually amongst traditional immunocompromised populations. Finally, 59,514 hospital admissions for chronic obstructive pulmonary disease (COPD) [9] account for approximately 214 IA cases, or 3.6 cases per 1,000 admissions [25], resulting in a total of 566 IA cases annually.

Over a 10-year period (2003-2012), 264 cases of *Pneumocystis pneumonia* (PCP) were diagnosed in single tertiary care centre in Montreal (personal communication, Dr M. Laskine,

Centre Hospitalier de l'Université de Montréal). Using this center's estimated catchment population of 3.4 million (41% of Quebec's population), one might expect as many as 252 cases per year across Canada.

A *C. gattii* outbreak has been ongoing in the province of British Columbia since 1999 [26] (Figure 1). Between 2010 and 2013, a mean of 23 *C. gattii* infections per year were reported to the British Columbia CDC [27]. During a 2-year period (1992-1994) prior to the *C. gattii* outbreak, investigators had identified 81 cases of *C. neoformans* disease across the country [1]. Assuming that *C. gattii* infections represent an increased burden of disease and that *C. neoformans* infection rates remain unchanged, then at least 63 cryptococcal infections are estimated to occur annually.

Endemic mycoses

Blastomycosis is endemic in the provinces of Ontario, Québec and Manitoba (Figure 1). It is considered hyperendemic in an area of northwestern Ontario [28]. In the former two provinces, recently published reports suggest that a mean of 44 culture-proven cases occur annually, [29,30]. In Manitoba where blastomycosis is a reportable infection, 19 cases were documented in 2013 [31]. Collectively, these data suggest an annual incidence of at least 63 cases of blastomycosis in these three provinces, hence in Canada.

Histoplasmosis occurs mainly along the Saint-Lawrence river valley, in Ontario and Québec (Figure 1). Histoplasmin reactivity studies have shown a prevalence of 9-27% in southern Québec [32-34]. The incidence of active histoplasmosis has been estimated at 27 cases per year in Canada in a study spanning the 2-year period 1992-1994 [1].

Coccidioides immitis/posadasii is not indigenous in Canada, though cases of active coccidioidomycosis are imported. Studies have estimated there are between 2 and 9 cases of coccidioidomycosis annually, largely in returning travelers [35,1].

Non-invasive pulmonary aspergillosis

Extrapolating from pulmonary tuberculosis 2014 annual incidence data (n=1,065) [11], the prevalence of cases of CPA following tuberculosis were estimated at 148 cases using a previously reported model [36]. If prior pulmonary tuberculosis is the predisposing factor in approximately 30% of cases of CPA [37], an estimated 492 new cases of CPA are expected.

ABPA burden was derived from the number of Canadian adults with asthma (n=2,444,804) [5] and the number of patients suffering from cystic fibrosis (CF; n=4,077) [12]. The proportion of ABPA among asthma patients was estimated at 2.5% [38] and 18% in those with CF [39], leading to a calculated national prevalence of 61,854 cases of ABPA. SAFS burden was derived from asthma prevalence and estimated at 73,344 cases, assuming a severe asthma prevalence of 10% of all cases of asthma and an *Aspergillus* sensitization prevalence of 30% [40].

Mucosal candidiasis

RVVC is reported to affect up to 9% of women of reproductive age [41]. Canadian prevalence of RVVC was calculated by applying 6% to the Canadian adult women's population (between 15 and 50 years old; n=8,311,477 [4]), yielding an estimated 498,688 cases present.

1
2 In a systematic review, the prevalence of OC in cancer patients was estimated 7.5%, 39.1%
3 and 32.6%, before, during and after the end of treatment, respectively [42]. In Canada,
4 196,900 new cases of cancer were estimated in 2015 [7]. Because the total number of patients
5 undergoing treatment and the survival rates are not known, only pre-treatment OC was
6 considered and the estimated prevalence was calculated at 14,767 cases of OC nationally.
7 HIV infection is responsible for an additional 752 cases, including approximately 1% of HIV-
8 positive patients without AIDS and 18% of those with AIDS [43], based on a prevalence of
9 71,300 HIV-positive patients in 2013 in Canada including 218 with AIDS [10].

10
11 EC is an AIDS-defining illness affecting nearly 20% of patients AIDS [44], thus at least 43
12 cases of EC are expected to occur annually in Canada. EC is also associated with other
13 underlying conditions (eg. receipt of corticosteroids), but data on infection rates in these
14 populations are not sufficient to inform accurate estimates.

15
16 Our study found that approximately 3,000 invasive fungal infections occur annually in
17 Canada, while over half a million people suffer from a chronic *Candida* or *Aspergillus*
18 infection. Overall, RVVC is likely the most frequent serious fungal disease in Canada, with
19 substantial quality of life impairment [45] and health costs [46], prompting the need for
20 specific guidelines addressing diagnostic and treatment. Education on current guidelines for
21 non-invasive forms of pulmonary aspergillosis is needed to increase awareness, diagnosis and
22 treatment proficiency among health care providers. Invasive fungal infections annual
23 incidence (8.3/100,000) is similar to that of invasive pneumococcal disease (8.9 in 2014) and
24 higher than invasive *S. pyogenes* disease (4.7 in 2013), infectious syphilis (5.1 in 2011),
25 tuberculosis (4.4 in 2014) and HIV (5.9 in 2013) all regarded as important in Canada [47].

1 Amongst invasive infections, candidiasis is by far the most common disease, followed by
2 aspergillosis. Cryptococcosis, pneumocystosis and endemic mycoses are less commonly
3 encountered, consistent with findings of a recent large registry across North America [48].
4 Invasive fungal infections data from two Canadian centers have confirmed 90-day fatality
5 rates of 41% and 36% for IC and IA, respectively [49], over a thousand deaths annually in
6 Canada.

7
8 Our estimates of serious fungal infection burden in Canada provide an opportunity for both
9 health care providers and public health agencies to improve diagnostics, surveillance and
10 formal epidemiological studies, including measures of health care utilization.

1 Tables & Figures

2 Table 1. Estimated annual burden of serious fungal diseases in Canada.

Serious fungal infection categories and types	Proportion affected (per 100,000 population)	# cases
Opportunistic invasive fungal infections	8,28	2,949
Invasive candidiasis (IC) without candidemia	2,91	1,034
Candidemia	2,91	1,034
Invasive aspergillosis (IA)	1,59	566
<i>Pneumocystis pneumonia</i> (PCP)	0,69	252
Cryptococcosis	0,18	63
Endemic mycoses	0,28	99
Blastomycosis	0,18	63
Histoplasmosis	0,08	27
Coccidioidomycosis	0,03	9
Non-invasive pulmonary aspergillosis	381,79	135,690
Severe asthma with fungal sensitization (SAFS)	206,37	73,344
Allergic bronchopulmonary aspergillosis (ABPA)	174,04	61,854
Chronic pulmonary aspergillosis (CPA)	1,38	492
Mucosal candidiasis	1,446,94	514,250
Recurrent vulvovaginal candidiasis (RVVC)	1,403,16	498,688
Oropharyngeal candidiasis (OC)	43,67	15,519
Esophageal candidiasis (EC)	0,12	43

3
4

Figure 1. Localization of geographically-confined mycoses in Canada. Colored areas represent the approximate geographic distribution of three fungal infections: Blue: *C. gattii*; Green: blastomycosis; Yellow: blastomycosis and histoplasmosis.

References

1. Nicolle L, Rotstein C, Bourgault A, St-Germain G, Garber G (1998) Invasive fungal infections in Canada from 1992 to 1994. *The Canadian journal of infectious diseases = Journal canadien des maladies infectieuses* 9 (6):347-352
2. Rodriguez-Tudela JL, Alastruey-Izquierdo A, Gago S, Cuenca-Estrella M, Leon C, Miro JM, Nunez Boluda A, Ruiz Camps I, Sole A, Denning DW (2015) Burden of serious fungal infections in Spain. *Clin Microbiol Infect* 21 (2):183-189. doi:10.1016/j.cmi.2014.07.013
3. Ruhnke M, Groll AH, Mayser P, Ullmann AJ, Mendling W, Hof H, Denning DW (2015) Estimated burden of fungal infections in Germany. *Mycoses* 58 Suppl 5:22-28. doi:10.1111/myc.12392
4. CANSIM Table 051-0001. Population by sex and age group (2014). Statistics Canada,
5. CANSIM Table 105-0502. Health indicator profile, two year period estimates, by age group and sex, Canada, provinces, territories, health regions (2012 boundaries) and peer groups (2013). Statistics Canada, Canadian Community Health Services,
6. Acute myelogenous leukemia statistics. (2015) Canadian Cancer Society. <http://www.cancer.ca/en/cancer-information/cancer-type/leukemia-acute-myelogenous-aml/statistics/?region=on>. Accessed April 10 2015
7. Canadian cancer statistics 2015 (2015). Toronto, Ontario
8. Canadian organ replacement register annual report: treatment of end-stage organ failure in Canada, 2003 to 2012 (2014). Canadian Institute for Health Information, Ottawa, Ontario
9. Health at a glance 2013: OECD Indicators (2013). OECD,
10. HIV and AIDS in Canada: Surveillance report to December 31, 2013. (2014). Minister of Public Works and Government Services Canada, Ottawa, Ontario
11. Tuberculosis country profiles -Canada. (2014) World Health Organization https://extranet.who.int/sree/Reports?op=Replet&name=%2FWHO_HQ_Reports%2FG2%2FPROD%2FEXT%2FTBCountryProfile&ISO2=CA&LAN=EN&outtype=html. Accessed August 16, 2016
12. The canadian cystic fibrosis registry: 2013 annual report (2015).
13. Laupland KB, Gregson DB, Church DL, Ross T, Elsayed S (2005) Invasive *Candida* species infections: a 5 year population-based assessment. *J Antimicrob Chemother* 56 (3):532-537. doi:10.1093/jac/dki258
14. St-Germain G, Laverdiere M, Pelletier R, Rene P, Bourgault AM, Lemieux C, Libman M (2008) Epidemiology and antifungal susceptibility of bloodstream *Candida* isolates in Quebec: Report on 453 cases between 2003 and 2005. *Can J Infect Dis Med Microbiol* 19 (1):55-62
15. Clancy CJ, Nguyen MH (2013) Finding the "missing 50%" of invasive candidiasis: how nonculture diagnostics will improve understanding of disease spectrum and transform patient care. *Clin Infect Dis* 56 (9):1284-1292. doi:10.1093/cid/cit006
16. Barkati S, Dufresne SF, Bélanger S, Vadnais B, Bergeron J, Labbe AC, Laverdiere M (2014) Incidence of invasive aspergillosis in remission-induction chemotherapy for acute leukemia: a retrospective cohort study in a single Canadian tertiary care center. *Canadian Medical Association Journal Open* 2 (2):E86-E93. doi:10.9778/cmajo.20130062
17. Pagano L, Caira M, Candoni A, Offidani M, Fianchi L, Martino B, Pastore D, Picardi M, Bonini A, Chierichini A, Fanci R, Caramatti C, Invernizzi R, Mattei D, Mitra ME, Melillo L, Aversa F, Van Lint MT, Falcucci P, Valentini CG, Girmenia C, Nosari A (2006) The epidemiology of fungal infections in patients with hematologic malignancies: the SEIFEM-2004 study. *Haematologica* 91 (8):1068-1075

18. Steinbach WJ, Marr KA, Anaissie EJ, Azie N, Quan SP, Meier-Kriesche HU, Apewokin S, Horn DL (2012) Clinical epidemiology of 960 patients with invasive aspergillosis from the PATH Alliance registry. *J Infect* 65 (5):453-464. doi:S0163-4453(12)00220-4 [pii] 10.1016/j.jinf.2012.08.003 [doi]
19. Pappas PG, Alexander BD, Andes DR, Hadley S, Kauffman CA, Freifeld A, Anaissie EJ, Brumble LM, Herwaldt L, Ito J, Kontoyiannis DP, Lyon GM, Marr KA, Morrison VA, Park BJ, Patterson TF, Perl TM, Oster RA, Schuster MG, Walker R, Walsh TJ, Wannemuehler KA, Chiller TM (2010) Invasive fungal infections among organ transplant recipients: results of the Transplant-Associated Infection Surveillance Network (TRANSNET). *Clin Infect Dis* 50 (8):1101-1111. doi:10.1086/651262 [doi]
20. Corzo-Leon DE, Satlin MJ, Soave R, Shore TB, Schuetz AN, Jacobs SE, Walsh TJ (2015) Epidemiology and outcomes of invasive fungal infections in allogeneic haematopoietic stem cell transplant recipients in the era of antifungal prophylaxis: a single-centre study with focus on emerging pathogens. *Mycoses* 58 (6):325-336. doi:10.1111/myc.12318
21. Harrison N, Mitterbauer M, Tobudic S, Kalhs P, Rabitsch W, Greinix H, Burgmann H, Willinger B, Presterl E, Forstner C (2015) Incidence and characteristics of invasive fungal diseases in allogeneic hematopoietic stem cell transplant recipients: a retrospective cohort study. *BMC Infect Dis* 15:584. doi:10.1186/s12879-015-1329-6
22. Wingard JR, Carter SL, Walsh TJ, Kurtzberg J, Small TN, Baden LR, Gersten ID, Mendizabal AM, Leather HL, Confer DL, Maziarz RT, Stadtmauer EA, Bolanos-Meade J, Brown J, Dipersio JF, Boeckh M, Marr KA (2010) Randomized, double-blind trial of fluconazole versus voriconazole for prevention of invasive fungal infection after allogeneic hematopoietic cell transplantation. *Blood* 116 (24):5111-5118. doi:blood-2010-02-268151 [pii] 10.1182/blood-2010-02-268151 [doi]
23. Montesinos P, Rodriguez-Veiga R, Boluda B, Martinez-Cuadron D, Cano I, Lancharro A, Sanz J, Arilla MJ, Lopez-Chulia F, Navarro I, Lorenzo I, Salavert M, Peman J, Calvillo P, Martinez J, Carpio N, Jarque I, Sanz GF, Sanz MA (2015) Incidence and risk factors of post-engraftment invasive fungal disease in adult allogeneic hematopoietic stem cell transplant recipients receiving oral azoles prophylaxis. *Bone Marrow Transplant* 50 (11):1465-1472. doi:10.1038/bmt.2015.181
24. Kontoyiannis DP, Marr KA, Park BJ, Alexander BD, Anaissie EJ, Walsh TJ, Ito J, Andes DR, Baddley JW, Brown JM, Brumble LM, Freifeld AG, Hadley S, Herwaldt LA, Kauffman CA, Knapp K, Lyon GM, Morrison VA, Papanicolaou G, Patterson TF, Perl TM, Schuster MG, Walker R, Wannemuehler KA, Wingard JR, Chiller TM, Pappas PG (2010) Prospective surveillance for invasive fungal infections in hematopoietic stem cell transplant recipients, 2001-2006: overview of the Transplant-Associated Infection Surveillance Network (TRANSNET) Database. *Clin Infect Dis* 50 (8):1091-1100. doi:10.1086/651263 [doi]
25. Guinea J, Torres-Narbona M, Gijon P, Munoz P, Pozo F, Pelaez T, de Miguel J, Bouza E (2010) Pulmonary aspergillosis in patients with chronic obstructive pulmonary disease: incidence, risk factors, and outcome. *Clin Microbiol Infect* 16 (7):870-877. doi:10.1111/j.1469-0691.2009.03015.x
26. Kidd SE, Hagen F, Tschärke RL, Huynh M, Bartlett KH, Fyfe M, Macdougall L, Boekhout T, Kwon-Chung KJ, Meyer W (2004) A rare genotype of *Cryptococcus gattii* caused the cryptococcosis outbreak on Vancouver Island (British Columbia, Canada). *Proc Natl Acad Sci U S A* 101 (49):17258-17263. doi:10.1073/pnas.0402981101
27. British Columbia Annual Summary of Reportable Diseases 2013 (2014). Communicable Disease Prevention and Control Services, Provincial Health Services Authority Vancouver, British Columbia

28. Dalcin D, Ahmed SZ (2015) Blastomycosis in northwestern Ontario, 2004 to 2014. *The Canadian Journal of Infectious Diseases & Medical Microbiology* 26 (5):259-262
29. Morris SK, Brophy J, Richardson SE, Summerbell R, Parkin PC, Jamieson F, Limerick B, Wiebe L, Ford-Jones EL (2006) Blastomycosis in Ontario, 1994-2003. *Emerg Infect Dis* 12 (2):274-279. doi:10.3201/eid1202.050849
30. Litvinov IV, St-Germain G, Pelletier R, Paradis M, Sheppard DC (2013) Endemic human blastomycosis in Quebec, Canada, 1988-2011. *Epidemiology and infection* 141 (6):1143-1147. doi:10.1017/s0950268812001860
31. Manitoba Annual Summary of Communicable Diseases 2013 (2014). Government of Manitoba. Manitoba Health, Healthy Living and Seniors. Public Health and Primary Health Care Division. Public Health Branch. Epidemiology and Surveillance.,
32. Guy R, Roy O, et al. (1949) Histoplasmin sensitivity; preliminary observations in a group of school children in the Province of Quebec. *Canadian journal of public health = Revue canadienne de sante publique* 40 (2):68-71
33. Leznoff A, Frank H, Taussig A, Brandt JL (1969) The focal distribution of histoplasmosis in Montreal. *Canadian journal of public health = Revue canadienne de sante publique* 60 (8):321-325
34. MacEachern EJ, McDonald JC (1971) Histoplasmin sensitivity in McGill University students. *Canadian journal of public health = Revue canadienne de sante publique* 62 (5):415-422
35. Sekhon AS, Isaac-Renton J, Dixon JM, Stein L, Sims HV (1991) Review of human and animal cases of coccidioidomycosis diagnosed in Canada. *Mycopathologia* 113 (1):1-10
36. Denning DW, Pleuvry A, Cole DC (2011) Global burden of chronic pulmonary aspergillosis as a sequel to pulmonary tuberculosis. *Bulletin of the World Health Organization* 89 (12):864-872. doi:10.2471/blt.11.089441
37. Smith NL, Denning DW (2011) Underlying conditions in chronic pulmonary aspergillosis including simple aspergilloma. *Eur Respir J* 37 (4):865-872. doi:10.1183/09031936.00054810
38. Denning DW, Pleuvry A, Cole DC (2013) Global burden of allergic bronchopulmonary aspergillosis with asthma and its complication chronic pulmonary aspergillosis in adults. *Med Mycol* 51 (4):361-370. doi:10.3109/13693786.2012.738312
39. Baxter CG, Dunn G, Jones AM, Webb K, Gore R, Richardson MD, Denning DW (2013) Novel immunologic classification of aspergillosis in adult cystic fibrosis. *J Allergy Clin Immunol* 132 (3):560-566.e510. doi:10.1016/j.jaci.2013.04.007
40. Agarwal R (2011) Severe asthma with fungal sensitization. *Current allergy and asthma reports* 11 (5):403-413. doi:10.1007/s11882-011-0217-4
41. Foxman B, Muraglia R, Dietz JP, Sobel JD, Wagner J (2013) Prevalence of recurrent vulvovaginal candidiasis in 5 European countries and the United States: results from an internet panel survey. *Journal of lower genital tract disease* 17 (3):340-345. doi:10.1097/LGT.0b013e318273e8cf
42. Lalla RV, Latortue MC, Hong CH, Ariyawardana A, D'Amato-Palumbo S, Fischer DJ, Martof A, Nicolatou-Galitis O, Patton LL, Elting LS, Spijkervet FK, Brennan MT (2010) A systematic review of oral fungal infections in patients receiving cancer therapy. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer* 18 (8):985-992. doi:10.1007/s00520-010-0892-z
43. Matee MI, Scheutz F, Moshy J (2000) Occurrence of oral lesions in relation to clinical and immunological status among HIV-infected adult Tanzanians. *Oral diseases* 6 (2):106-111
44. Buchacz K, Baker RK, Palella FJ, Jr., Chmiel JS, Lichtenstein KA, Novak RM, Wood KC, Brooks JT (2010) AIDS-defining opportunistic illnesses in US patients, 1994-2007: a cohort study. *Aids* 24 (10):1549-1559. doi:10.1097/QAD.0b013e32833a3967

- 1 45. Zhu YX, Li T, Fan SR, Liu XP, Liang YH, Liu P (2016) Health-related quality of life as
2 measured with the Short-Form 36 (SF-36) questionnaire in patients with recurrent
3 vulvovaginal candidiasis. *Health and quality of life outcomes* 14:65. doi:10.1186/s12955-016-
4 0470-2
- 5 46. Fong IW (1996) Clinical and cost considerations in the pharmacotherapy of vulvovaginal
6 candidiasis. *PharmacoEconomics* 9 (6):497-505
- 7 47. Public Health Agency of Canada - Surveillance. (2016) Public Health Agency of Canada.
8 <http://www.phac-aspc.gc.ca/surveillance-eng.php>. Accessed 2016-09-16 2016
- 9 48. Azie N, Neofytos D, Pfaller M, Meier-Kriesche HU, Quan SP, Horn D (2012) The PATH
10 (Prospective Antifungal Therapy) Alliance(R) registry and invasive fungal infections: update
11 2012. *Diagn Microbiol Infect Dis* 73 (4):293-300. doi:S0732-8893(12)00239-8 [pii]
12 10.1016/j.diagmicrobio.2012.06.012 [doi]
- 13 49. Haider S, Rotstein C, Horn D, Laverdiere M, Azie N (2014) The Prospective Antifungal
14 Therapy Alliance((R)) registry: A two-centre Canadian experience. *Can J Infect Dis Med*
15 *Microbiol* 25 (1):17-23
- 16