Sedimentology of the Essaouira Basin (Meskala Field) in context of regional sediment distribution patterns during upper Triassic pluvial events

DOI:

[10.1016/j.jafrearsci.2017.02.012](https://doi.org/10.1016/j.jafrearsci.2017.02.012)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Mader, N. K., Redfern, J., & El Ouataoui, M. (2017). Sedimentology of the Essaouira Basin (Meskala Field) in context of regional sediment distribution patterns during upper Triassic pluvial events. *Journal of African Earth Sciences*, 130, 293-318. <https://doi.org/10.1016/j.jafrearsci.2017.02.012>

Published in:

Journal of African Earth Sciences

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Accepted Manuscript

Sedimentology of the essaouira basin (meskala field) in context of regional sediment distribution patterns during upper triassic pluvial events

Nadine K. Mader, Jonathan Redfern, Majid El Ouataoui

PII: S1464-343X(17)30056-0
DOI: 10.1016/j.jafrearsci.2017.02.012
Reference: AES 2813
To appear in: *Journal of African Earth Sciences*

Received Date: 23 March 2016
Revised Date: 19 December 2016
Accepted Date: 06 February 2017

Please cite this article as: Nadine K. Mader, Jonathan Redfern, Majid El Ouataoui, Sedimentology of the essaouira basin (meskala field) in context of regional sediment distribution patterns during upper triassic pluvial events, *Journal of African Earth Sciences* (2017), doi: 10.1016/j.jafrearsci.2017.02.012

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**SEDIMENTOLOGY OF THE ESSAOUIRA BASIN (MESKALA FIELD) IN CONTEXT OF
REGIONAL SEDIMENT DISTRIBUTION PATTERNS DURING UPPER TRIASSIC PLUVIAL
EVENTS**

(Nadine K. Mader , Jonathan Redfern , Majid El Ouataoui)

HIGHLIGHTS

- This paper presents the first comprehensive depositional model for Upper Triassic sediments in the central Essaouira Basin.
- Results of this contribution add to the understanding of regional sedimentation patterns of (Upper) Triassic deposits in SW Morocco.
- The depositional model points towards much broader Triassic rift-basin geometries than previously expected, hence challenges previous representations of the evolution of the Atlantic rift during Triassic times.
- The study further provides new evidence for paleo-climatic reconstructions of the Moroccan Atlantic margin during Triassic times, presenting new data to support the southern extend of potential Carnian or ?Norian pluvial events evident in the investigated sequences of SW Morocco.
- The paper benefits from unpublished subsurface core data from the Essaouira Basin, including detailed core and lithofacies descriptions, data that is typically difficult to access.

SEDIMENTOLOGY OF THE ESSAOUIRA BASIN (MESKALA FIELD) IN CONTEXT OF REGIONAL SEDIMENT DISTRIBUTION PATTERNS DURING UPPER TRIASSIC PLUVIAL EVENTS

*Nadine K. Mader^{1,2}, Jonathan Redfern¹, Majid El Ouataoui³

¹North Africa Research Group, School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Williamson Building, Oxford Road, Manchester, M13 9 PL, United Kingdom.

²current Address: Maersk Oil Norge AS, Moseidsletta 122, 4068 Stavanger, Norway (madernadine@hotmail.com).

³ONHYM, 5 Avenue Moulay Hassan, B.P.99, 10050, Rabat, Morocco.

* CORRESPONDING AUTHOR

Keywords: Upper Triassic; Morocco; depositional reservoir model; regional sediment distribution; Triassic climate

ABSTRACT

Upper Triassic continental clastics (*TAGI: Trias Argilo-Greseux Inferieur*) in the Essaouira Basin are largely restricted to the subsurface, which has limited analysis of the depositional environments and led to speculation on potential provenance of the fluvial systems. Facies analysis of core from the Meskala Field onshore Essaouira Basin is compared with tentatively time-equivalent deposits exposed in extensive outcrops in the Argana Valley, to propose a process orientated model for local versus regional sediment distribution patterns in the continuously evolving Moroccan Atlantic rift during Carnian to Norian times. The study aims to unravel the climatic overprint and improve the understanding of paleo-climatic variations along the Moroccan Atlantic margin to previously recognised Upper Triassic pluvial events.

In the Essaouira Basin, four facies associations representing a progressive evolution from proximal to distal facies belts in a continental rift, detached from direct effects of sea-level changes were established. Early ephemeral braided river systems are followed by a wet aeolian sandflat environment with a strong arid overprint (FA1). This is followed by the onset of perennial fluvial deposits with extensive floodplain fines (FA2). The fluvial facies transitions to a shallow lacustrine or playa lake delta environment (FA3), accompanied by a distinct shift in fluvial style suggesting increase in discharge and related humidity, either locally in the basin or the catchment area. The succeeding deltaic environment exhibits cyclical abandonment and is progressively overlain by a terminal playa with extensive, mottled mudstones (FA4), interpreted to present a return from cyclical humid-arid conditions to prevailing aridity in the basin.

In terms of regional distribution and sediment source provenance, paleocurrent data from Carnian to Norian deposits (T5 to T8 member) in the Argana Valley suggest paleoflow focused towards the S and SW, not directed towards the Meskala area in the NW as previously suggested. A major depo-centre for fluvial

sediments is instead located in the southern Argana Valley, possibly the Souss Basin. To effectively source the reservoir sandstones found in the Meskala Field, a more local provenance area has hence to be envisaged. Despite this, the direct comparison of the genetic evolution of sedimentary sequences in the Argana Valley and Essaouira Basin shows a similar progression from dominantly arid ephemeral depositional environments to humid perennial sedimentation, returning to prominent arid conditions. This suggests climatic control in both regions, where an enhanced humid signal drives perennial fluvial flow in otherwise arid dominated sequences. On a regional scale, this is suggested to record the impact of strong Triassic pluvial events previously recognised in other basins along the Central Atlantic margin during the Carnian to Norian periods.

1. INTRODUCTION

Prior to the exploration efforts in SW Morocco, the development of new theories about global continent evolution during the initial stages of rifting along the Atlantic margin had already highlighted the importance of the investigated Moroccan basins for global pre-drift reconstructions (e.g. Van Houten, 1977; Manspeizer *et al.*, 1978; Jansa & Wiedmann, 1980; Stets & Wurster, 1981; Hinz *et al.*, 1981; Manspeizer, 1982; Laville & Petit, 1984; Manspeizer, 1988; Beauchamp, 1988; Medina, 1991, 1995; Piqué & Laville, 1996; Piqué *et al.*, 1998; Baudon *et al.*, 2012; Leleu *et al.*, 2016). With the discovery of hydrocarbons in the central part of the Agadir-Essaouira Basin in the early 1990's numerous studies evaluated the Triassic petroleum systems of Morocco (e.g. Broughton & Trépanier, 1993; Morabet *et al.*, 1998; Hafid *et al.*, 2000; Bouatmani *et al.*, 2004). Apart from a facies classification based on electrical log character of wells in the Meskala Field (Bouatmani *et al.*, 2004), most studies have a strong focus on the structural evolution with little focus on integrated sedimentological studies, leaving the regional facies distribution and provenance for the prolific Upper Triassic sandstone reservoirs within the greater Agadir-Essaouira Basin open to speculation. Broughton & Trépanier (1993) for example suggested the entire Triassic sequence of the Argana Valley as the eastward extension of the central Essaouira Basin, hence part of the sediment routing from the provenance area, while seismic interpretations and new dating of the overlying basalts led Hafid *et al.* (2000) to the conclusion, that only the upper part (Middle to Upper Triassic) of the Argana series was connected to the central Essaouira Basin, whereas the lower part was deposited in an isolated rift arm. To date, the relationship between Upper Triassic clastic sediments in the Argana Valley to Triassic reservoir sandstones of the central Essaouira Basin remains uncertain. In order to improve the regional understanding

of the spatial distribution of these extensive fluvio-aeolian reservoir facies in SW Morocco, this study presents new sedimentological analyses of core data from the Upper Triassic continental strata of the central Essaouira Basin (Meskala Field), integrated with results from sedimentological outcrop studies of the Argana Valley (Tadrart Ouadou Sandstone member, T6; Mader, 2005; Mader & Redfern, 2011), and published descriptions of additional Carnian-Norian deposits (T4 to T8) from the Argana Valley (Brown, 1980; Hofmann *et al.*, 2000). The study aims to integrate the sedimentological data within a regional climatic framework, improving the understanding of the control on sedimentation of extreme Triassic humid events as summarised by Preto *et al.* (2010) or proposed by Arche & Lopez (2014) for example.

2. GEOLOGICAL SETTING

Triassic sediments in Morocco outcrop within the Middle and High Atlas mountain chain, along the western margin of the African platform, and have also been encountered by wells in the Essaouira and Souss Basins, along the present offshore Moroccan continental margin (figure 1). The Triassic sequences of Northwest Africa typically overlay a major unconformity and are deposited on Palaeozoic rocks deformed by the Hercynian and older orogenies. Deposition of the Triassic sequences took place in a series of (half-) grabens or strike-slip basins related to rifting along the Atlantic margin during the break-up of the Pangaeon continent (Mattis, 1977; Van Houten, 1977; Manspeizer *et al.*, 1978; Laville & Petit, 1984; Medina, 1988, 1991, 1995) (figures 1, 2). Recent work suggests a change in how rift-basin geometry along the Central Atlantic Margin is viewed, and rather than the typical narrow rift-basin geometries more uncharacteristic broad sag-type basin styles have been reported, particularly for the Eastern provinces (Baudon *et al.*, 2012; Leleu *et al.*, 2016). The Triassic deposits infilling these basins commonly consist of coarse alluvial conglomerates, continental fluvial or aeolian sandstones, and alternations of fine siltstones, with evaporitic or non-evaporitic (playa) mudstones. In most parts of Morocco, the Triassic red beds are capped by the CAMP basalts (Central Atlantic Margin Province), which in turn are conformably transgressed, or interfinger, with limestones, dolomites, algal stromatolites, anhydrites, mudstones and halite deposits of an Early Jurassic sabkha environment (e.g. Courel *et al.*, 2003).

3. STRUCTURAL EVOLUTION

Mesozoic sequences in SW Morocco record the break-up of the super-continent Pangaea where crustal thinning and deep seated thermal activity led to the development of the Atlantic Ocean (Veevers, 1994). The Moroccan passive margin on the eastern side of the central Atlantic Ocean (e. g. Van Houten, 1977;

Manspeizer *et al.*, 1978; Piqué *et al.*, 1998) demonstrates at least four major tectonic phases influencing deposition of Latest Palaeozoic to Early Mesozoic strata (Manspeizer *et al.*, 1978; Laville & Petit, 1984; Piqué *et al.*, 1998). In a time period covering the Permian to the Middle Triassic, extensive uplift with related crustal thinning along the axis of the future Atlantic defines the initial phase of break-up (Van Houten, 1977). The second, late Middle Triassic tectonic phase, resulted in strike-slip faulting along E-trending fracture zones and initiated W trending Tethys transgressions (Piqué *et al.*, 1998). During the Late Triassic, continued rifting along the proto-Atlantic axis led to the deposition of clastic-evaporite sequences up to 5000 m thick in a relatively broad rift-setting with sag-geometries (Leleu *et al.*, 2016) (figure 1, 2B). Shearing occurred dominantly along E- to W-trending fracture zones inducing the decoupling of African and North American segments (Manspeizer *et al.*, 1978), creating a pathway for Tethyan transgressions from E to W. In a last major tectonic phase (Late Triassic to Early Jurassic) global plate reorganisation led to crustal extension and seafloor spreading, resulting in massive extrusions of olivine and quartz tholeiitic basalts and igneous rock emplacement across a large area of North America and West Africa described as the Central Atlantic Margin Province (CAMP) (Marzoli *et al.*, 1999; McHone, 2000; Olsen *et al.*, 2000; Youbi *et al.*, 2003; Knight *et al.*, 2004; Marzoli *et al.*, 2004; Verati *et al.*, 2007; Whiteside *et al.*, 2010; Deenen *et al.*, 2010). During this last phase, a strong transcurrent component with transtension is proposed by Laville & Petit (1984), while Piqué *et al.* (1998) suggest normal faulting during this final syn-rift stage in Morocco.

4. BASIN FILL

The Agadir-Essaouira Basin forms part of the El Jadida-Agadir rift segment (figure 1). It developed as a set of E-W oriented half-grabens and underlies an extensive area of the modern day eastern passive Atlantic margin (Piqué *et al.*, 1998). The nomenclature utilised to describe the Agadir-Essaouira Basin varies widely due to the complex basin segmentation and the lack of sufficient sub-surface or outcrop control (e.g. Société Chérifienne des Pétroles, 1966; Brown, 1980; Broughton & Trépanier, 1993; Piqué *et al.*, 1998; Hafid, 2000). For consistency, the sediments of the Meskala Field are described in this paper as clastics deposited within the “central Essaouira Basin”, and the term “Argana Valley” is adopted synonymously to describe clastic deposits exposed in the present day Argana physiographic depression due to the lack of a so far clearly defined outline or structural framework for a potential “Argana Basin”.

4.1 Central Essaouira Basin (Meskala Field)

The Essaouira Basin is situated in central western Morocco and extends towards the N and S of the city of Essaouira, offshore and onshore, along the present day coastline (figures 1, 2). The central basin is limited in the N by the Tensift-fault zone, and to the S by the Amsitten fault complex (figure 1). The basin contains an up to 4500 m thick sequence of Mesozoic strata (Broughton & Trépanier, 1993) (figure 3). Middle Triassic to Lower Jurassic continental red beds were deposited in an extensional rift setting, overlain by several kilometres of salt (figure 2C). Salt deposition was largely driven by marine incursions into the deepening basins from the Tethys during Upper Triassic to Lower Jurassic times (Courel *et al.*, 2003). Extensive emplacement of basalts reaching thicknesses of up to several hundred meters occurred around the Triassic-Jurassic boundary, and can be related to the CAMP magmatism (McHone, 2000; Knight *et al.*, 2004; Nomade *et al.*, 2007; Deenen *et al.*, 2010) (figure 2C). During the Atlasic (Alpine) tectonic phase from Late Eocene to Oligo-Miocene, the Essaouira Basin was moderately uplifted and tilted towards the W, which led to widespread erosion of overlying Cretaceous deposits. To the E, the onshore part of the Essaouira Basin is still buried below younger Mesozoic strata. Uncertainties exist in the thickness and distribution of the (Late) Triassic sequences towards the SE, in direction of the Argana Valley Triassic exposures due to the thick coverage of Jurassic and Cretaceous series and the lack of seismic control (e.g. Piqué *et al.*, 1998; Hafid *et al.*, 2000; LeRoy & Piqué, 2001).

4.2 Argana Valley

Extensive exposures of Permian and Triassic sediments are exhibited in the Argana Valley, located along the roughly N-S trending modern day physiographic depression, which runs parallel to the southern extents of the High Atlas mountain chain between Imi N'Tanoute in the N and Ameskrout in the S (figure 1). The valley developed during Cenozoic times, when rivers draining southward from the emergent High Atlas eroded through Cretaceous and Jurassic strata, resulting in the exposure of nearly 1500 km² of rocks. The Permian to Early Jurassic continental red bed sequence rests unconformably on Palaeozoic sediments, and attain an overall thickness of 4500 m (Duffaut *et al.*, 1966; Tixeront, 1973). The Argana basalt reaches thicknesses of up to 400 m and separates the Permo-Triassic lithostratigraphic members T1 to T8, from the Lower Jurassic units T9 and T10 (AitChayeb *et al.*, 1998; ElHachimi *et al.*, 2011) (figure 3). Radiometric ages have been recalculated for basalt flows to 197 ± 0.7 and 201.7 ± 2.4 Ma (Verati *et al.*, 2007) (table 1), suggesting a stratigraphic age of Sinemurian to Rhaetian calibrated to the timescale of Gradstein *et al.* (2012), associating them to the CAMP magmatism (Knight *et al.*, 2004; Marzoli *et al.*, 2004; Verati *et al.*,

2007; Deenen *et al.*, 2010). In contrast to the central Essaouira Basin, there is no direct evidence for marine incursions into the Argana basin during Upper Triassic or Lower Jurassic times (Hofmann *et al.*, 2000). Recent structural studies (Baudon *et al.*, 2012; Leleu *et al.*, 2016) highlight dominantly wide-rift or broad sag-basin styles for the eastern provinces, supporting earlier diagnostic work of Hofmann *et al.* (2000). During the Cenozoic, Alpine deformation caused high-angle faulting in the Central High Atlas, but only minor folding of the Mesozoic sedimentary rocks occurred. The tectonic overprint led to the formation of fault blocks, but in most cases no steep tilting took place and in general the dip of Triassic strata is less than 30 degrees (Van Houten, 1977; Manspeizer *et al.*, 1978; Manspeizer, 1988; Beauchamp, 1988; Manspeizer, 1994; Medina, 1995; Mader & Redfern, 2011; Baudon *et al.*, 2012).

5. UPPER TRIASSIC STRATIGRAPHY

Dating is a challenge in continental settings, which commonly lack direct biostratigraphic control. Palynological data is sparse, and the only conclusive age data available that is comparable between the two regions of investigation is presented by the basalt deposits related to the CAMP volcanism. Argana Valley basalt flows yield an absolute age of 197 ± 0.7 Ma and 201.7 ± 2.4 Ma. Within the central Essaouira Basin, dolerite samples from well JRP-1 have an absolute age of 210 Ma (Oujidi *et al.*, 2000) (figure 2C). No conclusive ages for the thick flow basalt sequences within the Meskala Field were however available (figure 2C). A summary of the stratigraphic framework calibrated by the sparse relative age data is provided for the two areas (figure 3, table 1) to allow for an integrated conclusion on the time-equivalence of the two areas and implications made in view of regional sedimentation patterns.

5.1 Central Essaouira Basin

A tectono-stratigraphic framework for the Essaouira Basin based on seismic interpretation was proposed by LeRoy & Piqué (2001) (figures 2B, 3). Within this tectono-stratigraphic framework the Mesozoic sequences were sub-divided into four sedimentary mega-sequences (LeRoy *et al.*, 1997; Manspeizer *et al.*, 1978; Piqué *et al.*, 1998)(figure 2B). The lowermost Upper Triassic to Lower Jurassic mega-sequence comprises red, silt-rich clastic deposits, thick evaporites and intercalated basalts, deposited during the syn-rift phase (Piqué *et al.*, 1998; LeRoy & Piqué, 2001). This mega-sequence is composed of three smaller-scale sedimentary sequences. The lowermost sequence sT1 is restricted to a westward tilted half-graben (Piqué *et al.*, 1998; LeRoy & Piqué, 2001), while the second sequence sT2 is thickest along the western border of the half-

graben (figure 2C). This sequence has been penetrated by well JPR-1, and comprises fluvial sandstones of Carnian age (Piqué *et al.*, 1998; LeRoy & Piqué, 2001) (figure 2C). The uppermost unit sT3 overlies an interpreted intercalated basalt and the maximum thicknesses of sT3 can be observed further towards the west (figures 2B,C). This highlights a general westward shift of depo-centres associated with increased thickness as suggested by Piqué *et al.* (1998) (figure 2B). Dolerite samples of the intercalated basalts were available from well JRP-1 (2768 m, 3163 m; 210 Ma; Oujidi *et al.*, 2000), suggesting a Norian or younger age (calibrated to Gradstein *et al.*, 2012) (figures 2B,C).

In the Meskala Field the Upper Triassic continental facies are overlain by reoccurring mudstones, and approximately 240 m of salt, eventually capped by an approximately 270 m thick basalt sequence (figures 2C, 4). A lower Liassic age has been assumed for the salt and mudstone sequences (Hafid *et al.*, 2000; Broughton & Trepanier, 1993; Youbi *et al.*, 2003). The thick basalt sequence can be traced regionally on seismic into the Jeer Field (Hafid *et al.*, 2000) (figure 2C) and has also been penetrated by wells in the Meskala Field (figure 4). Both, the extrusive basalts as well as the intrusive sills found within the Upper Triassic deposits (figure 2C) are expected to be related to the well documented CAMP event (McHone, 2000; Knight *et al.*, 2004; Nomade *et al.*, 2007; Deenen *et al.*, 2010). The limited radiometric ages from JRP-1 support this relationship, and the correlation is reinforced by more conclusive biostratigraphic data available for nearby wells MKL-1, where ostracodes of the genus *Darwinula* (between 3027-3300 m), and of the genus *Lymnocythere* in MKL-104 (2964 m) (Silmane & ElMostaine, 1997) were recorded, both supporting a Carnian to Norian age in line with the radiometric ages of the basalts. Since the CAMP event is documented to be very short-lived with an expulsion window as short as 1,5 Ma (McHone, 2000; Nomade *et al.*, 2007), some confidence with respect to the absolute timing is hence placed on the proposed chronostratigraphic framework.

5.2 Argana Valley

The exposed rift-sequences have formerly been divided into the Ikakern, Timezgadiouine and Bigoudine Formations, bound at the top by the Argana Basalt (Duffaut *et al.*, 1966; Tixeront, 1973) (figure 3). The formations have further been subdivided into ten lithostratigraphic units numbered T1 to T10 (Tixeront, 1973; Brown, 1980; Hofmann *et al.*, 2000). The Permian Ikakern Formation (T1 and T2) is characterised by alluvial fan and high-sinuosity meander-plain conglomerates with interbedded sandstone and siltstone alternations (Tixeront, 1973; Jones, 1975; Brown, 1980), the Timezgadiouine Formation (T3 to T5) consists

of braided or meandering fluvial channels, and cyclical playa deposits (Tixeront, 1973; Brown, 1980; Hofmann *et al.*, 2000). The base of the Bigoudine Formation rests conformably on underlying mudstones and fines, but can locally be unconformable and erosive exhibiting coarse ephemeral alluvial conglomerates (Mader, 2005; Mader & Redfern, 2011). The greatest part of the basal T6 member however consists of braided river deposits, perennial fluvial sandstones and aeolian dunes (Mader & Redfern, 2011), while the thickest part of the Bigoudine Formation (T7 to T10) displays highly cyclical and laterally extensive playa deposits (mudstones, siltstone, sandstone intercalations) (Tixeront, 1973; Brown, 1980; Hofmann *et al.*, 2000; Tourani *et al.*, 1999). A relatively well constrained biostratigraphic framework was established for the Permo-Triassic Argana Valley sequences combining the work of previous authors with new biostratigraphic data (Jones, 1975; Dutuit, 1976 & 1977; Manspeizer, 1978; Fiechtner *et al.*, 1992; Tourani *et al.*, 1999; 2010; Jalil & Peyer, 2007; Jalil *et al.*, 2009; Klein *et al.*, 2010, 2011; Verati *et al.*, 2007) (table 1). The chronostratigraphic framework suggests a Carnian to Norian age for the (uppermost?) T4, T5, T6, T7 and T8 deposits (figure 3). Consequently a review of T4/T5, T6 and T7/T8 members is provided in this study in view of major fluvial flow in the Argana Valley analogous to depositional ages anticipated for the Essaouira sequence in the Meskala Field.

6. CORE AND FACIES ANALYSIS: CENTRAL ESSAOUIRA BASIN, MESKALA FIELD

Core analysis of wells from the Meskala Field was undertaken (this study) and results were subsequently compared to outcrop data and literature from the Argana Valley. An almost fully cored section from well MKL101bis is presented and discussed in terms of the sedimentological evolution of the basin. MKL101bis was chosen as the example well, as it contains the most continuous Triassic section cored in the Meskala Field (figure 4). It hence offers the possibility to study a continuous vertical Triassic profile without major gaps. The generalised facies associations resulting from this work are under-pinned by additional wells. In total ten litho-sedimentary facies (*sub-facies*) were identified and combined into four facies associations (FAs) (table 2). Facies associations of well MKL101bis are discussed from base to top, the core is described with relative thickness in centimetres, and core gaps were noted where present (figures 5 to 10). The detailed facies description and interpretation for the core intervals is provided in tables 2 to 5, while in the following a brief summary of the facies associations is provided.

6.1 Summary facies associations

Facies Association 1 (FA1) consists of three *sub-facies* and is dominated by very fine to fine grained sandstones (table 3; figure 5). Finer material is sparse, but minor amounts of clay can be found in the form of thin (< 10 cm) mudstone horizons. Sedimentary structures include horizontal or low angle stratification (figure 9A), massive bedding or inverse grading. Secondary angular cavities (<1 cm in diameter) were abundant within this sandstone dominated unit. The facies association was interpreted to represent ephemeral fluvial braided river deposition in transition to a wet aeolian sandflat environment (table 3).

Dissimilar to FA1, FA2 comprises four *sub-facies* and is dominated by compound, well-sorted, very fine to fine-grained sandstone units interbedded with thick mudstone intervals (table 4; figure 6). The thickness of individual sandstone packages ranges from approximately 50 cm to up to 4 m, with interbedded dark red to purple mudstone units up to 2 m thick, and minor siltstones interspersed. The ratio between sandstone and mudstone dominated intervals is approximately 3:2, and sediments of FA2 were interpreted to represent perennial fluvial channels with adjacent floodplain fines (table 4).

For FA3 three *sub-facies* were distinguished (table 5; figures 7 to 8), which are repeated in three distinct fining-upward cycles. The two lower cycles, consist of 8 to 10 m thick sandstone dominated fining-upward sequences, while the upper cycle exhibits about 4-5 m thickness (figures 8, 10). Each fining upward cycle features a sharp, locally erosive basal contact, and the sandstones are interbedded with minor mud- to siltstone intervals (figures 7, 8, 10). Overall, FA3 is dominated by fine to very fine-grained, ripple cross-bedded to less abundant massive or destratified sandstones. Calcretes, or coaly interbeds are abundant, and the sequences is generally overprinted by root traces or mottling throughout. Dewatering structures and wavy horizontal lamination are abundant, and only remnant horizontal to low-angle lamination can be observed in the sandstones. Overall, a transition from fluvial delta top to shallow ?lacustrine or playa lake delta deposition is suggested (table 5).

In contrast to the sandstone dominated FA1 to FA3, FA4 is dominated by dark red to purple, mottled mudstones which can change progressively into dark red to purple silt- to mudstones with minor horizontal or wavy-horizontal lamination, regularly reverting back to cleaner mudstone intervals (table 5; figure 8). Minor very fine to fine grained sandstone intervals are interbedded in the form of small-scale (<1cm) adhesion ripple stratification. Dissimilar to the lower units and mudstones observed interbedded in the

sandstone dominated in FA2 and FA3, no intercalated dark grey to black mudstone intervals or thicker (>1 cm) sandstone intervals were observed. Full playa deposition with little clastic input to the system is hence anticipated (table 5).

7. DEPOSITIONAL MODEL ESSAOUIRA

Taking into account the climatic overprint inferred from sedimentary structures recorded in core, a progressive evolution from FA1 to FA4 suggests the following major depositional stages:

- **FA1:** ephemeral braided river deposition to wet aeolian sandflat environment (figure 11A);
- **FA2:** perennial fluvial deposits with extensive floodplain fines (figure 11B);
- **FA3:** fluvial deposits in transition to a shallow lacustrine delta environment with distributary mouthbars and cyclical abandonment of the fluvial facies (figure 11C);
- **FA4:** playa environment with extensive, highly mottled mudstones.

With the exception of FA1 and FA4, the main reservoir intervals comprise stacked fining-upward sequences interbedded with thick, occasionally organic-rich mudstones (figures 10, 11). The sandstones in FA1 differ from the overlying sandstone deposits of FA2/FA3 by being slightly coarser in grainsize, less well sorted, lacking fines and exhibiting a different fluvial style. The FA1 sandstones are typically horizontally laminated and imply sheet-like deposition with a high relatively well sorted sandstone content and little fines (figure 11A). The lack of fines is assumed to be related to proximity to source, but could also imply lateral confinement (e.g. Wright & Marriott, 1993). Overall, the sandstones of FA1 were interpreted to represent episodic discharge in a shallow braided river system (figure 11A), and in comparison to FA2 and FA3 a more proximal depositional setting is proposed. Within FA1 an obvious change from dark grey mudstones enriched with organic matter, to dark red to purple, often mottled mudstones was observed. While the presence of organic matter implies vegetation and humid conditions to transport and preserve the plant material, the shift from overall reducing to oxidising conditions is inferred from the colour change. More humid conditions, at least in the catchment area, are therefore proposed for the lower part of the system, but a shift in local climate to more arid (oxidising) conditions is apparent. Abandonment of the fluvial system is suggested, marked by an immature paleosol implying an extended time of exposure (*sub-facies 1b*). Above this internal bounding surface, a phase of amplified aridity is interpreted to have led to the development of a wet aeolian sandflat environment (*sub-facies 1c*). Increased winnowing of the sandstones, a dominance of

horizontally bedded and inverse graded deposits suggest augmented evaporation processes with fluctuating groundwater levels. This interpretation is supported by the recognition of a patchy anhydrite cement overprinting *sub-facies 1c*. This type of cementation has previously been associated with alternating groundwater levels, where above the groundwater table in the presence of saline waters and evaporation, the anhydrite is precipitated in form of patchy cements in an overall arid setting (Glennie, 1970).

A prominent change in depositional environment from sheet-like ephemeral flow to metre-scale fluvial channel deposits and abundant, thick floodplain fines follows FA1 (figure 11B). Several fining-upward sequences of well sorted, very fine grained sandstone packages associated with extensive dark red mudstones are documented (FA2). A fluvial setting with abundant crevasse splay and overbank deposition within laterally shifting facies belts associated with the migration of channels within an extensive floodplain environment is proposed (e.g. Miall, 1996) (figure 11B). The fining-upward profiles may reflect cycles of increased water availability and discharge, or simply lateral autocyclic migration and avulsion of the system. Extensive dewatering of the sediments and syn-sedimentary faulting observed in these units however implies relatively oversaturated sandstones, possibly related to more humid conditions. The increased influx of water to the area may have further led to a raised groundwater level, which favoured the dissolution of early calcites or evaporates indicated by the angular cavities prominent in FA1. The colour of the interbedded floodplain fines points towards overall oxidising conditions, however where deposition was rapid, preservation of dark black mudstones deposited under reducing conditions is possible. Alternatively, shallow semi-permanent lakes on the floodplain could be envisaged. The presence of reworked organic-rich to almost coaly reworked deposits in an overall oxidised floodplain environment is thus interpreted to suggest cyclical shifts between humid and more arid conditions, whereby the humidity could have been restricted to the catchment area.

The fluvial deposits become increasingly interdigitated with transgressive deposits, suggesting the expansion of an adjacent shifting shoreline (FA3) (figure 11C). Each of the fining-upward sequences exhibits deposition of dark grey or black organic-rich mudstones at the top (figure 10) interpreted to document the direct influence of a longer-lived shifting shoreline. These black organic-rich deposits change in composition throughout the interval, from organic-rich mudstones with organic wood-plant debris <40%, to coaly layers composed of 80 to 100% organic material, towards the top. The sandstones change from clearly defined fluvial channel deposits to dominantly ripple cross-stratified sandstones with abundant organic-rich to almost in-situ coal beds. The environment of deposition changes into a delta setting with distributary mouthbars and

abundant inter-pond areas (figure 11C), with more permanent water available. Simultaneous with the increase in organic matter to the system, the input of sandstones seems to decrease considerably. This decrease in clastic input could be an effect of numerous scenarios: less fluvial input related to decreased runoff and possibly an increase in aridity in the hinterland (Schumm, 1968b; Smoot, 1991; Shanley & McCabe, 1994; Howell & Mountney, 1997; Blum & Törnqvist, 2000; Nanson *et al.*, 1998), a less efficient hinterland source (e.g. Leleu & Hartley, 2010), or simply basin closure and back-filling or back-stepping of the system (e.g. Schumm, 1993). While an increase in humidity would be inferred from the high amounts of organic matter present, strong evidence for increased aridity is provided by the paleosols observed in the system. The formation of these nodular calcretes requires sufficient precipitation to translocate the carbonate, but equally prerequisites high evaporation rates (Reeves, 1976). Typically they form during arid to semi-arid intervals following humid periods (Wright, 1992; Alonzo-Zarza, 2003), or alternatively, calcretisation takes place in the vadose zone within the floodplain during arid climates (Wright, 1992; Wright & Marriot, 1996; Alonzo-Zarza, 2003; Khalaf, 2007). Both processes imply rather arid climates, which is also reflected in the increased destratification and related evaporation processes recognised in the host rock towards the top of the sequence. An increase in aridity, possibly within a system that still promotes high groundwater tables is therefore envisaged, and a combination of basin closure and changes to the prevailing climate back to more arid conditions is suggested.

Gradually, towards the top of the logged section, the input of coarser sandy material ceases entirely, and an extensive playa environment with dominantly reducing conditions, governs the last phase of deposition (FA4). Arche (2007) reviewed the appropriate use of the term playa, concluding that subsurface hydrogeology, evaporite geochemistry and tectonic regime are crucial to accurately define a particular playa or sabkha setting. With respect to the investigated deposits it remains open to discussion whether the term playa, i.e. deposits of continental dry depressions, versus sabkha defined as marine saline flats along arid shorelines is appropriate, as the exact nature of the shoreline could not be distinguished during this study. The sedimentological data suggests shifting shorelines, and only minor amounts of evaporite deposits were recognised. However, thick salt deposits of marine origin overlay the investigated deposits (figures 2B,C, 4), which certainly suggest marine incursion at a later stage. But since no direct marine indicators were recognised in the investigated deposits, a rechargeable or through flow playa system (Arche, 2007) with possibly a well-developed playa lake remains the preferred interpretation for the studied interval (FA4).

349 8. DISCUSSION: REGIONAL IMPACT

350 The Argana Valley is located approximately 80 km SE of the Meskala Field (figures 1, 12), where Carnian to
 351 Norian continental deposits (T4/5, T6, T7/T8) have been studied in outcrop by several authors (Tixeront,
 352 1973; Jones, 1975; Brown, 1980; Hofmann *et al.*, 2000; Tourani *et al.*, 2010; Mader & Redfern, 2011). In the
 353 past, the Argana sequence has been proposed to represent the proximal equivalent of the prolific reservoirs in
 354 the central Essaouira Basin (Brown, 1980; Broughton & Trépanier, 1993; Hafid *et al.*, 2000).

355 Owing to the lack of extensive subsurface and seismic data it is not possible to correlate directly between the
 356 Argana Valley exposures and the central Essaouira Basin subsurface section, thus these conclusions have
 357 been largely based on assumptions of structural trend predictions and limited paleocurrent data (e.g.
 358 Tixeront, 1973; Brown, 1980; Medina, 1991, 1995). Our study focuses on the sedimentological and climatic
 359 evidence from Carnian to Norian deposits in both areas, and conclusions that might be drawn from the
 360 sediments preserved. For the purpose of this comparative study, a synopsis of sedimentological work with
 361 emphasis on major fluvial flux in the Argana Valley as well as the concurrent climate is presented.

362 8.1 Literature review of Carnian to Norian deposits (?T4/T5, T6, T7/T8), Argana Valley

363 Based on Duffaut *et al.* (1966), Tixeront (1973) and Jones (1975) early work, Brown (1980) generated a
 364 synopsis on the tectono-stratigraphic evolution of the Argana Valley (figures 3, 12). In terms of facies
 365 distribution, Brown (1980) summarises the T4/T5 member as floodplain deposits with intercalated
 366 meandering river sandstones, T6 is described as a deltaic complex with westward prograding deltamouth
 367 bars, and T7/T8 were attributed to deltaplains and mud-flats with distributary channel deposition. Field
 368 measurements of flow directions led Brown (1980) to the conclusion that all sediments in the Argana Valley
 369 are initially derived from the E, and distributed towards the W following E-W trending basement faults
 370 (figure 12). He emphasises the predominance of normal faulting and suggests that some of the structures in
 371 the lower units functioned as tilted fault blocks, which influenced lateral sediment distribution throughout
 372 deposition of the entire Triassic sequences. His measurements for the Timezgadiouine (T4/T5) and
 373 Bigoudine Formation (T7/T8) led him to conclude that sediment dispersal is imposed by the Tirkou and
 374 Timezgadiouine horst (figure 12), diverting sediments S and N into the grabens respectively. Within the
 375 grabens paleocurrents are towards the W. Although not explicitly discussed, the measurements published by
 376 Brown (1980) clearly highlight a prominent S to SW component in addition to westerly trends, both in the

Timezgadiouine and the Bigoudine Formation (figure 12). Hofmann *et al.* (2000) presented detailed sedimentological work with a focus on paleoclimate in view of observed cyclicity, sediment accumulation history and basin evolution of the mudstone dominated sequences T4, and T7 to T10. Their work focuses on the mudstone dominated units and highlights a great degree of lateral continuity of cyclical mudstones deposits throughout the basin. The interbedded sandstone dominated facies has been described as unconfined sheetfloods, possibly related to ephemeral meandering streams (T4), a mixture of sheetflow deposits, often affected by haloturbation, sheet delta deposits with mud-draped wave ripple lamination, mixed fluvial aeolian massive deposits overprinted by haloturbation, and purely aeolian cross-bedded deposits interbedded with thick mudstone-claystone alteration of an arid saline shallow ephemeral lake environment (T7/T8). Hofmann *et al.* (2000) did not provide flow directions for any of the sandstone dominated facies, however highlights a general trend of progressive thickening and coarsening of the sandy facies. This coarsening upward trend is ascribed to a basinward migration pattern driven by a gradual climatic drying. Moreover, Hofmann *et al.* (2000) stresses the lateral continuity of T4, T7-T10, which they attribute to shifting facies belts between a marginal to a more central basinal position. A very flat basin topography is suggested based on the lateral extend of aeolian marker beds. Hofmann *et al.* (2000) further states that based on the cyclical stacking pattern, the extensive lateral extent of individual cycles and sheet-like character structural movement can be discounted. The observed cyclicity was assigned to oscillatory fluctuations in paleoclimate instead, and an overall drying upward cycle, which generated fluctuations in the playa basin driving lateral shifts in sediment distribution, was hence suggested.

8.2 Field study of T6, Argana Valley

An extensive field study (figure 12) focusing on the sandprone Tadrat Oudaou Sandstone member (T6) was carried out by Mader & Redfern (2011), who presented a comprehensive sedimentological model. Two main phases of sediment flux to the Argana Valley during T6 times were proposed, interrupted by a phase of pronounced aridity throughout the basin (figures 13-17). The early fluvial phase documented represents a coarse ephemeral alluvial fan to braidplain setting (figures 13, 14, 17A), characterised by high amounts of channel switching and restriction to the centre of the Argana Valley exposures (figure 12, locality 1, 2). Extensive flow directional measurements suggest dominantly E to W flow with sub-ordinate SE to NW flow directions (figures 12, 14). Remnant aeolian deposits as well as products of aeolian deflation are commonly interbedded throughout. Towards the W (figure 12, locality 2), in flow direction of the conglomeratic facies,

the fluvial deposits display a mere thickness of 2 m, which was attributed to reflect pinch-out of the system toward W and NW (figure 17A).

The second major fluvial phase, characterised by highly aggradational, and more perennial fluvial deposition, extensively distributed and exposed throughout the Argana Valley (figures 15, 16, 17B). The size and internal complexity of individual channels suggests a relatively long-lived fluvial system (figure 13A, 15). During this late fluvial phase flow directions of fine grained perennial channels are predominantly from N to S in both the northern most locality (figure 15A) and the southernmost locality (figure 15B). The southernmost locality (figures 12, locality 4) was interpreted to have acted as a major depo-centre for the basin, with maximum thicknesses of T6 preserved (up to 70 m) (figure 15B). E to W flow directions were established for the central locations (figure 12, locality 1), and a N-S running axial drainage system, fed by fluvial tributary systems from the E (for example in central Argana, locality 1) feeding the southern and southwestern Argana realm during this depositional stage was suggested (figures 14, 17B). Subtle changes in basin geometry affect spatial variations and facies variability throughout. The fluvial system is terminated along a sharp extensive upper bounding surface and the onset of aeolian sedimentation (Mader & Redfern, 2011), which is subsequently transgressed by lacustrine mudstones of the T7 member (Hofmann *et al.*, 2000).

8.3 Implications for regional sediment distribution

While growth faults are reported by Brown (1980) and Medina (1991; 1995) to be common, neither Hofmann *et al.* (2000), Mader & Redfern (2011) nor Baudon *et al.* (2012) recognise strong evidence for syn-sedimentary fault control during the Upper Triassic. Minor faults were attributed to dissolution of evaporites and linked collapse features (Hofmann *et al.*, 2000). Mader & Redfern (2011) highlight the possibility of limited fault controlled accommodation space only for the basal early fluvial phase in the centre of the Argana Valley (figure 12, locality 1), but point out that accommodation space equalised during the late fluvial phase of T6, and that facies distribution observed is a consequence of laterally shifting facies belts driven by the local climate rather than tectonic movement. Baudon *et al.* (2012) work suggest that the prominent NNE striking faults are post-depositional, and Hofmann *et al.* (2000) further concluded that sedimentary cycles observed within the Carnian to Norian T4, T5, T7 and T8 members related to a flat basin topography were correlatable with identical thickness over several 10s of kilometres. Baudon *et al.* (2012) who contributed the most recent structural review for the area, highlights the impact of the Massif Ancien as a

potential structural buffer between the Central High Atlas to explain the slower subsidence observed, and suggests a wide sag-basin type geometry for the Late Triassic section in the Argana Valley rather than narrow half-graben settings. This uncharacteristic broad basin morphology has further been documented on seismic lines in the Central Essaouira Basin (Hafid *et al.*, 2000), and is supported by a recent regional review of the Atlantic margin by Leleu *et al.* (2016). Both, the recent sedimentological and structural studies suggest that during Carnian to Norian times equalisation of accommodation space prevailed, and that the influence of deep seated structural highs as well as E-W trending basement faults had little to no influence on sedimentation as previously suggested (e.g. Brown, 1980, Laville & Petit, 1984; Medina, 1991, 1995). The main driving factor for the cyclicity observed were climate related, while lateral sediment distribution was attributed to autocyclic processes with fluctuating facies belts and compensational shifting (Hofmann *et al.*, 2000; Mader & Redfern, 2011).

Paleocurrent flow measurements from Brown (1980) and Mader & Redfern (2011), show two dominant directions of fluvial flow for the Timezgadiouine (T4-T5) and Bigoudine (T6-8) Formations:

- a) E to W flow direction;
- b) NE-SW and/or N to S.

Whilst an E-W paleocurrent component is prominent along the easternmost exposures, closest to the Eastern basin bounding margin, the Western exposures record a prominent N-S and NE-SW direction (figures 12, 17). A distributary system, feeding a southerly accommodation sink is suggested, not one focusing flow to the present day central Essaouira Basin in the NW (Meskala Field) (figure 1), casting further doubt on the model proposed by Brown (1980), Broughton & Trépanier (1993) and Hafid *et al.* (2000) who tentatively identified the Triassic sequence of the Argana Valley as the direct sediment provenance for the central Essaouira Basin.

Although the comparison of the Carnian to Norian sedimentary facies suit in the Argana Valley and the Essaouira Basin in context to common basin models (e.g. Matthews & Perlmutter, 1994) could suggest a direct proximal-distal relationship, regional sedimentological evidence is conflicting. For example, for deposition of T6 (Argana Valley) a model scenario could be envisaged, where T6 represent the proximal to mid-distal end member, and deposits from the central Essaouira Basin exhibit the transition from the mid-distal to distal play (lake) end member. To further use T6 fluvial deposits as a model scenario, a sediment

fairway with direct SE to NW fluvial flow from Argana to Essaouira would be required to specifically feed the accommodation space in the Meskala Field following this direct source to sink relationship (figure 1). While general facies distribution fits the model, field evidence from the Argana Valley of T4/T5, T6 and T7/T8 (Brown, 1980; Mader & Redfern, 2011) suggests, that direct SE to NW fluvial flow was only recognised close the easterly expected bounding fault entry points (Brown, 1980; Mader & Redfern, 2011), and subordinately for the early alluvial phase of T6, which is anticipated to be laterally restricted to central Argana Valley (figure 17A). Instead, S/SSW flow directions dominate along the western most exposures (figure 12), and both Brown (1980) and Mader & Redfern (2011) indicate increased accommodation space in the S. The onset of regionally extensive, isopachous lacustrine and playa strata of T7 to T10 (Hofmann *et al.*, 2000) would typically be interpreted as widespread post-rift sedimentation with major fluvial input to the system terminating in a closed basin setting. Leleu *et al.* (2016) point out that within the wide rift-basins and geometries, the onset of lacustrine strata could also fall into the late syn-rift stages of basin evolution, hence imply continued subsidence and accommodation space creation. A potential exit point for sediments in a fill and spill scenario would have hence had to be in the S, or SW/W (figure 1). The next ‘sink’ towards a southerly direction is the Souss Basin (figure 1), and Triassic deposits have been confirmed. Dependant on basin geometries towards the SW/W, interconnection and cross-basinal flow of sediment, changing flow direction from S to N in the newly developing western extensional basins, and then leading into the central Essaouira Basin area, would have to be envisaged (figure 1). This scenario would be analogous to fill and spill models for Triassic rift systems such as the Sherwood sandstone in the UK for example (Wills, 1970; Audley-Charles, 1970). With the progressive westward shift of active rift development in SW Morocco (LeRoy *et al.* 1997; 2001) a scenario where sediment is “stored” in the southern Argana or Souss Basin, later subsequently recycled and transported axially along the rift to the N through the newly developing western extensional basins, could be imagined. A certain diachronism between the source and sink would however have to be expected in this scenario.

The continuous evolution from proximal fluvial facies to distal delta or playa lake end member conditions in the Meskala Field presented in this study, are however anticipated to suggest a more local source to sink relationship. The Central Jebilet Massif located in the NW (figure 1) could present a more local hinterland source instead, or, similar to speculations for the Argana sediment routing (Mader & Redfern, 2011), the Anti Atlas could be envisaged as the easterly hinterland source, with a more northern point source than

Argana. Further regional provenance studies are required to reconstruct the Triassic hinterlands of Morocco with refined precision.

8.4 Regional climate

The climate-driven overprint on sedimentation in the two studied basins is considered the main force for sedimentary architectures, while higher-order changes in basin morphology and subsidence rates are assumed to have led to the sequence-scale basin fill. Sediments in both study areas were recognised to be strongly overprinted by the local climate and the climatic regime in the catchment-area (Hofmann *et al.*, 2000; Mader & Redfern, 2011; this study). Sedimentological evidence from the Argana Valley suggest a cyclical evolution from: (a) a dominance of arid climatic conditions with subordinate cyclical humid pulses, resulting in the deposition of stacked ephemeral channel deposits, and a documented direct effect on local groundwater tables affecting deposition-preservation of aeolianites within the fluvial sequences; (b) an increase in humidity (discharge) inferred from a change in fluvial architecture to perennial flow, with deeply incising channels reflecting an amplified climatic signal; (c) a return to aridity with full aeolian conditions and minor humid pulses towards the top of T6. In the Essaouira Basin, an analogous climatic evolution with (a) ephemeral braided river deposits followed by abandonment of the system shifting into a wet aeolian sandflat environment, is interpreted to highlight prevailing arid conditions with sub-ordinate humid pulses; (b) the onset of perennial fluvial deposits with extensive floodplain fines and subordinate cyclical aridity expressed by calcrete deposits, documents a pronounced shift related to increased discharge and prominent humidity, most likely in the catchment area; (c) playa environment with extensive, highly mottled mudstones and predominately arid overprint record a return to overall aridity.

An intriguing aspect comparing the Upper Triassic Essaouira with the Argana sequence is the observed well-defined shift from pronounced aridity, to fluvial sedimentation with a prominent perennial character, and a subsequent return to arid conditions. Moreover, the fluvial deposits exhibit sedimentary characteristics of long-lived, extensive perennial systems atypical for the overall arid Triassic period. In the Essaouira Basin, the perennial deposits are for example associated with organic-rich shales and coaly interbeds, while in Argana the perennial fluvial channels incise deeply into underlying strata, reflecting large river systems with exaggerate flow. This stands in direct contrast to the Triassic climate which is generally characterised by a “hot house” world with temperate warm conditions throughout (Kutzbach & Gallimore, 1989; Dubiel *et al.*, 1991; Robinson, 1973; Parrish, 1994). Moreover, various climate models have acknowledged the existence

of extreme climatic regimes, and a dominance of strong monsoonal circulation patterns has been recognised (e.g. Kutzbach & Gallimore, 1989; Crowley *et al.*, 1989; 1992; Dubiel *et al.*, 1991; Parrish, 1994, Parrish *et al.*, 1982). Analogous to the modern monsoons, the Pangaean monsoonal circulations are thought to have occurred due to the sensible heating of the vast Pangaean continent, the symmetry across the equator during the Triassic enhancing cross-equatorial thermal contrast and the existence of the warm Tethyan seaway in between maximising the pressure contrast (Young, 1987; Parrish, 1994; Kutzbach, 1994) (figure 18). Monsoonal circulation patterns can successfully be tested in the rock record (e.g. Nanson & Croke, 1992; Nanson *et al.*, 1998; Radis *et al.*, 2004; Sugai, 1993, Blum & Valastro, 1994; Holbrook, 2001, Gibling *et al.*, 2005), and the evidence of strong seasonality, paleosols (calcretes) or coaly layers observed in the studied sections, propose a direct relationship to Pangaean monsoonal patterns (see also Dubiel *et al.*, 1991; Parrish, 1994). Additionally, recent publications suggest particularly the Late Triassic to be interrupted by a series of regionally significant humid events (e.g. Preto *et al.*, 2010), the most prominent being the “Carnian pluvial event” (Simms & Ruffel, 1989, 90; Simms *et al.*, 1995; Parrish, 1994; Colombi & Parrish, 2008; Preto *et al.*, 2010; Arche & Lopez, 2014). The strong humid pulses preserved in the sedimentological record of Carnian to Norian strata in the Argana and Essaouira Basins could suggested that they also impacted these SW Moroccan basins. Arche & Lopez (2014) had already recognised the Argana Basin to fall within the regional trend of the Carnian pluvial event, and this study implies an even more regional extend to Essaouira. Similar to basins studied in Arche & Lopez’ (2014) regional approach, the Essaouira sequence exhibits a shift to arid condition sedimentation patterns, following the amplified humidity. In contrast to Arche & Lopes (2014) however, the Tethyan transgression is not assumed to have reached as far west as Essaouira during the Carnian inferred from the lack of clear marine indicators from this core study (figure 18). It further has to confirmed with more extensive biographic sampling, that the sub-surface data in Essaouira is indeed of Carnian age. Preto *et al.* (2010) summarised additional Upper Triassic humid events in the Norian, which have recently also been documented along the northernmost extends of the Pangaen equatorial regions in the Central North Sea Basin (e.g. McKie, 2014). With the sparse age dating from Eassouira it could be speculated, that there is a link to a later Norian humid event instead. Nevertheless, results from both Moroccan basins aid in redefining the paleogeographic and paleo-climatic setting along the south-eastern extends of the Tethyan margin (figure 18), placing them within the realm of mega-monsoonal oscillations during the Upper Triassic, and providing further evidence for the impact of regional pluvial events on the

stratigraphy of SW Morocco. In terms of the direct effect of monsoonal oscillation patterns, general circulation models (GCMs) for Pangaea during Upper Triassic times (Sellwood & Valdes, 2006) give rise to multiple assumption as to when strong rainfall could have affected the AOI (figures 18A,C): during the winter monsoon, as predicted for Arabia and NE Africa (figure 18B), or during the tropical summer monsoon (figure 18C). The predominantly E to W wind directions preserved in aeolianites of T6 in the Argana Valley led Mader & Redfern (2011) to believe that the climatic regime during the arid phases reflects the Pangaeian winter, while humid pulses are driven by the Pangaeian summer monsoon, however also point out the possible intra-mountainous position of the Argana Valley in a rainshadow position. Whether maximum rainfall affected the AOI during Pangaeian summers, or whether the Pangaeian winter monsoon modelled for Arabia and NE Africa reaches as far as SW Morocco remains speculative, however provides clear evidence for the eastern extends of monsoonal oscillations affecting the SW Moroccan basins in general.

9. CONCLUSIONS

9.1 Depositional model Essaouira Basin (Meskala Field)

Detailed core analysis from the main reservoir unit in the Meskala Field (central Essaouira Basin) has led to a defined suite of facies associations, which merit a new facies model for the Upper Triassic deposits. Four facies associations were defined and interpreted to represent a progressive evolution from ephemeral fluvial deposits characterised by episodic discharge in a shallow braided river system, evolving into a wet sandflat environment during a phase of pronounced aridity. Above a clear bounding surface a rapid change in depositional environment from sheet-like ephemeral flow to metre-scale fluvial sandstone packages with very fine-grained channel deposits and abundant, thick floodplain fines follows. Several fining-upward sequences were documented, and a fluvial perennial river setting is proposed. Subsequently, the fluvial deposits become increasingly overprinted by transgressive phases of an adjacent lacustrine or playa lake shifting shoreline. The environment of deposition changes into a delta setting with distributary mouthbars and abundant inter-pond areas on the delta-top. The sandstones change from clearly defined fluvial channel deposits to dominantly ripple cross-stratified sandstones with abundant organic-rich intercalations and paleosols preserved on the floodplain. Water availability and more perennial conditions were inferred. Towards the end of this transgressive phase, the overall input of sandstones decreases considerably and an

extensive playa environment returning to oxidising conditions in an arid climate is proposed.

9.2 Regional sediment distribution patterns SW Morocco

A comparison is made between the studied deposits in the Essaouira Basin and time-equivalent sections of Carnian to Norian age in the Argana Valley. A key question was, whether sediments exposed in the Argana Valley were directly linked to the Essaouira Basin, affording a direct sediment source for the prolific Upper Triassic reservoirs in the Meskala Field. Although a proximal to distal fluvial relationship between the Argana Valley and Essaouira Basin is apparent, SE to NW directed fluvial flow was only subordinately recognised in the Argana Valley questioning this direct source to sink relationship. Sedimentological data from the sandstone dominated facies indicate a dominantly W to SW to S paleoflow instead, which on a regional scale supports increased accommodation creation in the S/SW. The regional sedimentary evidence further suggests that sediment storage during the Carnian to Norian is likely to be in the southern Argana Valley or southern Souss Basins, gradually filled during the Upper Triassic, masking the previous inherent structural grain of the initial half-graben setting (Hofmann *et al.*, 2000; Mader & Redfern, 2011; Baudon *et al.*, 2012). No evidence for continuous, extensive NW flowing fluvial flow which would have been representative of a direct sediment fairway to the central Essaouira Basin was recognised, hence a more local provenance for the Meskala reservoirs is anticipated.

9.3 Paleo-climatic implications

Pronounced, possibly short but intensive wet seasons in an otherwise dominantly hot, arid climatic setting were documented in both studied localities. A link is made to regionally extensive Upper Triassic pluvial events such as the well document “Carnian pluvial event”. Results from this study aid in breaching the gap for paleogeographic and paleo-climatic reconstructions of the eastern Atlantic margin and the extends of Late Triassic Tethys transgressions.

ACKNOWLEDGMENTS

This work is part of a PhD thesis completed at the University of Manchester, UK. The North Africa Research Group and sponsoring companies provided the financial support necessary to carry out field work. ONHYM kindly provided access to the sub-surface data sets and allowed publication of the data. The reviewers S. Leleu (Université de Bordeaux Montaigne) and S. Jones (Durham University) provided valuable comments that improved the initial manuscript considerably.

REFERENCES

- Ait Chayeb, E. H., Youbi, N., El-Boukhari, A., Bouabdelli, M. & Amrhar, M. (1998). Permian–Mesozoic volcanism of the Argana Basin (western High Atlas, Morocco); intraplate magmatism associated with the opening of the Central Atlantic. *Journal of African Earth Sciences*, 26, 499–519.
- Alonso-Zarza, A.M. (2003). Palaeoenvironmental significance of palustrine carbonates and calcretes in the geological record. *Earth-Science Reviews*, 60, 261–298.
- Anderson, M.P., Aiken, J.S., Webb, E.K. & Mickelson, D.M. (1999). Sedimentology and hydrology of two braided stream systems. *Sedimentary Geology*, 129, 187–200.
- Arche, A. (2007). Some precisions on the use of the term playa in the geologic literature. *Journal of Iberian Geology*, 24, 5–10.
- Arche, A. & Lopez-Gomez, J. (2014). The Carnian Pluvial Event in Western Europe: New data from Iberia and correlation with the Western Neotethys and Eastern North America – NW Africa regions. *Earth-Science Reviews*, 128, 196–231.
- Audley-Charles, M. F. (1970). Triassic palaeogeography of the British Isles. *Quarterly Journal of the Geological Society London*, 126, 49–89.
- Baudon, C., Redfern, J. & Van Den Driessche (2012). Permo-Triassic structural evolution of the Argana Valley, impact of the Atlantic rifting in the High Atlas, Morocco. *Journal of African Earth Sciences*, 65, 91–104.
- Beauchamp, J. (1988). Triassic sedimentation and rifting in the High Atlas (Morocco). In: Manspeizer, W. (Ed), Triassic-Jurassic Rifting, *Developments in Geotectonics*, 22, 477–497.
- Blum, M. D. and Valastro, S. J. (1994). Late Quaternary sedimentation, lower Colorado River, Gulf Coastal Plain of Texas. *Geological Society of America Bulletin* 106(8), 1002–1016.
- Blum, M. D. and Törnqvist, T. E. (2000). Fluvial responses to climate and sea-level change: a review and look forward. *Sedimentology*, 47, 2–48.
- Bouatmani, R., Ahmamou, M., El Ouarghioui, A., Medina, F., El Mourabit, A. & Daoudi, L. (2004). Les environnements de dépôt des formations triasiques de la région de Meskala (bassin d’Essaouira, Maroc): apport de l’analyse des diagraphies et des carottes. *Bull. Inst. Sci., Rabat, section Sci. Terre*, 26, 49–67.
- Bridge, J.S. (1993). The interaction between channel geometry, water flow, sediment transport and

- deposition in braided rivers. In: J.L. Best, C.S. Bristow (Eds.), *Braided Rivers*, *Geological Society of London, Special Publications*, 75, 13–71
- Bridge, J.S. (2003).** Rivers and Floodplains. *Blackwell, Oxford* (2003).
- Bridge, J. S. (2006).** Fluvial facies models: recent developments. In: W. Posamentier, R.G. Walker (Eds), *Facies Models Revisited, SEPM Special Publication*, 84, 85–170.
- Bristow, C.S., Skelly, R.L. & Ethridge, F.G. (1999).** Crevasse splays from the rapidly aggrading, sandbed, braided Niobrara River, Nebraska: effect of base-level rise. *Sedimentology*, 46, 1029-1047.
- Broughton, P. & Trépanier, P. (1993).** Hydrocarbon generation in the Essaouira Basin of western Morocco. *AAPG Bulletin*, 77(6), 999-1015.
- Brown, R. H. (1980).** Triassic rocks of the Argana Valley, southern Morocco, and their regional structural implications. *AAPG Bulletin*, 64 (7), 988-1003.
- Cain, N.P. & Mountney, N.P. (2009).** Spatial and temporal evolution of a terminal fan system: the Permian Organ Rock Formation, SE Utah, USA. *Sedimentology*, 56, 1774-1800.
- Colombera, L., Mountney, N.P., & McCaffrey, W.D. (2015).** A meta-study of relationships between fluvial channel-body stacking pattern and aggradation rate: implications for sequence stratigraphy. *Geology*, 43, 283–286.
- Colombi, C.E. & Parrish, J.T. (2008).** Late Triassic environmental evolution in Southwestern Pangaea: plant taphonomy of the Ischigualasto Formation. *Palaios* 23, 778-795.
- Courel L., Ait Salem H., Benaouiss N.; Et-Touhami M., Fekirine B., Oujidi M.; Soussi M. & Tourani A. (2003).** Mid-Triassic to Early Jurassic clastic/evaporitic deposits over the Maghreb Platform. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 196/1, 157-176.
- Crowley, T. J., Hyde, W. T. and Short, D. A. (1989).** Seasonal cycle variations on the supercontinent of Pangaea: Implications for Early Permian vertebrate extinctions. *Geology*, 17, 457-460.
- Crowley, T. J., Kim, K. Y., Mengel, J. G. and Short, D. A. (1992).** Modelling 100,000-year climate fluctuations in pre-Pleistocene time series. *Science* 255(5045), 705-707.
- Deenen, M., Langreis, C; Krijgsman, W. ElHachimi, W. & Chellai, E.H. (2010).** Palaeomagnetic results from Upper Triassic red-beds and CAMP lavas of the Argana Basin, Morocco. In: Van Hinsbergen, D. J. J.,

- 661 Buiter, S. J. H., Torsvik, T. H., Gaina, C. & Webb, S. J. (Eds) The Formation and Evolution of Africa: A
662 Synopsis of 3.8 Ga of Earth History, *Geological Society, London, Special Publications*, 357, 195–209.
- 663 **Dubiel, R. F., Parrish, J. T., Parrish, M. and Good, S.C. (1991).** The Pangaeon Megamonsoon - Evidence
664 from the Upper Triassic Chinle Formation, Colorado Plateau. *Palaio*, 6, 347-370.
- 665 **Duffaut, M.F., Brun, L. and Plauchut, B. (1966).** *Notes mém. Serv. Géol. Maroc*, 85, 129-137.
- 666 **Dutuit, J.M. (1976).** Introduction à l'étude paléontologique du Trias continental marocain. Description des
667 premiers Stégocéphales recueillis dans le couloir d'Argana. *Mém. Mus. Natn. Hist. nat., Paris*, C 36, 253.
- 668 **Dutuit, J.M. (1977).** *Palaeorhinus magnoculus*, phytosaure du Trias supérieur de l'Atlas Marocain. *Géologie*
669 *Méditerranéenne*, 4, 225-238.
- 670 **Dutuit, J.M. (1988).** *Diplocaulus minimus* n. ps. (Amphibia; Nectridea), Lepospondyle de la formation
671 d'Argana, dans l'Atlas occidental marocain. *C.R. Acad. Sci. Paris*, II 307, 851-854.
- 672 **El Hachimi, H., Youbi, N., Madeira, J., Bensalah, M.K., Martins, L., Mata, J., Medina, F., Bertrand,**
673 **H., Marzoli, A., Munhá, J., Bellieni, J., Mahmoudi, A., Ben Abbou, M. & Assafar, H. (2011).**
674 Morphology, internal architecture, and emplacement mechanisms of lava flows from the Central Atlantic
675 Magmatic Province (CAMP) of Argana basin (Morocco). In: Van Hinsbergen, D.J.J. ; Buiter, S. ; Torsvik,
676 T.H. ; Gaina, C. & Webb, S. (Eds): The formation and evolution of Africa : A Synopsis of 3.8 Ga of Earth
677 History, *The Geological Society of London, Special Publication*, 357, 167-193.
- 678 **Fabuel-Pérez, I., Hodgetts, D., and Redfern, J. (2009a).** A new approach for outcrop characterization and
679 geostatistical analysis of a low-sinuosity fluvial-dominated succession using digital outcrop models: Upper
680 Triassic Oukaimeden Sandstone Formation, High Atlas, Morocco. *AAPG Bull.* 93, 795–827.
- 681 **Fabuel-Pérez, I., Redfern, J., and Hodgetts, D. (2009b).** Sedimentology of an intra-montane rift-controlled
682 fluvial-dominated succession: the Upper Triassic Oukaimeden Sandstone Formation, Central High Atlas,
683 Morocco. *Sediment. Geol.* 218, 103–140.
- 684 **Fiechtner, L., Friedrichsen, H. and Hammerschmidt, K. (1992).** Geochemistry and geochronology of
685 Early Mesozoic tholeiites from Central Morocco. *Int. Journal of Earth Sciences*, 81 (1), 45-62.
- 686 **Fisher, J.A., Nichols, G.J. & Waltham, D.A. (2008).** Unconfined flow deposits in distal sectors of fluvial
687 distributary systems: examples from the Miocene Luna and Huesca Systems, northern Spain. *Sedimentary*
688 *Geology*, 195 (2008), 55–77.

- 689 **Gibling, M. R., Tandon, S. K., Sinha, R. and Jain. M. (2005).** Discontinuity-bounded alluvial sequences
 690 of the southern Gangetic plains, India: aggradation and degradation in response to monsoonal strength.
 691 *Journal of Sedimentary Research*, 75, 369-385.
- 692 **Glennie, K. W. (1970).** Desert Sedimentary Environments. *Amsterdam, Elsevier.*
- 693 **Gradstein, F.; Ogg, J.; Schmitz, M. & Ogg, G. (2012).** The Geologic timescale. *2 Volumes, Elsevier.*
- 694 **Hafid, M. (2000).** Triassic - early Jurassic extensional systems and their Tertiary inversion, Essaouira Basin
 695 (Morocco). *Marine and Petroleum Geology*, 17, 409-429.
- 696 **Hafid, M., Ait Salem, A., and Bally, A.W. (2000).** The western termination of the Jebilet-High Atlas
 697 system (Offshore Essaouira Basin, Morocco). *Marine and Petroleum Geology*, 17, 431-443.
- 698 **Hinz, K., Dostmann, H. and Fritsch, J. (1981).** The continental margin of Morocco: seismic sequences,
 699 structural elements and geological development. *In: Von Rad, U., Hinz, K., Sarnthein, M. and Seibold, E.*
 700 (Eds), *Geology of the Northwest African Margin.*
- 701 **Hofmann, A., Tourani, A. and Gaupp, R. (2000).** Cyclicity of Triassic to Lower Jurassic continental red
 702 beds of the Argana Valley, Morocco: implications for palaeoclimate and basin evolution. *Palaeogeography,*
 703 *Palaeoclimatology, Palaeoecology*, 161, 226-266.
- 704 **Holbrook, J. (2001).** Origin, genetic interrelationships, and stratigraphy over the continuum of fluvial
 705 channel-form bounding surfaces: an illustration from middle Cretaceous strata, southeastern Colorado.
 706 *Sedimentary Geology*, 144, 179-222.
- 707 **Howell, J. and Mountney, N. (1997).** Climatic cyclicity and accommodation space in arid to semi-arid
 708 depositional systems: an example from the Rotliegend Group of the UK southern North Sea. *In: Ziegler, K.*
 709 *Turner, P. and Daines, S. R. (Eds), Petroleum Geology of the Southern North Sea: Future Potential,*
 710 *Geological Society London Special Publications*, 123, 63-86.
- 711 **Hminna, A., Voigt, S., Saber, H., Schneider, J.W. & Hmich, D. (2012).** On a moderately diverse
 712 continental ichnofauna from the Permian Ikakern Formation (Argana Basin, Western High Atlas, Morocco).
 713 *Journal of African Earth Sciences*, 68, 15–23.
- 714 **Hubert, J. F. and Forlenza, M.F. (1988).** Sedimentology of braided-river deposits in Upper Triassic
 715 Wolfville redbeds, southern shore of Cobequid Bay (sic), Nova Scotia, Canada. *In: Manspreizer, W. (Ed),*
 716 *Triassic-Jurassic Rifting: continental break-up and the origin of Atlantic ocean and passive margins.*

- Elsevier, Amsterdam, Netherlands, 231-247.
- Hunt, A.P. and Lucas, S.G. (1991).** The Palaeorhinus Biochron and the correlation of the non-marine Upper Triassic of Pangaea. *Palaeontology*, 34, 487-501.
- Jalil, N.-E. (1999).** Continental Permian and Triassic vertebrate localities from Algeria and Morocco and their stratigraphical correlations. *Journal of African Earth Sciences*, 29(1), 219-226.
- Jalil, N.-E. and Dutuit, J.-M. (1996).** Permian captorhinid reptiles from the Argana Formation, Morocco. *Palaeontology*, 39(4), 907-918.
- Jalil, N.E. and Janvier, P. (2005).** Les pareiasaures (Amniota, Parareptilia) du Permien supérieur du Bassin d'Argana, Maroc. *Geodiversitas*, 27, 35–132.
- Jalil, N.E. and Peyer, K. (2007).** A new rauisuchian (Archosauria, Suchia) from the Upper Triassic of the Argana Basin, Morocco. *Palaeontology*, 50 (2), 417–430.
- Jalil, N.-E, Janvier, P. and Steyer, J.-S. (2009).** A new cyclotosaurid (Amphibia, Temnospondyli) from the Triassic of Argana Basin (High Atlas Mountains, Morocco), biostratigraphic implications. *1st Int. Congr. N. African Vertebr. Palaeontol., Abstracts (2009)*, 36–37.
- Jansa, L. F. and Wiedmann, J. (1980).** Mesozoic-Cenozoic development of the E North American and Northwest African continental margins: a comparison. In: Von Rad, U., Hinz, K., Sarnthein, M. and Seibold, E. (Eds), *Geology of the Northwest African Margin*.
- Jones, D. F. (1975)** Stratigraphy, environments of deposition, petrology, age, and provenance of the basal red beds of the Argana Valley, Western High Atlas Mountains, Morocco. Socorro, New Mexico, *New Mexico Institute of Mining and Technology*, 144.
- Jopling, A.V. and Walker, R. (1968).** Morphology and origin of ripple-drift cross-lamination, with examples from the Pleistocene of Massachusetts. *Journal of Sedimentary Petrology*, 38, 971–98.
- Khalaf, I. F. (2007).** Occurrence and genesis of calcrete and dolocrete in the Mio-Pleistocene fluvial sequence in Kuwait, northeast Arabian Peninsula. *Sedimentary Geology*, 199, 129-139.
- Klein, H., Voigt, S., Hminna, A., Saber, H., Schneider, J. and D. Hmich (2010).** Early Triassic Archosaur-dominated footprint assemblage from the Argana Basin (Western High Atlas, Morocco). *Ichnos*, 17, 215–227.

- 744 Klein, H., Voigt, S., Hminna, A., Saber, H., Schneider, J., Fischer J., and Brosig, A. (2011). First
745 occurrence of a Middle Triassic tetrapod ichnofauna from the Argana Basin (Western High Atlas, Morocco).
746 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 307, 218–231.
- 747 Knight, K.B., Nomade, S., Renne, P.R., Marzoli, A., Bertrand, H. & Youbi, N. (2004). The Central
748 Atlantic magmatic province at the Triassic–Jurassic boundary: paleomagnetic and $^{40}\text{Ar}/^{39}\text{Ar}$ evidence from
749 Morocco for brief, episodic volcanism. *Earth and Planetary Science Letters*, 22, 143–160.
- 750 Kocurek, G. and Nielson, J. (1986). Conditions favourable for the formation of warm-climate aeolian sand
751 sheets. *Sedimentology*, 33, 795–816.
- 752 Kocurek, G. (1988). Late Palaeozoic and Mesozoic eolian deposits of the western interior of the United
753 States. *Sedimentary Geology-Special Issue*.
- 754 Kumar, R., Ghosh, S.K., Satish, J.S. (1999). Evolution of a Neogene fluvial system in a Himalayan
755 foreland basin, India. In: A. Macfarlane, R.B. Sorkhabi, J. Quade (Eds.), Himalaya and Tibet Mountain
756 Roots to Mountain Tops, *Geological Society of America, Special Papers*, 328, 239–256.
- 757 Kutzbach, J. E. (1994). Idealized Pangaeian climates: Sensitivity to orbital change. In: Klein, G. D.,
758 Pangaea: Palaeoclimate, tectonics, and sedimentation during accretion, zenith, and break-up of a super-
759 continent, Boulder, CO, *Geological Society of America*, 288, 41–55.
- 760 Kutzbach, J. E. and Gallimore, R. G. (1989). Pangaeian climates: Megamonsoons of the Megacontinent.
761 *Journal of Geophysical Research* 94(D3), 3341–3357.
- 762 Laville, E. and Petit, J.-P. (1984). Role of syn-sedimentary strike-slip faults in the formation of Moroccan
763 Triassic basin. *Geology*, 12, 424–427.
- 764 Leleu, S. & Hartley, A. J. (2010). Controls on the stratigraphic development of the Triassic Fundy basin,
765 Nova Scotia: Implications for the tectonostratigraphic evolution of Triassic Atlantic rift basins. *Journal of the*
766 *Geol. Soc. London*, 167, 437–454.
- 767 Leleu, S., Hartley, A. J., Van Oosterhout, C., Kennan, L., Ruckwied, K. and Gerdes, K. (2016).
768 Structural, stratigraphic and sedimentological characterisation of a wide rift system: The Triassic rift system
769 of the Central Atlantic Domain. *Earth-Science Reviews*, 158, 89–124.
- 770 LeRoy, P., LeGall, B., Ait Rahim, L., Morabet, A.M. and Demnati, A. (1997). Les bassins côtiers triasico-
771 liasiques du Maroc occidental et la diachronie du rifting intracontinental de l'Atlantique central. *Bulletin de*

- 772 *la Société géologique de France*, 168, 637-648.
- 773 **LeRoy, P. and Pique, A. (2001).** Triassic-Jurassic Western Moroccan synrift basins in relation to the
774 Central Atlantic opening. *Marine Geology*, 172, 359-381.
- 775 **Lucas, S. G. (1998).** Global Triassic tetrapod biostratigraphy and biochronology. *Palaeogeography*,
776 *Palaeoclimatology, Palaeoecology*, 143, 347-384.
- 777 **Mader, N.K. (2005).** Sedimentology and sediment distribution of Upper Triassic fluvio-aeolian reservoirs
778 on a regional scale (Central Algeria, SW Morocco, NE Canada): an integrated approach unravelling the
779 influence of climate versus tectonics on reservoir architecture. *Unpublished PhD thesis, Manchester*
780 *University, UK.*
- 781 **Mader, N. K. & Redfern, J. (2011).** A sedimentological model for the continental Upper Triassic Tadrart
782 Ouadou Sandstone Member : recording an interplay of climate and tectonics (Argana Valley; SW Morocco).
783 *Sedimentology*, 58(5), 1247-1282.
- 784 **Manspeizer, W. (1982).** Triassic-Jurassic Basins and Climate of the Atlantic Passive margins. *Geologische*
785 *Rundschau*, 71 (3), 895-917.
- 786 **Manspeizer, W. (1988).** A stratigraphic record from Morocco and North America of rifting, drifting and
787 Tethyan transgression of the Central proto-Atlantic. *Journal of African Earth Sciences*, 7(2), 369-373.
- 788 **Manspeizer, W. (1994).** The breakup of Pangaea and its impact on climate: consequences of Variscan-
789 Alleghanianide orogenic collapse. In: Klein (Ed). Pangaea: Palaeoclimate, Tectonics and Sedimentation
790 during Accretion, Zenith and Breakup of a Supercontinent. *Geol. Soc. Am. Spec. Papers*, 288, 169-185.
- 791 **Manspeizer, W., Puffer, J. H. and Cousminer, H. L. (1978).** Separation of Morocco and E North America:
792 a Triassic-Jurassic stratigraphic record. *Geol. Soc. Am. Bulletin*, 89, 901-920.
- 793 **Mariott, S.B. & Alexander, J (eds) (1999).** Floodplains: Interdisciplinary Approaches. *Geol. Soc. London*,
794 *Spec. Pub.*, 28, 271-281.
- 795 **Marzoli, A., Renne, P.R., Piccirillo, E.M., Ernesto, M., Belliene, G. & De Min, A. (1999).** Extensive 200-
796 Million-Year-Old Continental Flood Basalts of the Central Atlantic Magmatic Province. *Science* 23 April
797 1999, 284(5414), 616-618.

- 798 **Marzoli, A.; Bertrand, H.; Knight, K.B.; Cirilli, S.; N. Buratti, C. Vrati, S. Nomade,, P.R. Renne, N.**
 799 **Youbi, R., Martini, K. Allenbach, R. Neuwerth, C. Rapaille, L.Zaninetti, G. Bellieni (2004).** Synchrony
 800 of the Central Atlantic magmatic province and the Triassic-Jurassic boundary climatic and biotic crisis.
 801 *Geology*, 32, 973–976.
- 802 **Matthews, M.D. and Perlmutter, M.A. (1994).** Global cyclostratigraphy: an application to the Eocene
 803 Green River Basin. In: P.L. De Boer and D.G. Smith (Eds): *Orbital forcing and cyclical sequences.*,
 804 *Blackwell Scientific Publications, Oxford.*
- 805 **Mattis, A. F. (1977).** Nonmarine Triassic sedimentation, central High Atlas Mountains, Morocco. *Journal*
 806 *of Sedimentary Petrology*, 47(1), 107-119.
- 807 **McHone, G. (2000).** Non-plume magmatism and rifting during the opening of the central Atlantic Ocean.
 808 *Tectonophysics* 316, 287-296.
- 809 **McKie (2014).** Climatic and tectonic controls on Triassic dryland terminal fluvial system architecture,
 810 central North Sea. *Int. Assoc. Sedimentol. Spec. Publ.*, 46, 19-58.
- 811 **Medici, G., Boulesteix, Mountney, N.P., West, L.J. & Odling (2015).** Paleoenvironment of braided fluvial
 812 systems in different tectonic realms of the Triassic Sheerwood Sandstone Group, UK.. *Sedimentary Geology*,
 813 329, 188-210.
- 814 **Medina, F., (1988).** Tilted-blocks pattern, paleostress orientation and amount of extension, related to triassic
 815 early rifting of the central Atlantic in the Amzri area (Argana basin, Morocco). *Tectonophysics*, 148 (3–4),
 816 229–233.
- 817 **Medina, F. (1991).** Superimposed extensional tectonics in the Argana Triassic formations (Morocco),
 818 related to the early rifting of the central Atlantic. *Geological Magazine*, 128(5), 525-536.
- 819 **Medina, F. (1995).** Syn- and postrift evolution of the El Jadida-Agadir basin (Morocco): constraints for the
 820 rifting models of the central Atlantic. *Canadian Journal of Earth Sciences*, 32, 1273-1291.
- 821 **Miall, A. D. (1977).** A review of the braided-river depositional model environment. *Earth-Science Reviews*,
 822 13, 1–62.
- 823 **Miall, A. D. (1996).** The Geology of Fluvial Deposits. Sedimentary facies, Basin Analysis and Petroleum
 824 Geology. Heidelberg, Springer-Verlag.

- 825 **Miall, A.D. (2006).** The Geology of Fluvial Deposits. Sedimentary Facies, Basin Analysis and Petroleum
826 Geology. *Springer, Berlin (2006).*
- 827 **Morabet, A. M., Bouchta, R. and Jabour, H. (1998).** An overview of the petroleum systems of Morocco.
828 In: MacGregor, D. S., Moody, R. T. J. and Clark-Lowes, D. D., Petroleum Geology of North Africa,
829 *Geological Society London Special Publication, 132, 283-296.*
- 830 **Nanson, G.C., Rust, B.R. & Taylor, G. (1986).** Coexistent mud braids and anastomosing channels in arid
831 zone river: Cooper Creek, Central Australia. *Geology, 14, 175-178.*
- 832 **Nanson, G.C. & Croke, J.C. (1992).** A genetic classification of floodplains. *Geomorphology 4, 459-486.*
- 833 **Nanson, G.C., Callen, R.A. and Price, D.M. (1998).** Hydroclimatic interpretation of Quaternary shorelines
834 on South Australian playas. *Palaeogeography, Palaeoclimatology, Palaeoecology, 144, 281-305.*
- 835 **Newell, A.J., Tverdokhlebov, V.P. & Benton, M.J. (1999).** Interplay of tectonics and climate on a
836 transverse fluvial system, Upper Permian, Southern Uralian Foreland Basin, Russia. *Sedimentary Geology,*
837 *127, 11-29.*
- 838 **Nomade, S; Knight, K.B., Beutel, E. & Renne, P.R. (2007).** Chronology of the Central Atlantic Magmatic
839 Province: Implications for the Central Atlantic rifting processes and the Triassic-Jurassic biotic crisis.
840 *Palaeogeography, Palaeoclimatology, Palaeoecology, 244, 326-344.*
- 841 **Olsen, P.E., Kent, D.V., Fowell, S.J., Schlische, R.W., Withjack, M.O. & LeTourneau, P.M. (2000).**
842 Implications of a comparison of the stratigraphy and depositional environments of the Argana (Morocco) and
843 Fundy (Nova Scotia, Canada) Permian-Jurassic basins. In: Oujidi, M., Et-Touhami, M. (Eds): Le Permien et
844 le Trias du Maroc: Acte de la Premiere Reunion du Groupe Marocain du Permien et du Trias, *Oujda, 165-*
845 *183.*
- 846 **ONAREP (Office National de Recherches et d'Explorations Pétrolières, Morocco) (1995).** Onshore
847 Essaouira Basin: Exploration prospects Morocco. *ONAREP report (1995), 1-24.*
- 848 **Oujidi, M., Courel, L., Benaouiss, N, El Mostaine, M, El Touhami, M., Ouarhache, D., Sabaoui, A. &**
849 **Tourani, A.-I. (2000).** Triassic series of Morocco: stratigraphy, paleogeography and structuring of the
850 southwestern Peri-Tethyan Platform. An overview. In: Crasquin-Soleau, S. & Barrier, E. (Eds), Peri-Tethys
851 Memoir 5: new data on Peri-Tethyan sedimentary basins, *Mem. Mus. Natn. Hist. nat., 182: 23-38. Paris.*
- 852 **Owens, P.M., Walling, D.E. & Leeks, G.J.L (1999).** Deposition and storage of fine-grained sediment

- 853 within the main channel system of the River Tweed, Scotland. *Earth Surface Processes and Landforms*, 24,
854 1061–1076.
- 855 **Parrish, J. T. (1994).** Climate of the supercontinent Pangaea. *The Journal of Geology*, 101, 215-233.
- 856 **Parrish, J. T. and Curtis, R. L. (1982).** Atmospheric circulation, upwelling, and organic-rich rocks in the
857 Mesozoic and Cenozoic eras. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 40, 31-66.
- 858 **Piqué, A. and Laville, E. (1996).** The central Atlantic rifting: reactivation of Palaeozoic structures? *Journal*
859 *of Geodynamics*, 21, 235-255.
- 860 **Piqué, A., Le Roy, P. & Amrhar, M. (1998).** Transtensive synsedimentary tectonics associated with ocean
861 opening: the Essaouira-Agadir segment of the Moroccan Atlantic margin. *Journal of the Geological Society*,
862 *London*, 155, 913-928.
- 863 **Platt, N.H. & Keller, B. (1992).** Distal alluvial deposits in a foreland basin setting—the Lower Freshwater
864 (Miocene), Switzerland: sedimentology, architecture and palaeosols. *Sedimentology*, 39 (1992), 545–565. ’
- 865 **Preto, N, Kustatscher, E. & Wignall, P.B. (2010).** Triassic climates – state of the art and perspectives.
866 *Paleogeography, Paleoclimatology, Palaeoecology*, 290 , 1-10.
- 867 **Radis, D., Preusser, F., Matter, A. and Mange, M. (2004).** Eustatic and climatic controls on the
868 development of the Wahiba Sand Sea, Sultanate Oman. *Sedimentology*, 51, 1359-1385.
- 869 **Reeves Jr., C.C. (1976).** Caliche origin, classification, morphology and uses. *Estacado Books, Lubbock*.
870 *Schmid, S., Worden, R.H., Fisher, Q.J., 2006.*
- 871 **Retallack, G. J. (1988).** Field recognition of palaeosols. *Geological Society of America Special Paper*, 216,
872 1-20.
- 873 **Robinson, P.L. (1973).** Paleoclimatology and continental drift. In: *Tarling, D.H., Runcorn, S.K. (eds)*,
874 *Implications of continental drift to the Earth Sciences, Academic Press, London*, 449-476.
- 875 **Sahabi, M., Aslanian, D. and Olivet, J.L. (2004).** Un nouveau point de départ pour l’histoire de l’Atlantique
876 central. *C.R. Geoscience*, 336, 1041-1052.
- 877 **Schumm, S. A. (1968b).** Speculations concerning palaeohydrologic controls of terrestrial sedimentation.
878 *Geol. Soc. Am. Bull.*, 79, 1573-1588.
- 879 **Schumm, S. A. (1993).** River response to base-level changes: implications for sequence stratigraphy.

- Journal of Geology, 101, 279-294.
- Scotese, C.R. (2000).** Paleomap project. *Http://www.scotese.com*.
- Sellwood, B.C. and Valdes, P.J. (2006).** Mesozoic climates: General circulation models and the rock record. *Sedimentary Geology*, 190, 269-287.
- Shanley, K. W. & McCabe, P. J. (1994).** Perspectives on the sequence stratigraphy of continental strata. *The American Association of Petroleum Geologists Bulletin*, 78, 544-568.
- Silmane, A. & ElMostaine, M. (1997).** Observations biostratigraphique au niveau des formations rouge de la séquence synrift dans le bassins de Doukkla et Essaouira. In: 1^{ère} Réunion du Groupe marocain du Permien et du Trias, Oujda, Maroc. *Faculté des Science, Université Mohammed I, Oujda*: 54.
- Simms, M.J. & Ruffell, A.H. (1989).** Synchronicity of climatic change and extinctions in the Late Triassic. *Geology*, 17, 265-268.
- Simms, M.J. & Ruffell, A.H. (1990).** Climatic and biotic change in the late Triassic. *Journal of the Geological Society of London*, 147, 321-327.
- Simms, M.J., Ruffell, A.H. & Johnson, L.A. (1995).** Biotic and climatic changes in the Carnian (Triassic) of Europe and adjacent areas. In: Fraser, N.C., Sues, H.-D. (Eds) : In the shadow of the Dinosaurs: Early Mesozoic Tetrapods. *Cambridge University Press*, 352-365.
- Smoot, J.P. (1991).** Sedimentary facies and depositional environments of early Mesozoic Newark Supergroup basins, eastern North America. *Paleogeography, Paleoclimatology, Palaeoecology*, 84, 369-423.
- Societe Cherifienne des Petroles (1966).** Le bassin du sud-ouest marocain. In : *Sedimentary basins of the African coast* (Ed. D. Reyre). *Assoc. Geol. Soc. Africains, Paris*, 5-12.
- Stets, J. and Wurster, P. (1981).** Atlas and Atlantic-structural relations. In: *Geology of the Northwest African Margin*. Von Rad, U. Hinz, K., Sarnthein, M., and Seibold, E. (Eds), 68-84.
- Sugai, T. (1993).** River terrace development by concurrent fluvial processes and climatic changes. *Geomorphology*, 6, 243-252.
- Tixeront, M. (1973).** Lithostratigraphie et minéralisations cuprifères et uranifères stratiformes syngénétiques et familiaires des formations détritiques permotriassiques du couloir d'Argana, Haut Atlas occidental (Maroc). *Notes du service géologique du Maroc*, 33 (249), 147-177.

- 907 **Tourani, A., Lund, J. J., Benaouiss, N. and Gaupp, R. (1999).** Stratigraphy of Triassic syn-rift deposition
908 in Western Morocco. *Zentralblatt für Geologie und Paläontologie, Teil 1, Heft, 9-10, 1193-1215.*
- 909 **Tourani, A., Benaouiss, N., Gand, G., Bourquin, S., Jalil, N-E., Broutin, J., Battail, B., Germain, D.,**
910 **Khaldoune, F., Sebban, S., Steyer, J-S., and Vacant, R. (2010).** Evidence of an Early Triassic age
911 (Olenekian) in Argana Basin (High Atlas, Morocco) based on new chirotherioid traces. *C.R.Palevol (2010),*
912 *1-8.*
- 913 **Turnbridge, I.P. (1981).** Sandy high-energy flood sedimentation – some criteria for recognition, with an
914 example from the Devonian of SW England. *Sedimentary Geology, 28, 79-96.*
- 915 **Turner, P., Pilling, D., Walker, D., Exton, J., Binnie, J. and Sabaou, N. (2001).** Sequence stratigraphy
916 and sedimentology of the late Triassic TAGI (Blocks 401/402, Berkine Basin, Algeria). *Marine and*
917 *Petroleum Geology, 18, 959-981.*
- 918 **Turner, P. and Sherif, H. (2007).** A giant Late Triassic-Early Jurassic evaporitic basin on the Saharan
919 platform, North Africa. In: Schreiber, B.C., Lugli, S. & Babel, M. (Eds) : Evaporites through space and time.
920 *Geological Society, London, Special Publications, 285, 87-105.*
- 921 **Van Houten, F. B. (1977).** Triassic-Jurassic deposits of Morocco and E North America: comparison. *AAPG*
922 *Bulletin, 61(1), 79-99.*
- 923 **Veevers, J.J. (1994).** Pangaea: Evolution of the super-continent and its consequences for earth's
924 palaeoclimate and sedimentary environments. In: G.D. Klein (Ed): Pangaea: Palaeoclimate, Tectonics and
925 Sedimentation during Zenith and break-up of a super-continent, *Geological Society of America Spec. Pub.,*
926 *288.*
- 927 **Verati, C., Rapaille, C., Féraud G., Marzoli, A., Bertrand, H. & Youbi, N. (2007).** $^{40}\text{Ar}/^{39}\text{Ar}$ ages and
928 duration of the Central Atlantic Magmatic Province volcanism in Morocco and Portugal and its relation to
929 the Triassic–Jurassic boundary. *Palaeogeography, Palaeoclimatology, Palaeoecology, 244 (1–4), 308–325.*
- 930 **Voigt, S., Hminna, A., Saber, H., Schneider, J.W. and Klein, H. (2010).** Tetrapod footprints from the
931 uppermost level of the Permian Ikakern Formation (Argana Basin, Western High Atlas, Morocco). *Journal*
932 *of African Earth Sciences, 57, 470–478.*
- 933 **Wang, P.X. (2009).** Global monsoon in a geological perspective. *Chinese Science Bulletin, 54, 1113-1136.*

- 934 **Whiteside, J. H., Olsen, P. E., Eglinton, T., Brookfield, M. E. & Sambrotto, R. N. (2010).** Compound-
935 specific carbon isotopes from Earth's largest flood basalt eruptions directly linked to the end-Triassic mass
936 extinction. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 6721–6725
- 937 **Wills, L. J. (1970).** The Triassic successions in the Central Midlands in its regional setting. *Quarterly*
938 *Journal of the Geological Society London*, 126, 225-283.
- 939 **Wright, V.P. (1992).** Paleopedology: stratigraphic relationships and empirical models. *In*: Martini, I.P.,
940 Chesworth, W. (Eds), *Weathering, Soils and Paleosols. Elsevier, Amsterdam*, 475–499.
- 941 **Wright, V.P. and Mariott, S.B. (1993).** The sequence stratigraphy of fluvial depositional systems: the role
942 of floodplain sediment storage. *Sedimentary Geology*, 86, 203-210.
- 943 **Wright, V.P. and Marriot, S.B. (1996).** A quantitative approach to soil occurrence in alluvial deposits and
944 its application to Old Red Sandstone of Britain. *Quarterly Journal of the Geological Society of London*, 153,
945 907–913.
- 946 **Youbi, N., Martins, L.T., Munha, J. M., Ibouh, H., Madeira, J., Ait Chayeb & El Boukhari, A. (2003).**
947 The Late Triassic-Early Jurassic Volcanism of Morocco and Portugal in the Framework of the Central
948 Atlantic Magmatic Province: an Overview. *In*: Hames, W.E., McHone, J.G., Ruppel, C. (Eds): *The Central*
949 *Atlantic Province: Insights from fragments of Pangea, AGU, Geophys. Mon*, 136, 179- 207.
- 950 **Young, J. A. (1987).** Physics of monsoons: the current view. *In*: Fein, J. S. a. S. (Ed), *Monsoons. New York,*
951 *Wiley*, 211-243.

952

SEDIMENTOLOGY OF THE ESSAOUIRA BASIN (MESKALA FIELD) IN CONTEXT OF REGIONAL SEDIMENT DISTRIBUTION PATTERNS DURING UPPER TRIASSIC PLUVIAL EVENTS

*Nadine K. Mader^{1,2}, Jonathan Redfern¹, Majid El Ouataoui³

¹North Africa Research Group, School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Williamson Building, Oxford Road, Manchester, M13 9 PL, United Kingdom.

²current Address: Maersk Oil Norge AS, Moseidsletta 122, 4068 Stavanger, Norway (madernadine@hotmail.com).

³ONHYM, 5 Avenue Moulay Hassan, B.P.99, 10050, Rabat, Morocco.

* CORRESPONDING AUTHOR

Keywords: Upper Triassic; Morocco; depositional reservoir model; regional sediment distribution; Triassic climate

FIGURES

Figure 1 1A (inset) Location map of Morocco displaying the main structural domains with the WSW-ENE trending High Atlas, the SW-NE trending Middle Atlas, the Rif domain in the North, and the Moroccan and Oran Mesetas located in central Western and Eastern Morocco. The subsurface study was carried out in the central Essaouira Basin (1) on core of wells from the Meskala Field, while the outcrop study was undertaken in the Argana Valley (2), exposing extensive Permo-Triassic strata along the southern extends of the High Atlas mountain chain.

1B The map displays the structural framework of Mesozoic (Triassic to Early Jurassic) basins in SW Morocco (modified after LeRoy & Piqué, 2000; with additions for the Argana Valley modified after Brown, 1980). Triassic depo-centres either known from outcrop or the sub-surface are marked in orange, and key wells are marked in red (e.g. well MKL101bis). In the literature several of the SW-NE to WSW-ENE trending sub-basins and half-grabens displayed are summarised as the Agadir-Essaouira Basin (dark red dotted line). For the Argana Valley paleocurrent data for Carnian to Norian deposits (compilation from Brown, 1980; Mader, 2005; Mader & Redfern, 2011) are superimposed on the structural map of SW Morocco: black arrows are supported by field data, whereas the grey arrows mark anticipated transport routes towards the S, SW and W. Within the Argana Valley exposures, field data suggests main transport direction towards the S, locally to the W (see figure 12 for more detail). Only minor fluvial deposits show clear SE to NW flow directions, which could indicate a direct sediment transport link to the Meskala field.

Figure 2 The map in **figure 2A** shows the surface locations of key wells in the greater onshore Essaouira Basin as well as the approximate location and distance to the outcrop exposures in the Argana Valley to the SE (orange). The Meskala Field lies in the central part of the explored Essaouira Basin, and the red lines mark key cross-sections that are displayed and discussed in figures 2B, 2C and 4. **Figure 2B** The W-E running cross-section (A-A') exhibits the subsurface distribution of Triassic to Early Jurassic major depositional sequences. An overall broad rift-basin morphology with a west-ward shift of main depositional units is apparent, and Carnian to Norian sandstones (sT2) extend across the prominent Meskala-Zelten horst (see also figure 2C) (simplified cross-section is based on seismic interpretations by Piqué et al. (1998) and LeRoy & Piqué (2001)). **Figure 2C** The close-up integrated well and seismic correlation (B-B') across the Meskala Field into the Jeer Field to the E shows clear thinning of Upper Triassic reservoir units across the horst towards the W (modified after Broughton & Trepanier, 1993). Along the top, the Upper Triassic is bound by salt deposits and an extrusive basalts (*basalt superieur*) capping the reservoir units sub-regionally. The basalt is subsequently overlain by Early Jurassic deposits and younger sedimentary units (see also **2B**; sT3). In the E part towards the Jeer Field, sills can be intrusive into the Upper Triassic deposits (*basalt inferieur*). The key well discussed in this study Mkl101bis is located in a central to northern position of the Meskala Field (**2A**). Offset wells MKL1 and MKL4 (**2A,C**) contain Ostracodes (*Darwinula*, *Lymnocythere*) (Silmane & ElMostaine, 1997), suggesting a Carnian to Norian age for the investigated Triassic deposits. Absolute ages for intrusive dolerites from

adjacent well JRP-1 (2A,C) yield a Norian age (210 Ma).

Figure 3 Stratigraphic columns of the Essaouira Basin (left) and the Argana sequence (right), summarising major lithostratigraphic units for both study areas. For the Argana sequences lithostratigraphic members (T1-T10) and corresponding formations are displayed (after Tixeront, 1973; Brown, 1980). Members T5-T8 represent deposits of Carnian to Norian age that are considered to represent the time equivalent sequences in the Meskala Field in Essaouira (both highlighted in orange). Although dating is sparse for the Essaouira sequences, a Carnian to Norian age for the investigated sandstones is postulated based on biostratigraphic ages from offset wells as well as integration of stratigraphic considerations and the relative position to intercalated dolerites and overlying basalts (CAMP) (see also figure 2C).

Figure 4 Schematic cross-section across the Meskala Field, displaying key wells in relation to the reservoir interval (yellow), overlying salt (pink) and basalt (olive green) as well as core availability, location and total depth (TD) of the wells in meters below surface location (BSL) (the exploration wells are close to vertical) (modified after ONAREP, 1995).

Figure 5 Lower 8,65 m of cored section from MKL101bis. This part of the core exhibits facies association 1 (FA1) with detailed description of grain size, sedimentary structures, colour and cementation if apparent. In the interpretation column, the facies interpretation (*sub-facies* code) is displayed, whereas the detailed interpretations for this FA are summarised in table 3. The deposits of this facies association were interpreted to represent ephemeral shallow braided river deposits, with abandonment of the system towards the top and a transition into a wet aeolian sandflat environment. The legend provided for this figure is applicable for figures 5 to 10.

Figure 6 Above a 40 cm core gap to the basal part of the core in MKL101bis (figure 5), the 13,46 m described display facies association 2 with detailed description of grain size, sedimentary structures, colour and cementation if apparent. The facies interpretations for FA2 are summarised in table 4 and were interpreted to represent deposition in a perennial fluvial environment with extensive floodplains and associated crevasse splay deposition (for lithology legend see figure 5).

Figure 7 In these 13,83 m of core from MKL101bis, facies association 3 is described (continued in figure 8). The sediments of this facies association were interpreted to be deposited in transition from a fluvial environment to a delta environment with distributary mouthbars, characterised by the abundance of ripple cross-stratification and coaly interbeds. Delta top facies with organic-rich mudstones, paleosols and rooting is observed (for lithology legend see figure 5). The detailed interpretations for this FA are summarised in table 5.

Figure 8 Summary description of facies association 3 and 4 (15, 60 m). The sediments of facies association 3 were interpreted as delta top sediments with abundant organic-rich mudstones, paleosols, small channel or splay deposits and a high amount of rooting and mottling. The upper 7,5 m of core represent facies association 4, and are interpreted to exhibit full playa conditions under dominantly reducing conditions (note that FA 4 is not presented to scale) (for lithology legend see figure 5). The detailed interpretations for these FA are summarised in table 5.

Figure 9 Core pictures of MKL101bis, exhibiting a compilation of sedimentary features distinctive for facies associations FA2 (figures 9A, B) and FA3 (figures 9C-E).

Figure 10 Simplified summary litholog, documenting sedimentary facies associations and interpretation of the major sequences recognised across the Meskala Field using MKL101bis as an example. The core spans close to 60 m, with a minor core gap of 40 cm in the lower part of the well (see also figure 4 for position within the well). The continuous core was interpreted to represent a progressive evolution from an ephemeral fluvial to wet aeolian sandflat system. A marked change in fluvial depositional style and architecture from sandrich, shallow, sheet like ephemeral channels to larger, perennial fluvial channels with extensive rooted floodplains and paleosols is apparent. Reducing conditions are dominant in the accumulation area, even though the fluvial style and accompanying deposits suggest more perennial, possibly humid conditions. This more perennial fluvial system further develops into a shallow delta setting that is progressively followed by extensive playa deposition under reducing conditions.

Figure 11 Depositional model for sediments investigated in core from the Meskala Field. The figure displays the major depositional units summarised in FA 1 (A), FA2 (B) and FA3 (C).

Figure 12 Geological map of the Argana Valley displaying the lateral distribution of Permian to Lower Jurassic members and formations between Imi N'Tanoute in the North, and Ameskrout in the South. Structural elements (major faults as defined by Tixeront, 1973) are added. An overlay with flow directional measurements from Brown (1980), Mader (2005) and Mader & Redfern (2011) for Carnian to Norian sequences is provided. Study localities 1 to 4 of the outcrop work from Mader (2005) and Mader & Redfern (2011) are marked in red.

Figure 13 Overview on fluvial deposits exposed in the Argana Valley (figure 12, locality 1). **13A** Roughly N-S trending ridge exposes the early and late fluvial phases. The base of T6 (1 - white dashed line) in this locality is erosive and unconformable on underlying mudstones of T5. The early fluvial phase (white dashed line, interval between lines 1 and 2) thickens towards the N and pinches-out towards the S, thinning from 2 m to 0.5 m along this part of the outcrop over a short distance. The base (interval 3, white line) of the late fluvial phase incises deep into underlying floodplain fines of the early fluvial phase. The channels show large-scale cross-bed sets up to 2 m thick. **13B** Coarse pebble conglomerates with minor amounts of fine material, highly erosive, characterise the early fluvial phase (scale bar 2 m). **13C** Deposits of this early fluvial phase thicken from the exposure in 13A of 1 m, to more than 10 m within ~1 km distance. A sharp internal bounding surface (white arrow) can be recognised over several hundred metres, which could suggest some degree of confinement.

Figure 14 Simplified two stage depositional model for the early fluvial facies as recognised in locality 1 (Argana). **14A** Infill of early accommodation space by coarse alluvial fans and braided rivers. The deposits pinch-out to the S, and the main transport direction is towards the W, subordinately NW, with an easterly hinterland source. **14B** The figure highlights the progressive deposition of the low-sinuosity braided river system, sourced from the E. A likely hinterland source could be presented by the Anti Atlas or the Massiv Ancien further in the NE.

Figure 15 Late fluvial phase Argana Valley. **15A** The Amzri section (locality 3) exposes T6 along a roughly E-W trending ridge. White lines highlight the internal fluvial architectural elements of a major river system draining towards the S, SW. **15B** 3D view along a modern river bend in the southern Argana Valley close to the Anglou Dam (locality 4) shows an approximately 70 m thick stacked fluvial unit, composed of amalgamated fine grained sandstone sheets. The fluvial package is thinning towards the W, and exhibits transition into distal floodplain fines in this direction. The main flow direction is towards the S. The system shows a highly aggradational pattern (see also figure 17B).

Figure 16 Simplified depositional model for the late fluvial (perennial) phase (locality 1). Transport direction of fine grained, sand dominated fluvial sandstones is dominantly from East to West (WSW), and extensive floodplain fines to the North. In contrast to the early fluvial phase, the deposits are much cleaner, contain little to no extraformational clasts, and exhibit large (up to 2 m) bedforms of stacked fluvial dunes and bars. The fluvial system is laterally not confined as observed for the early phase.

Figure 17 Spatial sediment distribution of the early (**17A**) and late fluvial phases (**17B**) recognised for T6 in the Argana Valley. The distribution pattern is superimposed on the structural framework of Tixeront (1973). **17A** The limited extend of the early fluvial phase to the centre of the valley is highlighted, with a clear pinch-out mapped towards the S, N and NW. **17B** The spatial distribution of the late fluvial phase, with large, possibly longer-lived rivers extend throughout the entire valley exposures. Only in one locality (2) a dominance of aeolian deposits and only minor distal fluvial intercalations (away from main fairway) was recognised, why for this locality a fluvial margin position was postulated. The dominant transport direction of fluvial flow in locality 3 and 4 are towards the S (SW), whereas locality 1 exhibits an E-W (E-WSW) transport direction. Based on this facies distribution a roughly N-S running channel belt with extensive floodplain deposition, and a fluvial tributary systems from the E is postulated (green). Particularly for the N to S running channel belt, the hinterland source is more speculative and both the Anti Atlas, but also contribution from the Massie Ancien could be anticipated.

Figure 18 Summarises regional aspects of Upper Triassic sedimentation patterns and the regional distribution of prevalent dry climates, modelled for Upper Triassic times (Sellwood & Valdes, 2006).

18A Regional location map of Triassic basins situated around 25° N on the Pangaeon continent within the main corridor of mega-monsoonal oscillations (*sensu* Kutzback & Gallimore, 1989) (map modified after Turner *et al.*, 2001; Sahabi *et al.*, 2004; Turner & Sherif, 2007; Arche & Lopez, 2014; Leleu *et al.*, 2016). The black arrows highlight major transport directions of Upper Triassic alluvial/fluvial sediments investigated during this study, as well as from additional literature data (**1** this study; Mader & Redfern, 2011; **2** Fabuel-Perez *et al.*, 2009a,b; **3** Turner *et al.*, 2001 **4** Arche & Lopez, 2014; **5** Mader, 2005; Hubert & Forlenza, 1988). Super-imposed is the extend of the Tethyan ocean towards the W during Carnian times (after Arche & Lopez, 2014 with edits for the central Essaouira Basin). For the Essaouira basin the paleogeographic model was modified from the original as a result of this study, which suggests major marine incursions to be later than Carnian, possibly as late as Earliest Liassic.

18B & 18C General circulation models (GCMs) for the Upper Triassic (simplified after Sellwood & Valdes, 2016). **18B** shows the modelled simulation of mean seasonal precipitation for the northern hemisphere “winter” season (December- January-February) of the Upper Triassic Pangaeon continent. The red box positions the AOI and its surrounding basins in a zone of pronounced aridity (<0.5 mm/day rainfall), while during the summer months (June-July-August) mean seasonal precipitation is modelled to be < 8 mm/day in the equatorial top low latitudes surrounding the eastern Tethys (summerwet, tropical). The annual shift between these climatic extremes results in extensive evaporite and calcrete deposition in the Tethyan region (Scotese, 2000). While Argana exhibits a more intra mountainous year-round arid climate, possibly in a rain-shadow position, evidence from the Meskala Field of calcrete deposits as well as coaly interbeds points towards direct influence of monsoonal-oscillations with an overall hot and predominantly arid climate interspersed with short wet seasons.

1A MOROCCO

1B STRUCTURAL FRAMEWORK SW MOROCCO

Essaouira lithostratigraphic column modified after Broughton & Trepanier (1993) and Youbi et al. (2003)

Argana lithostratigraphic column modified after Tixeront (1973), Brown (1980) and Hofmann et al. (2000)

Legend

Facies Association 1																																							
Lithology	Scale	Description	Interpretation																																				
<div>Grainsize</div> <div>clay-silt vfs fs ms</div>																																							
<div>-log to scale-</div>	<div>~ 10cm</div> <div>~ 170cm</div> <div>~ 10cm</div> <div>~ 10cm</div> <div>~ 5cm</div> <div>~ 80cm</div> <div>~ 70cm</div> <div>~ 10cm</div> <div>~ 170cm</div> <div>~ 5cm</div> <div>~ 200cm</div> <div>~ 10cm</div> <div>~ 115cm</div>	<p>vfs-fs; dark-red; with abundant silt- mudstone component fs; orange-pink; less well cemented; dominated by horizontal bedding, but very often sedimentary structures overprinted/destratified by patchy, light pink cement s (anhydrite?); overall sandstones appear coarser and more friable (winnowing)</p> <p>vfs-fs; orange-pink; horizontal lamination; well cemented mudstone; dark grey; abundant reworked "wood" debris and coaly fragments (dissolved evaporites?) mudstone; dark grey; fine lamination</p> <p>vfs-fs; greyish-red; very well cemented; massive to remnant horizontal bedding, but no clear stratification or persistent lamination; minor (adhesion)ripples; abundant, packed cavities not coherent with structures</p> <p>vfs-fs; interbedded layers of a) light grey, horizontally laminated vfs-fs; b) dark-grey, horizontal layers with abundant small angular "cavities" (?dissolution); and c) red horizons with wavy to horizontal lamination; abundant mottling and rooting throughout organic-rich debris layer; black; abundant "wood" debris</p> <p>vfs-fs; grey to reddish; mainly massive</p> <p>mudstone; dark grey; fine horizontal lamination</p> <p>vfs-fs; grey to reddish colour; very well cemented; massive to horizontal, finely laminated; oftendestratified, but also abundant low-angle lamination to horizontal bedding preserved; abundant dissolution "cavities", but only in lower 50 cm of unit</p> <p>mudstone; black-dark grey; fine ripple-lamination</p> <p>vfs-fs; grey to reddish colour; very well cemented; massive (destratified?) to horizontal bedding; low angle lamination; abundant "cavities" randomly distributed (not related to sedimentary stratification)</p>	<p>WET AEOLIAN SANDFLAT (1c) arid</p> <p>--- bounding surface --- CREVASSE SPREAD (2c)</p> <p>--- bounding surface --- CHANNEL ABANDONMENT (1b) (rooting; paleosol)</p> <p>FLUVIAL CHANNEL (1a) (abandonment towards the top)</p> <p>--- FLUVIAL CHANNEL (1a)</p> <p>--- FLUVIAL CHANNEL (1a)</p>																																				
<div>Grainsize</div> <div>clay-silt vfs fs ms</div>																																							
<p>Legend: Lithology & Sedimentary Structures</p> <table border="0"> <tr> <td></td> <td>Red mudstone</td> <td></td> <td>Trough cross-bedding</td> <td></td> <td>Dissolution cavities</td> </tr> <tr> <td></td> <td>Black mudstone</td> <td></td> <td>Wavy lamination</td> <td></td> <td>Root traces</td> </tr> <tr> <td></td> <td>Distinct orange coloured sst</td> <td></td> <td>Horizontal lamination</td> <td></td> <td>Bioturbation</td> </tr> <tr> <td></td> <td>Sandstone (sst)</td> <td></td> <td>Low angle lamination</td> <td></td> <td>Water-escape structures</td> </tr> <tr> <td></td> <td></td> <td></td> <td>Moderate to steep cross-bedding</td> <td></td> <td>Ripple cross-stratification</td> </tr> <tr> <td></td> <td></td> <td></td> <td>Planar cross-bedding</td> <td></td> <td>Synsedimentary deformation</td> </tr> </table>					Red mudstone		Trough cross-bedding		Dissolution cavities		Black mudstone		Wavy lamination		Root traces		Distinct orange coloured sst		Horizontal lamination		Bioturbation		Sandstone (sst)		Low angle lamination		Water-escape structures				Moderate to steep cross-bedding		Ripple cross-stratification				Planar cross-bedding		Synsedimentary deformation
	Red mudstone		Trough cross-bedding		Dissolution cavities																																		
	Black mudstone		Wavy lamination		Root traces																																		
	Distinct orange coloured sst		Horizontal lamination		Bioturbation																																		
	Sandstone (sst)		Low angle lamination		Water-escape structures																																		
			Moderate to steep cross-bedding		Ripple cross-stratification																																		
			Planar cross-bedding		Synsedimentary deformation																																		

Legend: Lithology & Sedimentary Structures
Red mudstone		Trough cross-bedding		Dissolution cavities
Black mudstone		Wavy lamination		Root traces
Distinct orange coloured sst		Horizontal lamination		Bioturbation
Sandstone (sst)		Low angle lamination		Water-escape structures
		Moderate to steep cross-bedding		Ripple cross-stratification
		Planar cross-bedding		Synsedimentary deformation

Facies Association 2

Lithology					Scale	Description	Interpretation	
Grainsize	clay-silt	vfs	fs	ms				
- log to scale -					~ 40cm	mudstone ; black; minor wavy to fine ripple lamination	Lacustrine (2d)	dry
~ 80cm	mud-siltstone ; dark red-purple; minor wavy lamination; slickensides	Rooted Floodplain (2a)
~ 180cm	fs-vfs ; brownish-grey; very well sorted and very well cemented; base shows moderately steep cross-bedding	Abandoned Channel (2b)	dry
~ 5cm	fine conglomerate (components < 0.5cm); erosional base; angular basal
~ 3cm	mudstone ; dark grey; horizontal lamination
~ 50cm	conglomerate ; erosional base; light grey; with dark grey mudstone rip-up clasts
~ 3cm	mudstone ; dark grey; with vertical burrows down into breccia; burrows are infilled with dark mud	SHEET FLOODS (flashy) (2c)	wet
~ 50cm	conglomerate ; erosional base; medium-dark grey; clast dominated; clasts consists of red or dark grey mudstone rip-up clasts, minor "wood" fragments and organic debris
~ 180cm	silt-mudstone ; minor vfs-fs interbedded in form of ripples; dark red to purple; horizontally laminated; towards the top, increased destratification and minor fine roots	rooted Flood plain (Playa-lake margin?) (2a)	dry
~ 140cm	vfs-silt ; minor fs; dominated by fine ripple cross-bedding/ lamination; ripple-size (height) decreases towards the top of the interval	Channel abandonment or crevasse splay (2c)
~ 70cm	vfs ; increase in silt- and mudstones; still very well sorted sands; red to orange-red; top of fining upward sequence	Exposure bounding surface (2b)	dry
~ 360cm	vfs-fs (0.088-0.177mm); minor silt; very well sorted; grey-light pink-withish; very well cemented; fine horizontal or low-angle lamination; often massive; fining upward sequence (drying out towards the top)	FLUVIAL CHANNEL (2a)
~ 5cm	mud-siltstone ; dark-light grey; abundant fine "wood" debris and organic material interbedded		wet
~ 130cm	(vfs)-silt-mudstone ; dark red-purple; wavy to horizontal lamination; minor sand input in form of very small scale ripples; minor roots; water-escape structures; overall destratified	FLOODPLAIN (2a)	dry
Grainsize	clay-silt	vfs	fs	ms				

Facies Association 3

Lithology					Scale	Description	Interpretation	
Grainsize	clay-silt	vfs	fs	ms
~400cm	vfs-siltstone ; pinkish; dominated by ripple cross-bedding; sharp basal contact - no transition from underlying black mudstone into vfs; ripples often feature black mud drapes; ripple size <1cm; overall ripple dominated, but as well massive intervals, which are either greyish or pinkish in colour (more clay?); massive intervals can show remnant syn-sedimentary faulting or water escape structures, or in places ripple lamination; ripple lamination often almost merges into horizontal laminated strata	(3a)	DELTA FRONT/ MOUTHBAR in transition to ?LACUSTRINE shoreline
					~ 30cm	mudstone ; black; well cemented; burrows		
					~ 30cm	mudstone ; purple	(3b)	LAKE (pond?) wet dry
					~100cm	vfs-siltstone ; dominated by fine horizontal or ripple lamination; fining upward; increase in mud, and in red-purple colour upwards		
					~ 5cm	mudstone ; black-purple; organic-rich layer	TRANSITION to DELTA CHANNEL Abandonment Delta top	wet
					~100cm	vfs-siltstone ; dominated by fine horizontal to ripple lamination		
					~ 3cm	mud-siltstone ; dark grey; horizontal layer		
					~100cm	vfs-siltstone ; dominated by fine horizontal or ripple lamination		
					~ 5cm	mud-siltstone ; dark grey-black; horizontal lamination dominates		
					~100cm	vfs-fs ; well sorted; dominated by ripple cross-bedding and planar trough-cross bedded sets (~4 cm thick sets); well cemented		
					~500cm	fs-vfs ; light grey- light pink; very well sorted and very well cemented; some horizontal to wavy lamination, but mainly massive	(3a)	FLUVIAL CHANNEL (destratified) wet
						<i>towards top of interval</i> : planar tabular, or trough cross-bed set ~4 cm in thickness; in places fine horizontal lamination and ripple lamination with dark grey to black mud drapes		
						<i>base</i> : low angle to moderately steep cross-bedding		
					~ 10cm	conglomerate ; erosional base; clasts are dominated by orange coloured pedogenic calcrete (dolocrete?) nodules; matrix sediment is moderately steep cross-bedded		
Grainsize	clay-silt	vfs	fs	ms				

Facies Associations 3 in transition to Facies Association 4

A Finely horizontal laminated sst characteristic for fluvial facies of FA2 (scale in cm).

B Dark grey conglomerate, with crude cross-bedding, dominated by mdst rip-up clasts. Clasts are sub-rounded to angular. Abundant organic debris interbedded (scale in cm).

C Dark grey, organic-rich mdsts. Darker layers (black arrows) indicate thin coal layers.

D Well-defined, well cemented ripple-lamination with small scale planar cross-bed sets and s-shaped foresets. **E** Reworked orange to pink calcrite nodules (diameter <1 cm). In the upper part (white dotted lines) moderately steep cross-bedding (scale in cm).

Figure 11 Core pictures of MKL101bis, exhibiting a compilation of sedimentary features distinctive for facies associations FA2 (figures 9A, B) and FA3 (figures 9C-E).

Evolution of the sand dominated reservoir facies (FA1 to FA3) in the Meskala Field

- A FA1** Ephemeral braided river to sheetflood environment
- B FA2** Perennial fluvial channel deposits with extensive floodplain deposition
- C FA3** Shallow (?lacustrine/playa) delta with mouthbars and channel deposits

LEGEND

- 9** Ripple cross-bedded delta sandstones
- 8** Delta mouthbar sandstones
- 7** Channelised delta sandstones
- 6** Interpond mudstones (oxidising) & coaly deposits
- 5** Floodplain mudstones (reducing) & paleosols
- 4** Crevasse splay deposit (channels & splays)
- 3** Anastomosing fluvial channels
- 2** Overbank and abandonment fines
- 1** Stacked channels and sheetfloods

Geological Map of the Argana Valley (after Tixeront, 1973)

8 km

Imi N'Tanoute

Locality 3

Amzri

Locality 2

Ait Khetab Ouzedder

Locality 1

Central Argana

Locality 4

Anglou Dam

Jurassic mudstones (T9-10) & limestones

ARGANA SEQUENCE

T9-10 & Jurassic limestones

Argana Basalt

Bigoudine Fm

Hasseine Mudstone

Sidi Mansour Mudstone

T6 Tadrart Ouadou Sandstone

Timezgadiouine Fm

T5 Irohalene Mudstone

T4 Aglegale Sandstone

T3 Tanameurt Volcaniclastics

Ikakaern Fm

T2 Tourbihine Sandstone

T1 Driss River Conglomerate

LEGEND: Flow directional measurements

Timezgadiouine Fm
(Brown, 1980)

T6 Tadrart Ouadou sst
(Mader, 2005)

Bigoudine Fm
(Brown, 1980)

Faults

Village

A Early Fluvial Phase - stage 1

Central Argana (locality 1)

B Early Fluvial Phase - stage 2

Central Argana (locality 1)

main transport
directions

W WSW

SEDIMENTOLOGY OF THE ESSAOUIRA BASIN (MESKALA FIELD) IN CONTEXT OF REGIONAL SEDIMENT DISTRIBUTION PATTERNS DURING UPPER TRIASSIC PLUVIAL EVENTS

*Nadine K. Mader^{1,2}, Jonathan Redfern¹, Majid El Ouataoui³

¹North Africa Research Group, School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Williamson Building, Oxford Road, Manchester, M13 9 PL, United Kingdom.

²current Address: Maersk Oil Norge AS, Moseidsletta 122, 4068, 4033 Stavanger, Norway (madernadine@hotmail.com).

³ONHYM, 5 Avenue Moulay Hassan, B.P.99, 10050, Rabat, Morocco.

* CORRESPONDING AUTHOR

Keywords: Upper Triassic; Morocco; depositional reservoir model; regional sediment distribution; Triassic climate

TABLES

Member	Author	Vertebrates	Palynology	Radeometric Ages	Chronostratigraphic Age*
Argana Basalt	Verati et al. (2007)			197.8 ± 0.7 Ma	Rhaetian to Sinemurian
	Fiechter et al. (1992)			2.4 Ma	Norian to Pliensbachian
	Manspeizer (1978)			K-Ar 205 ± 17 Ma	Rhaetian to Norian
T8	Tourani et al. (1999)		Assemblages similar to T7 with very good <i>Camerosporites secatus</i> and absence of <i>Corollia/Classopollis</i> (11)	Ar-Ar 200 to 210 Ma	11 Earliest Norian (or younger)
T7	Tourani et al. (1999)		<i>Patinasporites densus</i> (6) <i>Paracirculina quadruplicis</i> (10) <i>Camerosporites secatus</i> (9) <i>Spiritisporites spirabilis</i> (8)		10/9 Late Carnian to Early Norian 8 Late Carnian (Tuvanian)(or younger)
T6	Tourani et al. (1999)		<i>Pseudozonasporites sumus</i> (7) <i>Patinasporites densus</i> (6)		7 Late Carnian (Tuvanian) to Norian 6 Carnian to Middle Carnian
T5	Dutuit (1976, 1977); Jalil & Dutuit (1996); Hunt & Lucas (1991); Lucas (1998)	Phytosaurs (A) <i>Paleorhinus magnolucis</i> (5), <i>Angistorhinus talanti</i> Metoposaurs (B) <i>Metoposaurus azerouali</i> <i>Dicynodonts</i> (C) <i>Moghreberia rinachouensis</i>	Oligochalkian Landfauna		5 Late Triassic; 5 Paleorhinus-biochron (Pangaea): Late Carnian (Tuvanian) A/B/C Assemblage: Carnian to Upper Carnian
	Tourani et al. (1999)			undated; poor data recovery	
	Jalil (1999)	Phytosaurs, Metoposaurs, Aetosaurs and Placerias-like forms			Carnian (Middle to Late Carnian) resembles Upper Triassic European and American assemblages: no distinct dating!
T4	Jalil (1999)	<i>Capitosaurus</i> remains <i>Cyclotus</i> (x)			
T3	Jones (1975); Tourani et al. (2010)	Chirotheriid traces (1) (<i>Synaptichnium</i> , <i>Chirotherium</i> , <i>Brachychirotherium</i> , <i>Isochirotherium</i>)		Volcanics: 247 ± 15 Ma (2); 280 ± 15 Ma (3), 300 ± 15 Ma (4)	Olenekian (1) (Middle Triassic (2); Permian (3) Upper Carboniferous (4))
T2	Dutuit (1988); Jalil & Dutuit (1996); Jalil (1999); Jalil & Janvier (2005); Voigt et al. (2010); Hminna et al. (2012)	Diploid amphibians (A) <i>Diplocolus minimus</i> Chaptorinid reptiles (B) <i>Moradisaurinae</i> <i>Diplocaulida</i> , <i>Capthorinida</i> , <i>Pareiasauria</i> , <i>Hylodichnus</i> , <i>Pachypes</i> , <i>Amphisauroposus</i> ;			Upper Permian; Permian (Ufimian-Kazanian)(A/B)
T1	Jones (1975)			Volcanics in conglomerates 247 ± 15 Ma	Upper Permian to Middle/Low er Triassic

*calibrated to Gradstein et al. (2012)

Table 1 Compilation of biostratigraphic data and absolute ages available for the Argana sequence. Carnian to Norian deposits time-equivalent to the Meskala field sediments are represented in Argana by members (T4) T5 to T8 (highlighted by yellow stripy pattern). Where contradicting datums were available, the more unreliable dates are greyed out in the figure and only the best data sets are highlighted in black font for the chronostratigraphic age column.

Facies Association (FA)	Sub-Facies Code (sfa)	Description	Depositional Environment / Interpretation sub-facies		Climatic Implications
			Sfa	FA	
FA 1	1a	Vfs; minor sltst-mdst; horizontal to low angle stratification; minor ripple stratification; massive; colour change from grey to red; angular dissolution	Ephemeral fluvial channels	Ephemeral fluvial braided river deposition transition to wet aeolian sandflats	periodic arid-humid changing to dominantly arid
	1b	Mixed grey vfs-fs, vfs; finely laminated to adhesion ripple cross-stratified and mottled (rooted?); winnowing; reworked material; abundant secondary angular dissolution	Channel abandonment facies (paleosols, rooting, deflation)		
	1c	Fs; well sorted; horizontally laminated; well cemented; patchy pink cement (anhydrite?); winnowed; distinctively orange coloured	Wet aeolian sandflats		arid
FA 2	2a	Intercalated mdst-sltst; horizontal lamination or highly destratified; dark red to purple; minor vfs-fs in form of ripples interbedded	Floodplain deposits	Perennial fluvial channels with floodplain deposition	humid in hinterland, periodic arid to humid in depositional setting
	2b	Vfs-fs; horizontal lamination to low-angle; or massive to destratified; well cemented; pink colour; fining upward sequences	Fluvial channels (perennial)		
	2c	Clast dominated conglomerate; dark grey or red/purple mudstone rip-up clasts; vfs-fs sst matrix; interbedded with dark-grey mudstones	Sheet flood/splay deposits		
	2d	Mdst; black; enriched w organic material; minor sltst interbedded (ripples)	?Lacustrine/Playa lake shales		
FA 3	3a	Vfs-fs; medium steep to massive destratified; remnant horizontal, wavy laminations; abundant water escape structures and syn-sedimentary faulting; erosive bases with reworked material	(Fluvial) Channels /Delta top (destratified)	Fluvial delta plain to delta top deposition	humid in hinterland, periodic arid to humid in depositional setting
	3b	Vfs-sltst; fining-upward cycles; well preserved horizontal lamination and ripple-cross bedding; thin mudstone horizons and clay drapes are intercalated; locally minor bioturbation	Delta plain/Delta top with delta mouthbars		
	3c	Sltst-mdst; red to purple; massive; alternating with black mdst enriched with organic debris; interbedded vfs ripple cross-stratification; abundant water escape structures; calcretes	Delta top transition to playa (floodplain)		
FA 4	undivided	Mdst; dark red to purple; minor sltst; mottled; minor horizontal lamination	Playa mudstones	Playa deposition	arid

Table 2 Summary of facies associations and related litho-sedimentary *sub-facies*. Both, description and interpretation of the main depositional units are provided in the table. More detailed descriptions and depositional environment descriptions are summarised in tables 3 to 5, and illustrated in figures 5 to 10.

FA1 Sub-Facies Code (<i>sfa</i>)	Description	Interpretation: Ephemeral fluvial braided river deposition in transition to wet aeolian sandflats
1a Ephemeral fluvial channels	Sub-facies 1a consists of moderately well-sorted, very fine-grained, horizontal to low-angle stratified reddish grey sandstones (Figure 5). They often appear massive or destratified and are very well cemented. Typically up to three 1 m to 2 m thick sandstones were recognised throughout the cores. There is an apparent increase in destratification up section. The sandstones are often capped at the top by a thin (<10 cm) black to dark grey mudstone horizon. Silt or finer sandstones can be interbedded with the mudstones in form of thin ripple- or horizontal lamination. The intercalated mudstones show a distinct colour change from dark grey and black to dark red up section. Angular cavities seemingly unrelated to primary stratification can be abundant.	The dominance of horizontal to low-angle stratification, the sheet-like character with sharp top and basal contacts and the overall lack of fine material are interpreted to represent sheetflood fluvial deposition during repeated discharge events (Miall, 1977, 1996, 2006; Fisher <i>et al.</i> , 2008; Anderson <i>et al.</i> , 1999). No clear erosive bases or higher angle cross-bed sets which would point to deep, confined channels were recognised, and deposition in a non-confined braided river system is inferred (Miall, 1977, 1996, 2006; Bridge, 1993; Bridge, 2003, 2006). Occasional interbedded thin mudstone horizons that cap these well sorted sandstones are interpreted to be genetically linked and record deposition from suspension during low-energy flow conditions of waning flow (Miall, 1977; Turnbridge, 1981; Kumar <i>et al.</i> , 1999; Newell <i>et al.</i> , 1999; Platt & Keller, 1992; Owens <i>et al.</i> , 1999; Bridge, 2003, 2006). Massive sandstones defined by prominent destratification and minor remnant faint patchy lamination often associated with these deposits, are hence interpreted to represent post-depositional modification and dewatering during abandonment. Finer, often ripple-laminated sandstones, were mainly recognised at the tops of individual packages, and associated to bedform migration by unconfined tractional flow while lower flow regime conditions prevail (e.g. Jopling and Walker, 1968).
1b Channel abandonment facies (paleosols, rooting, deflation, etc.)	This sub-facies consists of alternations of very fine to fine-grained sandstones that are either a) light grey and horizontally laminated, b) greyish to red with adhesion ripple stratification, c) dark grey, horizontally laminated, with abundant angular cavities, or d) dark red with wavy to horizontal lamination and a mottled (rooted?) appearance. Additionally, thin (<10 cm) dark grey mudstones with abundant organic debris and coaly fragments may be interbedded. Individual bed thickness is centimetre to decimetre scale, and the sediments appear generally very mottled, disrupted and destratified. The sub-facies is further characterised by abundant secondary dissolution cavities.	The mixed interval of <i>sub-facies 1b</i> is characterised by sedimentary structures related to exposure, where highly mottled sediments as a result of rooting or disrupted stratification due to evaporation processes are common. Adhesion ripple stratification as well as interbedded sandstones with sharp top and basal surfaces suggest periodic deposition. Reworked organic matter and organic clasts are also present, and the development of a weakly developed immature paleosol is hence proposed (Retallack, 1988).
1c Wet aeolian sandflats	<i>Sub-facies 1c</i> consists of fine-grained, very well-sorted, exclusively horizontally laminated sandstones, which are less well cemented in comparison to the deposits of <i>sub-facies 1a</i> and <i>1b</i> . The stratification is very well preserved, and distinct inverse grading is characteristic. The very well sorted sandstones lack any fine material, and they are defined by a discrete colour change to orange-pink, that corresponds with the occurrence of patchy pink (?anhydrite) cements.	These deposits above an internal bounding surface are generally slightly coarser and appear winnowed, displaying inverse grading. The sandstones show a marked colour change to orange-pink above the immature paleosol. Concurrently, the interbedded mudstones exhibit a change in colour from grey-black (suggesting reducing conditions) to red (oxidising conditions). The dominance of horizontally bedded inverse stratification in the sandstones is interpreted to highlight the transition from a shallow braided fluvial system (<i>sfa1a</i>) to a wet aeolian sand-flat environment (<i>sfa1c</i>), dominated by fluctuating water tables in a semi-arid to arid environment (e.g. Kocurek & Nielson, 1986; Kocurek, 1988).
Secondary processes	Angular cavities are distinctive in this FA and overprint a variety of <i>sub-facies</i> . The cavities are angular and typically <2 cm in diameter. In the sandstone dominated part of <i>sub-facies 1a</i> , they initially don't appear to be related to depositional sedimentary structures. However, they show an association with facies where primary lamination is strongly altered by dewatering processes. Considered in context to the overall depositional environment, the cavities could represent ghosts of clay rip-up clasts along the bases of braided fluvial sheet flood units or shallow channels. In <i>sub-facies 1b</i> similar dissolution cavities are concentrated to specific horizons. These deposits were instead interpreted as immature paleosols, and the cavities are proposed to record dissolution of an early calcite cement, or gypsum layers related to evaporation and subsequent capillary groundwater processes (Retallack, 1988).	

Table 3 Detailed descriptions and summary interpretation for Facies Association 1 with sub-ordinate *sub-facies 1a* to *1c*. The deposits of this FA are interpreted to report ephemeral fluvial, braided river deposition in transition to a wet aeolian sandflat environment and abandonment of the system (figure 11A). A summary log with typical lithological characteristics of this facies association is in addition displayed in figure 5.

FA2 Sub-Facies Code (sfa)	Description	Interpretation: Perennial fluvial channels with floodplain deposition
2a Floodplain deposits	<i>Sub-facies 2a</i> consists of dark red to purple mudstones with minor siltstone or fine grained sandstone intercalations. Particularly where silt- and sandstones are interbedded, horizontal or wavy as well as ripple lamination is observed. The fine grained sandstones often consist of centimetre-scale single ripple trains forming beds up to 2 cm . In general the mudstones appear highly destratified. Small rootlets were recognised as well as dewatering structures. The mudstone dominated intervals are up to 2 m thick. Towards the top of the section slickensides were observed.	Above a major bounding surface, sedimentary facies changes (figure 6) into several meter thick compound sandstone intervals (<i>sfa2b</i>), interbedded with several meter thick, dark red to purple mud-siltstone units (<i>sfa2a</i>). The upward-fining sandstone units are interpreted to represent fluvial channels (<i>sub-facies 2b</i>) within an extensive floodplain environment (<i>sub-facies 2a</i>) (e.g. Miall, 1995; Nanson & Croke, 1992; Kumar <i>et al.</i> , 1999; Newell <i>et al.</i> , 1999; Colombero <i>et al.</i> , 2015). The intercalated very fine-grained sandstone deposits are up to 4 m thick, and are suggested to represent both amalgamated sheet-like sandstone elements, and more confined (crevasse) channel deposits (Nanson <i>et al.</i> , 1986; Miall, 1996; Bristow <i>et al.</i> , 1999; Bridge, 2003; Medici <i>et al.</i> , 2015). In general it can be noted that silt- to sandstone material is commonly interbedded with the mudstones, suggesting frequent low-energy flood events with non-confined flow during times when fluvial discharge exceeds the systems capacity (Bridge, 2003, 2006; Cain & Mountney, 2009).
2b Perennial fluvial channels	This sub-facies comprises of very fine to fine grained (0.088 – 0.177 mm), very well sorted, well cemented, horizontal to low-angle laminated grey to light pink coloured sandstones (Figure 9A). The sandstones are also massive or destratified in part. Locally, moderately steep cross-bedding, particularly along the basal contacts of individual sandstone packages are apparent. These thick clastic units range between 150 to 400 cm , exhibiting fining-upward from fine to very fine sst and sltst with an increasing mud component. The upper parts of the fining-upward sequences can be dominated by ripple laminations and are often mottled and destratified along their top boundaries.	
2c Sheet flood or splay deposits	This lithofacies comprises of clast-rich conglomerates consisting of dark grey or red mudstone rip-up clasts and minor organic debris within a very fine silt- to very fine to fine grained sandstone matrix (Figure 9B). Individual beds are up to 50 cm thick , have an erosive base and sharp top, and can locally be interbedded with dark grey mudstone layers. In the mudstones abundant mud-filled simple burrows were recognised. The burrows are mainly vertical from the top surfaces down, with diameters of <0,5 cm and vertical length of 1 to 2 cm.	Conglomeratic, sharp based and possibly sheet-like deposits of <i>sub-facies 2c</i> are interpreted to represent overbank, splay and sheetflood deposition. They become more prominent up section, which could suggest progradation of the crevasse splays.
2d ?Lacustrine/ Playa shales	Up to 40 cm thick organic-rich black mudstone horizons, interbedded with thin siltstone intercalations featuring minor wavy to fine ripple lamination.	The top bounding surface of this sequence is marked by the deposition of a 40 cm thick black mudstone horizon (<i>sub-facies 2d</i>). A transgressive event, related to expansion of a shallow (playa) lake, or alternatively the development of laterally extensive inter-pond area on the floodplain with reducing conditions is anticipated (e.g. Wright & Mariott, 1993; Mariott & Alexander, 1999; Bridge, 2003). For both scenarios high groundwater tables are a prerequisite.

Table 4 Detailed descriptions and summary interpretation for Facies Association 2 with sub-ordinate *sub-facies 2a* to *2d*. FA2 was interpreted to represent perennial fluvial conditions with extensive floodplain deposition and associated crevasse splay deposition (figure 11A). A summary log with typical lithological characteristics of this facies association is in addition displayed in figure 6.

FA3 Sub-Facies Code (sfa)		Description	Interpretation: Fluvial to delta plain and delta top deposition
FA 3	3a (fluvial) Channels to delta top (destratified)	<i>Sfa3a</i> is on first account very similar to <i>sub-facies 2b</i> , characterised by moderately steep cross-bedded or massive, destratified very fine to fine grained sandstones which are light grey in colour. Thicknesses of individual sandstone units are comparable (up to ~4 m), but in contrast to <i>sub-facies 2b</i> , the basal contacts of <i>sub-facies 3a</i> are often erosive and basal contacts contain reworked material, such as calcrete (?dolocrete) nodules (figure 9C). The reworked nodules are orange to whitish in colour and typically <1 cm in diameter, aligning with the foresets of the medium steep cross-bedding.	Above a black shale marker horizon, two to three fining-upward sequences were recognised that comprise 70-80% of FA3 (figures 8, 10). Both fining-upward sequences consist of moderately steep cross-bedded to massive deposits (<i>sub-facies 3a</i>) in the lower part, transforming into ripple cross-stratification interbedded with thin organic-rich mudstone intercalations in the upper part (<i>sub-facies 3b</i>). Sedimentary evidence is ambiguous, and sedimentary features could record similar facies previously described in crevasse splay deposits and lacustrine deltas (Miall, 1996; Bridge, 2003). Certainly in the lowermost sequence investigated in the core, the erosive based, moderately steep cross-bedding with abundant reworked coarse material suggests an erosive channel base and large scale dune migration from traction currents. As flow wanes, abandonment of the system would lead to extensive ripple laminated strata as observed. However, the entire second fining upward cycle is dominated by ripple-cross stratification throughout, featuring abundant climbing ripple trains above a sharp but non erosive basal contact. The abundance of organic material interbedded further suggests transition into a more lacustrine delta depositional environment, with subaqueous levees or delta-mouthbar deposits (e.g. Miall, 1996; Bridge, 2003).
	3b Delta plain or delta top and delta front mouthbars	<i>Sub-facies 3b</i> is dominated by very fine grained sandstones to siltstones with very well-defined and well-preserved ripple cross-stratification, planar trough-cross bed sets and minor horizontal lamination or massive appearance. The ripple facies consist of stacked, small scale (2-4 cm) planar cross-bed sets with s-shaped foresets, and finer material lining the latter (Figure 9D). Individual ripples are low relief, often < 1 cm, almost merging into horizontal lamination. The ripple cross-stratified sandstones are either intercalated with thin (~5 cm thick), dark grey mudstone-siltstone horizons, or abundant thin, black, organic-rich mudstone layers with a coaly character (figure 9C). Locally, small vertical burrows and rootlets intersect the mudstones. Water-escape structures and syn-sedimentary faulting appear throughout. The upper 2 m to 4 m of each fining upward cycle are dominated by this sub-facies.	
	3c Delta top in transition to playa (floodplain)	This sub-facies shows similarities to <i>sub-facies 2a</i> , with heterolithic deposits characterised by up to 3 m thick mudstone-siltstone-sandstone packages. The mudstone intervals are overprinted by an abundance of rootlets and highlight an overall mottled appearance as in <i>sub-facies 2a</i> , but the reddish background facies can in contrast be interbedded with dark grey to black mudstones enriched with organic debris (up to 5 cm large clasts). In addition, up to 30 cm thick layers packed with orange coloured calcrete (?dolocrete) nodules (<1 cm) were observed (Figure 9D). The interbedded coarser material is defined by sharp basal contacts, that are typically ripple cross-stratified and massive to destratified with visible water-escape structures towards their top contacts. The orange coloured nodules do not follow any obvious bedding, but are concentrated in packed layers with little matrix surrounding them.	At the base of the lower fining-upward sequence (figures 8, 10), the occurrence of reworked orange-coloured pedogenic nodules suggest the development of palaeosols (calcrete/?dolocrete) (Retallack, 1988; Khalaf, 2007) within the adjacent floodplain or delta top. In the upper fining-upward sequence in-situ calcrete nodules were recognised. In these sediments primary stratification of the host sediment is distorted and sedimentary structures indicative of reworking and flash-flood deposition are lacking, confirming the preservation of more mature paleosols in the system. In addition, the upper part of each cycle contains abundant organic-rich layers, adhesion ripples, mottling and rooting, interpreted to represent periodic exposure, following either a shifting lake shoreline or simply lateral channel switching on the flood or delta plain.
FA4 Sub-Facies Code (sfa)		Description	Interpretation: Playa Fines
FA 4	Playa deposits	The FA is dominated by dark red to purple, mottled mudstones which can change progressively into dark red to purple silt- to mudstones with minor horizontal or wavy-horizontal lamination, regularly reverting back to cleaner mudstone intervals. Minor very fine to fine grained sandstone intervals are interbedded in the form of small-scale (<1cm) adhesion ripple stratification. In contrast to the lower units and similar mudstones observed in FA2 and FA3, no intercalated dark grey to black mudstone intervals or thicker (>1 cm) sandstone intervals were observed.	The dominance of dark red mudstones is interpreted to represent full playa conditions (e.g. Hofmann <i>et al.</i> , 2000; Arche, 2007). Repeating shallowing and subsequent drying of the playa is indicated by an increase in mottling and rooting and the abundance of adhesion ripple stratification. Destratification and water-escape structures are common and the deposits show a high degree of exposure (e.g. Nanson <i>et al.</i> , 1986; Hofmann <i>et al.</i> , 2000).

Table 5 Detailed descriptions and summary interpretation for Facies Association 3 and 4 with subordinate sub-facies 3a to 3c. The deposits of FA3 were interpreted to consist of fluvial delta plain to delta top deposits in a shallow (playa) lake environment (figure 11C), whereas the mudstone dominated FA4 was interpreted to highlight full playa conditions. A summary log with typical lithological characteristics of these facies associations are in addition displayed in figure 7 and 8.