Primary microglia maintain capacity to function despite internalisation and intracellular loading with carbon nanotubes

DOI:

[10.1039/C7NH00055C](https://doi.org/10.1039/C7NH00055C)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Bussy, C., Bianco, A., Prato, M., & Kostarelos, K. (2017). Primary microglia maintain capacity to function despite internalisation and intracellular loading with carbon nanotubes. *Nanoscale Horizons*, 2. <https://doi.org/10.1039/C7NH00055C>

Published in:

Nanoscale Horizons

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Nanoscale Horizons

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: C. bussy, A. Bianco, M. Prato and K. Kostarelos, *Nanoscale Horiz.*, 2017, DOI: 10.1039/C7NH00055C.

This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Primary microglia maintain capacity to function despite
internalisation and intracellular loading with carbon nanotubes

Cyrill Bussy^{1,*}, Alberto Bianco², Maurizio Prato^{3,4,5} and Kostas Kostarelos^{1,*}

¹ *Nanomedicine Lab, Faculty of Biology, Medicine and Health & National Graphene Institute, University of Manchester, AV Hill Building, Manchester M13 9PT, United Kingdom*

² *CNRS, Institut de Biologie Moléculaire et Cellulaire, UPR 3572, Immunopathologie et Chimie Thérapeutique, Strasbourg, France*

³ *Center of Excellence for Nanostructured Materials, Department of Chemical and Pharmaceutical Sciences, University of Trieste, Trieste, Italy*

⁴ *Carbon Nanobiotechnology Laboratory, CIC biomaGUNE, Paseo de Miramón 182, 20009 Donostia-San Sebastian, Spain*

⁵ *Basque Science Foundation, Ikerbasque, Bilbao 48013, Spain*

* Corresponding to: cyrill.bussy@manchester.ac.uk; kostas.kostarelos@manchester.ac.uk

Abstract

Biomedical applications of functionalised carbon nanotubes (*f*-CNTs) for imaging as well as drug or gene delivery in the brain have recently gained interest. Several studies have demonstrated the potential of *f*-CNTs to offer treatment options for neurological conditions with success. However, there is also evidence that *f*-CNTs accumulate preferentially within microglial cells when introduced in the brain. Considering the key immunologic role of these cells in the brain and the limited knowledge regarding the interaction of CNTs with microglial cells, it is imperative to understand whether accumulation of CNTs in microglial cells can alter their physiological functions or trigger pro-inflammatory signalling. The aim of the present study was to investigate the basic physiological functions of isolated primary microglial cells over time, following their exposure to multi-walled carbon nanotubes functionalized via different surface chemistries. We rationalized that some chemical strategies may be more deleterious for microglial cell functions than others. We used rat primary microglial cells that can be maintained in cell culture for a long period of time without undergoing cell division. Cell viability, phagocytosis, migration, pro-inflammatory factor release were studied over one month, following a single exposure to a non-cytotoxic concentration of three different *f*-CNTs (carboxylated, aminated and both carboxylated and aminated). We show that *f*-CNTs do not induce inflammation and are not affecting basic functions of microglial cells under these conditions. Importantly, this was the case from day 1 to day 28 post-exposure, even though microglial cells had internalized *f*-CNTs in subcellular compartments and retained the nanotube load in their cytoplasm over time.

Key words

nanomedicine; brain; biocompatibility; central nervous system; phagocytosis; neurotoxicity; nanotoxicology

1 **Introduction**

2
3 In recent years, numerous nanomaterials have been utilised in an attempt to
4 overcome the great challenge of drug delivery to the central nervous system ¹⁻³. Among
5 them, carbon nanotubes (CNTs) have lately shown great promise ⁴ due to their inherent
6 properties, but also a unique ability to cross cellular barriers ⁵. For instance, acetylcholine
7 adsorbed onto single-walled CNTs (SWNTs) was efficiently delivered to cells in a kainic acid
8 induced Alzheimer's Disease (AD) mouse model ⁶. Amine functionalized multi-walled CNTs
9 (MWNTs) have also been used successfully to deliver Caspase 3 siRNA into neurons with
10 the aim to reduce the ischemic injuries in the motor cortex of an endothelin-1 induced stroke
11 model ⁷. Similarly, amine functionalized SWNTs were injected in mouse brain ventricles in
12 order to decrease the effects of ischemia and inflammation in an artery occlusion model of
13 stroke ⁸. The SWNT treated animals showed better histopathological and behavioural profiles
14 after injury, indicating that aminated SWNTs could achieve a neuroprotective effect even
15 without carrying any therapeutic agent. More recently, PEG-SWCNTs have also been
16 proposed as solutions to alleviate glutamate excitotoxicity of neurons by increasing astrocytic
17 glutamate uptake ⁹.

18
19 Once localised in the brain parenchyma, knowledge of the fate and biological effects
20 of those functionalised nanomaterials remains however limited. This raises the need for a
21 better understanding of the interactions of carbon nanomaterials with the different cell
22 subpopulations in the central nervous system (CNS). While neurons might be a preferred
23 target for therapeutic purposes, nanomaterial in the brain parenchyma are likely to interact
24 with other cell types including the resident macrophages of the brain, namely the microglial
25 cells. Even in the case of nanoscale vectors bearing neuron-targeting moieties, microglial
26 cells have indeed been shown to be the cell type in which a significant fraction of the
27 nanomaterials were accumulating ¹⁰. Considering that microglial cells are the main cellular
28 component of the immune system in the brain, any nanomaterial that is intended for
29 intracerebral applications should therefore have a controlled and well-understood interaction
30 with those cells.

31 In terms of our current understanding of CNT-microglia interactions, microglial cells
32 have been reported to play a major role in internalising functionalised CNTs (*f*-CNTs) in
33 comparison to other cell types following stereotactic administration of *f*-CNT in the cortex of
34 mice ^{7, 11, 12}. Similar results were obtained *in vitro* in mixed cell culture of primary glial cells,
35 whereby microglial cells were displaying higher uptake of functionalised MWNTs (*f*-MWNTs)
36 compared to astrocytes ¹³. Interestingly, following internalisation in microglial cells, we have
37 observed the partial degradation of *f*-MWNTs over time, both *in vivo* over 14 days ¹² and *in*

1 *vitro* in a primary cell culture model up to 90 days^{14, 15}. Alteration within 24h of the graphitic
2 structure of *f*-MWNTs in the immortalised N9 microglial cell line offered independent
3 validation¹⁶.

4 These degradation results suggested that despite the internalisation of *f*-CNTs,
5 microglial cells were able to maintain one of their primary functions over time: their ability to
6 degrade foreign materials^{12, 14, 17}. In contrast, MWNTs dispersed in horse serum were shown
7 *in vitro* to alter microglial functions such as migration and phagocytosis, and also cell division
8 in BV2 immortalised microglial cell model after their internalisation¹⁸. The main hypothesis to
9 explain these alterations was the direct interaction of MWNTs with the cytoskeleton of BV2
10 cells leading to an impairment of all cytoskeleton related functions including migration and
11 phagocytosis. The present study was therefore designed to interrogate the impact of
12 internalisation and retention of *f*-CNTs in microglial cells on their capacity to perform normal
13 cellular and macrophage-specific functions, other than their degradation ability that was
14 previously demonstrated^{14, 15}.

15
16 In addition, using different chemical derivatives of *f*-CNTs, surface properties have
17 been revealed as essential determinants that could modulate both CNT-microglia interaction
18 and bioreactivity of CNTs towards microglial cells. Firstly, the type of surface functionalisation
19 was demonstrated to influence the uptake¹⁹. Phosphatidyl-serine coated SWNTs were
20 engulfed by primary rat microglia while phosphatidyl-choline coated and non-coated SWNTs
21 were not. Similarly, microglial cells were shown to be very efficient at internalising oxidised
22 MWNTs but were not fully able to break apart agglomerates of pristine MWNTs and to
23 internalise them¹⁶. Secondly, the type of functionalisation was shown to modulate the
24 response of microglial cells. While both pristine and oxidised MWNTs were not altering cell
25 viability, pro-inflammatory cytokine release or nitric oxide (NO) production, a significant
26 increase of reactive oxygen species production was measured after exposure to pristine
27 MWNTs, suggesting possible impact on cell viability at long term¹⁶. Accordingly, using an *in*
28 *vivo* cortical stereotactic injection model, brain cells and microglial ones in particular were
29 reported to differently respond to *f*-CNTs exposure depending on the surface
30 functionalisation of the materials injected¹¹. Functionalisation was not only impacting the
31 localisation and distribution pattern of the injected materials within the brain, but also
32 modulating the inflammatory profile of the different *f*-CNTs. Oxidised MWNTs were shown to
33 induce and sustain a microglial cell mediated inflammatory response while aminated MWNTs
34 were only transiently inducing inflammation.

35 Taken together, these studies demonstrate the major importance of material surface
36 properties and in particular the type of surface functionalisation on the biological outcomes of
37 the material-cell interaction. Moreover, ability by CNTs to induce inflammation and impact on

1 microglial cell physiological functions are two key endpoints that should be systematically
2 investigated and correlated to the surface chemistry of the materials in order to determine the
3 best biocompatibility profile for putative CNT-based brain delivery systems.
4

5 The objectives of the present study were therefore to investigate whether microglial
6 cell functions can be affected by the internalisation of *f*-CNTs, whether varying the surface
7 chemistry of *f*-CNTs can modulate the cell response, and whether the impact of these
8 different *f*-CNTs on microglial cell functions may evolve over time. Oxidation of CNTs has
9 been shown to increase the bioreactivity of those materials for macrophages^{11, 20}, we
10 therefore hypothesised that the degree of oxidation/carboxylation is determinant to modulate
11 the biological responses and can affect microglial cell functions while not inducing cell death.
12 To assess this hypothesis we selected three MWNTs with different carboxylation degree
13 (purely carboxylated, purely aminated and one type bearing both carboxyl and amine groups).
14 We additionally developed a primary rat microglial cell model that demonstrated a long
15 lifespan *in vitro* without undergoing division to monitor over time the evolution of microglial
16 cell functions after internalisation of the materials. Cytotoxicity, phagocytosis, migration and
17 pro-inflammatory mediator release were studied at different time points and up to 28 days
18 following a single exposure to a non-cytotoxic concentration of the three selected *f*-CNTs.
19 We report that *f*-MWNTs irrespective of their chemical functionalisation did not affect
20 microglial cell viability under our conditions and did not induce release of pro-inflammatory
21 mediators at any time points. Moreover, despite an important perinuclear internalisation of
22 materials, microglial cells maintained their abilities to phagocytose and to migrate.
23 Interestingly, oxidation of MWNTs stimulated the migration ability of microglial cells that have
24 engulfed those CNTs in comparison to aminated MWNT laden cells.

25

26 Results and Discussion

27 **Chemical structure and characterisation of MWNT suspension.** The selection of
28 MWCNTs over other CNT types such as single walled- or double walled- CNTs was
29 motivated by our previous works on the interactions of *f*-MWNTs with brain cells^{7, 11-13, 21} and
30 microglial cells in particular^{14, 15}. We have also demonstrated that these functionalised
31 nanomaterials can be used as delivery platforms for drugs²², genes^{7, 23} or contrast agents²⁴.
32 The type of chemical functionalisation was previously identified by some of us as a major
33 determinant of the *in vivo* ability of *f*-MWNTs to induce inflammation in the brain¹¹.
34 Comparing two types of *f*-MWNTs, one that has been pre-oxidised and then aminated with
35 one that has been aminated only, oxidation/carboxylation was revealed as causing
36 inflammation, even though the effects of MWNTs bearing only carboxyl groups (i.e. negative

charges) were not directly investigated. Here, three chemically functionalised MWNTs with different level of oxidation were synthesised from the same starting materials in order to limit batch-to-batch variations. Their chemical structure is presented in **Figure 1** and they include: (i) carboxylated MWNTs (ox-MWNTs) prepared by treatment of the starting materials in strong acid conditions²⁵; (ii) amino-functionalized oxidised MWNTs (ox-MWNT-NH₃⁺) prepared by 1,3 dipolar-cycloaddition reaction after an initial oxidation as described previously²⁶; and (iii) amino-functionalized MWNTs (MWNT-NH₃⁺) prepared following the 1,3 dipolar-cycloaddition reaction on the starting materials as described previously^{27, 28}. Length of MWNT-NH₃⁺ remained similar to the one of the starting materials (between 0.5-2 µm) while it was reduced for materials undergoing oxidation (shorten to 300-400nm for ox-MWNT and ox-MWNT-NH₃⁺). Suspensions of MWNTs were prepared by hydrating dry powder with a 5% dextrose water solution, which was then sonicated until MWNTs reached a good dispersion (about 40 min). It was essential to achieve a dispersion of similar quality for the three materials to prevent non-intrinsic differences amongst them. Difference in dispersion quality is indeed known to induce differences in biological responses especially inflammation^{29, 30} and could therefore lead to biases in experiment outcomes and misinterpretation in regards to the role of chemical functionalisations. To assess the stability of the 3 MWNT dispersions in 5% dextrose water solution, analysis of the absorbance (used as an index of dispersion, well dispersed materials absorbing more than agglomerated ones) was performed at 0 h and 24 h (**Figure S1**, first column). Despite the intrinsic differences between the 3 materials (highest absorbance: ox-MWNT-NH₃⁺ > MWNT-NH₃⁺ > ox-MWNT), no difference between the absorbances at 0 h and 24 h could be observed for any materials. All 3 materials were stable over 24 h in 5% dextrose. Similar stability results over 24 h were observed when the materials initially suspended in 5% dextrose were diluted in cell culture medium supplemented with 12% FBS (**Figure S1**, second column). However, when materials initially suspended in 5% dextrose were then diluted in cell culture medium without FBS, the UV-Vis spectrometric analysis evidenced a clear sedimentation for all 3 materials, regardless of their surface chemistries (**Figure S1**, third column). These results suggested that MWNT suspensions were not stable in cell culture medium without the addition of FBS, and would sediment at the bottom of the cell culture wells under these conditions, increasing the potential of materials to interact with the monolayer of cells. In contrast the addition of FBS would not be favourable to interaction with cells as most materials would stay in suspension, above the cell monolayer.

Functionalised nanotubes had no effect on microglial cell viability. Microglial cells were revealed as the main cell population in which f-MWNTs are accumulating when infused in the brain parenchyma^{7, 11, 12}, or when exposing mixed glial cells cultures (containing mainly

1 astrocytes and microglial cells)^{13, 14}. However, the impact of this preferential accumulation
2 remains elusive and contradictory in the literature. We therefore first questioned whether the
3 exposure to different *f*-MWNTs using a non-overload concentration will affect the viability of
4 microglial cells over time. To answer that question, a model of primary microglial cell cultures
5 that can be maintained over long period of time without cell division was used. It was
6 rationalised that once in the brain parenchyma, *f*-MWNTs will diffuse throughout the tissue as
7 observed for other nanomaterials³¹ and therefore each microglial cell will likely be exposed
8 to a limited amount of nanotubes. To ensure that microglial cells interacted with the MWNTs,
9 we favoured the CNT sedimentation by exposing cells to MWNTs in a serum free
10 environment for 2 h, following which the cell culture medium was supplemented with 12%
11 FBS. Following this single exposure to a unique dose of 10 µg/mL, we used a modified LDH
12 assay developed in our lab³² to assess the impact of *f*-MWNTs on microglial cell viability at
13 1, 7, 14 and 28 days after the initial exposure (**Figure 2**). No effect on the viability of exposed
14 microglial cells compared to untreated cells was observed irrespective of *f*-MWNTs used or
15 time after exposure. However, a statistically significant difference was observed at day 1 for
16 ox-MWNT-NH₃⁺ treated cells when compared to their respective control (untreated cells). This
17 could be ascribed to the primary nature of the cell cultures used here (more dying cells in
18 untreated cells than in ox-MWNT-NH₃⁺ treated cells)¹⁴ or to a stimulation of viability as
19 primary microglial cells preserve the ability to divide under stimulatory conditions³³. Further
20 investigations will be needed to address this question. Overall, these results suggested that
21 microglial cells are tolerant to *f*-MWNTs exposure under the conditions used here, and that
22 this tolerance is not dependent on the type of chemical functionalisation for this
23 concentration. These results were in agreement with previous observations of a dose
24 dependent cytotoxicity appearing after 24 h above 10 µg/mL¹³.

25

26 **Primary microglial cells internalised and kept nanotubes over time.** CNTs found
27 primarily free in the cytoplasm were reported to reduce proliferation and induce apoptosis in
28 BV2 cells, an immortalised model of microglial cells¹⁸. In order to explain the difference
29 between this study and the viability results of the present study (**Figure 2**), optical
30 microscopy was used to investigate the internalisation pattern and evolution of CNT loading
31 over time (**Figure 3a, S2 and S3**). For all three *f*-MWNTs and irrespective of the time point
32 considered, the distribution of nanotubes observed as black material in bright-field mode
33 (**Figure S2 and S3**) was characteristic, with an accumulation of materials around the nucleus
34 (perinuclear region), absence of black material in the nucleus, and no apparent accumulation
35 in the cytoplasm periphery close to the cell boundaries (**Figure S3**). No obvious difference in
36 terms of CNT loading from 1 day to 28 days was observed for all three *f*-MWNTs, suggesting

that exocytosis was limited. In addition, using immunocytochemistry (anti-CD68), a singular pattern of CD68 immunolabelling was observed, with black materials appearing as entrapped in vesicles positive for CD68 immunoreactivity (**Figure S4**). The vesicular pattern was enhanced and contrasted by the black materials contained within the CD68 positive vesicles. CD68 is predominantly an intracellular glycosylated transmembrane protein with minimal surface expression in unactivated macrophages³⁴. The perinuclear position of CNT accumulation, the absence of diffuse CNT-related black signal across the cytoplasm, and the vesicular pattern of CD68 immunoreactivity suggested a vesicular accumulation of *f*-MWNTs.

In order to confirm that the punctuated black halo around the nucleus (**Figure S5b**) was indeed due to localised accumulation of carbon nanotubes, Raman spectroscopy analysis was performed at a single cell level and in different places of the considered cell (**Figure 3 b**). The characteristic peaks at 1360 cm⁻¹ (D band), 1585 cm⁻¹ (G band) and 1620 cm⁻¹ (D' band) were used to confirm the CNT nature of the accumulation^{35, 36}. Following exposure to *f*-CNTs, all three microglial cell cultures exhibited the specific Raman signature of MWNTs (*i.e.* D, G, D' bands) in the perinuclear region (**Figure S5b**). For each *f*-CNTs, the Raman profile found in cells (**Figure 3b**) was similar to the Raman profile of the starting materials used to treat the cells (**Figure S5a**), confirming the adequacy and specificity of Raman fingerprints to characterise carbon nanomaterials. In addition, Raman signature of MWNTs was observed at each time points tested (*i.e.* 1, 7, 14 and 28 days), confirming the persisting perinuclear presence of *f*-CNTs in each case (**Figure 3b**). Taken together, these results show that the type of chemical functionalisation had no influence on the internalisation of *f*-MWNTs, their distribution within the cells, and their overtime retention in microglial cells. However, both endocytosis and exocytosis of *f*-MWNTs in primary microglial cells will need further investigations to fully describe the mechanisms and kinetic of processes at stake. In particular, further studies investigating the amount of materials internalised and maintained in cells over time for each *f*-MWNT considered will be important, as over time degradation in microglial cells in a surface chemistry dependent manner has been recently confirmed using two different methodological approaches^{14, 15}. Importantly, as mentioned above, the persistent retention of *f*-MWNTs over 28 days around the nucleus of microglial cells had no effect on viability, which demonstrated an absence of correlation between intracellular accumulation of materials and cytotoxicity under the considered conditions.

Uptake of *f*-MWNTs did not induce pro-inflammatory reaction in microglial cells. Since internalisation of CNTs had no effect on microglial cells viability over time, it was then questioned whether the CNT uptake could induce the release of pro-inflammatory factors. The Griess assay was therefore used to measure the amount of NO released by MWNT

1 exposed microglial cells at 1, 7, 14 and 28 days after the initial exposure (**Figure 4**). NO is
2 one of the main markers for microglial cell activation and pro-inflammation induction, and is
3 also a toxic agent for neurons^{17, 37}. A statistically significant release of NO compared to
4 untreated cells was observed 24 h after the initial exposure to CNTs for all three *f*-MWNTs.
5 However, this positive release of pro-inflammatory markers was largely limited in comparison
6 to LPS stimulated microglial cells. At later time points, the release of NO became not
7 significant compared to untreated cells and disappeared at 28 days. These data showed that
8 *f*-MWNTs irrespective of their surface functionalisation have a limited ability to induce
9 inflammation *in vitro* under the conditions used here, and that their internalisation and
10 persistence in microglial cells over time did not lead to chronic pro-inflammatory reaction, as
11 would have been expected if microglial cells were activated and inducing an immune
12 response, as previously observed with other MWNTs¹⁶.

13 **Uptake of *f*-MWNTs did not alter internalisation or migration abilities.** The presence of *f*-
14 MWNTs inside microglial cells for 28 days did not seem to affect their viability or to induce an
15 inflammatory reaction. The next question was then to know whether other basic physiological
16 functions characteristic of resident macrophages would stay unimpaired by *f*-MWNT
17 exposure and over-time persistence in cells.

18 Firstly, the internalisation ability of CNT-laden microglial cells was tested, using 0.5
19 µm fluorescently labelled polystyrene beads. It was rationalised that if the internalisation
20 ability was maintained, the uptake of polystyrene beads for each microglial cell considered
21 would be limited only by the amount of materials (*i.e.* CNTs) already inside these MWNT-
22 preloaded cells. The internalisation of polystyrene beads inside microglia previously exposed
23 to MWNTs was assessed over 8 h and compared to naïve microglial cells (*i.e.* non-exposed
24 to MWNTs) and activated microglial cells (*i.e.* non-exposed to MWNTs but stimulated by LPS
25 (200 ng/mL)) (**Figure 5 and S6**). The internalisation assay was performed only at day 7 after
26 exposure to CNTs because it was observed that engulfment of polystyrene beads was
27 leading to acute cytotoxicity at later time points (14 and 28 days) for all cells, naïve
28 microglial cells included. This was attributed to the age of the primary cells, which have been
29 isolated from brain tissue about 1.5 months before day 14 time point. Similarly, the test was
30 not performed at day 1 after exposure to MWNTs because the cells were exposed to
31 MWNTs the day before and some free nanotubes might still be in the supernatant and would
32 be in direct competition with the polystyrene beads for internalisation. At the earliest time
33 point (1 h after the bead treatment), there was no major difference between groups, including
34 untreated cells or LPS stimulated cells (**Figure 5c and S6**). While non-statistically significant,
35 the amount of polystyrene beads (*i.e.* fluorescence) in microglial cells previously exposed to
36 oxidised MWNTs (*i.e.* both ox-MWNT and ox-MWNT-NH₃⁺) was however lower compared to

untreated cells (**Figure 5c and S6**). After 2 h, the LPS stimulation greatly enhanced the polystyrene beads internalisation compared to the other groups (**Figure 5a and 5c**). In addition, the internalisation of polystyrene beads was slightly higher for microglia exposed to aminated MWNTs (*i.e.* both ox-MWNT-NH₃⁺ and MWNT-NH₃⁺) compared to naïve microglia or microglia exposed to ox-MWNT. After 8 h, there was no significant difference between the various groups suggesting that all groups of cells had reached their highest loading in polystyrene beads possible (**Figure 5b and 5c**). It also suggested that despite their initial CNT loading, CNT-laden microglial cells were able to internalise polystyrene beads to the same extend as cells not pre-loaded with CNTs. These results showed that the internalisation ability of microglial cells was not perturbed by uptake of *f*-MWNTs, irrespective of the type of chemical functionalisation tested. These results also underlined the relative tolerance of cells to *f*-MWNTs exposure, especially when considering non-overload non-toxic doses (*i.e.* 10 µg/mL).

The migration ability of CNT-laden microglial cells was then tested. Following the observation that unintended scratch on the plastic cell culture surface during the renewal of cell culture medium was inducing migration of microglial cells toward the scratch, a specific migration assay based on “scratch stimulated” migration was developed. The migration assay was performed at day 7, 14 and 28 days after the initial exposure to one of the three *f*-MWNTs and compared to the results with naïve microglial cells and microglial cells stimulated with LPS (200 ng/mL) (**Figure 6 and S7**). The assay was not performed on the day following the exposure to CNTs as free CNTs might still be present in the supernatant and perturb the assay. Unexpectedly, migration of microglial cells stimulated by LPS was inhibited at all three time points tested compared to untreated cells (**Figure S7 and 6**). LPS is indeed known to stimulate migration and should have acted as a positive control/stimulator of migration³⁸. It was previously reported that the amount of LPS used during an assay is a critical parameter that can influence the motility of primary microglial cells³⁹. While low (0.1 ng/mL) or high (100 ng/mL) concentrations were shown to have no effect on promoting migration, 10 ng/mL was reported as the adjusted dose which induce higher migration compared to control cells. In our case, the reduce motility of microglial cells under LPS condition can possibly be explained by the deviation of normal macrophage functionalities toward an activation profile dedicated to pro-inflammatory factors secretion due to overload LPS dose (*i.e.* 200 ng/mL was used for both NO release and migration assay). This will require further investigations to fully understand the reason of this unusual behaviour of LPS stimulated microglial cells and confirming the optimal concentration of 10 ng/mL for promoting migration³⁹.

As regards to *f*-MWNTs, a chemical functionalisation dependent modification of the microglial cell migration ability was observed (**Figure 6**). At any time point, microglial cells

1 exposed to ox-MWNT had higher migration ability than cells exposed to aminated MWNTs
2 (*i.e.* both ox-MWNT-NH₃⁺ and MWNT-NH₃⁺) with MWNTs bearing both chemical functions
3 (*i.e.* ox-MWNT-NH₃⁺) in between ox-MWNT and MWNT-NH₃⁺. At day 7, microglial cells
4 exposed to ox-MWNT migrated to the same extend as untreated cells whereas microglial
5 cells exposed to aminated MWNTs (*i.e.* both ox-MWNT-NH₃⁺ and MWNT-NH₃⁺) moved less
6 than untreated cells. At day 14, surprisingly, microglial cells exposed to any type of MWNTs
7 migrated more than untreated cells. At day 28, only microglial cells exposed to oxidised
8 MWNTs (*i.e.* both ox-MWNT and ox-MWNT-NH₃⁺) migrated more than untreated cells or
9 cells exposed to MWNT-NH₃⁺, which both show similar migration ability. These results
10 showed that internalisation of *f*-MWNTs did not alter microglial cell migration ability and that
11 exposure to (and uptake of) *f*-MWNTs can actually stimulate migration ability in a chemical
12 functionalisation dependent fashion. In contrast to amination, oxidation of MWNTs seemed to
13 stimulate the migration ability. Similar enhance migration ability in comparison to untreated
14 cells has been recently observed when peritoneal macrophages were exposed to nano
15 graphene oxide flakes and to a higher extend when exposed to nano graphene oxide
16 functionalised with PEG groups ⁴⁰. In this study, enhanced cell migration was attributed to the
17 activation of the cells towards a pro-inflammatory profile upon exposure to graphene oxide
18 flakes. A correlation between enhanced cell migration and activation was not identified in the
19 present study (**Figure 4 and 6**) and would therefore require further investigations to reveal
20 the mechanism at the origin of the enhanced migration ability of cells exposed to oxidised
21 MWNTs.

23 In the present *in vitro* study, no activation of pro-inflammation reaction following the
24 exposure of primary microglial cells to different types of *f*-MWNTs was observed, despite the
25 over-time persistence of the nanomaterials in the cytoplasm, in vesicles (**Figure S4 and**
26 **inset of Figure S5**), and regardless of the type of surface functionalisation. These results
27 are in agreement with other *in vitro* studies performed using the BV2 microglial cell line
28 exposed to non-functionalised MWNTs, in which either cytotoxicity nor induction of
29 expression of major inflammation markers were observed despite CNT internalisation at 48 h
30 ⁴¹. Similarly, no cytotoxicity or pro-inflammatory cytokine release or NO production were
31 observed *in vitro* in the N9 microglial cell line exposed to oxidised MWNTs despite
32 internalisation ¹⁶. On the contrary, using similar MWNTs and similar surface functionalisation
33 (ox-MWNT-NH₃⁺ and MWNT-NH₃⁺), *in vivo* cortical injection of *f*-MWNTs was demonstrated
34 to induce a neuroinflammatory response that was surface chemical functionalisation
35 dependent ¹¹. One month after injection, the degree of oxidation of the *f*-MWNTs was
36 suggested to be a major factor driving the immune tissue response to explain the higher

amount of CD11b positive (microglial cell marker) and GFAP positive (astrocyte marker) cells around the site of injection for animals injected with *f*-MWNTs compared to vehicle injected animals.

In contrast to previous reports, no effects due to interaction of *f*-MWNTs with the cytoskeleton were observed here. It was indeed previously speculate that if CNTs are free in the cytoplasm and interacting with the cytoskeleton, eventually forming bundles with microtubules and microfilaments, they could possibly rigidify cells and consequently reduce both migration and phagocytosis/internalisation abilities of the cells¹⁸. However, both internalisation and migration abilities were here found to be normal in *f*-MWNTs exposed microglial cells, despite a high internalisation of materials illustrated by a black halo around the nucleus region and confirmed to be made of CNTs by Raman spectroscopy. This suggested that *f*-MWNTs were likely not free in the cytoplasm but in vesicular compartments, which is in agreement with the microscopic results presented in **Figure 3a, S2, S3, S4 and S5**. Secondly, this also meant that although their cytoplasm were relatively congested with CNT loading, cells were still elastic enough to perform basic cytoskeleton-mediated functions and especially cell migration, one of the most important cell function *in vivo* for clearance of foreign materials^{17, 37}. This led us to believe that *in vivo* the tested materials would probably undergo brain tissue emigration through microglial cell-mediated drainage finishing in the lymphatic system without harming microglial cells. However, further studies are needed to address that specific hypothesis and to assess whether free *f*-MWNTs (not engulfed in any brain cells) would be able to use the brain solutes elimination system (*i.e.* glymphatic clearance system)^{42, 43}.

Theoretically, CNT-cytoskeleton interactions can also hamper the cytoskeleton dynamics required during cell division, and therefore modify the cell cycle, which would eventually lead to apoptosis. Using the dividing BV2 microglial cell model, MWNTs were indeed shown to induce apoptosis at the relatively low concentration levels of 0.6 and 6 µg/mL (35% and 66%, respectively)¹⁸. Induction of apoptotic pathway was also explained by cell cycle arrest due to CNT interactions with cytoskeleton molecules^{18, 44}. In contrast to these studies, no cell death at 10 µg/mL regardless of the time point considered was observed here. Seclusion of *f*-MWNT loading in vesicular compartments over time is likely the best explanation for the difference between the present results and previous reports. This will require complementary investigations on the subcellular fate of *f*-MWNTs, notably to confirm the nature of the vesicular compartmentalisation of the CNT content and its potential dynamics (endocytosis/exocytosis) over time. Moreover, a complete dose-response relationship correlated with the physicochemical features of the materials should be performed in order to identify the full extent of microglial tolerance to *f*-MWNTs and its limitations.

Degradation and clearance of cell debris and foreign materials are also a fundamental role of microglial cells for maintaining brain homeostasis¹⁷. After stereotactic administration of aminated MWNTs in the cortex of mice, resident microglial cells were reported to internalise and then alter the structural integrity of the injected materials¹². Analysis of tissue sections via Raman spectroscopy and electron microscopy gave evidence of the initiation of a degradative process within two days following the injection. Similarly, it was shown that after 24 h, the N9 microglial cells were able to alter the graphitic structure of oxidised MWNTs¹⁶. And finally, but more importantly using the same model as described in the present study (same primary cells, same *f*-MWNTs, same 10 µg/mL dose), primary microglial cells were showed to degrade *f*-MWNTs over time, irrespective of the surface functionalisation of MWNTs^{14, 15}. These results demonstrate that in addition to conserving their internalisation and migration abilities as reported here, microglial cells had preserved their degradation functions despite their CNT loading. Taken together, these results confirm that microglial cells exposed to 10 µg/mL of the *f*-MWNTs used here are still fully functional cells.

The agglomeration and colloidal properties of the three CNT suspensions were not found to be different (**Figure S1**). Similarly, variations in the surface chemical functionalisation were not found to be deterministic in the impact of the three CNTs apart from the migration ability that seemed to be stimulated by the two oxidised CNTs, in an oxidation degree dependent manner (ox-MWNT >> ox-MWNT-NH₃⁺; **Figure 6**). But alongside surface chemical functionalisation and agglomeration, dimensions (both length and diameter) are important parameters for CNT biological impact⁴⁵⁻⁴⁸. In this respect, the type of CNTs (multi-walled rather than single-walled or double-walled) could also be important as dimensions together with agglomeration will highly differ between CNT types. Therefore, the present results obtained with MWNTs are likely to be different if SWNTs would have been used.

27

Conclusion

In summary, primary microglial cells when exposed to a low concentration of *f*-CNTs maintained their basic resident macrophage functions (phagocytosis/internalisation, migration, degradation) and did not initiate an inflammatory reaction despite a large internalisation of materials in cytoplasm vesicles. Considering the key role of microglial cells within the brain, the demonstration of the relative absence of bioreactivity of *f*-CNTs upon interaction with microglial cells is an important step towards their development as nanovectors for drug delivery purpose in the central nervous system.

Experimental

Materials, MWNTs and preparation of MWNT suspensions for cell exposure. Chemicals and solvents were obtained from Merck / Sigma-Aldrich (UK). Cell culture reagents (PBS, trypsin, foetal bovine serum, DMEM:F12) were purchased from Gibco (ThermoScientific, UK). Multi-walled carbon nanotubes (MWNT) were purchased from Nanostructured and Amorphous Materials Inc. (Houston, TX, USA; Lot # 1240XH, 95%). The different chemical functionalisation of the MWNTs were performed as previously reported^{25, 27, 28, 32}. The amount of COOH (on both ox-MWNT and ox-MWNT-NH₃⁺) after the oxidation reaction has been previously calculated using thermogravimetric analysis, corresponding to 1.7 $\mu\text{mol}\cdot\text{g}^{-1}$ ⁴⁹ while the amount of amino groups measured by Kaiser's test was 170 $\mu\text{mol}\cdot\text{g}^{-1}$ for ox-MWNT-NH₃⁺ and 84 $\mu\text{mol}\cdot\text{g}^{-1}$ for MWNT-NH₃⁺. Preparations of *f*-MWNT dispersion for cell exposure were performed as previously reported¹⁴.

Stability of *f*-MWNT suspension in different media assessed by UV-Vis spectrophotometry.

Suspension of the three *f*-MWNTs originally prepared in 5% dextrose water solution (1 mg/mL) were diluted to 10 $\mu\text{g}/\text{mL}$ in 5% dextrose water solution, cell culture medium (DMEM:F12) supplemented with 12% FBS or cell culture medium (DMEM:F12) without FBS. Absorbance spectra were acquired at 0 h and 24 h after preparing the different 10 $\mu\text{g}/\text{mL}$ suspensions using a Cary 50 Bio UV/Vis spectrophotometer (Varian Inc., Agilent Technologies, UK). For measurement, the prepared suspensions were transferred from a microtube (0.5 mL) to a 1 mL quartz cuvette with 1 cm path length. Measurements were performed at room temperature, using a dual beam mode with baseline correction based on 5% dextrose water solution, cell culture medium with 12% FBS or cell culture medium without FBS, as blank controls to create the baseline.

Primary microglial cell cultures and exposure to functionalised CNT. The preparation of primary microglial cell cultures were performed as previously reported¹⁴ and according to a previously described method, based on mild trypsinisation of mixed glial cell culture⁵⁰ prepared from rat embryonic striatal extracts. Microglial cell cultures were then exposed for 2 h to a non-cytotoxic 10 $\mu\text{g}/\text{mL}$ CNT suspension prepared in serum free cell culture medium (DMEM:F12 medium). After 2 h incubation at 37°C without serum, the cell culture medium was completed with heat inactivated foetal bovine serum (12%). After 24 h incubation at 37°C, the supernatant containing CNTs was removed, cells were washed twice with pre-warmed PBS and then incubated with CNT free complete medium (DMEM:F12 with 12% serum) for 1, 7, 14 and 28 days (recovery periods). CNT-free complete medium was changed every 3 days.

Cell viability assess by modified LDH assay. Cell viability was assessed by a modified LDH assay as previously described³². The Promega Cytotox 96 © Non-radioactive cytotoxicity assay (Promega, UK) was modified to avoid interference between CNTs and LDH. DMSO 10% was used as positive control of the assay. Microglial cell cultures were generated in 24 well plates. At the desired time points (1, 7, 14 and 28 days), cells were lysed with 10 μL of lysis buffer (0.9% triton X100 in distilled water) per 100 μL serum free media and left for 45 min at 37°C. After centrifuging (13000 rpm, 5 min), 50 μL of cell lysate supernatant were mixed with 50 μL of substrate mixture in a new microtiter plate and incubated for 6 min at room temperature. LDH content was determined by reading the absorbance at 490 nm using a multiwell plate reader (Fluostar Omega, BMG Labtech Ltd.). The amount of LDH detected after cell lysis indicated the number of cells that have survived to CNT exposure. The percentage cell survival is expressed as [LDH released from tested cells - Blank (media alone) / LDH released control cells- Blank (media alone)] $\times 100$. Each assay was performed at least 3 times, using 3 wells per conditions at each considered times.

Raman spectroscopy. A Thermo Scientific DXR Raman microscope, equipped with an Olympus microscope (100x objective lens) and a 633 nm LASER (1 mW of LASER power) was used to perform Raman spectroscopy. Collection of Raman spectra was performed as previously described¹⁴.

Immunocytochemistry of microglial cells. Microglial cell cultures were prepared on poly-L-lysine coated plastic tissue-culture treated dishes (35 mm diam.). At the end of exposure, cells were fixed with pre-cooled methanol (-20°C) for 10 min, air-dried and then washed twice with PBS containing 0.1% triton (TPBS, pH 7.4) before staining. Staining with primary antibody was performed after 1 h incubation (i.e. blocking) with 5% of normal goat serum in TPBS. Primary antibodies used were: mouse monoclonal anti-CD11b/c (1:200)(Ox42, Abcam) or mouse monoclonal anti-CD68 (ED1, Abcam, UK); both diluted in 2% BSA in TPBS and incubated for 2 h at room temperature. The secondary antibody (goat anti-mouse Cy3 labelled) was used at the dilution of 1:250 in 2% BSA in TPBS for 1 h at room temperature. After washing, cell preparations were mounted in appropriate medium containing anti-fading agent and DAPI (Vectashield, Vector laboratories LTD). Immunoreactivity was assessed on an inverted epifluorescence microscope (Carl Zeiss Axio Observer A1).

Inflammation markers (NO release) assess by Griess reagent assay. Inflammation reaction was based on the release of NO, a well-known biomarker of inflammation for microglial cells. As for the cytotoxicity assay, microglial cell cultures were generated in 24 well plates. Cells were then exposed to 10 µg/mL of the three different CNT suspensions or to CNT free medium. LPS (E. Coli, O26:B6, 200 ng/mL) was used as positive control for inflammation and was applied 24 h before performing the measurement on cell supernatants. At day 1, 7, 14 and 28 after CNT exposure, the amount of nitrite (stable breakdown products of NO) was measured using the Griess reagent system assay (Promega, UK) following the manufacturer's instructions. In brief, 50 µl of each experimental sample was mixed with 50 µl of the Sulfanilamide Solution. Following 8 min incubation, 50 µl of the NED Solution were added to all wells for 8 min incubation before reading absorbance at 530 nm. The amount of nitrite was determined by comparison to a nitrite standard reference curve, prepared and measured in parallel to each sample measurement.

Phagocytosis/internalisation ability assess by fluorescent polystyrene beads assay. To assess the internalisation ability of CNT-laden microglial cells, we used 0.50µm Fluoresbrite® YellowGreen Microspheres (Polysciences, Inc.) and reduced the cell density of microglial cell cultures (1/2 of usual cell density when preparing mixed glia cell culture). No assay was performed at day 1 after the CNT exposure as some CNTs might still be present in the cell culture medium and compete with the beads for internalisation, despite precautions washings. Assays were therefore performed at day 7 of the recovery period. At this time point, 2 µL of the initial beads suspension was diluted in 2 mL of complete cell culture medium. This bead diluted suspension was used to treat CNT-laden microglial cells prepared and cultured in 35 mm cell culture dishes. The phagocytosis of the fluorescently labelled beads was followed on an inverted epifluorescence microscope tuned on the FITC channel (Carl Zeiss Axio Observer A1) and recorded at 1, 2 and 8 h after bead treatment. Green fluorescence present in each picture was measured using ImageJ image processing and analysis software (NIH, USA).

Migration ability assess by scratch assay. To assess the migration ability of CNT-laden microglial cells, we developed a migration assay based on the well-known wound-healing assay and using the formation of scratches at the bottom of the plastic cell culture support. Using reduced cell density of microglial cell cultures (1/2 of usual mixed glia cell density), we first exposed cells to 10 µg/mL of the different f-MWNT suspension for 24 h for loading purpose. As for the internalisation assay, no migration assay was performed at day 1 after CNT exposure because it was considered too soon after exposure and the presence of free CNTs on the cell culture surface might bias the migration assessment of CNT-laden microglial cells. On day 7, 14 and 28 of the recovery period, we generated straight scratches on the plastic surface of 35 mm cell culture dishes by gently drawing straight lines using a sterile plastic pipette tip. To allow a maximum number of cells to move towards the scratches, migration ability was assessed 72 h after generation of the scratches. Pictures were centred on the scratch and a ROI was drawn around the scratch to count the cells present in the ROI of each picture (ImageJ image processing and analysis software (NIH, USA)) as illustrated in supplementary information **Figure S7**.

Statistics. Results are expressed as mean $\pm$ SD. Statistical differences were analysed using the Student's t test, and $p < 0.05$ (*) were considered as significantly different. Each assay was performed independently at least 3 times. One way ANOVA, Kruskal-Wallis test with Dunn's multiple comparison test was used to analyse the differences in the viability assay $p < 0.05$ (*) were considered as significantly different.

Acknowledgments

This work was supported by the European Commission, under the FP-7 *Marie Curie* actions (Career Development Intra-European Fellowship, PIEF-GA-2010-276051, project NANONEUROHOP). AB wishes to thank the CNRS financial support from PICS (Project for International Scientific Cooperation). MP, as the recipient of the AXA Chair, is grateful to the AXA Research Fund for financial support. MP was also supported by the Spanish Ministry of Economy and Competitiveness MINECO (project CTQ2016-76721-R), by the University of Trieste and by Diputación Foral de Gipuzkoa program Red (101/16).

Figure Legends

Figure 1: Characteristics of the different types of functionalised MWNTs. (a) Chemical structures of the different chemically functionalised MWNTs; (b) Table providing the abbreviations used in the manuscript, the length and the amount of amino groups for the different MWNTs.

Figure 2: Cytotoxic effects of different MWNTs on primary microglial cell cultures evaluated by modified LDH assay. Microglial cell cultures were exposed during 24 h to a unique dose of different MWNTs (10 µg/mL). Following different period of time after the exposure, the LDH assay was performed. No significant effect was observed compared to untreated cells or positive control (DMSO 10%). One way ANOVA, Kruskal-Wallis test with Dunn's multiple comparison test was used to analyse the differences in the viability assay $p < 0.05$ (*) were considered as significantly different.

Figure 3: Imaging of primary microglial cells exposed to different functionalised MWNTs via optical microscopy and confirmation of CNT via Raman spectroscopy. Primary microglial cell cultures were exposed during 24 h to a unique dose of different MWNTs (10 µg/mL) and then leaved for different period of time before imaging. (a) Live imaging (phase contrast) of primary microglial cells at day 1 to 28 for the different exposure. LPS activation (200 ng/mL) was used as a control that induced a change in cell shape (cells flattened and spreaded). No significant difference in terms of shape or number of cells was observed between the non-treated cells and the cells exposed to MWNTs at any time of recovery after the initial exposure. (b) At the end of each desired time of recovery, primary microglial cells were fixed with -20°C pre-cooled methanol and then air dried for at least one hour before further analysis. The Raman analysis of regions randomly selected within the cell boundaries was performed (as presented in **Figure S5c**). Intensities were normalised to the intensity of G band ($I_G = 1$). For all MWNTs and for all time points considered, the typical Raman signature of MWNTs constituted of D, G and D' bands was obtained in regions which appeared black under bright-field microscopy. No Raman signature of CNTs was detected in white regions (nucleus and cell periphery). All scale bars shown are 100 µm.

Figure 4: Inflammation reaction assessed by a measurement of NO release using the Griess reagent assay. Microglial cell cultures were exposed during 24 h to a unique dose of different MWNTs (10 µg/mL). Following different period of time after the exposure, the Griess reagent system assay was performed. No significant effect was observed compared to untreated cells or positive control (LPS 200 ng/mL). Only at day 1 after the initial CNT exposure, the NO release was higher than the untreated cells but minimal compared to the NO release induced by LPS stimulation.

Figure 5: Internalisation ability assessed by a fluorescently labelled microsphere assay. Microglial cell cultures were exposed during 24 h to a unique dose of different MWNTs (10 µg/mL). On day 7 after this initial exposure, microglial cells were treated with fluorescently labelled microspheres (0.5 µm). The internalisation of the fluorescent microspheres was recorded at 2 h (a) and 8 h (b) after the treatment. For the LPS control, untreated cells were exposed to LPS (200 ng/mL) 24 h before the microsphere treatment in order to separate LPS stimulation and microsphere exposure, as for the MWNT exposure. A quantitative measurement of the fluorescence obtained in each pictures was analysed via ImageJ and is presented in (c). For 8 h, results presented have been divided by a factor 10 for all conditions. After 1 and 8 h, no significant internalisation was observed compared to untreated or LPS stimulated cells. At 2 h following microsphere exposure, LPS stimulated cells internalised more microsphere than untreated cells. All scale bars shown are 100 µm.

Figure 6: Migration ability assessed by a scratch assay. Microglial cell cultures were exposed during 24 h to a unique dose of different MWNTs (10 µg/mL). On day 7, 14 and 28 days after this initial exposure, the plastic surface of microglial cell culture was submitted to scratches. Seventy two hours after the scratch, the amount of cells that have migrated toward the scratch was imaged and

counted using ImageJ. For the LPS control, untreated cells were exposed to LPS (200 ng/mL) 24 h before the formation of scratch on the plastic surface. The migration is presented as the percentage compared to untreated cells. LPS treatment was founded to inhibit the migration of microglial cells. The exposure to ox-MWNTs or ox-MWNT-NH₃⁺ were found to significantly stimulate microglial cell migration at both 14 and 28 days, with ox-MWNTs having a higher stimulation than ox-MWNT-NH₃⁺. In contrast, MWNT-NH₃⁺ showed a limited effect on the microglial cell migration ability compared to untreated cells.

Supplementary Figure Legends

Figure S1: Stability of *f*-MWNT suspension in different media assessed by UV-Vis

spectrophotometry. Stability of the three *f*-MWNTs (10 µg/mL) suspended in 5% dextrose water solution (first column), in cell culture medium (DMEM:F12) supplemented with 12% FBS (second column) and in cell culture medium without FBS (third column) are presented at 0 h (red spectra) and 24 h (blue spectra). Absorbance spectra for the suspension media are also displayed (bottom row). Each *f*-MWNT presented specific spectra in each medium. All three *f*-MWNTs showed no difference in absorbance between 0 h (blue spectra) and 24 h (red spectra) when suspended in 5% dextrose or in cell culture medium with 12% FBS, red spectra overlapped with blue spectra. In contrast, a clear decrease in absorbance between 0 h and 24 h was observed for all three *f*-MWNTs when suspended in cell culture medium without FBS. While the blue spectra were specific to each MWNT tested, a similar red spectra below the blue one was observed at 24 h for all three *f*-MWNTs, suggesting instability of suspensions and sedimentation of the three *f*-MWNTs when suspended in this medium.

Figure S2: Optical microscopy of live primary microglial cells after exposure to *f*-MWNTs.

Primary microglia cell cultures were imaged at 1 (a), 7 (b), 14 (c) or 28 (d) days after a 24 h treatment with different *f*-MWNTs (10 µg/mL or 0 µg/mL). The same field of view is displayed after imaging under either phase contrast or bright field illumination. Bright field was used to highlight black regions in cells indicating the presence of carbon nanotubes. For all microglial cell cultures (exposed and non-exposed), cells with either elongated, spread or round shapes were found. All scale bars shown are 100 µm.

Figure S3: Optical microscopy of live primary microglia exposed to *f*-MWNTs (higher magnification).

Primary microglia cell cultures were imaged at 1 (a), 14 (b) or 28 (c) days after a 24 h treatment with different *f*-MWNTs (10 µg/mL) using a 20x objective. Higher magnification at 40x are provided for MWNT-NH₃⁺ after 7 days (d). The same field of view is displayed after imaging under either phase contrast or bright field illumination. Bright field was used to highlight black regions in cells indicating the presence of carbon nanotubes. All scale bars shown are 100 µm.

Figure S4: Immunocytochemistry of *f*-MWNTs laden microglia.

Microglial cells cultures were fixed with pre-cooled methanol, washed and immunostained with mouse anti-rat CD11b/c (microglial cell marker) antibodies in combination with secondary anti-mouse Cy3 antibodies or with mouse anti-rat CD68 (macrophage marker) antibodies in combination with secondary anti-mouse Cy3 antibodies. (a) Merged images of DAPI stain with CD11b/c or CD68 immunoreactivities for the different conditions. (b) Close up on *f*-MWNT laden microglial cells for the 3 different types of MWNTs tested highlighting the vesicular pattern of the MWNT loading inside the cytoplasm of microglial cells. All scale bars shown are 100 µm.

Figure S5: Raman spectroscopy of starting materials before cell exposure and imaging of fixed

cells before Raman analysis. (a) Raman spectroscopy of the three different materials used before cell exposure was performed by depositing a drop of the starting 5% dextrose dispersion of MWNTs (1

Published on 15 June 2017. Downloaded on 15/06/2017 17:31:58.

1 µg MWNT/mL) on a glass slide. (b) Representative pictures of cells exposed to MWNT-NH₃⁺ showing
2 the typical distribution of nanotubes around the nucleus and a granular accumulation pattern
3 suggesting vesicular accumulation, both in phase contrast and bright-field mode. (c) Representative
4 pictures of untreated cells or cells exposed to MWNT-NH₃⁺ in bright-field mode at different levels of
5 magnification. Also presented is a typical map for point and shoot Raman analysis.

6
7 **Figure S6: Internalisation ability assessed by a fluorescently labelled microsphere assay.**

8 Microglial cell cultures were exposed during 24 h to a unique dose of different MWNTs (10 µg/mL). On
9 day 7 after this initial exposure, microglial cells were treated with fluorescently labelled microspheres
10 (0.5 µm). The internalisation of the fluorescent microspheres was recorded at 1 h after the treatment.
11 For the LPS control, untreated cells were exposed to LPS (200 ng/mL) 24 h before the microsphere
12 treatment in order to separate LPS stimulation and microsphere exposure, as for the MWNTs
13 exposure. A quantitative measurement of the fluorescence obtained in each pictures was analysed via
14 ImageJ and is presented in Figure 3c. All scale bars shown are 100 µm.

15
16 **Figure S7: Migration ability assessed by a scratch assay.** Microglial cell cultures were exposed
17 during 24 h to a unique dose of different MWNTs (10 µg/mL). On day 7, 14 and 28 days after this
18 initial exposure, the plastic surface of cell culture was submitted to scratches (see schemes). After the
19 scratch, cells that have migrated towards the scratch was imaged (representative pictures, 72 h after
20 the scratch) and counted using ImageJ in the ROI highlighted in yellow. For the LPS control, untreated
21 cells were exposed to LPS (200 ng/mL) 24 h before the formation of scratch on the plastic surface. All
22 scale bars shown are 100 µm.

References

1. P. Blasi, S. Giovagnoli, A. Schoubben, M. Ricci and C. Rossi, *Adv Drug Deliv Rev*, 2007, **59**, 454-477.
2. T. Patel, J. Zhou, J. M. Piepmeier and W. M. Saltzman, *Adv Drug Deliv Rev*, 2012, **64**, 701-705.
3. S. B. Pehlivan, *Pharm Res*, 2013, **30**, 2499-2511.
4. J. T. Wang and K. T. Al-Jamal, *Nanomedicine (Lond)*, 2015, **10**, 2639-2642.
5. K. Kostarelos, L. Lacerda, G. Pastorin, W. Wu, S. Wieckowski, J. Luangsivilay, S. Godefroy, D. Pantarotto, J. P. Briand, S. Muller, M. Prato and A. Bianco, *Nat Nanotechnol*, 2007, **2**, 108-113.
6. Z. Yang, Y. Zhang, Y. Yang, L. Sun, D. Han, H. Li and C. Wang, *Nanomedicine-Uk*, 2010, **6**, 427-441.
7. K. T. Al-Jamal, L. Gherardini, G. Bardi, A. Nunes, C. Guo, C. Bussy, M. A. Herrero, A. Bianco, M. Prato, K. Kostarelos and T. Pizzorusso, *Proc Natl Acad Sci U S A*, 2011, **108**, 10952-10957.
8. H. J. Lee, J. Park, O. J. Yoon, H. W. Kim, D. Y. Lee, D. H. Kim, W. B. Lee, N. E. Lee, J. V. Bonventre and S. S. Kim, *Nature Nanotechnology*, 2011, **6**, 120-124.
9. M. K. Gottipati, E. Bekyarova, R. C. Haddon and V. Parpura, *Amino acids*, 2015, **47**, 1379-1388.
10. G. Tosi, A. Vilella, R. Chhabra, M. J. Schmeisser, T. M. Boeckers, B. Ruozzi, M. A. Vandelli, F. Forni, M. Zoli and A. M. Grabrucker, *J Control Release*, 2014, **177**, 96-107.
11. G. Bardi, A. Nunes, L. Gherardini, K. Bates, K. T. Al-Jamal, C. Gaillard, M. Prato, A. Bianco, T. Pizzorusso and K. Kostarelos, *PLoS One*, 2013, **8**, e80964.
12. A. Nunes, C. Bussy, L. Gherardini, M. Meneghetti, M. A. Herrero, A. Bianco, M. Prato, T. Pizzorusso, K. T. Al-Jamal and K. Kostarelos, *Nanomedicine (Lond)*, 2012, **7**, 1485-1494.
13. C. Bussy, K. T. Al-Jamal, J. Boczkowski, S. Lanone, M. Prato, A. Bianco and K. Kostarelos, *ACS Nano*, 2015, **9**, 7815-7830.
14. C. Bussy, C. Hadad, M. Prato, A. Bianco and K. Kostarelos, *Nanoscale*, 2016, **8**, 590-601.
15. J. Russier, L. Oudjedi, M. Piponnier, C. Bussy, M. Prato, K. Kostarelos, B. Lounis, A. Bianco and L. Cognet, *Nanoscale*, 2017.
16. A. E. Goode, D. A. Gonzalez Carter, M. Motskin, I. S. Pienaar, S. Chen, S. Hu, P. Ruenraroengsak, M. P. Ryan, M. S. Shaffer, D. T. Dexter and A. E. Porter, *Biomaterials*, 2015, **70**, 57-70.
17. L. C. Davies, S. J. Jenkins, J. E. Allen and P. R. Taylor, *Nat Immunol*, 2013, **14**, 986-995.
18. J. C. Villegas, L. Alvarez-Montes, L. Rodriguez-Fernandez, J. Gonzalez, R. Valiente and M. L. Fanarraga, *Adv Healthc Mater*, 2014, **3**, 424-432.
19. N. V. Konduru, Y. Y. Tyurina, W. Feng, L. V. Basova, N. A. Belikova, H. Bayir, K. Clark, M. Rubin, D. Stolz, H. Vallhov, A. Scheynius, E. Witas, B. Fadeel, P. D. Kichambare, A. Star, E. R. Kisin, A. R. Murray, A. A. Shvedova and V. E. Kagan, *Plos One*, 2009, **4**, e4398.
20. C. Bussy, M. Pinault, J. Cambedouzou, M. J. Landry, P. Jegou, M. Mayne-L'hermite, P. Launois, J. Boczkowski and S. Lanone, *Part Fibre Toxicol*, 2012, **9**, 46.
21. J. T. Wang, N. Rubio, H. Kafa, E. Venturelli, C. Fabbro, C. Menard-Moyon, T. Da Ros, J. K. Sosabowski, A. D. Lawson, M. K. Robinson, M. Prato, A. Bianco, F. Festy, J. E. Preston, K. Kostarelos and K. T. Al-Jamal, *J Control Release*, 2016, **224**, 22-32.

- 1 22. H. Ali-Boucetta, K. T. Al-Jamal, D. McCarthy, M. Prato, A. Bianco and K.
2 Kostarelos, *Chem Commun (Camb)*, 2008, 459-461.
- 3 23. R. Singh, D. Pantarotto, D. McCarthy, O. Chaloin, J. Hoebeke, C. D. Partidos, J. P.
4 Briand, M. Prato, A. Bianco and K. Kostarelos, *J Am Chem Soc*, 2005, **127**, 4388-
5 4396.
- 6 24. A. Servant, I. Jacobs, C. Bussy, C. Fabbro, T. da Ros, E. Pach, B. Ballesteros, M.
7 Prato, K. Nicolay and K. Kostarelos, *Carbon*, 2016, **97**, 126-133.
- 8 25. S. P. Li, W. Wu, S. Campidelli, V. Sarnatskaia, M. Prato, A. Tridon, A. Nikolaev, V.
9 Nikolaev, A. Bianco and E. Snezhkova, *Carbon*, 2008, **46**, 1091-1095.
- 10 26. C. Gaillard, G. Cellot, S. P. Li, F. M. Toma, H. Dumortier, G. Spalluto, B. Cacciari,
11 M. Prato, L. Ballerini and A. Bianco, *Advanced Materials*, 2009, **21**, 2903-+.
- 12 27. V. Georgakilas, K. Kordatos, M. Prato, D. M. Guldi, M. Holzinger and A. Hirsch,
13 *Journal of the American Chemical Society*, 2002, **124**, 760-761.
- 14 28. V. Georgakilas, N. Tagmatarchis, D. Pantarotto, A. Bianco, J. P. Briand and M. Prato,
15 *Chem Commun (Camb)*, 2002, 3050-3051.
- 16 29. P. Wick, P. Manser, L. K. Limbach, U. Dettlaff-Weglikowska, F. Krumeich, S. Roth,
17 W. J. Stark and A. Bruinink, *Toxicol Lett*, 2007, **168**, 121-131.
- 18 30. L. Dong, C. M. Witkowski, M. M. Craig, M. M. Greenwade and K. L. Joseph,
19 *Nanoscale Res Lett*, 2009, **4**, 1517-1523.
- 20 31. E. A. Salegio, H. Streeter, N. Dube, P. Hadaczek, L. Samaranch, A. P. Kells, W. San
21 Sebastian, Y. Zhai, J. Bringas, T. Xu, J. Forsayeth and K. S. Bankiewicz, *Frontiers in*
22 *neuroanatomy*, 2014, **8**, 9.
- 23 32. H. Ali-Boucetta, K. T. Al-Jamal, K. H. Muller, S. Li, A. E. Porter, A. Eddaoudi, M.
24 Prato, A. Bianco and K. Kostarelos, *Small*, 2011, **7**, 3230-3238.
- 25 33. S. Ganter, H. Northoff, D. Mannel and P. J. Gebickeharter, *J Neurosci Res*, 1992, **33**,
26 218-230.
- 27 34. M. P. Ramprasad, V. Terpstra, N. Kondratenko, O. Quehenberger and D. Steinberg,
28 *Proc Natl Acad Sci U S A*, 1996, **93**, 14833-14838.
- 29 35. M. S. Dresselhaus, G. Dresselhaus, R. Saito and A. Jorio, *Physics Reports*, 2005, **409**,
30 47-99.
- 31 36. M. S. Dresselhaus, A. Jorio, M. Hofmann, G. Dresselhaus and R. Saito, *Nano Lett*,
32 2010, **10**, 751-758.
- 33 37. K. Saijo and C. K. Glass, *Nature reviews. Immunology*, 2011, **11**, 775-787.
- 34 38. T. Tajima, T. Murata, K. Aritake, Y. Urade, H. Hirai, M. Nakamura, H. Ozaki and M.
35 Hori, *J Pharmacol Exp Ther*, 2008, **326**, 493-501.
- 36 39. H. Scheiblich, F. Roloff, V. Singh, M. Stangel, M. Stern and G. Bicker, *Brain Res*,
37 2014, **1564**, 9-21.
- 38 40. N. Luo, J. K. Weber, S. Wang, B. Luan, H. Yue, X. Xi, J. Du, Z. Yang, W. Wei, R.
39 Zhou and G. Ma, *Nat Commun*, 2017, **8**, 14537.
- 40 41. B. Kateb, M. Van Handel, L. Zhang, M. J. Bronikowski, H. Manohara and B. Badie,
41 *Neuroimage*, 2007, **37 Suppl 1**, S9-17.
- 42 42. L. Xie, H. Kang, Q. Xu, M. J. Chen, Y. Liao, M. Thiyagarajan, J. O'Donnell, D. J.
43 Christensen, C. Nicholson, J. J. Iliff, T. Takano, R. Deane and M. Nedergaard,
44 *Science*, 2013, **342**, 373-377.
- 45 43. J. J. Iliff, M. Wang, Y. Liao, B. A. Plogg, W. Peng, G. A. Gundersen, H. Benveniste,
46 G. E. Vates, R. Deane, S. A. Goldman, E. A. Nagelhus and M. Nedergaard, *Science*
47 *translational medicine*, 2012, **4**, 147ra111.
- 48 44. L. Rodriguez-Fernandez, R. Valiente, J. Gonzalez, J. C. Villegas and M. L. Fanarraga,
49 *ACS Nano*, 2012, **6**, 6614-6625.

- 1 45. S. Lanone, P. Andujar, A. Kermanizadeh and J. Boczkowski, *Adv Drug Deliv Rev*,
2 2013, **65**, 2063-2069.
- 3 46. H. Nagai, Y. Okazaki, S. H. Chew, N. Misawa, Y. Yamashita, S. Akatsuka, T.
4 Ishihara, K. Yamashita, Y. Yoshikawa, H. Yasui, L. Jiang, H. Ohara, T. Takahashi, G.
5 Ichihara, K. Kostarelos, Y. Miyata, H. Shinohara and S. Toyokuni, *Proc Natl Acad Sci*
6 *U S A*, 2011, **108**, E1330-1338.
- 7 47. C. A. Poland, R. Duffin, I. Kinloch, A. Maynard, W. A. Wallace, A. Seaton, V. Stone,
8 S. Brown, W. Macnee and K. Donaldson, *Nat Nanotechnol*, 2008, **3**, 423-428.
- 9 48. C. Muhlfeld, C. A. Poland, R. Duffin, C. Brandenberger, F. A. Murphy, B. Rothen-
10 Rutishauser, P. Gehr and K. Donaldson, *Nanotoxicology*, 2012, **6**, 867-879.
- 11 49. J. Russier, C. Menard-Moyon, E. Venturelli, E. Gravel, G. Marcolongo, M.
12 Meneghetti, E. Doris and A. Bianco, *Nanoscale*, 2011, **3**, 893-896.
- 13 50. J. Saura, J. M. Tusell and J. Serratosa, *Glia*, 2003, **44**, 183-189.
- 14

b

Short name	Nature of MWCNT	Amount of amino groups (Kaiser test)	Length (TEM)
ox-MWNT	carboxylated	-	200 – 300 nm
ox-MWNT-NH ₃ ⁺	carboxylated and aminated (cycloaddition)	170 μmol/g	200 – 300 nm
MWNT-NH ₃ ⁺	aminated (cycloaddition)	84 μmol/g	0.5 – 2 μm

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6