Radiative light dark matter

DOI:

[10.1103/PhysRevD.95.115037](https://doi.org/10.1103/PhysRevD.95.115037)

Document Version

Final published version

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Dedes, A., Karamitros, D., & Pilaftsis, A. (2017). Radiative light dark matter. *Physical Review D - Particles, Fields, Gravitation and Cosmology*, 95, [115037]. <https://doi.org/10.1103/PhysRevD.95.115037>

Published in:

Physical Review D - Particles, Fields, Gravitation and Cosmology

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Radiative light dark matterA. Dedes,^{1,*} D. Karamitros,^{1,†} and A. Pilaftsis^{2,‡}¹*Department of Physics, Division of Theoretical Physics, University of Ioannina, Ioannina, GR 45110, Greece*²*Consortium for Fundamental Physics, School of Physics and Astronomy, University of Manchester, Manchester M13 9PL, United Kingdom*

(Received 11 April 2017; published 29 June 2017)

We present a Peccei-Quinn (PQ)-symmetric two-Higgs doublet model that naturally predicts a fermionic singlet dark matter in the mass range 10 keV–1 GeV. The origin of the smallness of the mass of this light singlet fermion arises predominantly at the one-loop level, upon soft or spontaneous breakdown of the PQ symmetry via a complex scalar field in a fashion similar to the so-called Dine-Fischler-Sredniki-Zhitnitsky axion model. The mass generation of this fermionic radiative light dark matter (RLDM) requires the existence of two heavy vectorlike SU(2) isodoublets, which are not charged under the PQ symmetry. We show how the RLDM can be produced via the freeze-in mechanism, thus accounting for the missing matter in the Universe. Finally, we briefly discuss possible theoretical and phenomenological implications of the RLDM model for the strong *CP* problem and the CERN Large Hadron Collider (LHC).

DOI: 10.1103/PhysRevD.95.115037

I. INTRODUCTION

Ongoing searches for the elusive missing matter component of the Universe, the so-called dark matter (DM), have offered no conclusive evidence so far. From analyses of the cosmic microwave background power spectrum and from pertinent astronomical studies, we now know that about one quarter of the energy budget of our Universe should be in the form of DM, and so many candidate theories have been put forward to address this well-known DM problem [1]. Among the suggested scenarios, those predicting weakly interactive massive particles (WIMPs) constitute one class of popular models that may not only account for the DM itself, but also leave their footprints in low-energy experiments, or even at high-energy colliders, such as the LHC [2]. In particular, for WIMPs near the electroweak scale, the WIMP-nucleon scattering cross section is estimated to be somewhat below 10^{-46} cm² as measured by LUX [3].

Projected experiments that lie not very far ahead in future will be capable of reaching sensitivity in the ballpark $10^{-47} - 10^{-48}$ cm² [4], and so they will be getting closer to the neutrino-nucleon background cross section, the infamous “neutrino floor,” where disentangling neutrino signals from those of WIMPs will become almost an impossible task [5]. Therefore, DM models have to be constructed (or revisited) to avoid such severe constraints, e.g. by contemplating scenarios that either sufficiently suppress the WIMP-nucleon interaction, or move the DM mass to the sub-GeV or ultra-TeV region.

Several models have been proposed featuring a light DM in the mass range $\mathcal{O}(\text{keV}) - \mathcal{O}(\text{GeV})$, such as sterile

neutrino DM [6–10], light scalar DM [11] and milli-charged DM [12], including their possible implications for future DM searches [13,14]. However, one central problem of such models is the actual origin of the small mass for the light DM, which could be more than 6 orders of magnitude below the electroweak scale.

In this paper we address this mass hierarchy problem, by presenting a new radiative mechanism that can predominantly account for the smallness in mass for the light DM. The so-generated radiative light dark matter (RLDM) is a fermionic singlet *S* and can naturally acquire a mass in the desired range: 10 keV–1 GeV. A minimal realization of this radiative mechanism requires the extension of the Standard Model (SM) by one extra scalar doublet, resulting in a Peccei-Quinn (PQ)-symmetric two-Higgs doublet model [15,16], augmented by two fermionic heavy vectorlike SU(2) isodoublets *D*₁ and *D*₂, which are not charged under the PQ symmetry. The mass of the RLDM is predominantly generated at the one-loop level, upon soft or spontaneous breakdown of the PQ symmetry via a complex scalar field, e.g. Σ , in close analogy to the so-called Dine-Fischler-Sredniki-Zhitnitsky (DFSZ) axion model that addresses the strong *CP* problem [17,18].

We analyse the production mechanisms of the RLDM in the early universe, and show that it can account for its missing matter component via the so-called freeze-in mechanism [19]. In fact, we illustrate how the freeze-in mechanism remains effective in the RLDM model, without the need to resort to suppressed Yukawa couplings. In this context, we investigate two possible scenarios of both theoretical and phenomenological interest. In the first scenario, we consider the breaking of the PQ scale f_{PQ} to be comparable to the one required for the DFSZ model to solve the strong *CP* problem, i.e. $f_{\text{PQ}} \sim 10^9$ GeV.

*adedes@cc.uoi.gr

†dkaramit@cc.uoi.gr

‡Apostolos.Pilaftsis@manchester.ac.uk

We find that such PQ scale can exist within this realization, provided an appropriate isodoublet mass M_D and reheating temperature T_{RH} is considered. In the second scenario, we relax the constraint of the strong CP problem on f_{PQ} , and investigate its possible lower limit, with the only requirement that T_{RH} be larger than the critical temperature T_C of the SM electroweak phase transition, thus allowing for the $B + L$ -violating sphaleron processes to be in thermal equilibrium. This requirement is introduced here, so as to leave open the possibility of explaining the cosmological baryon-to-photon ratio η_B via low-scale baryogenesis mechanisms, such as electroweak baryogenesis [20,21] and resonant leptogenesis [22–25]. In this second scenario, we find that the heavy Higgs bosons of the two-Higgs doublet model (2HDM) may have masses as low as a few TeV, which are well within reach of the LHC.

The layout of the paper is as follows. In Sec. II, we first introduce the PQ-symmetric 2HDM, augmented with a singlet fermion S and a fermionic pair of vectorlike doublets $D_{1,2}$. Then, we describe the radiative mechanism for the RLDM, once the PQ symmetry is broken softly, and show that a radiative mass in the range 10 keV–1 GeV can be naturally generated. In Sec. III, we outline the relevant Boltzmann equation for computing the relic abundance of the RLDM. Utilizing the freeze-in mechanism, we present in Sec. IV numerical estimates for the allowed parameter space of our RLDM model. Based on these results, we explore the possibility whether our model can account for the strong CP problem within a scenario similar to the DFSZ axion model. Moreover, we investigate whether an absolute lower limit exists for the heavy Higgs-boson masses in our effective 2HDM. Indeed, we find that our RLDM model may allow for heavy Higgs bosons at the TeV scale, whose existence can be probed at the LHC. Finally, Sec. V summarizes our conclusions and outlines possible new directions for further research.

II. RADIATIVE MECHANISM

In this section we present a minimal extension of the SM, in which the small mass of the light DM, in the region 10 keV–1 GeV, can have a radiative origin, generated at the one-loop level. This radiative mechanism is minimally realized within the context of a constrained 2HDM obeying a Peccei-Quinn symmetry. In addition, the model under study contains a singlet fermion S charged under the PQ symmetry and a fermionic pair of massive isodoublets $D_{1,2}$ with zero PQ charges. Finally, we delineate the parameter space for which a viable scenario of radiative light dark matter can be obtained consistent with the observed relic abundance.

A. The model

In the 2HDM under consideration, we impose a global PQ symmetry $U(1)_{\text{PQ}}$, which forbids the appearance of a bare mass term for the singlet fermion S at the tree level.

This PQ symmetry will be broken softly or spontaneously which in turn triggers a radiative mass for S at the one-loop level. The fermion S is stable and receives naturally a small sub-GeV mass, leading to a RLDM scenario. On the other hand, we note that a candidate for a light DM would probably be relativistic at its freeze-out, resulting in an extremely large relic abundance (similar to [26]) for the allowed range of DM masses that are larger than about 3 keV, e.g. see [27,28]. Therefore, the DM should be produced out of thermal equilibrium in the early universe. The mechanism that we will be utilizing here is the so-called freeze-in mechanism [19], which assumes that the DM particles were absent initially and are produced only later from the plasma.

The relevant Yukawa and potential terms of our model are given by

$$-\mathcal{L}^Y = Y_1 \epsilon^{ab} \Phi_{1a} D_{1b} S + Y_2 \Phi_2^{\dagger a} D_{2a} S + M_D \epsilon^{ab} D_{1a} D_{2b} + \text{H.c.}, \quad (1)$$

$$V(\Phi_1, \Phi_2) = m_{11}^2 \Phi_1^{\dagger a} \Phi_{1a} + m_{22}^2 \Phi_2^{\dagger a} \Phi_{2a} - m_{12}^2 (\Phi_1^{\dagger a} \Phi_{2a} + \text{H.c.}) + \frac{\lambda_1}{2} (\Phi_1^{\dagger a} \Phi_{1a})^2 + \frac{\lambda_2}{2} (\Phi_2^{\dagger a} \Phi_{2a})^2 + \lambda_3 \Phi_1^{\dagger a} \Phi_{1a} \Phi_2^{\dagger b} \Phi_{2b} + \lambda_4 |\Phi_1^{\dagger a} \Phi_{2a}|^2, \quad (2)$$

where $a, b = 1, 2$ are $SU(2)_L$ -group indices (with $\epsilon^{12} = -\epsilon^{21} = +1$), S is a Weyl-fermion SM singlet, $D_{1,2}$ are two Weyl-fermion $SU(2)_L$ doublets, and $\Phi_{1,2}$ are two scalar $SU(2)_L$ doublets. A complete list of the PQ and hypercharge quantum numbers of the aforementioned particles is given in Table I, including a $\mathbb{Z}_2$ parity which excludes the mixing of dark-sector particles with those of the SM. For simplicity, we assume that the new dark-sector interactions are CP invariant and so take their respective couplings to be real in the physical mass basis.

As can be seen from (2), we have assumed that the PQ symmetry is broken by the lowest dimensionally possible mass operator in the scalar potential V , namely by allowing only the dimension-2 mixing term m_{12}^2 between Φ_1 and Φ_2 . This dimension-2 operator breaks softly the $U(1)_{\text{PQ}}$ symmetry in the potential, but could result from spontaneous

TABLE I. Quantum number assignments of particles pertinent to the RLDM model.

	$SU(2)_L$	$U(1)_Y$	$U(1)_{\text{PQ}}$	$\mathbb{Z}_2$
S	1	0	−1	Odd
D_1	2	−1	0	Odd
D_2	2	1	0	Odd
Φ_1	2	1	1	Even
Φ_2	2	1	−1	Even

breaking of the $U(1)_{\text{PQ}}$ by a scalar Σ , which acquires a vacuum expectation value (VEV) $\langle \Sigma \rangle \equiv f_{\text{PQ}} \sim m_{12}$ (see Sec. IV). If the PQ-breaking scale f_{PQ} is high enough, one may neglect, to a good approximation, the potential quartic couplings $\lambda_{1,2,3,4}$, as they do not affect much the radiative mass mechanism and the DM production rates which we will be discussing in the next section.

The mass parameters m_{11}^2 and m_{22}^2 of the scalar potential V in Eq. (2) may be eliminated in favor of the VEVs $v_{1,2}$ of the Higgs doublets $\Phi_{1,2}$, by virtue of the minimization conditions on V (for a review on 2HDMs, see [29]). These VEVs are related to the SM Higgs VEV v , through: $v^2 = v_1^2 + v_2^2$. In the kinematic region where $m_{12}^2 \gg v^2$, the mass parameters m_{11}^2 and m_{22}^2 are approximately given by

$$m_{11}^2 \approx m_{12}^2 t_\beta + \mathcal{O}(v^2), \quad (3)$$

$$m_{22}^2 \approx m_{12}^2 t_\beta^{-1} + \mathcal{O}(v^2), \quad (4)$$

where $t_\beta \equiv \tan \beta = v_2/v_1$.

B. One-loop radiative mass

Having introduced the minimal model under investigation, we can now discuss the radiative mechanism responsible for the generation of a mass of dimension 3 for the singlet fermion S . We assume that $m_{12} \gtrsim 1$ TeV, such that the main contribution to the mass of the S particle comes from the diagram shown in Fig. 1. In addition, there will be a tree-level mass M_S^{tree} generated after the SM electroweak phase transition, given by $M_S^{\text{tree}} \approx Y_1 Y_2 v^2 / M_D$. Under the assumption that M_D is very large, i.e. $M_D \gg v$, the tree-level contribution turns out to be subdominant compared to the radiatively induced mass M_S^{rad} , and hence it can be ignored for most of the parameter space. We will return to this point at the end of this section.

After evaluating the relevant one-loop self-energy graph shown in Fig. 1 at zero external momentum ($p \rightarrow 0$), we obtain

$$M_S^{\text{rad}} = -2Y_1 Y_2 M_D m_{12}^2 I(M_D, m_{11}, m_{22}), \quad (5)$$

where

FIG. 1. One-loop diagram responsible for the mass generation of the singlet fermion S .

$$I(M_D, m_{11}, m_{22}) = \int \frac{d^4 q}{(2\pi)^4} \frac{1}{(q^2 - M_D^2)(q^2 - m_{11}^2)(q^2 - m_{22}^2)}. \quad (6)$$

Employing the approximate relations given in (3) and (5), the one-loop radiative mass of S is finite and may conveniently be expressed as follows:

$$M_S^{\text{rad}} = \frac{2y^2}{(4\pi)^2} \frac{M_D}{t_\beta - t_\beta^{-1}} \left[\frac{t_\beta \ln(t_\beta/r^2)}{t_\beta - r^2} - \frac{t_\beta^{-1} \ln(t_\beta^{-1}/r^2)}{t_\beta^{-1} - r^2} \right], \quad (7)$$

with $y^2 \equiv Y_1 Y_2$ and $r \equiv M_D/m_{12}$. Observe that the interchange $t_\beta \leftrightarrow t_\beta^{-1}$ leaves M_S^{rad} unchanged. Assuming that $t_\beta = 1$ for different kinematic regimes of the ratio r , the following simplified forms for M_S^{rad} are obtained:

$$M_S^{\text{rad}} \approx \frac{2y^2}{(4\pi)^2} M_D \quad \text{for } r \ll 1, \quad (8)$$

$$M_S^{\text{rad}} \approx \frac{y^2}{(4\pi)^2} M_D \quad \text{for } r \sim 1, \quad (9)$$

$$M_S^{\text{rad}} \approx \frac{2y^2}{(4\pi)^2} \frac{M_D \ln r^2}{r^2} \quad \text{for } r \gg 1. \quad (10)$$

Note that for $M_D \gg m_{12}$ (corresponding to $r \gg 1$), the radiative mass M_S^{rad} of the singlet fermion S is suppressed by the square of the hierarchy factor r . The latter allows for scenarios, for which the Yukawa couplings are of order 1, i.e. $y^2 = Y_1 Y_2 = \mathcal{O}(1)$, for $10 \text{ keV} \leq M_S^{\text{rad}} \leq 1 \text{ GeV}$. On the other hand, for $r \sim 1$ and $r \ll 1$, one needs either a low M_D of order TeV and $y \approx 0.1$, or $M_D \approx 10^8 - 10^9 \text{ GeV}$ and $y \approx 10^{-3} - 10^{-4}$.

In Fig. 2, we display the values of the coupling parameter $y = \sqrt{Y_1 Y_2}$, as a function of M_D , which yield a radiatively induced mass M_S^{rad} for the singlet fermion S in the region

FIG. 2. Predicted values for M_D versus $y = \sqrt{Y_1 Y_2}$ as obtained from (7), for $M_S \approx M_S^{\text{rad}}$ ranging from 10 keV to 1 GeV, after setting $t_\beta = 1$ and $r = 10^{-2}$.

$10 \text{ keV} \leq M_S^{\text{rad}} \leq 1 \text{ GeV}$, for $t_\beta = 1$ and $r = 10^{-2}$. In particular, we see that for every set of M_S^{rad}, M_D, r , there is an acceptable range of perturbative values for y . However, if $r \gg 1$, the desirable value of y may exceed 10 according to (10), and our perturbative results do no longer apply. Such nonperturbative values of y are excluded from our numerical estimates for the determination of the relic abundance of S which we perform in the next section.

In a similar context, we note that a large mass for m_{11}, m_{12}, m_{22} and M_D might seem to be a huge fine-tuning for generating a light sub-GeV radiative mass for S . However, we may easily convince ourselves that this is not the case. The absence of fine-tuning can be seen in an easier way, if we rotate from the general weak basis spanned by Φ_1 and Φ_2 to the so-called *Higgs basis* [29,30], H_1 and H_2 , where H_1 contains the SM VEV v and H_2 has exactly no VEV. Note that in the Higgs basis, the assignment of the PQ charges to the fields H_1 and H_2 is not canonical. Moreover, in this rotated Higgs basis, one has that the new Higgs-mass parameters obey the relation: $\tilde{m}_{22}^2 \gg \tilde{m}_{11}^2, \tilde{m}_{12}^2$. In addition, the analogue of the diagram in Fig. 1 is now represented by a set of two self-energy graphs, where the fields H_1 and H_2 are circulating in the loop. The ultraviolet (UV) infinities cancel, after the contributions from these two diagrams are added. For $t_\beta = 1$ and $r = 1$, we then obtain the same result as the one stated in (9). Hence, we observe that a small mass for the singlet fermion S arises naturally in a SM+S effective field theory. This effective field theory results from integrating out the heavy $D_{1,2}$ and H_2 fields from (1) in the Higgs basis.

Besides the radiative mass M_S^{rad} of S which violates the PQ symmetry by two units (cf. Table I), there will be a tree-level contribution to the mass of S after the SM electroweak phase transition. For most r values of interest here, the relative size of the two contributions can naively be estimated to be

$$\frac{M_S^{\text{tree}}}{M_S^{\text{rad}}} \sim \frac{8\pi^2 v^2}{M_D^2}. \quad (11)$$

Thus, for $M_D \gg \sqrt{8\pi}v \approx 2.2 \text{ TeV}$, the tree-level contribution can be safely ignored. In our numerical estimates, the tree-level mass term M_S^{tree} is always less than 10% of the radiative mass term M_S^{rad} . Hence, the total mass M_S of the stable fermion S is given predominantly by the radiative mass term, implying that $M_S \approx M_S^{\text{rad}}$ to a very good approximation.

We conclude this section by commenting on the possibility of considering a radiative model alternative to the one discussed here. For instance, one may envisage a scenario that instead of the single S , one of the neutral components of the doublets $D_{1,2}$ becomes the RLDM. In this case, however, the charged component $D^\pm$ from $D_{1,2}$ will be almost degenerate with the light sub-GeV DM

particle, which is excluded experimentally. The general SM + $D_{1,2}$ effective theory has been studied in [31].

III. DARK MATTER ABUNDANCE

In this section we first describe the relevant effective Lagrangian that governs the production of the stable fermions S in the early universe. We then solve numerically the Boltzmann equation that determines the yield $Y_S \equiv n_S/s$ of these fermions S , where n_S is the number density of S particles and s is the entropy density of the plasma. Having thus estimated the value of Y_S , we can then use it to deduce the respective relic abundance $\Omega_S h^2$ of the S particles in the present epoch. Finally, we present approximate analytic results for $\Omega_S h^2$ and compare these with the observationally favored value: $\Omega_{\text{DM}} h^2 \approx 0.12$.

As mentioned in the previous section, the stable fermions S will play the role of the DM, which are produced via the freeze-in mechanism [19]. The key assumption is that the DM fermions S were absent (i.e. their number density was suppressed) in the early universe and were produced later from annihilations and decays of plasma particles, e.g. from $\Phi_{1,2}$ and $D_{1,2}$, according to the model discussed in Sec. II. Furthermore, we will assume that $D_{1,2}$ were also absent in the early universe, so as to avoid overclosure of the Universe, unless the Yukawa couplings $Y_{1,2}$ are taken to be extremely suppressed, such that decays of the sort $D_1^0 \rightarrow hS$ are made slow and inefficient. The latter results in a contrived scenario, in which obtaining a viable DM parameter space requires a good degree of fine-tuning. In order for the $\text{SU}(2)_L$ -doublet fermions $D_{1,2}$ to be absent, we take their bare mass M_D to be above the reheating temperature T_{RH} of the Universe. This simplifies considerably our analysis, as the heavy fermions $D_{1,2}$ can be integrated out.

The effective Lagrangian that determines the production rate of S particles after reheating is given by

$$-\mathcal{L}_{\text{eff}}^{d=5} = \frac{1}{2\Lambda} (\Phi_1^\dagger \Phi_1 + \tilde{a} \Phi_2^\dagger \Phi_2 + \tilde{b} \Phi_1^\dagger \Phi_2 + \tilde{c} \Phi_2^\dagger \Phi_1) S S + \text{H.c.}, \quad (12)$$

where $\tilde{a}$, $\tilde{b}$ and $\tilde{c}$ denote the Wilson coefficients of the dimension-5 operators. The calculation of the relic abundance is not straightforward in this basis, since $\Phi_{1,2}$ mix and the identification of the physical fields is obscured, especially after spontaneous symmetry breaking where further mixing between the scalar fields is introduced. Therefore, according to our discussion at the end of Sec. II B, it would be more convenient to rotate the scalars to the so-called Higgs basis [29], where only one doublet H_1 develops a VEV and is identified with the SM Higgs doublet. To further simplify calculations, and without much loss of generality, we assume that the Higgs basis is also the mass eigenstate basis. This assumption is well justified for

relatively large values of m_{12} , as it leads to the so-called *alignment limit* of the 2HDM [32–36], which is favored in the light of global analyses of experimental constraints [37,38]. In the Higgs basis, the dimension-5 effective Lagrangian reads

$$-\mathcal{L}_{\text{eff}}^{d=5} = \frac{y^2}{M_D} \frac{t_\beta}{1+t_\beta^2} (H_1^\dagger H_1 - H_2^\dagger H_2 - t_\beta H_1^\dagger H_2 + t_\beta^{-1} H_2^\dagger H_1) SS + \text{H.c.}, \quad (13)$$

where H_1 is the SM Higgs doublet and H_2 is the heavy scalar doublet with $\langle H_2 \rangle = 0$.

A. Boltzmann equation for Y_S

In order to determine the relic abundance of S particles, we need to solve the Boltzmann equation for their yield Y_S . Since we assume that the singlets S remained out of equilibrium throughout the history of the Universe (at least up to the phase of reheating), our only concern will then be their production. The main production channels, depending on the plasma temperature T , are the following:

$$\begin{aligned} H_1^\dagger H_1, H_1^\dagger H_2, H_2^\dagger H_1 &\rightarrow SS \quad \text{for } T_C \leq T < T_{\text{RH}}, \\ H_2^\dagger H_2 &\rightarrow SS \quad \text{for } T < T_{\text{RH}}, \\ h &\rightarrow SS \quad \text{for } T < T_C, \end{aligned} \quad (14)$$

where h is the Higgs field with mass $m_h \approx 125$ GeV and $T_C \approx 130$ GeV is the critical temperature of the SM electroweak phase transition. For $T < T_C$, one has to add new channels, for instance $W^+ W^- \rightarrow SS$, but their contribution to the production of the DM particles is negligible compared to $h \rightarrow SS$.

Following [19], the Boltzmann equation for the yield Y_S becomes

$$\begin{aligned} sH \frac{dY_S}{dT} = & -\frac{1}{512\pi^5} \sum_{i,j=H_1,H_2} \left[\int_{(m_i+m_j)^2}^{\infty} d\hat{s} P_{ij} |M_{ij}|^2 K_1 \left(\frac{\sqrt{\hat{s}}}{T} \right) \right] \\ & + \left(\frac{t_\beta}{1+t_\beta^2} \right)^2 \frac{y^4}{2\pi^3} \frac{m_h^3 v^2}{M_D^2} K_1 \left(\frac{m_h}{T} \right), \end{aligned} \quad (15)$$

where T is the temperature of the plasma, H is the Hubble parameter, K_1 is the first modified Bessel function of the second kind, $P_{ij} \equiv \sqrt{\hat{s} - (m_i + m_j)^2} \sqrt{\hat{s} - (m_i - m_j)^2} / \sqrt{\hat{s}}$ is a kinematic factor, and $|M_{ij}|^2$ is the squared matrix element, summed over internal degrees of freedom, for the $2 \rightarrow 2$ annihilation processes: $H_i^\dagger H_j \rightarrow SS$. The last term on the rhs of (15) arises from the decay $h \rightarrow SS$, upon ignoring the mass of the S particles. Also, upon ignoring M_S , the squared matrix elements $|M_{ij}|^2$ for the various $2 \rightarrow 2$ processes are

$$\begin{aligned} |M_{H_1^\dagger H_1 \rightarrow SS}|^2 &= |M_{H_2^\dagger H_2 \rightarrow SS}|^2 = 16 \left(\frac{t_\beta}{1+t_\beta^2} \right)^2 y^4 \frac{\hat{s}}{M_D^2}, \\ |M_{H_1^\dagger H_2 \rightarrow SS}|^2 &= |M_{H_2^\dagger H_1 \rightarrow SS}|^2 = 8(t_\beta^2 + t_\beta^{-2}) \left(\frac{t_\beta}{1+t_\beta^2} \right)^2 y^4 \frac{\hat{s}}{M_D^2}. \end{aligned} \quad (16)$$

The solution to the Boltzmann equation is obtained by integrating (15) over the temperature T . The limits of integration for the various channels are the ones shown in (14). However, before doing that, we have to make an assumption for the critical temperature and the thermal corrections to the masses of the scalar fields. In what follows, we assume that the critical temperature T_C and the thermal effects on the masses (for $T > T_C$) are similar to the pure SM Higgs sector and they are given by [39]

$$T_C \sim m_h, \quad m_{H_1}^2 \approx m_h^2 + \frac{1}{2} T^2, \quad m_{H_2}^2 \approx \frac{1+t_\beta^2}{t_\beta} m_{12}^2 + \frac{1}{2} T^2. \quad (17)$$

Under these assumptions and restricting T_{RH} to be above T_C , we can compute the yield Y_S at $T \approx 0$, which in turn implies the relic abundance [40]

$$\Omega_S h^2 \approx 0.12 \times \left(\frac{M_S}{1 \text{ GeV}} \right) \left(\frac{Y_S(T=0)}{4.3 \times 10^{-10}} \right). \quad (18)$$

B. Approximate results for $\Omega_S h^2$

In general, the yield Y_S cannot be calculated analytically, but depending on the reheating temperature T_{RH} , we are able to present approximate analytic results. We find that for decoupled $D_{1,2}$, i.e. $T_{\text{RH}} > M_D$, the relic abundance $\Omega_S h^2$ derived from Y_S in (18) takes on the form

$$\begin{aligned} \Omega_S h^2 \approx & 0.12 \times \left(\frac{M_S}{10^{-5} \text{ GeV}} \right) \left(\frac{2 \times 10^8 \text{ GeV}}{M_D} \right)^2 \\ & \times \left(\frac{y}{4.7 \times 10^{-2}} \right)^4 \left[\left(\frac{t_\beta}{1+t_\beta^2} \right)^2 + \left(\frac{T_{\text{RH}}}{10^4 \text{ GeV}} \right) \right], \end{aligned} \quad (19)$$

for $T_{\text{RH}} \gg m_{12}$, and

$$\begin{aligned} \Omega_S h^2 \approx & 0.12 \times \left(\frac{M_S}{10^{-3} \text{ GeV}} \right) \left(\frac{2 \times 10^5 \text{ GeV}}{M_D} \right)^2 \\ & \times \left(\frac{y}{4.7 \times 10^{-4}} \right)^4 \left(\frac{t_\beta}{1+t_\beta^2} \right)^2 \left[1 + \left(\frac{T_{\text{RH}}}{10^4 \text{ GeV}} \right) \right], \end{aligned} \quad (20)$$

for $T_{\text{RH}} \ll m_{12}$. Equations (19) and (20) are accurate up to 1%, except for $T_{\text{RH}} \sim m_{H_2}$, where the deviation from the

exact result is about 20%. Note that in both the regimes of T_{RH} , there are two contributions to $\Omega_S h^2$, given by the two terms contained in the last factors of (19) and (20). The first contribution does not depend on the reheating temperature T_{RH} and arises from the decay $h \rightarrow SS$, while the second one is proportional to T_{RH} . This second contribution is a result of the decoupling of the heavy fermionic doublets $D_{1,2}$ and indicates that for $T_{\text{RH}} \gtrsim 10^4$ GeV, the production of S particles is dominated by $2 \rightarrow 2$ annihilation processes given in (14). As discussed in [19,41], the latter is a general result for the freeze-in production mechanism via non-renormalizable operators. Finally, it is worth pointing out that $\Omega_S h^2$ is symmetric under $t_\beta \rightarrow t_\beta^{-1}$, as is the expression for M_S in (7).

IV. RESULTS

In Sec. II B, we have shown that the mass of the singlet S can be generated at the one-loop level, if the PQ symmetry is softly broken, and in Sec. III we have calculated the relic abundance of the S particles. In this section, we will be exploring the validity of the parameter space of our minimal model. To this end, one may consider the parameters,

$$T_{\text{RH}}, \quad M_D, \quad y^2, \quad t_\beta \quad \text{and} \quad m_{12},$$

as being independent. However, we prefer to solve the mass formula M_S^{rad} in (7) for y^2 and replace it with a physical observable, the S -particle mass M_S which is taken in our numerical estimates to be in the region: $10 \text{ keV} \leq M_S \leq 1 \text{ GeV}$. Consequently, the parameters that we allow to vary independently are

$$T_{\text{RH}}, \quad M_D, \quad M_S, \quad t_\beta \quad \text{and} \quad m_{12}. \quad (21)$$

We perform a scan over this parameter space, while imposing the perturbativity constraint on the Yukawa couplings: $Y_{1,2} < \sqrt{4\pi}$. In this way, we find the values of these parameters that satisfy the observed DM relic abundance [42]:

$$\Omega_S h^2 = \Omega_{\text{DM}} h^2 = 0.1198 \pm 0.0026. \quad (22)$$

In Fig. 3 we present contour lines on the $T_{\text{RH}} - m_{12}$ plane for discrete values of the S -particle mass M_S in the region: $1 \text{ keV} \leq M_S \leq 1 \text{ GeV}$, for $t_\beta = 1$ and $r = 10^{-2}$, which give the DM relic abundance (22). For $m_{12} \approx 10^{10}$ GeV, the reheating temperature T_{RH} can vary between the critical temperature $T_C \approx 130 \text{ GeV}$ and 10^8 GeV . This upper bound on T_{RH} may be as high as 10^{14} GeV , if the parameter $r = M_D/m_{12}$ is increased to the value $r = 10^5$, as depicted in Fig. 4. Yet, at the same time, m_{12} increases by 1 order of magnitude or so. On the other hand, for $m_{12} \approx 1 - 10 \text{ TeV}$, an acceptable DM relic abundance is reached only for large r and for $M_S \approx 1 \text{ keV}$, as can be seen from Fig. 4. Most remarkably, we notice that the predicted

FIG. 3. T_{RH} versus m_{12} for $r = M_D/m_{12} = 10^{-2}$, $t_\beta = 1$ and several RLDM masses M_S .

values for $\Omega_S h^2$ are compatible with the observed DM relic abundance $\Omega_{\text{DM}} h^2$, for a wide range of values for the parameters m_{12} , M_D and T_{RH} . Interestingly enough, the required Yukawa couplings $Y_{1,2}$ for a viable RLDM are sizeable, and always larger than the electron Yukawa coupling.

We recall here that we explore only regions where the fermion doublets $D_{1,2}$ are decoupled after the reheating of the Universe, i.e. we assume $M_D \gg T_{\text{RH}}$. As a working hypothesis, we assume the decoupling condition: $M_D > 3T_{\text{RH}}$. This condition is motivated by the fact for $T \approx M_D/3$, the $D_{1,2}$ particles become nonrelativistic and, as a consequence, its number density is exponentially suppressed by a Boltzmann factor. Correspondingly, for the scenario considered in Fig. 3, the heavy scalar H_2 will be also decoupled, because $m_{H_2} \gg M_D$.

Furthermore, we observe that for $T_{\text{RH}} \gtrsim 10^4 \text{ GeV}$, m_{12} becomes linearly dependent on the reheating temperature, as expected from the approximate analytic expression in (20). We also obtain a similar behavior in Fig. 4. In this case, however, the heavy scalar doublet H_2 is no longer constrained to be decoupled. As a result, there is an interface region at $T_{\text{RH}} \sim m_{12}$ that lies between the two

FIG. 4. The same as in Fig. 3, but for $r = M_D/m_{12} = 10^5$.

linear regimes, $T_{\text{RH}} \ll m_{12}$ and $T_{\text{RH}} \gg m_{12}$. At the interface region, there is a transition caused by the contribution of the heavy scalar doublet H_2 to the production of singlet fermions S [cf. (14)], which can reach equilibrium with the plasma when $T_{\text{RH}} \gg m_{12}$.

A. Solving the strong CP problem

It is known that in the SM there is an explicit breaking of CP (and P) discrete symmetry due to the instanton-induced term

$$\mathcal{L}_\theta = \frac{\theta}{32\pi^2} \text{Tr}(G_{\mu\nu} \tilde{G}^{\mu\nu}). \quad (23)$$

In the above, θ is a CP -odd parameter which can be absorbed into the quark masses. However, this θ parameter cannot be fully eliminated, since the combination: $\bar{\theta} = \theta - \text{ArgDet}M_q$, where M_q is the quark mass matrix, becomes a physical observable. It contributes to the neutron dipole moment and experimentally, it is severely bounded to be $|\bar{\theta}| \lesssim 10^{-11}$ [43]. The problem of why $\bar{\theta}$ is much smaller than all other CP -violating parameters, such as the well-known parameter $\epsilon_K \sim 10^{-3}$ from the $K^0 \bar{K}^0$ system, introduces another hierarchy problem in the SM known as the *strong problem*. A possible solution, suggested by Peccei and Quinn [15,16], is to promote the θ parameter into a dynamical field which naturally minimizes the energy. This dynamical field, called *the axion* [44,45], is a pseudo-Goldstone boson of the global anomalous PQ symmetry.

The SM has no global anomalous $U(1)_{\text{PQ}} n$ symmetry. One possible way to realize such a symmetry is to non-trivially extend its Higgs sector by adding a second Higgs doublet, resulting in the PQ-symmetric 2HDM. However, charging simply the field doublets Φ_1 and Φ_2 under the PQ symmetry as done in Table I does not lead to a healthy model. Such a model predicts a visible keV-axion with PQ-breaking scale $f_{\text{PQ}} \sim 100$ GeV, which is already excluded by the experiment. A minimal extension suggested by DFSZ [17] is to add a SM singlet Σ with charge +1 under $U(1)_{\text{PQ}}$ symmetry such that the scalar potential term,

$$\lambda_\Sigma \Sigma^2 \Phi_1^\dagger \Phi_2 + \text{H.c.} \subset V(\Phi_1, \Phi_2, \Sigma), \quad (24)$$

is invariant. Then, such a Σ -dependent term that occurs in the DFSZ potential $V(\Phi_1, \Phi_2, \Sigma)$ breaks the PQ symmetry spontaneously, when the electroweak singlet field Σ receives a large VEV $\langle \Sigma \rangle$ which is not necessarily tied in with that of the electroweak scale v . For this reason, in this paper we have made the identification

$$\langle \Sigma \rangle \equiv f_{\text{PQ}} \approx m_{12}, \quad (25)$$

with $\lambda_\Sigma \approx 1$. From experimental constraints and astrophysical considerations, the PQ-breaking scale f_{PQ} must be

typically larger than 10^9 GeV [46]. Interestingly, within the RLDM scenario, there are values for m_{12} satisfying this constraint and at the same time are compatible with the observed $\Omega_{\text{DM}} h^2$, as discussed in the previous section. An example is shown in Fig. 3 for $M_S = 1$ GeV and $T_{\text{RH}} \ll m_{12}$. In addition, values where $m_{12} \gtrsim 10^9$ GeV can be also obtained for other hierarchies e.g. $r \sim 1$ and $r \gg 1$, as shown in Fig 4. This seems to be a rather generic feature of the RLDM realization.

Although the above is a strong indication that the DFSZ solution to the strong CP problem is consistent with the RLDM scenario, a detailed analysis of the UV-complete DSFZ-extended model lies beyond the scope of this article. In particular, for $f_{\text{PQ}} \sim 10^{11}$ GeV [47], the axion becomes a sizeable DM component resulting in a two-component DM, consisting of the axion and the S particle, and so a more careful treatment will be required.

B. Detection of RLDM

We observe that for small enough reheating temperatures, $T_{\text{RH}} \sim 1$ TeV, the fermion doublets $D_{1,2}$, as well as the heavy scalar doublet H_2 , can lie at the TeV scale, provided that M_S is of order $\mathcal{O}(10 \text{ keV})$. This is shown in Figs. 3 and 4 for light M_S , where M_D and m_{12} lie in the vicinity of the TeV scale. As a result, the DM particle S can be probed indirectly by looking for its associated “partners” of the heavy Higgs doublet H_2 . In general, we expect that at the LHC, the heavy sector of the 2HDM will be efficiently explored up to the TeV scale [34,48]. For the RLDM scenario at hand, however, such exploration may be somehow challenging, when looking for charged Higgs bosons with masses larger than ~ 1 TeV for a wide range of t_β values [34,48].

On the other hand, direct detection experiments for sub-GeV DM particles focus on their interactions with atomic electrons, e.g. see [49,50]. However, in the RLDM scenario, such a detection of S particles is practically unattainable, because S interacts feebly with the SM Higgs boson with a coupling proportional to $v/M_D \ll 1$ yielding a cross section for $Se \rightarrow Se$, which is highly suppressed by fourth powers of the electron-to-Higgs-mass ratio, i.e.

$$\begin{aligned} \bar{\sigma}_{Se} &\approx \frac{y^4}{\pi} \frac{t_\beta^2}{(1+t_\beta^2)^2} \left(\frac{m_e}{m_h}\right)^4 \frac{1}{M_D^2} \\ &\approx 10^{-50} \times \frac{y^4 t_\beta^2}{(1+t_\beta^2)^2} \left(\frac{1 \text{ GeV}}{M_D}\right)^2 \text{ cm}^2. \end{aligned} \quad (26)$$

Hence, a simple estimate shows that $\bar{\sigma}_{Se}$ is much smaller than its current experimental reach: $\bar{\sigma}_{Se}^{\text{exp}} \simeq 10^{-38} \text{ cm}^2$.

Another potentially observable effect could originate from the invisible Higgs-boson decay, $h \rightarrow SS$. Current LHC analyses report the upper bound [51]: $\text{Br}(h \rightarrow \text{inv}) < 0.28$, which for the RLDM scenario translates into

$$M_D \gtrsim 10^4 \times y^2 \frac{t_\beta}{1 + t_\beta^2} \text{ GeV}. \quad (27)$$

Note that this constraint is comfortably satisfied for the entire range of our parameter space.

In summary, at least for the foreseeable future, the RLDM particle S proposed here will remain elusive. This leaves only a window for the LHC to find indirectly a second heavy Higgs doublet H_2 and/or a pair of heavy fermion doublets $D_{1,2}$.

V. CONCLUSIONS

One central problem of most electroweak scenarios that require the existence of very light DM particles in the keV-to-GeV mass range is the actual origin of this sub-GeV scale. To address the origin of such a small scale, we have presented a novel radiative mechanism that can naturally generate a sub-GeV mass for a light singlet fermion S , which is stable and can successfully play the role of the DM.

In order to minimally realize such a radiative light dark matter, we have considered a Peccei-Quinn symmetric two-Higgs doublet model, which was extended with the addition of a singlet fermion S and a pair of massive vectorlike SU(2) isodoublets $D_{1,2}$ that are not charged under the PQ symmetry. Instead, the singlet fermion S is charged under the PQ symmetry and so it has no bare mass at the tree level. However, upon soft breaking of the PQ symmetry, we have shown how the singlet fermion S receives a nonzero mass at the one-loop level. The so-generated radiative mass for the singlet fermion S lies naturally in the cosmologically allowed region of $\sim 10 \text{ keV} - 1 \text{ GeV}$.

We have computed the relic abundance of the RLDM S , for different plausible heavy mass scenarios. Specifically, for all scenarios we have been studying, we have assumed that the S particles were absent in the early universe, while the fermion isodoublets $D_{1,2}$ stay out of equilibrium

through the entire thermal history of the Universe, because their gauge-invariant mass M_D is taken to be well above the reheating temperature T_{RH} . Then, we have found that the observationally required relic abundance for the RLDM S can be produced via decays and annihilations of Higgs-sector particles.

We have analyzed a heavy mass scenario where the PQ-breaking scale f_{PQ} can reach values $\sim 10^9 \text{ GeV}$ as required by the Dine-Fischler-Sredniki-Zhitnitsky axion model to explain the strong CP problem. We have found that for appropriate isodoublet masses (e.g. in Fig. 3 $M_D \sim 10^{-2} f_{\text{PQ}}$), the RLDM particle S in such a scenario can successfully account for the missing matter component of the Universe. In addition, we have investigated whether a lower mass limit exists for the heavy Higgs scalars, within the context of a viable RLDM scenario. We have found that the masses of the heavy scalars can be as low as TeV, which allows for their possible detection at the LHC in the near future.

The PQ-symmetric scenario we have studied here generates a viable RLDM at the one-loop level. However, one may envisage other extensions of the SM, in which the required small mass for the light DM could be produced at two or higher loops. For instance, if the SM is extended by two scalar triplets, a small DM mass can be generated through their mixing at the two-loop level, in a fashion similar to the Zee model. In this context, it would be interesting to explore possible models where both the tiny mass of the SM neutrinos and the small mass of the light DM have a common radiative origin and study their phenomenological implications.

ACKNOWLEDGMENTS

The work of A. P. is supported in part by the Lancaster-Manchester-Sheffield Consortium for Fundamental Physics, under Science and Technology Facilities Council (STFC) research Grant No. ST/L000520/1.

-
- [1] G. Bertone, D. Hooper, and J. Silk, Particle dark matter: Evidence, candidates and constraints, *Phys. Rep.* **405**, 279 (2005).
 - [2] F. Kahlhoefer, Review of LHC dark matter searches, *Int. J. Mod. Phys. A* **32**, 1730006 (2017).
 - [3] D. S. Akerib *et al.* (LUX Collaboration), Results from a Search for Dark Matter in the Complete LUX Exposure, *Phys. Rev. Lett.* **118**, 021303 (2017).
 - [4] E. Aprile *et al.* (XENON Collaboration), Physics reach of the XENON1T dark matter experiment, *J. Cosmol. Astropart. Phys.* **04** (2016) 027.
 - [5] J. Billard, L. Strigari, and E. Figueroa-Feliciano, Implication of neutrino backgrounds on the reach of next generation dark matter direct detection experiments, *Phys. Rev. D* **89**, 023524 (2014).
 - [6] T. Asaka, S. Blanchet, and M. Shaposhnikov, The nuMSM, dark matter and neutrino masses, *Phys. Lett. B* **631**, 151 (2005).
 - [7] A. Kusenko, Sterile neutrinos: The dark side of the light fermions, *Phys. Rep.* **481**, 1 (2009).
 - [8] N. E. Mavromatos and A. Pilaftsis, Anomalous Majorana neutrino masses from torsionful quantum gravity, *Phys. Rev. D* **86**, 124038 (2012).
 - [9] R. Adhikari *et al.*, A white paper on keV sterile neutrino dark matter, *J. Cosmol. Astropart. Phys.* **01** (2017) 025.
 - [10] L. Heurtier and D. Teresi, Dark matter and observable lepton flavor violation, *Phys. Rev. D* **94**, 125022 (2016).

- [11] C. Boehm and P. Fayet, Scalar dark matter candidates, *Nucl. Phys.* **B683**, 219 (2004).
- [12] J.-H. Huh, J. E. Kim, J.-C. Park, and S. C. Park, Galactic 511 keV line from MeV milli-charged dark matter, *Phys. Rev. D* **77**, 123503 (2008).
- [13] R. Essig *et al.*, Dark sectors and new, light, weakly coupled particles, [arXiv:1311.0029](#).
- [14] J. Alexander *et al.*, Dark sectors 2016 workshop: Community report, [arXiv:1608.08632](#).
- [15] R. D. Peccei and H. R. Quinn, Constraints imposed by conservation in the presence of instantons, *Phys. Rev. D* **16**, 1791 (1977).
- [16] R. D. Peccei and H. R. Quinn, Conservation in the Presence of Instantons, *Phys. Rev. Lett.* **38**, 1440 (1977).
- [17] A. R. Zhitnitsky, On possible suppression of the axion hadron interactions, *Yad. Fiz.* **31**, 497 (1980) [*Sov. J. Nucl. Phys.* **31**, 260 (1980)].
- [18] M. Dine, W. Fischler, and M. Srednicki, A simple solution to the strong problem with a harmless axion, *Phys. Lett.* **104B**, 199 (1981).
- [19] L. J. Hall, K. Jedamzik, J. March-Russell, and S. M. West, Freeze-in production of FIMP dark matter, *J. High Energy Phys.* **03** (2010) 080.
- [20] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, On the anomalous electroweak baryon number nonconservation in the early universe, *Phys. Lett.* **155B**, 36 (1985).
- [21] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, Progress in electroweak baryogenesis, *Annu. Rev. Nucl. Part. Sci.* **43**, 27 (1993).
- [22] A. Pilaftsis, *CP* violation and baryogenesis due to heavy Majorana neutrinos, *Phys. Rev. D* **56**, 5431 (1997).
- [23] A. Pilaftsis and T. E. J. Underwood, Resonant leptogenesis, *Nucl. Phys.* **B692**, 303 (2004).
- [24] P. S. B. Dev, P. Millington, A. Pilaftsis, and D. Teresi, Flavour covariant transport Equations: An application to resonant leptogenesis, *Nucl. Phys.* **B886**, 569 (2014).
- [25] A. Achelashvili and Z. Tavartkiladze, Calculable cosmological violation and resonant leptogenesis, [arXiv:1611.07956](#).
- [26] B. W. Lee and S. Weinberg, Cosmological Lower Bound on Heavy Neutrino Masses, *Phys. Rev. Lett.* **39**, 165 (1977).
- [27] V. Iršič *et al.*, New constraints on the free-streaming of warm dark matter from intermediate and small scale Lyman- α forest data, [arXiv:1702.01764](#).
- [28] M. Viel, G. D. Becker, J. S. Bolton, and M. G. Haehnelt, Warm dark matter as a solution to the small scale crisis: New constraints from high redshift Lyman- α forest data, *Phys. Rev. D* **88**, 043502 (2013).
- [29] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, and J. P. Silva, Theory and phenomenology of two-Higgs-doublet models, *Phys. Rep.* **516**, 1 (2012).
- [30] H. Georgi and D. V. Nanopoulos, Suppression of flavor changing effects from neutral spinless meson exchange in gauge theories, *Phys. Lett.* **82B**, 95 (1979).
- [31] A. Dedes, D. Karamitros, and V. C. Spanos, Effective theory for electroweak doublet dark matter, *Phys. Rev. D* **94**, 095008 (2016).
- [32] J. F. Gunion and H. E. Haber, The *CP* conserving two Higgs doublet model: The Approach to the decoupling limit, *Phys. Rev. D* **67**, 075019 (2003).
- [33] I. F. Ginzburg and M. Krawczyk, Symmetries of two Higgs doublet model and *CP* violation, *Phys. Rev. D* **72**, 115013 (2005).
- [34] P. S. B. Dev and A. Pilaftsis, Maximally symmetric two Higgs doublet model with natural standard model alignment, *J. High Energy Phys.* **12** (2014) 024; Erratum: Maximally symmetric two Higgs doublet model with natural standard model alignment, *J. High Energy Phys.* **11** (2015) 147.
- [35] A. Pilaftsis, Symmetries for standard model alignment in multi-Higgs doublet models, *Phys. Rev. D* **93**, 075012 (2016).
- [36] B. Grzadkowski, O. M. Ogreid, and P. Osland, Spontaneous *CP* violation in the 2HDM: physical conditions and the alignment limit, *Phys. Rev. D* **94**, 115002 (2016).
- [37] M. Gorbahn, J. M. No, and V. Sanz, Benchmarks for Higgs effective theory: Extended Higgs sectors, *J. High Energy Phys.* **10** (2015) 036.
- [38] H. Bêlusca-Maitô, A. Falkowski, D. Fontes, J. C. Romão, and J. P. Silva, Higgs EFT for 2HDM and beyond, *Eur. Phys. J. C* **77**, 176 (2017).
- [39] Z. Fodor and A. Hebecker, Finite temperature effective potential to order g^4 , λ^2 and the electroweak phase transition, *Nucl. Phys.* **B432**, 127 (1994).
- [40] J. Edsjo and P. Gondolo, Neutralino relic density including coannihilations, *Phys. Rev. D* **56**, 1879 (1997).
- [41] F. Elahi, C. Kolda, and J. Unwin, Ultraviolet freeze-in, *J. High Energy Phys.* **03** (2015) 048.
- [42] P. A. R. Ade *et al.* (Planck Collaboration), Planck 2013 results. XVI. Cosmological parameters, *Astron. Astrophys.* **571**, A16 (2014).
- [43] J. E. Kim and G. Carosi, Axions and the strong *CP* problem, *Rev. Mod. Phys.* **82**, 557 (2010).
- [44] S. Weinberg, A New Light Boson?, *Phys. Rev. Lett.* **40**, 223 (1978).
- [45] F. Wilczek, Problem of Strong *P* and *T* Invariance in the Presence of Instantons, *Phys. Rev. Lett.* **40**, 279 (1978).
- [46] G. G. Raffelt, Astrophysical axion bounds, *Lect. Notes Phys.* **741**, 51 (2008).
- [47] M. S. Turner, Cosmic and local mass density of invisible axions, *Phys. Rev. D* **33**, 889 (1986).
- [48] K. Pedersen and Z. Sullivan, Probing the two Higgs doublet wedge region with charged Higgs boson decays to boosted jets, *Phys. Rev. D* **95**, 035037 (2017).
- [49] A. Dedes, I. Giomataris, K. Suxho, and J. D. Vergados, Searching for secluded dark matter via direct detection of recoiling nuclei as well as low energy electrons, *Nucl. Phys.* **B826**, 148 (2010).
- [50] R. Essig, T. Volansky, and T.-T. Yu, New constraints and prospects for sub-GeV dark matter scattering off electrons in xenon, [arXiv:1703.00910](#).
- [51] G. Aad *et al.* (ATLAS Collaboration), Search for invisible decays of a Higgs boson using vector-boson fusion in *pp* collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *J. High Energy Phys.* **01** (2016) 172.