Evolutionary history resolves global organization of root functional traits

DOI:

[10.1038/nature25783](https://doi.org/10.1038/nature25783)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Ma, Z., Guo, D., Xu, X., Lu, M., Bardgett, R., Eissenstat, D. M., McCormack, M. L., & Hedin, L. O. (2018). Evolutionary history resolves global organization of root functional traits. *Nature*, 555(7694), 94-97. <https://doi.org/10.1038/nature25783>

Published in:

Nature

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

1 **Evolutionary history resolves global organization of root functional traits**

2 **Short title:** Root trait biogeography

3

4 Zeqing Ma¹, Dali Guo¹, Xingliang Xu¹, Mingzhen Lu², Richard D. Bardgett³, David
5 M. Eissenstat⁴, M. Luke McCormack^{1,5}, Lars O. Hedin²

6 ¹Center for Forest Ecosystem Studies and Qianyanzhou Ecological Station, Key
7 Laboratory of Ecosystem Network Observation and Modeling, Institute of
8 Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences,
9 Beijing 100101, China

10 ²Department of Ecology and Evolutionary Biology, Princeton University, NJ 08544,
11 USA

12 ³School of Earth and Environmental Sciences, The University of Manchester,
13 Manchester, M13 9PT, UK

14 ⁴Department of Ecosystem Science and Management, The Pennsylvania State
15 University, University Park, PA 16802, USA

16 ⁵Department of Plant and Microbial Biology, University of Minnesota, St Paul, MN
17 55108, USA

18

19 **E-mail addresses of all authors:**

20 Z.M.: mazq@igsnrr.ac.cn; D.G.: guodl@igsnrr.ac.cn; X.X.: xuxingl@hotmail.com;

21 M.L.: mingzhen@princeton.edu; R.D.B.: richard.bardgett@manchester.ac.uk;

22 D.M.E.: dme9@psu.edu; M.L.M.: mltmcc@gmail.com; L.O.H.:

23 lhedin@princeton.edu

24

25 **Keywords:** plant functional traits, belowground, root traits, root diameter, root
26 economics spectrum, phylogeny, root biogeography, biomes, mycorrhizal fungi,
27 nutrient foraging, root lifespan, root nitrogen

28

29 **Article Type:** Letters

30 **Number of words in the abstract:** 282

31 **Number of words in the main text:** 1737 body + 349 method summary

32 **Number of references:** 28 Main Text including Methods Summary + 34 Methods +
33 43 Extended Data + 30 Supplementary Information (total 135)

34 **Number of figures:** 2, **tables:** 0

35 **Number of text box:** 0

36 **Author for correspondence:** Dr. Dali Guo, Center for Forest Ecosystem Studies and
37 Qianyanzhou Ecological Station, Key Laboratory of Ecosystem Network Observation
38 and Modeling, Institute of Geographic Sciences and Natural Resources Research,
39 Chinese Academy of Sciences, Beijing 100101, China.

40 Phone: 86-10-64889027, Fax: 86-10-64888153.

41 E-mail: guodl@igsnr.ac.cn

42

Abstract

Plant roots have diversified greatly in both form and function since the first land plants emerged^{1,2}, but the global organization of root functional traits remains poorly understood^{3,4}. We analyzed a new global dataset of ten functionally important traits, compiled from metabolically active first-order roots collected from 369 species distributed across natural plant communities of seven biomes globally. Our results identify a high degree of organization of root traits across species and biomes, but also that the observed pattern differs from expectations based on studies of leaf traits. Root diameter exerted the strongest influence on root trait variation across plant species, growth forms, and biomes. Our analysis suggests that plants have evolved thinner roots since they first emerged in land ecosystems, which, in turn, has allowed them to dramatically improve the efficiency of soil exploration per unit of carbon invested and reduce dependence on symbiotic mycorrhizal fungi. We also found that diversity in root morphological traits is greatest in the tropics where plant diversity is highest and many ancestral phylogenetic groups are preserved, but declines sharply from the tropics to temperate regions to desert biomes, presumably due to changes in resource supply caused by seasonally inhospitable abiotic conditions. Our results suggest that root traits have evolved along a spectrum bounded by two contrasting strategies of root life: from an ancestral ‘conservative’ strategy in which plants with thick roots depend on symbiosis with mycorrhizal fungi for soil resources, to a derived ‘opportunistic’ strategy in which thin roots allow

65 **plants to more efficiently leverage photosynthetic carbon for soil exploration.**
66 **These findings imply that innovations of belowground traits have been key for**
67 **preparing plants to colonize new habitats, and for generating biodiversity within**
68 **and across biomes.**

Recent efforts to understand how functional traits are organized across land plants have revealed striking patterns across the leaf economic spectrum^{5,6}, but whether such high degree of organization is reflected also in root traits remains controversial^{4,7}. A key factor limiting progress has been the paucity of data on root traits across plant species and biomes, as roots are difficult to sample and characterize^{8,9}. Yet, roots are vital for the ability of plants to acquire nutrients and water – two functions of fundamental importance to whole-plant performance and for predicting how plants respond to elevated CO₂ and climate change¹⁰⁻¹².

Roots face ecological and physiological challenges that differ fundamentally from leaves. Roots must compete for, and acquire, nutrients and water in environments that vary greatly across the world's biomes, with biophysical conditions ranging from relatively stable (*e.g.*, tropical rainforests) to highly seasonal (*e.g.*, deserts or boreal forests). The high diversity that exists in root form and function, and in the degree of association with symbiotic mycorrhizal fungi, raises a fundamental question: How are root traits organized across the diverse taxa that inhabit different ecological conditions worldwide?

Here we propose a new model of root trait organization that is functionally decoupled from the leaf economic spectrum, and that derives from the phylogenetic history of root diameter and its evolutionary consequences for plant resource acquisition.

Specifically, we evaluated an unprecedented species- and biome-specific dataset of 10 root traits in three major categories^{3,13} (morphology, physiology, and

mycorrhizal association, *SI* note 1), from >1,200 individual plants distributed across 369 species (210 genera, and 79 families), seven major biomes, and three continents of the world (Extended Data Table 1). This dataset is unique in that our observations: (i) derive solely from native plant communities with natural soil and nutrient conditions; (ii) focus on first-order roots (the most distal and absorptive roots of the branching system), which are subject to strong selection by the local environment^{8,9,14}; (iii) accurately identify species and root order (*i.e.*, measure of branching hierarchy⁸) in mixed-species ecosystems, by tracing roots to parent trees¹⁵; and (iv) apply consistent analytical methods to trait measures across all species and biomes (94% of observations were collected by ourselves; Methods).

We first evaluated whether the first-order root traits are globally organized in a manner analogous to the leaf economic spectrum^{5,6}, a composite axis of trait variation that ranges from nitrogen-rich leaves with high specific leaf area and short leaf lifespan, to nitrogen-poor leaves with low specific leaf area and long leaf lifespan. In roots, nitrogen supports metabolic activity including nutrient and water transport, enzyme functioning, and mycorrhizal symbiosis¹⁶. As a result, nitrogen has been proposed to serve a similarly central role in the trait organization of roots, with high root nitrogen occurring in species with high leaf nitrogen, rapid growth, and short root lifespan^{4,17}.

Our results do not support the idea of an analogous organizing role of nitrogen in a global root economic spectrum (*c.f.*, similar conclusion from taxonomically and geographically smaller datasets; refs.4,18,19). First, a principal component analysis

failed to identify root nitrogen (analogous to leaf nitrogen) as a significant contributor to the primary axis of trait variation (Extended Data Fig.1 and Table 2). Instead, root traits were most strongly (46%) explained by root diameter and a group of traits associated with root construction and mycorrhizal association (axis1 in Extended Data Fig.1).

Second, root nitrogen was not correlated to specific root length in a manner analogous to the relationship between specific leaf area and leaf nitrogen (Extended Data Fig.2). Third, *in situ* ($n=73$) and hydroponic-based ($n=119$) measures showed no systematic relationship between root nitrogen uptake (analogous to leaf photosynthetic capacity) and root diameter, specific root length, or plant growth form (Extended Data Figs.3b,d and 5). Combined, these results suggest that nitrogen is less important in belowground nutrient foraging than in aboveground light and CO₂ capture (*SI* note 3). Moreover, root lifespan (analogous to leaf lifespan) was correlated with root diameter and specific root length, but explained only 14% and 17% of the respective variance ($P<0.01$ for both, linear model; Extended Data Fig.3a,c).

We next analyzed the organizing role of root diameter in determining trait variation across plants. We found that the length of root per unit biomass invested – or specific root length (*SRL*) – increases non-linearly with decreasing root diameter (D) following the allometric relationship $SRL = \frac{16.8}{\pi D^2}$ (Fig.1a, red line; *SI* note 1). This relationship indicates that as roots get thinner, plants can explore dramatically greater volumes of soil per unit carbon invested. Albeit with some overlap, we also

found that woody and herbaceous plants occupy different parts of the specific root length vs. root diameter relationship: woody plants (Fig. 1a, brown points) occur in a region where differences in root diameter have limited effect on specific root length, while herbaceous plants (green points) reside in a region where even small diameter variations cause large changes in specific root length.

We further found that – in thin-rooted species – even a modest evolutionary change in tissue density of first-order roots can greatly alter the soil length explored per unit carbon invested. The dashed red lines in Fig. 1a indicate the sensitivity of the *SRL*-diameter relationship to changes in root tissue density across a physiologically relevant range (0.1 to 1 g cm⁻³). For example, low root density allows the grass *Agropyron cristatum* to explore ~350m more soil per gram biomass than the shrub *Rhaphiolepis indica*, despite similar (~0.2mm) root diameter (Fig. 1a, red arrows and cross section photos). We infer that, over evolutionary time, plants can use both root diameter (Fig. 1a, x-axis) and tissue density (Fig. 1a, dashed lines) to influence specific root length: thin and soft first-order roots have the advantage of efficient soil exploration, but incur the cost tradeoff of sacrificing water conduction, tissue permanence, and the ability to penetrate the soil matrix.

We next examined the role of evolutionary history in structuring the differences in root diameter across all major vascular plant families in our dataset. We found that (Fig. 1b), on average, thick roots are associated with evolutionarily ancient taxa (*e.g.*, *Magnoliaceae*) while thin roots are increasingly common in taxa that have recently diverged from their ancestral lineage (*e.g.*, *Betulaceae*)(weighted linear regression:

157 $r^2=0.54$; $P<0.001$). Herbaceous plants evolved more recently (Fig.1b, green circles)
158 and – with the exception of *Amaryllidaceae* and *Boraginaceae* – are thus
159 characterized by thin roots and specific root lengths that exceed woody plants (Fig.
160 1a). Together, these patterns broadly characterize an evolutionary transition from
161 ancient tree taxa, defined by thick first-order roots, to more recently radiated^{20,21}
162 woody and herbaceous plants with thin roots that can explore dramatically greater
163 lengths of soil per carbon invested.

164 The trend towards thinner roots has had major consequences for the symbiosis
165 between plant roots and mycorrhizal fungi. We found that mycorrhizal colonization
166 (*i.e.*, % root length colonized) declines as roots get thinner (Fig.1c), but also that
167 herbaceous roots have ~30% less colonization than woody plants at the same root
168 diameter (linear model; $r^2=0.63$ with difference between herbaceous and woody
169 plants at $P<0.001$). In addition, herbaceous plants have on average 33% lower root
170 tissue density than woody plants (Fig.1d; unequal variance t-test; $P<0.001$), though
171 considerable unexplained variation exists across taxa. These differences suggest that
172 first-order roots have become less dependent on mycorrhizae as they have evolved
173 thinner diameter, but also that the innovation of the short-lived herbaceous growth
174 form has fundamentally changed the relationship between root diameter and
175 mycorrhizal colonization.

176 A phylogenetic independence contrasts (PICs) analysis¹⁷ confirmed that variation
177 in root diameter, specific root length, and mycorrhizal colonization are strongly
178 influenced by evolutionary history (Blomberg's K value in Extended Data Table1).

In contrast, root chemical traits did not display a clear phylogenetic signal, indicating that, for these traits ecological variation overshadows evolutionary constraints²².

When combined, our results identify a general evolutionary trend from thick roots that rely on mycorrhizal fungi for resource acquisition, to thin roots that can explore the soil at high carbon use efficiency but with less reliance on mycorrhizae. The observed root trait combinations imply selection for two contrasting plant strategies: (i) a ‘*conservative*’ strategy, in which carbon allocation to mycorrhizae enhances the ability of plants to compete in environments with stable resources and intense plant-plant competition; and (ii) an ‘*opportunistic*’ strategy, in which thin roots benefit plants in less predictable environments (*e.g.*, seasonal drought or cold), where rapid root growth response to fluctuating resource supply is rewarded.

It is less clear, however, why herbaceous plants have lower mycorrhizal colonization than woody plants at similar diameter (Fig.1c), although softer tissue may cause roots to be less permanent (Extended Data Fig.2), and therefore less able to maintain stable mycorrhizal relationships.

We next evaluated whether the distribution of root diameter changed across biomes that may differ in the pattern and stability of resource supply. First, we found an overall trend of decreasing variance in root diameter of woody plants from the more stable conditions of tropical and sub-tropical forests, to the highly seasonal boreal and desert biomes (Fig.2, Levene’s test, Extended Data Table 3). Second, while woody plants were limited to thin-rooted species in the most seasonal biomes,

the diameter of herbaceous plant roots did not differ systematically across biomes
(Extended Data Fig.4a,b).

These patterns are consistent with biome-specific differences in both evolutionary history and stability of resource supply and abiotic conditions. The tropical forest biome is ancient²³, characterized by seasonally stable supplies of soil resources, and holds species that range from ancestral thick-rooted to more derived thin-rooted taxa. In contrast, boreal and desert biomes are evolutionarily young²⁴, and have been colonized mainly by thin-rooted species that, in theory, can rapidly respond²⁵ to fluctuating soil resources and seasonally inhospitable conditions. The coexistence of thin-rooted plants with more ancient thick-rooted strategies suggests that heterogeneity within the tropical biome is sufficient to maintain a range of niche conditions for plant belowground strategies.

Our findings suggest that – at the timescale of plant evolution – innovations of belowground traits have been key for preparing plants to colonize new habitats, and for the rich generation of biodiversity within and across biomes. The dominant dimension of trait evolution for first-order roots has been a decrease in diameter, which, in turn, has reduced the dependence on mycorrhizal fungi, increased the efficiency of root growth, and thus elevated the ability of plants to leverage photosynthetic carbon for soil exploration. An improved functional understanding of root traits is critical for comprehending the history and distribution of plant life, and may help to predict the risk of species extinction and to conserve biodiversity in the face of environmental change.

METHODS SUMMARY

We collected roots from natural plant communities across seven major biomes and three continents (Asia, Europe and North America) during 2004 to 2016. We adopted the root-order based methodology (established in 2002; ref.8). At each sampling site, we selected common indigenous species that are representative of the local plant community. For each species, we sampled at least three individual plants to derive mean trait values for each species. We analyzed four morphological traits (diameter, specific root length, root tissue density, and root length), three physiological-chemical traits (root nitrogen content, root carbon content, and root carbon to nitrogen ratio), and the extent of mycorrhizal colonization (Detailed in Methods). We calculated percent mycorrhizal colonization by sampling first-order roots and determining by microscope the presence of either arbuscular mycorrhizal or ectomycorrhizal fungal structures within an individual root segment. For each species, percent colonization was calculated across 20-150 first-order root segments as detailed in Methods. We enhanced coverage of some biomes by including literature data (~5% of final dataset), but only if methodologies were consistent with our methods. In total, we gathered traits from 369 species (281 woody, 88 herbaceous, Extended Data Table 1, SI Fig.S3).

We collected data on plant root lifespan for 40 species using *in situ* minirhizotrons across boreal and temperate forests, with individual measures spanning at least one year. We acquired additional lifespan data from published literature identified via Web of Science and Google Scholar. We measured per-

biomass root nitrogen uptake rates in 36 plant species and acquired data for an additional 101 species through published literature (Extended Data Table 4).

Principal component analysis (PCA) and variance partitioned among traits in our database were analyzed by the FactorR package of the R platform. Linear models, linear mixed effects models (package lme4) and Levene's test were also conducted in R. Data were log transformed to correct for deviations from normality. We followed the APG III phylogenetic system in all analyses²⁶ and used PHYLOCOM²⁷ to construct phylogenetic trees (*SI Fig.S4*). Following Wikstrom, et al.²⁸, we defined the divergence time of a plant family using the earliest diverging genus within that family.

Full Methods and any associated references are available in the online version of the paper at www.nature.com/nature.

Methods

Sampling approach. We collected roots from natural plant communities across seven major biomes and three continents (Asia, Europe, and North America) during 2004 to 2016. Our sampling sites range from -1.4°C to 22.4°C in mean annual temperature, and from 35mm to 2651mm in mean annual precipitation. At each site, we selected common indigenous species that are representative of local plant communities. We sampled multiple root branches or segments from at least three individual plants from each species to derive the mean species trait value. For species

that occupied more than one sample location, we merged the local means into one species trait value. Eleven species occurred in more than a single biome; for these we calculated a mean value for each biome.

We identified roots to species level in mixed-species ecosystems by tracing a root to its parent tree. During the growing season, we selected mature individuals and excavated the surface soil (0–20cm) around the plant stem to expose lateral roots. We then sampled multiple intact root branches and gently cleared the attached soil. Sampled roots were bagged and immediately placed in a cooler, and then either transferred to a refrigerator for processing within the next few days or kept frozen until later laboratory analyses.

Laboratory analyses of root functional traits. We adopted a root branching-order based approach, where absorptive fine roots are sorted based on their position in the branching architecture (established in 2002, details in *SI* note 2; ref. 8). We dissected root branches according to standard methodology¹⁵ and determined root diameter, root length, specific root length, root tissue density, root nitrogen and carbon concentration, and mycorrhizal colonization (*SI* note 1).

Root diameter and length were measured using stereomicroscope with an ocular micrometer. Specific root length was determined by dividing root length by the dry biomass weight. We calculated the volume of root segments from root diameter and length, assuming segments are cylinders. Root tissue density was then calculated using dry mass and volume. Sampled roots were oven-dried at 60°C for 48h, ground

to fine powder with a ball-mill for subsequent measurements of carbon and nitrogen on an elemental analyzer (Vario EL Cube; Elementar, Hanau, Germany).

We measured the length of root colonized by mycorrhizal fungi in 137 species from sub-tropical¹⁹ forests, temperate forests¹⁵, and temperate grasslands²⁹. We calculated percent mycorrhizal colonization by sampling first-order roots and determining by microscope the presence of either arbuscular mycorrhizal or ectomycorrhizal fungal structures within an individual root segment. For arbuscular mycorrhizal fungi we used a standard staining technique to identify coils and arbuscules^{19,29}; no stain was needed to identify ectomycorrhizal fungal sheaths¹⁹. For each individual plant, we selected at least 10 root branches (containing multiple order of roots). We next randomly selected 20-150 first-order root segments^{19,29} for each species, ensuring that each segment length was consistent across all roots sampled. We then calculated the species-specific percent length colonization as the ratio of the sum of infected root segments over all root segments examined. We used two different techniques: one based on cross-sectional analysis (MC_1 ; $n=110$ species) and one based on scanning the root surface (MC_2 ; $n=27$ species); both allowed us to quantify fungal association within a standardized root area. We kept the effective area examined per root segment approximately the same for both methods (173 vs. 169 mm²), such that the results are equivalent ($MC_2 = 1.02 \times MC_1 - 0.02$; $r^2=0.988$).

Since distal fine roots (*i.e.*, first-order roots) are primarily responsible for plant nutrient acquisition^{8,9}, we focused our analyses on first-order root traits. We accumulated 480 species-specific observations, of which 187 are unpublished and 256

309 published by our groups^{14,15,19,25,29}. To cover a broader range of biomes (*e.g.*, boreal
 310 and Mediterranean), we added 37 observations from the literature^{8,30-32} into our
 311 dataset, taking care to only include studies of first-order roots and consistent methods.
 312 In total, we compiled 480 species-specific observations, covering 210 genera and 79
 313 families (Extended Data Table 1).

314 **Root lifespan.** We collected multi-year root lifespan data of 40 species using *in situ*
 315 minirhizotrons across subtropical forest, tropical forest and temperate forest of
 316 Europe³³, Asia^{34,35} and North America^{36,37}. We further added data from the published
 317 literature to develop a global dataset. We carefully selected observations only from
 318 studies of first-order roots or distal roots, using *in situ* minirhizotrons or root
 319 windows³⁸⁻⁵⁹. When corresponding root traits (*e.g.*, diameter, *SRL*) were not
 320 available, we used species-specific observations from our own dataset to match the
 321 life span data. In total, we obtained 70 species-specific observations and 13
 322 community-observations across 5 biomes.

323 **Root nitrogen uptake rates.** We measured root biomass-specific nitrogen uptake
 324 rates using two standard approaches: (i) by isolating an intact living root branch and
 325 exposing it to a hydroponic solution labeled with isotopically labeled ammonium
 326 nitrate (intrusive approach, elevated nitrogen concentration; see ref. 60); and (ii) by
 327 applying nutrient solution to soil and allowing plant roots to take up nutrients *in situ*
 328 (non-intrusive, low nitrogen concentration; see ref. 61). The first approach allows an
 329 estimation of the maximum uptake rate of absorptive roots, while the second approach
 330 more accurately reflects the uptake rate of roots in natural conditions. We then

331 supplemented our dataset with 43 published measurements (keyword search for “root
 332 nitrogen uptake” using the Web of Science and Google Scholar). This dataset
 333 included 210 species-specific and 34 community-specific observations across major
 334 five biomes (Extended Data Table 4).

335 **Species and phylogeny.** Our root trait dataset represented a wide range of taxa,
 336 covering 210 genera and 79 families, with species names confirmed in The Plant List
 337 (<http://www.theplantlist.org>). We constructed the plant phylogenetic relationship
 338 using PHYLOCOM²⁷ (<http://phylodiversity.net/phyloomatic>) and determined the
 339 divergence time of plant families based on the earliest diverging genus within that
 340 family²⁸. We calculated Blomberg’s *K*-statistic⁶² using the “Picante” package in R
 341 and evaluated the strength of the phylogenetic signal for each traits; a large
 342 Blomberg’s *K* value is thought to indicate phylogenetic conservatism. We performed
 343 phylogenic independent contrasts analyses (PICs) to correct for shared evolutionary
 344 histories among traits and look for the pure effect of environmental influences
 345 (Extended Data Table 5).

346 **Statistical analyses.** Shapiro-Wilk tests revealed that all of our traits were
 347 significantly non-normal ($P < 0.05$), which we corrected by log10 transforming our
 348 data. We performed the Principal component analysis (PCA) using R package
 349 FactorR. The linear regression between root diameter vs. divergence time was
 350 weighted by the number of species within each family. We used linear regression to
 351 test the effect of root diameter and growth form (woody vs. herbaceous) on root
 352 mycorrhizal colonization, and tested equality of variance in root diameter among

353 biomes using Levene's test. We used linear mixed effects model to compare the
354 difference of root diameter across biomes. All statistical analyses were performed
355 using the R software, version 2.15.0.

356 **Code availability.** The R scripts used in Fig.1 and Fig.2 are available from the
357 corresponding author upon reasonable request.

358 **Data availability.** The data that support the findings of this study are available from
359 the corresponding authors upon request.

References

- 1 Kenrick, P. & Crane, P. R. The origin and early evolution of plants on land. *Nature* **389**, 33-39 (1997).
- 2 Field, K. J., Pressel, S., Duckett, J. G., Rimington, W. R. & Bidartondo, M. I. Symbiotic options for the conquest of land. *Trends Ecol. Evol.* **30**, 477-486 (2015).
- 3 Bardgett, R. D., Mommer, L. & De Vries, F. T. Going underground: root traits as drivers of ecosystem processes. *Trends Ecol. Evol.* **29**, 692-699 (2014).
- 4 Weemstra, M. *et al.* Towards a multidimensional root trait framework: a tree root review. *New Phytol.* **211**, 1159-1169 (2016).
- 5 Diaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167-171 (2016).
- 6 Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428**, 821-827, (2004).
- 7 Roumet, C. *et al.* Root structure-function relationships in 74 species: evidence of a root economics spectrum related to carbon economy. *New Phytol.* **210**, 815-826 (2016).
- 8 Pregitzer, K. S. *et al.* Fine root architecture of nine North American trees. *Ecol. Monogr.* **72**, 293-309 (2002).
- 9 McCormack, M. L. *et al.* Redefining fine roots improves understanding of below-ground contributions to terrestrial biosphere processes. *New Phytol.* **207**, 505-518 (2015).
- 10 Chen, W. L. *et al.* Root morphology and mycorrhizal symbioses together shape nutrient foraging strategies of temperate trees. *P. Natl. Acad. Sci. USA* **113**, 8741-8746 (2016).
- 11 Norby, R. J., Ledford, J., Reilly, C. D., Miller, N. E. & O'Neill, E. G. Fine-root production dominates response of a deciduous forest to atmospheric CO₂ enrichment. *P. Natl. Acad. Sci. USA* **101**, 9689-9693 (2004).
- 12 Jackson, R. B., Mooney, H. A. & Schulze, E. D. A global budget for fine root

- biomass, surface area, and nutrient contents. *P. Natl. Acad. Sci. USA* **94**, 7362-7366 (1997).
- 13 McCormack, M. L. *et al.* Building a better foundation: improving root-trait measurements to understand and model plant and ecosystem processes. *New Phytol.* **215**, 27-37 (2017).
- 14 Chen, W. L., Zeng, H., Eissenstat, D. M. & Guo, D. L. Variation of first-order root traits across climatic gradients and evolutionary trends in geological time. *Global Ecol. Biogeogr.* **22**, 846-856 (2013).
- 15 Guo, D. L. *et al.* Anatomical traits associated with absorption and mycorrhizal colonization are linked to root branch order in twenty-three Chinese temperate tree species. *New Phytol.* **180**, 673-683 (2008).
- 16 Reich, P. B. *et al.* Scaling of respiration to nitrogen in leaves, stems and roots of higher land plants. *Ecology Letters* **11**, 793-801 (2008).
- 17 Valverde-Barrantes, O. J., Freschet, G. T., Roumet, C. & Blackwood, C. B. A worldview of root traits: the influence of ancestry, growth form, climate and mycorrhizal association on the functional trait variation of fine-root tissues in seed plants. *New Phytologist* **215**, 1562-1573 (2017).
- 18 Iversen, C. M. *et al.* A global Fine-Root Ecology Database to address below-ground challenges in plant ecology. *New Phytol.* **215**, 15-26 (2017).
- 19 Kong, D. L. *et al.* Leading dimensions in absorptive root trait variation across 96 subtropical forest species. *New Phytol.* **203**, 863-872 (2014).
- 20 Feild, T. S. & Arens, N. C. Form, function and environments of the early angiosperms: merging extant phylogeny and ecophysiology with fossils. *New Phytol.* **166**, 383-408 (2005).
- 21 Comas, L. H. *et al.* Evolutionary Patterns and Biogeochemical Significance of Angiosperm Root Traits. *Int J Plant Sci* **173**, 584-595 (2012).
- 22 Ackerly, D. D. & Reich, P. B. Convergence and correlations among leaf size and function in seed plants: A comparative test using independent contrasts. *American Journal of Botany* **86**, 1272-1281 (1999).
- 23 Wing, S. L. *et al.* Late Paleocene fossils from the Cerrejón Formation, Colombia,

- 419 are the earliest record of Neotropical rainforest. *P. Natl. Acad. Sci. USA* **106**,
 420 18627-18632 (2009).
- 421 24 Woodward, F. I., Lomas, M. R. & Kelly, C. K. Global climate and the distribution
 422 of plant biomes. *Philos. T. Roy. Soc. B.* **359**, 1465-1476 (2004).
- 423 25 Eissenstat, D. M., Kucharski, J. M., Zadworny, M., Adams, T. S. & Koide, R. T.
 424 Linking root traits to nutrient foraging in arbuscular mycorrhizal trees in a
 425 temperate forest. *New Phytologist* **208**, 114-124 (2015).
- 426 26 Bremer, B. *et al.* An update of the angiosperm phylogeny group classification for
 427 the orders and families of flowering plants: APG III. *Bot. J. Linn. Soc.* **161**, 105-
 428 121 (2009).
- 429 27 Webb, C. O., Ackerly, D. D. & Kembel, S. W. Phylocom: software for the
 430 analysis of phylogenetic community structure and trait evolution. *Bioinformatics*
 431 **24**, 2098-2100 (2008).
- 432 28 Wikstrom, N., Savolainen, V. & Chase, M. W. Evolution of the angiosperms:
 433 calibrating the family tree. *P. Roy. Soc. B-Biol. Sci.* **268**, 2211-2220 (2001).
- 434 29 Li, H., Liu, B., McCormack, M. L., Ma, Z. & Guo, D. Diverse belowground
 435 resource strategies underlie plant species coexistence and spatial distribution.
 436 *New Phytol.*, doi:10.1111/nph.14710 (2017).
- 437 30 Hummel, I. *et al.* Relating root structure and anatomy to whole-plant functioning
 438 in 14 herbaceous Mediterranean species. *New Phytol.* **173**, 313-321 (2007).
- 439 31 Ostonen, I. *et al.* Fine root foraging strategies in Norway spruce forests across a
 440 European climate gradient. *Global Change. Biol.* **17**, 3620-3632 (2011).
- 441 32 Leppalammi-Kujansuu, J. *et al.* Fine root longevity and carbon input into soil
 442 from below- and aboveground litter in climatically contrasting forests. *Forest*
 443 *Ecol. Manag.* **326**, 79-90 (2014).
- 444 33 Withington, J. M., Reich, P. B., Oleksyn, J. & Eissenstat, D. M. Comparisons of
 445 structure and life span in roots and leaves among temperate trees. *Ecol. Monogr.*
 446 **76**, 381-397 (2006).
- 447 34 Sun, K., McCormack, M. L., Li, L., Ma, Z. Q. & Guo, D. L. Fast-cycling unit of
 448 root turnover in perennial herbaceous plants in a cold temperate ecosystem. *Sci.*

- 449 *Rep-Uk* **6**, doi:10.1038/srep19698 (2016).
- 450 35 Xia, M., Guo, D. & Pregitzer, K. S. Ephemeral root modules in *Fraxinus*
 451 *mandshurica*. *New Phytol.* **188**, 1065-1074 (2010).
- 452 36 McCormack, M. L., Adams, T. S., Smithwick, E. A. H. & Eissenstat, D. M.
 453 Predicting fine root lifespan from plant functional traits in temperate trees. *New*
 454 *Phytol.* **195**, 823-831 (2012).
- 455 37 Valenzuela-Estrada, L. R., Vera-Caraballo, V., Ruth, L. E. & Eissenstat, D. M.
 456 Root anatomy, morphology, and longevity among root orders in *Vaccinium*
 457 *Corymbosum* (Ericaceae). *Am. J. Bot.* **95**, 1506-1514 (2008).
- 458 38 Liu, B., He, J. X., Zeng, F. J., Lei, J. Q. & Arndt, S. K. Life span and structure of
 459 ephemeral root modules of different functional groups from a desert system. *New*
 460 *Phytol.* **211**, 103-112 (2016).
- 461 39 Hansson, K., Helmisaari, H.-S., Sah, S. P. & Lange, H. Fine root production and
 462 turnover of tree and understorey vegetation in Scots pine, silver birch and
 463 Norway spruce stands in SW Sweden. *Forest Ecol. Manag.* **309**, 58-65 (2013).
- 464 40 Huang, J. X. *et al.* Fine root longevity and controlling factors in subtropical
 465 *Altingia grilipes* and *Castanopsis carlesii* forests. *Acta Ecologica Sinica* **32**,
 466 1932-1942 (2012).
- 467 41 Yavitt, J. B., Harms, K. E., Garcia, M. N., Mirabello, M. J. & Wright, S. J. Soil
 468 fertility and fine root dynamics in response to 4 years of nutrient (N, P, K)
 469 fertilization in a lowland tropical moist forest, Panama. *Austral. Ecol.* **36**, 433-
 470 445 (2011).
- 471 42 Ling, H. *et al.* Influencing factors of fine root lifespans in two Chinese fir
 472 plantations in subtropical China. *Acta Ecologica Sinica* **31**, 1130-1138 (2011).
- 473 43 Stover, D. B., Day, F. P., Drake, B. G. & Hinkle, C. R. The long-term effects of
 474 CO₂ enrichment on fine root productivity, mortality, and survivorship in a scrub-
 475 oak ecosystem at Kennedy Space Center, Florida, USA. *Environ. Exp. Bot.* **69**,
 476 214-222 (2010).
- 477 44 Krasowski, M. J., Lavigne, M. B., Olesinski, J. & Bernier, P. Y. Advantages of
 478 long-term measurement of fine root demographics with a minirhizotron at two

479 balsam fir sites. *Can. J. Forest Res.* **40**, 1128-1135 (2010).

480 45 Huang, G., Zhao, X. Y., Zhao, H. L., Huang, Y. X. & Zuo, X. A. Linking root
481 morphology, longevity and function to root branch order: a case study in three
482 shrubs. *Plant Soil* **336**, 197-208 (2010).

483 46 Girardin, C. A. J. *et al.* Net primary productivity allocation and cycling of carbon
484 along a tropical forest elevational transect in the Peruvian Andes. *Global Change*
485 *Biol.* **16**, 3176-3192 (2010).

486 47 Espeleta, J. F., West, J. B. & Donovan, L. A. Tree species fine-root demography
487 parallels habitat specialization across a sandhill soil resource gradient. *Ecology*
488 **90**, 1773-1787 (2009).

489 48 Vargas, R. & Allen, M. F. Dynamics of fine root, fungal rhizomorphs, and soil
490 respiration in a mixed temperate forest: integrating sensors and observations.
491 *Vadose Zone J.* **7**, 1055-1064 (2008).

492 49 Graefe, S., Hertel, D. & Leuschner, C. Fine root dynamics along a 2,000-m
493 elevation transect in South Ecuadorian mountain rainforests. *Plant Soil* **313**, 155-
494 166 (2008).

495 50 Green, I. J., Dawson, L. A., Proctor, J., Duff, E. I. & Elston, D. A. Fine root
496 dynamics in a tropical rain forest is influenced by rainfall. *Plant Soil* **276**, 23-32
497 (2005).

498 51 Baddeley, J. A. & Watson, C. A. Influences of root diameter, tree age, soil depth
499 and season on fine root survivorship in *Prunus avium*. *Plant Soil* **276**, 15-22
500 (2005).

501 52 Kern, C. C., Friend, A. L., Johnson, J. M. F. & Coleman, M. D. Fine root
502 dynamics in a developing *Populus deltoides* plantation. *Tree Physiol.* **24**, 651-660
503 (2004).

504 53 Wells, C. E., Glenn, D. M. & Eissenstat, D. M. Changes in the risk of fine-root
505 mortality with age: a case study in peach, *Prunus persica* (Rosaceae). *Am. J. Bot.*
506 **89**, 79-87 (2002).

507 54 King, J. S. *et al.* Below - ground carbon input to soil is controlled by nutrient
508 availability and fine root dynamics in loblolly pine. *New Phytol.* **154**, 389-398

(2002).

55 Gill, R. A., Burke, I. C., Lauenroth, W. K. & Milchunas, D. G. Longevity and turnover of roots in the shortgrass steppe: influence of diameter and depth. *Plant Ecol.* **159**, 241-251 (2002).

56 Wells, C. E. & Eissenstat, D. M. Marked differences in survivorship among apple roots of different diameters. *Ecology* **82**, 882-892 (2001).

57 Tierney, G. L. & Fahey, T. J. Evaluating minirhizotron estimates of fine root longevity and production in the forest floor of a temperate broadleaf forest. *Plant Soil* **229**, 167-176 (2001).

58 Majdi, H., Damm, E. & Nylund, J. E. Longevity of mycorrhizal roots depends on branching order and nutrient availability. *New Phytol.* **150**, 195-202 (2001).

59 Coleman, M. D., Dickson, R. E. & Isebrands, J. G. Contrasting fine-root production, survival and soil CO₂ efflux in pine and poplar plantations. *Plant Soil* **225**, 129-139 (2000).

60 Chapin, F. S., Moilanen, L. & Kielland, K. Preferential use of organic nitrogen for growth by a nonmycorrhizal Arctic sedge. *Nature* **361**, 150-153 (1993).

61 McKane, R. B. *et al.* Resource-based niches provide a basis for plant species diversity and dominance in arctic tundra. *Nature* **415**, 68-71 (2002).

62 Blomberg, S. P., Garland, T. & Ives, A. R. Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution* **57**, 717-745 (2003).

63 Aanderud, Z. T. & Bledsoe, C. S. Preferences for ¹⁵N-ammonium, ¹⁵N-nitrate, and ¹⁵N-glycine differ among dominant exotic and subordinate native grasses from a California oak woodland. *Environ. Exp. Bot.* **65**, 205-209 (2009).

64 Andersen, K. M. & Turner, B. L. Preferences or plasticity in nitrogen acquisition by understorey palms in a tropical montane forest. *J. Ecol.* **101**, 819-825 (2013).

65 Andresen, L. C., Michelsen, A., Jonasson, S. & Strom, L. Seasonal changes in nitrogen availability, and root and microbial uptake of ¹⁵N¹³C₉-phenylalanine and ¹⁵N-ammonium in situ at a temperate heath. *Appl. Soil Ecol.* **51**, 94-101 (2011).

66 Averill, C. & Finzi, A. C. Increasing plant use of organic nitrogen with elevation

538 is reflected in nitrogen uptake rates and ecosystem $\delta^{15}\text{N}$. *Ecology* **92**, 883-891
539 (2011).

540 67 Bardgett, R. D., Streeter, T. C. & Bol, R. Soil microbes compete effectively with
541 plants for organic-nitrogen inputs to temperate grasslands. *Ecology* **84**, 1277-
542 1287 (2003).

543 68 Boczulak, S. A., Hawkins, B. J. & Roy, R. Temperature effects on nitrogen form
544 uptake by seedling roots of three contrasting conifers. *Tree Physiol.* **34**, 513-523
545 (2014).

546 69 Cheng, X. M. & Bledsoe, C. S. Competition for inorganic and organic N by blue
547 oak (*Quercus douglasii*) seedlings, an annual grass, and soil microorganisms in a
548 pot study. *Soil Biol. Biochem.* **36**, 135-144 (2004).

549 70 Dunn, R. M., Mikola, J., Bol, R. & Bardgett, R. D. Influence of microbial activity
550 on plant-microbial competition for organic and inorganic nitrogen. *Plant Soil*
551 **289**, 321-334 (2006).

552 71 Finzi, A. C. & Berthrong, S. T. The uptake of amino acids by microbes and trees
553 in three cold-temperate forests. *Ecology* **86**, 3345-3353 (2005).

554 72 Gallet-Budynek, A. *et al.* Intact amino acid uptake by northern hardwood and
555 conifer trees. *Oecologia* **160**, 129-138 (2009).

556 73 Harrison, K. A., Bol, R. & Bardgett, R. D. Preferences for different nitrogen
557 forms by coexisting plant species and soil microbes. *Ecology* **88**, 989-999 (2007).

558 74 Henry, H. A. L. & Jefferies, R. L. Interactions in the uptake of amino acids,
559 ammonium and nitrate ions in the Arctic salt-marsh grass, *Puccinellia*
560 phryganeodes. *Plant Cell Environ.* **26**, 419-428 (2003).

561 75 Jin, V. L. & Evans, R. D. Microbial ^{13}C utilization patterns via stable isotope
562 probing of phospholipid biomarkers in Mojave Desert soils exposed to ambient
563 and elevated atmospheric CO_2 . *Global Change Biol.* **16**, 2334-2344 (2010).

564 76 Jin, V. L., Romanek, C. S., Donovan, L. A. & Sharitz, R. R. Soil nitrogen
565 availability and in situ nitrogen uptake by *Acer rubrum* L. and *Pinus palustris*
566 Mill. in the southeastern U. S. Coastal Plain. *J. Torrey Bot. Soc.* **137**, 339-347
567 (2010).

- 568 77 Kahmen, A., Livesley, S. J. & Arndt, S. K. High potential, but low actual, glycine
569 uptake of dominant plant species in three Australian land-use types with
570 intermediate N availability. *Plant Soil* **325**, 109-121 (2009).
- 571 78 Kastovska, E. & Santruckova, H. Comparison of uptake of different N forms by
572 soil microorganisms and two wet-grassland plants: a pot study. *Soil Biol.*
573 *Biochem.* **43**, 1285-1291 (2011).
- 574 79 Li, C. C. *et al.* Inorganic and organic nitrogen uptake by nine dominant
575 subtropical tree species. *Iforest* **9**, 253-258 (2015).
- 576 80 McFarland, J. W. *et al.* Cross-ecosystem comparisons of in situ plant uptake of
577 amino acid-N and NH_4^+ . *Ecosystems* **13**, 177-193 (2010).
- 578 81 Metcalfe, R. J., Nault, J. & Hawkins, B. J. Adaptations to nitrogen form:
579 comparing inorganic nitrogen and amino acid availability and uptake by four
580 temperate forest plants. *Can. J. Forest Res.* **41**, 1626-1637 (2011).
- 581 82 Mozdzer, T. J., Zieman, J. C. & McGlathery, K. J. Nitrogen uptake by native and
582 invasive temperate coastal macrophytes: importance of dissolved organic
583 nitrogen. *Estuar Coast* **33**, 784-797 (2010).
- 584 83 Nie, M. *et al.* Plants' use of different nitrogen forms in response to crude oil
585 contamination. *Environ. Pollut.* **159**, 157-163 (2011).
- 586 84 Nordin, A., Hogberg, P. & Nasholm, T. Soil nitrogen form and plant nitrogen
587 uptake along a boreal forest productivity gradient. *Oecologia* **129**, 125-132
588 (2001).
- 589 85 Öhlund, J. & Näsholm, T. Growth of conifer seedlings on organic and inorganic
590 nitrogen sources. *Tree Physiol.* **21**, 1319-1326 (2001).
- 591 86 Ouyang, S. *et al.* Nitrogen competition between three dominant plant species and
592 microbes in a temperate grassland. *Plant Soil* **408**, 121-132, (2016).
- 593 87 Paulding, E. M., Baker, A. J. M. & Warren, C. R. Competition for nitrogen by
594 three sympatric species of Eucalyptus. *Ann. For. Sci.* **67**, 406-406 (2010).
- 595 88 Persson, J. *et al.* Nitrogen acquisition from inorganic and organic sources by
596 boreal forest plants in the field. *Oecologia* **137**, 252-257 (2003).
- 597 89 Persson, J. & Näsholm, T. Regulation of amino acid uptake in conifers by

- exogenous and endogenous nitrogen. *Planta* **215**, 639-644 (2002).
- 90 Pfautsch, S., Rennenberg, H., Bell, T. L. & Adams, M. A. Nitrogen uptake by
Eucalyptus regnans and *Acacia* spp. preferences, resource overlap and energetic
costs. *Tree Physiol.* **29**, 389-399 (2009).
- 91 Rains, K. C. & Bledsoe, C. S. Rapid uptake of ^{15}N -ammonium and glycine- ^{13}C ,
 ^{15}N by arbuscular and ericoid mycorrhizal plants native to a Northern California
coastal pygmy forest. *Soil Biol. Biochem.* **39**, 1078-1086 (2007).
- 92 Schmidt, S. & Stewart, G. R. Waterlogging and fire impacts on nitrogen
availability and utilization in a subtropical wet heathland (wallum). *Plant Cell
Environ.* **20**, 1231-1241 (1997).
- 93 Schmidt, S. & Stewart, G. R. Glycine metabolism by plant roots and its
occurrence in Australian plant communities. *Aust. J. Plant Physiol.* **26**, 253-264
(1999).
- 94 Scott, E. E. & Rothstein, D. E. Amino acid uptake by temperate tree species
characteristic of low- and high-fertility habitats. *Oecologia* **167**, 547-557 (2011).
- 95 Simon, J. *et al.* Competition for nitrogen between adult European beech and its
offspring is reduced by avoidance strategy. *Forest Ecol. Manag.* **262**, 105-114
(2011).
- 96 Simon, J., Waldhecker, P., Bruggemann, N. & Rennenberg, H. Competition for
nitrogen sources between European beech (*Fagus sylvatica*) and sycamore maple
(*Acer pseudoplatanus*) seedlings. *Plant Biol.* **12**, 453-458 (2010).
- 97 Stoelken, G., Simon, J., Ehling, B. & Rennenberg, H. The presence of amino
acids affects inorganic N uptake in non-mycorrhizal seedlings of European beech
(*Fagus sylvatica*). *Tree Physiol.* **30**, 1118-1128 (2010).
- 98 Wallander, H., Arnebrant, K., Ostrand, F. & Karen, O. Uptake of N^{15} -labelled
alanine, ammonium and nitrate in *Pinus sylvestris* L. ectomycorrhiza growing in
forest soil treated with nitrogen, sulphur or lime. *Plant Soil* **195**, 329-338 (1997).
- 99 Wanek, W., Arndt, S. K., Huber, W. & Popp, M. Nitrogen nutrition during
ontogeny of hemiepiphytic *Clusia* species. *Funct. Plant Biol.* **29**, 733-740 (2002).
- 100 Warren, C. R. Potential organic and inorganic N uptake by six *Eucalyptus*

species. *Funct. Plant Biol.* **33**, 653-660 (2006).

101 Warren, C. R. Does nitrogen concentration affect relative uptake rates of nitrate,
ammonium, and glycine? *J. Plant Nutr. Soil Sc.* **172**, 224-229 (2009).

102 Warren, C. R. & Adams, P. R. Uptake of nitrate, ammonium and glycine by plants
of Tasmanian wet eucalypt forests. *Tree Physiol.* **27**, 413-419 (2007).

103 Wei, L. L., Chen, C. R. & Yu, S. Uptake of organic nitrogen and preference for
inorganic nitrogen by two Australian native Araucariaceae species. *Plant Ecol.*
Divers. **8**, 259-264 (2015).

104 Weigelt, A., Bol, R. & Bardgett, R. D. Preferential uptake of soil nitrogen forms
by grassland plant species. *Oecologia* **142**, 627-635 (2005).

105 Wu, J. R. *et al.* Mycorrhizas alter nitrogen acquisition by the terrestrial orchid
Cymbidium goeringii. *Ann. Bot-London* **111**, 1181-1187 (2013).

Supplementary Information is linked to the online version of the paper at
www.nature.com/nature.

Acknowledgments We thank C. Ma, L. Li, Y. Yue, M. Liu, F. Ma, H. Li, D. Kong,
B. Liu, K. Sun for collecting data, X. Liu, X. Deng for their assistance in field
sampling. We also thank all field research stations of the Chinese Academy of
Sciences for their warm support. This study was funded by the Natural Science
Foundation of China (NSFC Grants 31325006, 31530011, and 41571130041).

Author Contributions Z.M., D.G. and L.O.H. developed the overall conceptual
approach and analysis. Z.M. compiled and analyzed the data, and X.X. provided
nitrogen uptake rate data. R.D.B., D.M.E. and M.L.M. contributed to the formulation
of research questions and interpretations. Z.M., D.G, M.L and L.O.H. wrote the

652 paper. Z.M., D.G. and L.O.H contributed equally to this work and all authors
653 contributed to revisions.

654 **Author Information** Reprints and permissions information is available at
655 www.nature.com/reprints. The authors declare no competing financial interests.

656 Readers are welcome to comment on the online version of the paper. Correspondence
657 and requests for materials should be addressed to D.G. (guodl@igsnrr.ac.cn) or
658 L.O.H. (lhedin@princeton).

Figure 1 | Root trait dimensions organized by root diameter and growth form.

At the species level, the diameter of first-order roots is inversely correlated with specific root length (*SRL*) (**a**), positively related to the evolutionary time of divergence of major taxonomic groups (**b**), and positively related to the length of root (in %) colonized by mycorrhizal fungi (**c**). Moreover, root tissue density (RTD) differs across plant growth form (**d**), with herbaceous plants (green points) displaying more constrained variation than woody plants (brown points)(F-test, $P < 0.001$; note logarithmic scale on Y-axis). The solid red line in (**a**) identifies the relationship (*SI* note 1) between specific root length and root diameter assuming a root tissue density of 0.25 g cm^{-3} ; dashed red lines identify tissue densities of 0.1 and 1.0 g cm^{-3} (upper vs. lower line, respectively). We used a linear regression weighted by number of species in panel (**b**) and a linear regression with woody and non-woody growth forms as categorical variables in panel (**c**). Root cross-section images in (**a**) are from the low-density grass *Agropyron cristatum* (upper right) and the high-density woody shrub *Rhaphiolepis indica* (lower left), as discussed in main text.

Figure 2 | Density distributions of first-order root diameter across seven biomes.

The variance in root diameter declines from biomes with equable conditions (*e.g.*, tropical forests) to biomes with pronounced seasonality in soil resource supplies (*e.g.*, deserts). Numbers in brackets identify species-specific observations, and letters identify significant pairwise differences in a Levene's variance test (Extended Data Table 3). Woody biomes are identified as shades of tan to yellow and non-woody biomes as shades of green.