Platinum-group element remobilisation and concentration in the Cliff chromitites of the Shetland Ophiolite Complex, Scotland

DOI:

[10.1180/minmag.2017.081.108](https://doi.org/10.1180/minmag.2017.081.108)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

O'Driscoll, B., Garwood, R., Day, J. M. D., & Wogelius, R. A. (2018). Platinum-group element remobilisation and concentration in the Cliff chromitites of the Shetland Ophiolite Complex, Scotland. *Mineralogical Magazine*, 82(3), 471-490. <https://doi.org/10.1180/minmag.2017.081.108>

Published in:

Mineralogical Magazine

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Platinum-group element remobilisation and concentration in the Cliff chromitites of the Shetland Ophiolite Complex, Scotland

Brian O'Driscoll¹, Russell Garwood^{1,2}, James M.D. Day³, Roy Wogelius¹

¹School of Earth and Environmental Sciences, Oxford Road, University of Manchester, M13 9PL, UK

²Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

³Geosciences Research Division, Scripps Institution of Oceanography, UCSD, La Jolla, CA 92093-0244, USA

Abstract

The ~492 Ma Shetland Ophiolite Complex (SOC) contains an extensive mantle section, within which numerous podiform chromitite bodies formed during melt percolation in a supra-subduction zone setting. One of the SOC chromitite localities has an unusual style of platinum-group element (PGE) mineralisation. Specifically, the Cliff chromitite suite has relatively high (>250 ppm) Pt plus Pd, compared to other SOC chromitites. In this study, we apply petrographic observation, mineral chemistry and novel X-ray microtomography data to elucidate the petrogenesis of PGE-bearing phases at Cliff. The combined datasets reveal that the PGE at Cliff have likely been fractionated by an As-rich fluid, concentrating Pt and Ir into visible (0.1-1 µm) platinum-group minerals (PGM) such as sperrylite and irarsite, respectively. The high (>1 ppm) bulk rock concentrations of the other PGE (e.g., Os) in the Cliff chromitites suggests the presence of abundant fine-grained unidentified PGM in the serpentinised groundmass. The spatial association of arsenide phases and PGM with alteration rims on Cr-spinel grains suggests that the high Pt and Pd abundances at Cliff result from a late-stage low-temperature (e.g., 200-300°C) hydrothermal event. This conclusion highlights the potential effects that secondary alteration processes can have on modifying and upgrading the tenor of PGE deposits.

Introduction

The platinum-group metals are high-value industrial commodities that concentrate naturally in select geological environments. In particular, ore deposits of the platinum-group elements (PGE: Os, Ir, Ru, Rh, Pt, Pd) are known from layered mafic-ultramafic intrusions and from the mantle sections of supra-subduction zone ophiolites. In many - but not all - of these occurrences, the PGE are hosted in chromitite seams (>60 vol.% Cr-spinel). A significant body of work demonstrates the petrogenetic link between the concentration of the PGE and the formation of chromitite (e.g., Irvine 1977; Melcher et al. 1997; O'Driscoll et al. 2009; González-Jiménez et al. 2014a, 2014b; Latypov

et al. 2013; Godel 2015). Layered intrusion-bearing ‘stratiform’ chromitite seams are associated with two of the largest deposits (reefs) of the PGE on Earth: the Merensky Reef and the UG-2 chromitite, both situated in the Critical Zone of the ~2 Ga Bushveld Complex (South Africa; cf. Schouwstra et al. 2000). However, podiform chromitites hosted in ophiolitic mantle may also contain significant enrichments of the PGE, up to economically viable grades (>20 ppm Pt, Pd). In contrast to layered intrusion-hosted PGE deposits which typically show high PPGE (Pt, Pd)/IPGE (Os, Ir, Ru, Rh) ratios, ophiolite PGE-rich deposits typically have low PPGE/IPGE (e.g., Page and Talkington 1984, Oshin and Crocket 1986, Melcher et al. 1997, O’Driscoll and González-Jiménez 2016; **Fig. 1**).

The mechanism(s) by which the PGE concentrate in the upper mantle, and the relationship between the petrological processes that form chromitite and the platinum-group minerals (PGM), are poorly constrained. In the majority of cases, base-metal sulphides are thought to play an important role by scavenging the PGE in upper mantle chromitite-forming magma conduits (cf. Holwell and McDonald 2010). This is because the PGE exhibit strongly chalcophile tendencies in terrestrial magmatic systems (i.e., partition coefficients up to 10^6 ; Mungall and Brenan, 2014). However, evidence suggests that within cooling magmatic systems, the PGE only exist transiently in solid-solution in base-metal sulphides, and the PGM form during a complex down-temperature crystallisation process. Based on experimental data from the Fe-Ni-Cu-S system that have been used to examine PGE partitioning behaviour, Holwell and McDonald (2010) presented a model that proposed the partitioning of Os, Ir and Ru into Fe-Ni-S monosulphide solution (mss) that crystallised from base-metal sulphide liquid at high temperature (~1000°C). Platinum and Pd concentrate in the Cu-rich liquid fraction that remains (referred to as intermediate solid solution; iss), with As, Bi, Te. Unmixing of the iss leads to the crystallisation of the platinum-group minerals (PGM), which are a diverse group of (~135) alloys, sulphides, arsenides and other phases, that form as accessory minerals in these systems (O’Driscoll and González-Jiménez 2016).

64 Critical gaps remain in our understanding of PGM petrogenesis. In particular, the degree to
65 which low-temperature processes may modify the distribution of the PGE and form new PGM as
66 subsolidus reaction products is unknown (cf. Hanley 2005). Some authors have suggested that
67 desulphurisation may be a relevant process (e.g., Godel and Barnes 2008). Others have argued more
68 generally for metasomatism as an important mechanism - if not in the primary concentration of the
69 PGE in a rock, then through the dissolution, mobilisation and reprecipitation of the PGM. For
70 example, Godel and Barnes (2008) highlight that PGE mobilisation has occurred in sulphides of the
71 ~2.7 Ga JM Reef (Stillwater Complex), during the Laramide Orogeny at 70-80 Ma. Similarly,
72 studies carried out on the Platinova Reef of the Skaergaard suggest that its PGM assemblage
73 equilibrated at temperatures as low as 300°C (Karup-Møller et al. 2008). Coggon et al. (2011)
74 inferred that PGM compositions in the Merensky Reef were modified by a hydrothermal fluid after
75 the crystallisation age of the reef (the latter age is constrained by U-Pb in zircon). Thus, the whole
76 rock budget of the PGE can potentially be modified and enhanced during metasomatism, so there
77 are important economic implications for a better understanding of relatively low-temperature, as
78 well as magmatic, ore-forming processes. Although ophiolite chromitites are not a significant
79 contributor to the global PGE supply, these deposits offer different and potentially valuable
80 perspectives on the petrogenesis of the PGM to that gained from the study of mid-upper crustal
81 layered intrusions.

82 In this study, an anomalous upper mantle chromitite deposit from the Shetland Ophiolite
83 Complex (SOC; **Fig. 2**) in northern Scotland was targeted to further interrogate the petrogenesis of
84 PGM formation in upper mantle environments. The chromitite, from the locality of Cliff on the
85 island of Unst (**Fig. 2**), is known for its exceptionally high levels of enrichment in the PPGE (~250
86 ppm Pt+Pd, O'Driscoll et al. 2012; see also Prichard and Lord 1993). Unusually for ophiolite
87 chromitite deposits, the PPGE are enriched by an order of magnitude over the IPGE (**Fig. 1**) such
88 that the distribution of PGE overall is more similar to that from layered intrusions. Another
89 intriguing feature of the Cliff PGE mineralisation is that it is associated with As-rich phases; there

is little or no base metal sulphide in these rocks. Here, we combine X-ray microtomography (μ CT) with petrography and mineral chemistry to visualise the 3D distribution and microstructure of the base- and precious metal-rich phases and link this to their chemical signatures. Establishing the microstructural relationships between Cr-spinel, base-metal phases and the PGM in chromitite is pivotal to understanding the petrogenesis of these deposits; distinguishing between these high density phases is an objective particularly well-addressed by μ CT (e.g., Ketchum and Carlson 2001; Godel et al. 2010; Godel et al. 2013; Godel, 2015; Prichard et al. 2017). The key goal in this study is to elucidate the distribution of the PGE between specific carrier phases and link this to the chromitite microstructure and mineral chemistry to formulate a comprehensive petrogenetic interpretation.

Geological setting

The SOC is a ~492 Ma ophiolite (Prichard 1985; Spray and Dunning 1991), thought to have been obducted onto Laurentia at ~470 Ma (Flinn et al. 1991; Chew et al. 2010). Although overlying components of the ophiolite are missing or not well-exposed (e.g., the sheeted dyke complex) on the islands of Unst or Fetlar, the mantle section is extensive and well-preserved (Derbyshire et al., 2013). Notably, a portion of mantle peridotite associated with pillow lavas, sediments and cross-cutting dykes has been recognised on mainland Shetland. This has been linked to the SOC to the north by gravity and aeromagnetic anomalies, possibly representing an extensive sliver of Iapetus Ocean mantle (Day et al., 2017). At Unst and Fetlar, the mantle section predominantly comprises serpentinised harzburgite, with numerous dunite layers and lenses, and less abundant chromitites and pyroxenites (**Fig. 2**; Flinn 2001). Chromitite seams occur encased in dunite sheets and pods; a relationship that has previously been invoked to explain the formation of chromitite due to high degrees of melt-rock reaction in zones (channels) of focused melt flow (O'Driscoll et al. 2012; Derbyshire et al. 2013). This melt percolation is likely to have occurred in a mantle wedge

115 environment, related to subduction of Iapetus lithosphere in a forearc setting (Crowley and Strachan
116 2015).

117 The petrological Moho is well-exposed on Unst, and the Cliff chromitite suite lies ≤ 1 km
118 stratigraphically below its position, close to the trace of a structure (to the west) that probably
119 excised the ophiolite sole thrust (Cutts et al. 2011). Harzburgites that host the Cliff chromitites yield
120 mantle-like $^{187}\text{Os}/^{188}\text{Os}$ initial isotopic compositions at 492 Ma (O'Driscoll et al. 2012). The Cliff
121 chromitites are between 0.1-2 m thick and are structurally complex, occurring as stock-work veins
122 and laterally discontinuous pods and lenses, mostly oriented sub-parallel to the petrological Moho
123 (e.g., Derbyshire et al. 2013). We sampled material from pits dug within the Cliff chromitite suite in
124 the 1800s, which were used for producing yellow pigment from the chromitite.

125

126 **Analytical techniques**

127 *Mineral chemistry*

128 A JEOL JXA-8900RL electron microprobe at the University of Göttingen (Germany) was
129 employed to make the back-scattered electron images of the Cliff chromitites, using a 20.0 kV
130 acceleration voltage and a 20 nA beam current. The instrument settings and protocols have
131 otherwise been previously outlined in Derbyshire et al. (2013), but the details of the arsenide and
132 PGM measurements are summarised here. For the arsenide analyses, peak and background count
133 times were 15 s and 5 s, respectively, for S, Fe and Zn, 30 s and 15 s peak and background,
134 respectively, for As, Cu, Ni, Pb, Sb and Co, and 60 s and 30 s, respectively, for Se and Te. A range
135 of standards including ZnS, AsGa, galena, gahnite and native metals and semi-metals Te, Cu, Ni,
136 Se, Sb, Fe and Co. For the PGM, count times of 15 s and 5 s were used for S, As, Fe, Si, Ni,
137 respectively; 30 s and 15 s for Sb, Cu, Co, Ru, Ir, Rh and Pt, respectively; 60 s and 30 s for Pd and
138 Os, respectively. The standards used for the PGM analyses include ZnS, AsGa, wollastonite and a
139 range of native metals for Sb, Fe, Cu, Co, Ru, Pd, Ni, Ir, Rh, Os and Pt calibration. Detection limits

140 for PGE in the PGM are as follows: Ir: 592 ppm, Os: 2196 ppm, Ru: 237 ppm, Rh: 276 ppm, Pd:
141 276 ppm, Pt: 649 ppm (Derbyshire et al. 2013).

142 The laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) data were
143 collected at the time of the analytical campaign reported in O'Driscoll et al. (2012). In the Cliff
144 samples, LA-ICP-MS analyses were carried out on Cr-spinel, arsenide and across host serpentinitised
145 matrix material (but not PGM, which are typically too small to be identified). The measurements
146 were performed at the University of Maryland (USA) using a *New Wave* 213 nm wavelength laser-
147 ablation system attached to a *Thermo Fisher* Element 2 ICP-MS. Trace-element abundances were
148 determined on individual spots and raster paths using a variable laser beam diameter (6-55 μm) and
149 a laser repetition rate of 7 Hz, at 1.4 to 2.5 J/cm². Argide and Ni/Cu interferences were monitored
150 during analysis and were found to be negligible. Ablation was conducted under a helium
151 atmosphere to enhance production of fine aerosols and the ablated sample was mixed with argon
152 prior to transport to the ICP-MS. Each analysis consisted of 60 s of data collection. Backgrounds
153 on the ICP-MS sample gas were collected for approximately 20 s followed by approximately 40 s
154 of laser ablation of the sample. Data were collected in time-resolved mode so that effects of
155 inclusions, mineral zoning, and possible penetration of the laser beam to underlying phases could be
156 evaluated for each analysis. Plots of counts per second versus time were examined for each
157 analysis and integration intervals for the gas background and the sample analysis were selected
158 manually. Each LA-ICP-MS analysis was normalised to a major-element oxide (typically Fe),
159 measured by electron microprobe, as an internal standard to account for variable ablation yield.
160 This internal normalisation is supported by the electron microprobe data, which allowed no more
161 than ~4% relative difference in Ni or Cr content over a 50 μm spot and the LA-ICP-MS time-
162 resolved patterns having shown that the ablated volumes were generally homogeneous. For all
163 data, the NIST 610 glass standard was used for calibration of relative element sensitivities, using
164 values from a compiled database of preferred values from GEOREM. Replicate LA-ICP-MS
165 analyses of the University of Toronto JB Sulphide standard run at intervals during the analytical

166 session, yielded an external precision of 1% (2σ relative standard deviation) for all trace- and
167 major-element compositions.

168

169 *3D image analysis by X-ray microtomography*

170 Hand specimen rock samples were investigated with laboratory-based microtomography in the
171 Manchester X-ray Imaging Facility (MXIF, University of Manchester). Scans were conducted using
172 a Nikon Metris 225/320 kV X-ray CT, within a customised bay, and with a 2000x2000 Perkin
173 Elmer 1621-16-bit amorphous silicon flat-panel detector. All scans used 3142 projections, a
174 tungsten reflection target, with the following variable settings:

- 175 • 1000 ms exposure, unfiltered source at 115kV and 60 μ A, giving a voxel size of 32.8 μ m.
- 176 • 1000 ms exposure, source at 200 kV and 75 μ A with a 0.5mm copper filter, giving a voxel
177 size of 35.2 μ m.
- 178 • 708 ms exposure, source at 130 kV and 120 μ A with a 0.5mm copper filter, giving a voxel
179 size of 45.1 μ m.

180 In order to interrogate the distribution of the small volumes of arsenide and PGM inclusions,
181 higher resolution data were collected at the Diamond Light Source (DLS) synchrotron (UK). The
182 high flux - in part achieved by using a pink beam - of beamline I13 facilitated successful scans at
183 1.1 micron voxel size, despite relatively X-ray dense samples. Approximately 1 mm diameter cores
184 were drilled from hand specimens. These were mounted using glue to the tip of the metal pin on a
185 magnetic sample holder. Specimens were scanned using a pink beam, with a 3 mm Al filter to
186 reduce the flux of lower energy X-rays, and hence minimise beam hardening (the undulator gap was
187 reduced to a point where there were sufficient counts through the densest parts of a specimen). A
188 PCO4000 camera with scientific grade 4008×2672 pixel CCD sensor was used to collect 1801 0.3
189 second projections on a 4 $\times$ objective for lower resolution scans (providing voxel sizes of 1.1 μ m),
190 and 3600 0.5 second projections using a 10 $\times$ objective (providing voxel sizes 0.45 μ m respectively).
191 Further scans were collected with a 28 keV monochromatic source, and 10 $\times$ objective, reduced Al

192 filter (0.2mm thickness), and 2 second exposure to collect data without beam hardening. All
193 projections employed an X-ray sensitive scintillator emitting visible light (cadmium tungstate), and
194 volumes were reconstructed using in-house filtered back projection reconstruction algorithm
195 (Titarenko et al. 2010).

196 Volumes were visualised using both volume and surface rendering. For both forms of
197 visualisation, tiff stacks (DLS) or VGI/VOL files (MXIF) were loaded using Drishti import, the
198 histograms examined and correctly windowed prior to conversion to 8 bit grayscale data. These
199 were either exported as Drishti volumes (.pvl.nc), or as PNG stacks for the SPIERS software suite.
200 Volume rendering was conducted in Drishti (Limaye 2012), which can render data based on a two
201 dimensional (value-gradient magnitude) histogram. A transfer function was used to select, and then
202 render, different regions of this histogram in three dimensions: for this work, different phases were
203 selected using multiple transfer functions - typically to pick out the rock exterior, oxide phases of
204 medium X-ray attenuation, as well as the densest arsenide and PGM inclusions. These transfer
205 functions were copied between datasets and modified as required. Visualisations were also
206 conducted in SPIERS following the methods of Garwood and Sutton (2010) and Sutton et al.
207 (2012), and subsequently rendered using the open source Raytracer Blender (see Garwood and
208 Dunlop 2014).

209

210 **Results**

211 *Petrographic observations*

212 Aspects of the petrography and mineralogy of the Cliff chromitites have been previously described
213 by Prichard and Tarkian (1988), Prichard et al. (1994) and Derbyshire et al. (2013), and we focus
214 here on new observations that bear directly on the microstructural distribution of Ni-arsenide and
215 PGM. The Cliff chromitites contain 50-90 vol.% Cr-spinel, in a groundmass composed
216 predominantly of lizardite (**Figs. 3a, 3b**). The Cr-spinel forms subhedral-anhedral crystals and
217 aggregates of crystals that are heavily fractured (**Fig. 4a**). Many of the larger fractures continue into

218 the groundmass serpentinite (**Fig. 3c**). Chrome-spinel crystal edges and the interiors of intra-Cr-
219 spinel fractures are typically lined with selvages of sieve-textured ferrian chromite (although some
220 of this material is also likely to be non-stoichiometric ferritchromite, as noted by Derbyshire et al.
221 2013), with extensive further (exterior) rimming of ferrian chromite rims by a chloritic mineral
222 (**Figs. 3a, 3b**). The chlorite-like rims are typically about 10-20 μm thick, and in addition to
223 occurring on grain edges, they line serpentinitised inclusions within Cr-spinel grains (e.g., **Fig. 3b**)
224 and fill intra-Cr-spinel fractures (**Fig. 3b**). The Fe-rich rims and ferrian chromite deposited within
225 intra-crystal fractures are compositionally heterogeneous (composite) (**Fig. 4b**), suggesting that the
226 alteration of primary Cr-spinel occurred in multiple stages. Base-metal sulphides are rare to
227 completely absent in Cliff chromitite samples.

228 Nickel-arsenide is present in accessory abundances, with grain sizes that are typically 20 to
229 50 μm in diameter. Grains of Ni-arsenide $<5 \mu\text{m}$ in size are not common. The Ni-arsenides are
230 consistently situated close to Cr-spinel grain boundaries and within intra-Cr-spinel fractures and
231 alteration zones (**Figs. 4b, 4c**), often in small clusters or linear trains of crystals comprising less
232 than 10 visible grains (e.g., **Fig. 4c**). Thus, there is a close textural association between Ni-arsenide
233 and the ferrian chromite. Isolated larger grains of Ni-arsenide are also less commonly observed
234 within the groundmass (**Fig. 4a**). A critically important microstructural observation is that the
235 visible PGM in the Cliff chromitite are spatially correlated with the Ni-arsenide and ferrian
236 chromite (**Figs. 4d, 4e, 4f**). The PGM are typically $\leq 5 \mu\text{m}$ in diameter and EDS analysis suggests
237 that many of the 'bright' phases analysed below this size threshold are PGM. However, the majority
238 of grains are much smaller than this, meaning that quantitative determination of their compositions
239 using the electron microprobe is not possible. The PGM observed may occur either as inclusions
240 within Ni-arsenides, attached to the edges of Ni-arsenides (e.g., **Fig. 4f**), or in close proximity to
241 them.

242

243 *Mineral compositional data*

244 The Ni-arsenides at Cliff predominantly comprise maucherite ($\text{Ni}_{11}\text{As}_8$), with subsidiary orcelite
 245 ($\text{Ni}_{5-x}\text{As}_2$). Prichard et al. (2004) and Derbyshire et al. (2013) have also reported nickeline (NiAs),
 246 as well as accessory godlevskite ($[\text{Ni,Fe}]_9\text{S}_8$), millerite (NiS) and heazlewoodite (Ni_3S_2). The
 247 maucherite and orcelite have ~1 wt.% level impurities of S, Fe, Sb and Co. Systematic
 248 compositional variation is not observed in the minor elements present in the Ni-arsenides, with the
 249 exception of S, which exhibits a weak negative correlation with Ni content (**Fig. 5**). The Cliff PGM
 250 assemblage comprises sperrylite (PtAs_2), stibiopalladinite (Pd_5Sb_2) and irarsite ($[\text{Ir,Ru,Rh,Pt}]\text{AsS}$),
 251 with subordinate laurite (RuS_2), confirming and extending the observations of Tarkian and Prichard
 252 (1987), Prichard and Tarkian (1988) and Prichard et al. (1994). Other PGM reported from Cliff by
 253 Tarkian and Prichard (1987) and Prichard and Tarkian (1988) include hongshiite (PtCu), potarite
 254 (PdHg) and hollingworthite ($\text{Rh,Pt,Pd}]\text{AsS}$), but these phases were not observed here. The Ru-rich
 255 pentlandite reported by the latter authors was also not observed here. The compositional variation
 256 of the Cliff PGM population analysed in the current study is illustrated in **Figure 6**. The sperrylite
 257 at Cliff is characterised by S, Fe, Sb and Ir impurities. Stibiopalladinite shows some intergrain
 258 compositional variation, with Fe, As, Cu and Pt present in up to wt.% concentrations. Irarsite
 259 contains sub wt.% levels of Cu, Fe and Sb. The single grain of laurite analysed has a relatively high
 260 As concentration (~3 wt.%), with measurable Os (>1 wt.%) and Rh (~4 wt.%) concentrations.
 261 Covariation of certain elements is observed between different PGM groups (e.g., Rh versus S; **Fig.**
 262 **6a**), that might suggest that the PGM behaved as a single paragenesis, or a sequence of parageneses,
 263 and there is overlap between irarsite, sperrylite and stibiopalladinite in Ni versus Fe space (**Fig. 6b**).
 264 In other respects (e.g., Fe versus Pt, **Fig. 6c**; Ir versus As, **Fig. 6d**), the different phases are clearly
 265 distinguished from one another and compositional variation within mineral groups is also observed.
 266 Stibiopalladinite shows a weak trend of sympathetically increasing Cu with Pt (**Fig. 6e**) and
 267 sperrylite and irarsite show a negative correlation between Sb and As (**Fig. 6f**).

268 The LA-ICP-MS data indicate that the PGE are not hosted within the fresh Cr-spinel.
 269 Concentrations of Co, Zn, As and Pb are relatively high in Cr-spinel (maximum concentrations of

270 278, 428, 49 and 1.5 ppb, respectively). A positive correlation is observed between Co and Zn (**Fig.**
271 **7a**), as well as weaker, but sympathetic covariation of As with Pb and Co (**Figs. 7b, 7c**). Maucherite
272 grains show weak positive correlations between base and precious metals in some cases (e.g., Au,
273 Pt versus Cu; **Figs. 8a, 8b**), but not in others (e.g., Co versus Rh; **Fig. 8c**). Platinum and Pd are also
274 positively correlated in maucherite (**Fig. 8d**). The LA-ICP-MS traverses from Cr-spinel through the
275 ferrian chromite rims and into the groundmass give insights into the distribution of the PGE in the
276 various mineral components, as well as other trace metals and semi-metals (**Figs. 9a-c**). The ferrian
277 chromite rims contain elevated Co and Ni (and Cu and Zn, not shown in **Fig. 9**), relative to fresh
278 Cr-spinel. Arsenic is also enriched in the sieve-textured rims, and follows the behavior of Ni
279 particularly well. Cobalt (and Zn) concentrations are considerably lower in the serpentinite
280 groundmass, but Ni and As are higher. The traverse shown in **Figure 9c** suggests that the
281 distribution of Ni and As in the serpentinite groundmass is relatively uniform, indicating either
282 abundant pervasively disseminated fine-grained Ni-arsenides, or another mode of occurrence. In
283 most cases, the PGE are indistinguishable from background levels within Cr-spinel grains, with the
284 exception of obvious inclusions of Ni-arsenide (e.g., **Fig. 9a**). However, in some samples the
285 altered Cr-spinel rims contain elevated Os and Pd (**Figs. 9b, 9c**). The serpentinite groundmass
286 around some Cr-spinel crystals contains concentrations of Os, Pd and to a lesser extent, Ru, which
287 are significantly higher than background levels. **Figure 9c** indicates elevated levels of Os relatively
288 homogeneously distributed throughout the groundmass. The distribution of Pd is more irregular,
289 though given the lack of covariation of base-metals it is not clear if this is a mineralogical control
290 (**Fig. 9c**). Platinum and Ir in the Cr-spinel, the altered rims and the groundmass serpentinite are
291 generally indistinguishable from background along the traverses analysed.

292

293 *X-ray microtomography*

294 The X-ray microtomography analyses provide insights into the spatial distribution of Ni-arsenides
295 in the Cliff chromitites (**Figs. 10, 11**). In this study, Ni-arsenides have not been thresholded from

PGM due to the similarities in grey levels of both mineral groups and the fact that the PGM are typically <5 µm in diameter. On the basis of the detailed textural observations, it is evident that they are distributed similarly. As such, conclusions drawn from X-ray microtomography on the microstructural relationships of the Ni-arsenide are broadly applied to the PGM too. The microtomography highlights that clusters of Ni-arsenide (and PGM) are typically sited at the grain boundaries of Cr-spinel crystals (**Figs. 11a,b,c**), often at points where fractures in the crystals intersect the groundmass serpentinite. Such fractures extend into the serpentinite, and may control the distribution of Ni-arsenide grains that appear to be isolated in the groundmass (**Figs. 11a,b**). The microtomography data also highlight the considerable density of microfractures that cut through the Cr-spinel crystals (**Fig. 11b**), and subtly suggest that olivine pseudomorphs survive in the groundmass (**Fig. 10c**), hinting at the protolith texture and mineralogy prior to serpentinisation.

Discussion

A low temperature Ni-arsenide-PGM paragenesis at Cliff

The formation of ferrian chromite (plus ferritchromit) in ophiolite and komatiite chromitites has been attributed to low grade (greenschist-lower amphibolite facies) metamorphism of Cr-spinel-bearing serpentinite (Kimball, 1990; Barnes and Roeder 2001; Mellini et al. 2005; González-Jiménez et al. 2009; Derbyshire et al. 2013). Barnes (2000) described the effects of metamorphic processes on the chemical composition of Cr-spinels from komatiites. In particular, Barnes concluded that Fe and Zn enrichment begins at greenschist facies, but continues on and is enhanced at amphibolite facies as magnetite replacement advances its encroachment on Cr-spinel crystal cores. There is a corresponding depletion in Ni during this process which occurs mostly at amphibolite grade. The stabilisation of chlorite (at ~550°C) plays a significant role in the mobility of Al³⁺ cations from Cr-spinel above this temperature. At relatively low metamorphic grades (greenschist facies), Zn, Mn and Co enrichment in Cr-spinel is observed. The Cliff chromitites have sieve-textures and alteration occurring both along grain boundaries and intragrain fractures. This

322 strongly suggests that a reactive fluid was responsible for the metamorphism of the Cr-spinels. A
323 good correlation is observed between Zn and Co (**Fig. 7a**) in ferrian chromite, suggesting that these
324 metals were mobile during metamorphism. In addition, As correlates positively with Pb and Co,
325 with the highest concentrations of these elements present in the ferrian chromite rims. This suggests
326 that the fluids responsible for alteration of the Cr-spinel had a high As activity. The production of
327 ferrian chromite is an oxidizing reaction, and has clearly also resulted in extensive precipitation of
328 the chlorite mineral on Cr-spinel grain boundaries (**Figs. 3a, 3b**). Derbyshire et al. (2013) suggested
329 that this phase was septechlorite, an aluminous serpentine, but Cr-rich chlorite is also possible along
330 the lines of a reaction such as: Cr-spinel + Antigorite $\rightarrow$ Fe-rich Cr-spinel + Cr-chlorite (Merlini et
331 al. 2009). Further analyses of this mineral are required for a definitive identification.

332 The Ni-arsenide in the Cliff chromitites is dominated by maucherite, with subordinate
333 orcelite and nickeline (Prichard and Lord, 1993; Derbyshire et al. 2013). Nickel-sulphide is
334 conspicuous by its absence; by contrast, other Unst chromitites and their host dunites typically
335 contain Ni-sulphide. The visible PGE mineralogy is dominated by arsenide phases, with a
336 preponderance of sperrylite controlling the distribution of Pt in the Cliff chromitites. The Ni-
337 arsenides and PGM are concentrated in the alteration zones and intra-crystal fractures of Cr-spinel
338 crystals, especially where Cr-spinel is altered to sieve-textured ferrian chromite. We suggest this
339 relationship implies that the reaction responsible for the formation of the ferrian chromite also
340 involved the Ni-arsenide, or perhaps the Ni-arsenide was a product of that reaction. In addition,
341 sulphide melts wet Cr-spinel surfaces much better than they do silicate mineral surfaces (Brenan
342 and Rose 2002), so it seems reasonable to suggest that arsenide might behave similarly. Such
343 behavior would explain the proximity of the bulk of the Ni-arsenide grains to Cr-spinel in these
344 samples. The novel X-ray tomography data presented here demonstrates that the Cliff chromitites
345 are pervasively deformed; the constituent Cr-spinels preserve evidence for complex fracturing and
346 alteration at sub-grain length-scales. Fluids were clearly focused along these intra-grain fractures
347 leading to precipitation of the PGM (**Fig. 4d**). This observation conclusively demonstrates a late-

stage (and presumably low-temperature, compared to chromitite formation) origin for the style of precious metal enrichment at Cliff. The three major PGM observed here (sperrylite, irarsite and stibiopalladinite) do not occur in different microstructural positions to one another, i.e., all are observed as inclusions in Ni-arsenide, sited along Cr-spinel fractures and at Cr-spinel grain boundaries (e.g., **Fig. 4c**). This suggests that all are part of the same paragenesis, and that they are coeval with the Ni-arsenide and the ferrian chromite. The broad variability of mineral compositions measured by electron microprobe (both in terms of base and precious metal abundances; **Fig. 6**) suggests that the processes responsible for their crystallisation occurred over either a range in temperature or from fluids with varying compositions. Maucherite is stable over a wide temperature interval (**Fig. 12**; Yund 1961), down to ~200°C, as is sperrylite. It therefore seems plausible that several generations of hydrothermal fluid were involved, to account for the complex microstructural relationships, intra-grain textures (e.g., **Fig. 4f**) and the variation in PGM chemistry observed.

Evidence for groundmass-hosted PGE enrichments

The LA-ICP-MS traverses across Cr-spinel grain boundaries in the Cliff samples yield intriguing insights into the distribution of the PGE. The serpentinite groundmass contains concentrations of Os and Pd that are significantly higher than background, and that are impossible to spatially resolve as discrete PGM grains (e.g., Os in the traverse illustrated in **Fig. 9c**). The PGE data along these traverses might potentially be explained by abundant very fine grained Os-rich phases disseminated throughout the serpentinite. The whole rock Os concentration of the Cliff chromitites is ~3 ppm (O'Driscoll et al. 2012). As shown in **Figure 9**, this Os is not in the Cr-spinel, so there is apparently a significant (unidentified) Os carrier present, potentially a nuggeting or clustering effect of Os in the groundmass. Relatively high levels of Pd are also present throughout the groundmass serpentinite. The traverses also indicate that Pt levels in the serpentinite groundmass and the Cr-spinel are indistinguishable from background levels. This is satisfactorily explained by the relatively large grain size of sperrylite crystals throughout the samples, which presumably act as

374 high concentration nodes for Pt and potentially account for the bulk rock concentration of up to
 375 ~100 ppm (O'Driscoll et al. 2012).

376

377 *Arsenide as a collector of PGE at Cliff*

378 The potential importance of arsenide as a collector of the PGE in magmatic and hydrothermal
 379 systems has been well-documented (Hanley, 2007; Dare et al. 2010; Godel et al. 2012; Le Vaillant
 380 et al. 2015; Prichard et al. 2013; Helmy et al. 2013; Helmy and Bragagini 2017), and though
 381 arguably not completely understood, it is known that As-rich deposits can exhibit very high PGE
 382 grades. Arsenic is enriched in intermediate solid solution (iss) during cooling (Holwell and
 383 McDonald, 2010), and there is a broad spectrum of Pd and Pt-rich arsenide phases that attest to the
 384 ease at which bonding with the metals occurs. Indeed, although experimental evidence for the
 385 system is not extensive, Pt and Pd may have greater affinity for arsenide than for sulphide (Fleet et
 386 al. 1993; Wood 2003; Helmy et al. 2013). Helmy et al. (2013) conclude that Pt and Pd associate
 387 with As^{n-} in preference to S^{2-} before saturation of arsenide in sulphide melts (perhaps as nano-metre
 388 sized As-metal clusters). The latter authors propose that the affinities of the precious and base-
 389 metals for an immiscible arsenide melt are: $\text{Pt} > \text{Pd} > \text{Ni} > \text{Fe} \approx \text{Cu}$. Pina et al. (2013) also reported that
 390 maucherite was strongly enriched in most chalcophile elements (including the PGE) relative to
 391 coexisting (co-paragenetic) sulphides (by factors of 10^{-1} to 10^2). Many of the studies listed above have
 392 attributed As mobilisation to magmatic processes, such as incorporation of an As-rich assimilate
 393 into magma (Godel et al. 2012), or leaching of As from crustal rocks by hydrothermal fluids (Le
 394 Vaillant et al. 2015). Kiefer et al. (2017) have also attributed the formation of complex sulfarsenide
 395 and arsenide mineralisation in the Dobšiná deposit (Slovakia) to the activity of fluids operating at
 396 temperatures less than 200°C. In the Bou Azzer ophiolite (southern Morocco), Ahmed et al. (2009)
 397 suggested mobilisation of As and Co from serpentinite to acidic (magmatic) fluids, at high fluid-
 398 rock ratios.

399 Generally, enrichments in As are not positively correlated with the PGE in the SOC
400 (Prichard and Lord 1993). The exception is the Cliff locality, where there are multiple potential
401 sources for the As. Derbyshire et al. (2013) reported sulphides with hundreds to thousands of ppm
402 of As in dunites and chromitites of other parts of the SOC. Heazlewoodite (more typically found in
403 chromitite, such as at the Harold's Grave or Hagdale localities) has higher concentrations (>1000
404 ppm) of As than dunite (in which pentlandite is dominant, <500 ppm). Thus, reworking (alteration)
405 of chromitite-hosted heazlewoodite could account for the As in the Cliff chromitites; the
406 observation that Cliff is the only chromitite locality in the SOC where little or no sulphide is
407 observed in most samples is intriguing and does suggest that local desulphurisation may have
408 occurred in the chromitite. However, a similar process operating on sulphide-bearing mantle dunites
409 such as those from The Viels (Derbyshire et al. 2013) might also be reasonable. In support of a
410 desulphurisation process being important at Cliff, Helmy et al. (2013) note that sperrylite is stable
411 relative to platarsite at relatively low sulphur activities.

412 It is also worth noting that Cliff is located close to the contact between the SOC and the
413 underlying Neoproterozoic metasediments (Dalradian Supergroup). Immediately across the
414 ophiolite contact (which is not the sole thrust, but a later structure referred to as the Burra Firth
415 Lineament that represents a younger tectonic break; Cutts et al. 2011) are graphitic schists and
416 limestones. Prichard and Lord (1993) reported high As concentrations in serpentinites along the
417 Burra Firth Lineament (that they mistakenly refer to as the basal thrust). It is possible that the
418 mineralisation at Cliff occurred during the reworking of the basal thrust that gave rise to the Burra
419 Firth Lineament. Indeed, Hanley (2007) has previously suggested that graphitic schists might have
420 acted as the source of As in the PGE-rich arsenide ores at Dundonald Beach South (Ontario,
421 Canada). The Os isotope data of O'Driscoll et al. (2012) indicate that Cliff chromitites are more
422 radiogenic ($^{187}\text{Os}/^{188}\text{Os}_{\text{initial}}$ of ~0.1292) than other SOC chromitites (~0.125), potentially supporting
423 the role of a hydrothermal fluid that had interacted with the metasediments in the provision of the
424 As. In fact, Prichard et al. (2017) recently reported variable Os isotope compositions for two PGM

425 grains from Cliff ($^{187}\text{Os}/^{188}\text{Os}$ between 0.1270-0.1300), suggesting the presence of >1 PGM
426 paragenesis, i.e., that the more radiogenic Os isotope compositions observed at Cliff may be
427 controlled by a specific PGM paragenesis. In this scenario, which is our preferred interpretation, we
428 envisage that As-rich fluids altered the existing sulphide assemblage in the Cliff chromitites, such
429 that the high PGE concentrations probably reflect the original chromitite PGE budgets, albeit with
430 reworking and upgrading of the PPGE in particular.

431

432 **Conclusions**

433 The SOC Cliff chromitite PGE occurrence is characterised by high PPGE relative to IPGE. Detailed
434 investigations of the rocks reveal that unlike other SOC chromitites, there is little to no Ni-sulphide
435 present. Instead, the primary ligand responsible for concentrating the PGE appears to have been As,
436 as evidenced by the dominant PGM present (e.g., sperrylite and irarsite). Abundant Ni-arsenide and
437 PGM are sited close to the edges of Cr-spinel crystals, and lining intra-crystal fractures. Together
438 with observations that ferrian chromite and septechnorite formed coevally with the As-rich PGM
439 paragenesis, it seems reasonable to suggest that a late-stage hydrothermal event has been
440 responsible for remobilizing and upgrading the primary PGE abundances and distribution within the
441 chromitites. The hydrothermal event could have resulted from movement on the Burra Firth
442 Lineament, a structure that excised the ophiolite basal thrust in the region of Cliff. The source of the
443 As is unknown, but may have come from Neoproterozoic metasediments on the opposite side of the
444 Burra Firth Lineament, including graphitic schists and impure limestones. Recent studies that report
445 relatively radiogenic chromitite Os isotope compositions from the Cliff area support the idea that a
446 late-stage hydrothermal fluid, derived from crustal rocks, was involved.

447

448 **Acknowledgements**

449 BO'D would like to acknowledge support from the Manchester X-ray Imaging Facility (MXIF;
450 Project #405) at the University of Manchester, and the Manchester-Diamond collaboration is also

acknowledged for beamtime at I13 (Project # MT12201). B.O'D. also acknowledges support from a Natural Environment Research Council (NERC) New Investigator grant NE/J00457X/1 and Standard Grant NE/L004011/1 during the writing of this paper. Nick Edwards, Jennifer Anne, and Joan Vila-Comamala are thanked for assistance at the Diamond Light Source Facility, and Tristan Lowe for hosting the authors at MXIF. MXIF is funded in part by the EPSRC (grants EP/F007906/1, EP/F001452/1 and EP/I02249X/1).

References

- Ahmed, A.H., Arai, S. and Ikenne, M. (2009) Mineralogy and Paragenesis of the C-Ni arsenide ores of Bou Azzer, Anti-Atlas, Morocco. *Economic Geology*, **104**, 249-266.
- Barnes, S.J. (2000) Chromite in Komatiites II. Modification during greenschist to mid-amphibolite facies metamorphism. *Journal of Petrology*, **41**, 387-409.
- Barnes, S.J. and Roeder, P.L. (2001) The range of spinel compositions in terrestrial mafic and ultramafic rocks. *Journal of Petrology*, **42**, 2279-2302.
- Brenan, J.M., and Rose, L.A. (2002) Experimental constraints on the wetting of chromite by sulfide liquid. *The Canadian Mineralogist*, **40**, 1113-1126.
- Chew, D.M., Daly, J.S., Magna, T., Pagé, L.M., Kirkland, C.L., Whitehouse, M.J. and Lam, R. (2010) Timing of ophiolite obduction in the Grampian orogen. *Geological Society of America Bulletin*, **122**, 1787-1799.
- Coggon, J.A., Nowell, G.M., Pearson, D.G., Lorand, J.-P., Oberthür, T. and Parman, S.W. (2011) The ¹⁹⁰Pt-¹⁸⁶Os decay system applied to dating platinum-group element mineralization of the Bushveld Complex, South Africa. *Chemical Geology*, **302-303**, 48-60.
- Crowley, Q.G. and Strachan, R.A. (2015) U-Pb constraints on obduction initiation of the Unst Ophiolite: an oceanic core complex in the Scottish Caledonides? *Journal of the Geological Society, London*, **172**, 279-282.
- Cutts, K.A., Hand, M., Kelsey, D.E., and Strachan, R.A. (2011) P-T constraints and timing of Barrovian metamorphism in the Shetland Islands, Scottish Caledonides: implications for the structural setting of the Unst ophiolite. *Journal of the Geological Society, London*, **168**, 1265-1284.
- Dare, S.A.S., Barnes, S.-J., Prichard, H.M. and Fisher, P.C. (2010) The timing and formation of platinum-group minerals from the Creighton Ni-Cu-platinum-group element sulfide deposit, Sudbury, Canada: Early crystallization of PGE-rich sulfarsenides. *Economic Geology*, **105**, 1071-1096.
- Day, J.M.D., O'Driscoll, B., Strachan, R.A., Daly, J.S. and Walker, R.J. (2017) Identification of mantle peridotite as a possible Iapetan ophiolite sliver in south Shetland, Scottish Caledonides. *Journal of the Geological Society, London*, **174**(1), 88-92.

496 Derbyshire, E.J., O'Driscoll, B., Lenaz, D., Gertisser, R. and Kronz, A. (2013) Compositionally
 497 heterogeneous podiform chromitite in the Shetland Ophiolite Complex (Scotland): implications for
 498 chromitite petrogenesis and late-stage alteration in the upper mantle portion of a supra-subduction
 499 zone ophiolite. *Lithos*, **162-163**, 279-300.
 500
 501 Fleet, M.E., Chrysosoulis, S.L., Stone, W.E., and Weisener, G. (1993) Partitioning of platinum-group
 502 elements and Au in the Fe-Ni-Cu-S system: Experiments on the fractional crystallization of sulfide
 503 melt. *Contributions to Mineralogy and Petrology*, **115**, 36-44.
 504
 505 Flinn, D. (2001) The basic rocks of the Shetland Ophiolite Complex and their bearing on its
 506 genesis. *Scottish Journal of Geology*, **37**, 79-96.
 507
 508 Flinn, D., Miller, J.A. and Roddom, D. (1991) The age of the Norwick hornblendic schists of Unst
 509 and Fetlar and the obduction of the Shetland ophiolite. *Scottish Journal of Geology*, **27**, 11-19.
 510
 511 Garwood, R.J. and Sutton, M.D. (2010) X-ray micro-tomography of Carboniferous stem-
 512 Dictyoptera: New insights into early insects. *Biology Letters*, **6**, 699-702.
 513
 514 Garwood, R.J. and Dunlop, J.A. (2014) The walking dead: Blender as a tool for palaeontologists
 515 with a case study on extinct arachnids. *Journal of Palaeontology*, **88(4)**, 735-746.
 516
 517 Godel, B. (2013) High resolution X-ray computed tomography and its application to ore deposits:
 518 case studies from Ni-Cu-PGE deposits. *Economic Geology*, **108**, 2005-2019.
 519
 520 Godel, B. (2015) Platinum-Group Element Deposits in Layered Intrusions: Recent Advances in the
 521 Understanding of the Ore Forming Processes. In Charlier, B., Namur, O., Latypov, R. and Tegner,
 522 C., eds., *Layered Intrusions*: Springer Netherlands, p. 379-432.
 523
 524 Godel, B. and Barnes, S.-J. (2008) Image analysis and composition of platinum-group minerals in
 525 the J-M Reef, Stillwater Complex. *Economic Geology*, **103**, 637-651.
 526
 527 Godel, B., Barnes, S.J., Barnes, S.-J. and Maier, W.D. (2010) Platinum ore in 3D: Insights from
 528 high-resolution X-ray computed tomography. *Geology*, **38(12)**, 1127-1130.
 529
 530 Godel, B., Gonzalez-Alvarez, I., Barnes, S.J., Barnes, S.-J., Parker, P. and Day, J. (2012) Sulfides
 531 and sulfarsenides from the Rosie Nickel Prospect, Duketon Greenstone Belt, Western Australia.
 532 *Economic Geology*, **107**, 275-294.
 533
 534 González-Jiménez, J.M., Kerestedjian, T., Proenza, J.A. and Gervilla, F. (2009) Metamorphism of
 535 chromite ores from the Dobromirski Ultramafic Massif, Rhodope Mountains (SE Bulgaria).
 536 *Geologica Acta*, **7**, 413-429.
 537
 538 González-Jiménez, J.M., Griffin, W.L., Gervilla, F., Proenza, J.A., O'Reilly, S.Y. and Pearson, N.J.
 539 (2014a) Chromitites in ophiolites: how, where, when, why? Part I. Origin and significance of
 540 platinum-group minerals. *Lithos*, **189**, 127-139.
 541
 542 González-Jiménez, J.M., Griffin, W.L., Proenza, J.A., Gervilla, F., O'Reilly, S.Y., Akbulut, M.,
 543 Pearson, N.J. and Arai, S. (2014b) Chromitites in ophiolites: how, where, when, why? Part II. The
 544 crystallisation of chromitites. *Lithos*, **189**, 140-158.
 545
 546

547 Hanley, J.J. (2005) The Aqueous Geochemistry of the Platinum-Group Elements (PGE) in Surficial,
548 Low-T Hydrothermal and High-T Magmatic Hydrothermal Environments. In Mungall, J.E., ed.,
549 *Exploration for Platinum-Group Element Deposits*, Mineral Association of Canada, Quebec, QC,
550 Canada, P. 35-56.
551

552 Hanley, J.J. (2007) The role of arsenic-rich melts and mineral phases in the development of high
553 grade Pt-Pd mineralization within komatiite-associated magmatic Ni–Cu sulfide horizons at
554 Dundonald Beach South, Abitibi Subprovince, Ontario, Canada. *Economic Geology*, **102**, 305-317.
555

556 Helmy, H.M., Ballhaus, C., Fonseca, R.O.C., and Nagel, T.J. (2013) Fractionation of platinum,
557 palladium, nickel, and copper in sulfide-arsenide systems at magmatic temperature. *Contributions*
558 *to Mineralogy and Petrology*, **166**, 1725-1737.
559

560 Helmy, H.M. and Bragagini, A. (2017) Platinum-group elements fractionation by selective
561 complexing, the Os, Ir, Ru, Rh-arsenide-sulfide systems above 1020°C. *Geochimica et*
562 *Cosmochimica Acta*, in press.
563

564 Holwell, D.A. and McDonald, I. (2010) A Review of the Behaviour of Platinum Group Elements
565 within Natural Magmatic Sulfide Ore Systems. *Platinum Metals Review*, **54**, 26-36.
566

567 Irvine, T.N. (1977) Origin of chromitite layers in the Muskox intrusion and other stratiform
568 intrusions: A new interpretation. *Geology*, **5**, 273-277.
569

570 Karup-Møller, S., Mackovicky, E. and Barnes, S.-J. (2008) The metal-rich portions of the phase
571 system Cu–Fe–Pd–S at 1000 °C, 900 °C and 725 °C: implications for mineralization in the
572 Skaergaard intrusion. *Mineralogical Magazine*, **72**, 941-951.
573

574 Ketcham, R. A. and Carlson, W. D. (2001) Acquisition, optimization and interpretation of X-ray
575 computed tomographic imagery: applications to the geosciences. *Computers and Geosciences*, **27**,
576 381-400.
577

578 Kiefer, S., Majzlan, J., Chovan, M. and Števko, M. (2017) Mineral compositions and phase
579 relations of the complex sulfarsenides and arsenides from Dobšiná (Western Carpathians,
580 Slovakia). *Ore Geology Reviews*, **89**, 894-908.
581

582 Kimball, K. (1990) Effects of hydrothermal alteration on the composition of chromian spinels.
583 *Contributions to Mineral Petrology*, **105**, 337-346.
584

585 Latypov, R., O’Driscoll, B. and Lavrenchuk, A. (2013) Towards a model for an in situ origin for
586 PGE reefs in layered intrusions: insights from the chromitite seams of the Rum Eastern Layered
587 Intrusion, Scotland. *Contributions to Mineral Petrology*, **166**, 309-327.
588

589 Le Vaillant, M., Barnes, S.J., Fiorentini, M.L., Miller, J., McCuaig, T.C. and Muccilli, P. (2015) A
590 hydrothermal Ni-As-PGE geochemical halo around the Miitel komatiite-hosted nickel sulfide
591 deposit, Yilgarn Craton, Western Australia. *Economic Geology*, **110**, 505-530.
592

593 Limaye, A. (2012) Drishti: a volume exploration and presentation tool. In: SPIE Optical
594 Engineering+ Applications. International Society for Optics and Photonics, pp. 85060X-85060X.
595

596 Melcher, F., Grum, W., Simon, G., Thalhammer, T.V. and Stumpfl, E.F. (1997) Petrogenesis of
597 the ophiolitic giant chromite deposits of Kempirsai, Kazakhstan: a study of solid and fluid
598 inclusions in chromite. *Journal of Petrology*, **38**, 1419-1458.

599
600 Mellini, M., Rumori, C. and Viti, C. (2005) Hydrothermally reset magmatic spinels in retrograde
601 serpentinites: formation of ‘ferritchromit’ rims and chlorite aureoles. *Contributions to Mineralogy
602 and Petrology*, **149**, 266-275.
603
604 Merlino, A., Grieco, G. and Diella, V. (2009) Ferritchromit and chromium-chlorite formation in
605 melange-hosted Kalkan chromitite (Southern Urals, Russia). *American Mineralogist*, **94**, 1459-
606 1467.
607
608 Mungall, J.E. and Brenan, J.M. (2014) Partitioning of platinum-group elements and Au between
609 sulfide liquid and basalt and the origins of mantle–crust fractionation of the chalcophile elements.
610 *Geochimica et Cosmochimica Acta*, **125**, 265-289.
611
612 Naldrett, A.J. and Duke, J.M. (1980) Platinum metals in magmatic sulfide ores. *Science*, **208**, 1417-
613 1424.
614
615 O’Driscoll, B., Day, J.M.D., Daly, J.S., Walker, R.J. and McDonough, W.F. (2009) Rhenium-
616 osmium isotope and platinum-group elements in the Rum Layered Suite, Scotland: Implications for
617 Cr-spinel seam formation and the composition of the Iceland mantle anomaly. *Earth and Planetary
618 Science Letters*, **286**, 41-51.
619
620 O’Driscoll, B., Day, J.M.D., Walker, R.J., Daly, J.S., McDonough, W.F. and Piccoli, P.M. (2012)
621 Chemical heterogeneity in the upper mantle recorded by peridotites and chromitites from the
622 Shetland Ophiolite Complex, Scotland. *Earth and Planetary Science Letters*, **333-334**, 226-237.
623
624 O’Driscoll, B. and González-Jiménez, J.M. (2016) Petrogenesis of the platinum-group minerals.
625 *Reviews in Mineralogy and Geochemistry, Mineralogical Society of America*, **81**, 489-578.
626
627 Oshin, I.O. and Crocket, J.H. (1986) Noble metals in the Thetford Mines ophiolites, Quebec,
628 Canada. II. Distribution of gold, silver, iridium, platinum and palladium in the Lac de l’Est volcano-
629 sedimentary section. *Economic Geology*, **81**, 931-945.
630
631 Page, N.J. and Talkington, R.W. (1984) Palladium, platinum, rhodium, ruthenium, and iridium in
632 peridotites and chromites from ophiolite complexes in Newfoundland. *Canadian Mineralogist*, **22**,
633 137-149.
634
635 Pina, R., Gervilla, F., Barnes, S.-J., Ortega, L. and Lunar, R. (2013) Partition coefficients of
636 platinum group and chalcophile elements between arsenide and sulfide phases as determined in the
637 Beni Bousera Cr-Ni mineralization (North Morocco). *Economic Geology*, **108**, 935-951.
638
639 Prichard, H.M. (1985) The Shetland Ophiolite. In Gee, D.G. and Sturt, B.A. Eds., *The Caledonide
640 Orogen - Scandinavia and related areas*. Wiley, New York, p. 1173-1184.
641
642 Prichard, H. and Tarkian, M. (1988) Platinum and Palladium minerals from two PGE-rich localities
643 in the Shetland Ophiolite Complex. *The Canadian Mineralogist*, **26**, 979-990.
644
645 Prichard, H.M. and Lord, R. (1993) An overview of the PGE concentrations in the Shetland
646 Ophiolite Complex. *Geological Society of London Special Publications*, **76**, 273-294.
647
648 Prichard, H.M., Ixer, R., Lord, R.A., Maynard, J. and Williams, N. (1994) Assemblages of
649 platinum-group minerals and sulphides in silicate lithologies and chromite-rich rocks within the
650 Shetland Ophiolite. *The Canadian Mineralogist*, **32**, 271-294.

651
652 Prichard, H.M., Fisher, P.C., McDonald, I., Knight, R.D., Sharp, D.R. and Williams J.P. (2013) The
653 distribution of PGE and the role of Arsenic as a collector of PGE in the Spotted Quoll nickel ore
654 deposit in the Forresteria Greenstone Belt, Western Australia. *Economic Geology*, **108**, 1903-1921.
655
656 Prichard, H.M., Barnes, S. J., Godel, B., Halfpenny, A., Neary, C.R. and Fisher, P.C. (2015) The
657 structure of and origin of nodular chromite from the Troodos ophiolite, Cyprus, revealed using
658 high-resolution X-ray computed tomography and electron backscatter diffraction. *Lithos*, **218-219**,
659 87-98.
660
661 Prichard H.M., Barnes S.J., Dale C.W., Godel B., Fisher P.C., Nowell, G.M. (2017) Paragenesis of
662 multiple platinum-group mineral populations in Shetland ophiolite chromitite: 3D X-ray
663 tomography and in situ Os isotopes. *Geochimica et Cosmochimica Acta*, in press.
664
665 Schouwstra, R.P., Kinloch, E.D. and Lee, C.A. (2000) A short geological review of the Bushveld
666 Complex. *Platinum Metals Review*, **44**, 33-39.
667
668 Spray, J.G. and Dunning, G.R. (1991) A U/Pb age for the Shetland Islands oceanic fragment,
669 Scottish Caledonides: evidence from anatectic plagiogranites in 'layer 3' shear zones. *Geological*
670 *Magazine*, **128**, 667-671.
671
672 Sutton, M.D., Garwood, R.J., Siveter, D.J. and Siveter, D.J. (2012) Spiers and VAXML; A software
673 toolkit for tomographic visualisation, and a format for virtual specimen interchange. *Palaeontologia*
674 *Electronica*, **15(2)**, 14p.
675
676 Tarkian, M. and Prichard, H.M. (1987) Irarsite-hollingworthite solid-solution series and other
677 associated Ru-, Os-, Ir- and Rh-bearing PGM's from the Shetland ophiolite complex. *Mineralium*
678 *Deposita*, **22**, 178-184.
679
680 Titarenko, V., Bradley, R., Martin, C., Withers, P. J. and Titarenko, S. (2010) Regularization
681 methods for inverse problems in x-ray tomography. In: *Developments in X-Ray Tomography VII*
682 (ed. Stock, S. R.) 7804, 40Z.
683
684 Wood, M. (2003) Arsenic in igneous systems: An experimental investigation: Unpublished B.A.Sc.
685 thesis, Toronto, Ontario, Canada, University of Toronto, 32 p.
686
687 Yund, R.A. (1961) Phase relations in the system Ni-As. *Economic Geology*, **56**, 1273-1296.
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702

Figure Captions

Figure 1. (a) Chondrite-normalised PGE patterns of various Shetland Ophiolite Complex chromitites, including Cliff, as well as other ophiolite chromitites. The chondrite normalization data are from Naldrett and Duke (1980).

Figure 2. Geological map of the Shetland Ophiolite Complex, illustrating the major components of the ophiolite as well as the chromitite localities (red dots) referred to in the text. Inset shows the regional position of the Shetland archipelago in the North Sea, relative to Scotland and Norway.

Figure 3. Cross-polarised light photomicrographs of Cliff chromitite, illustrating the development of chlorite rims on Cr-spinel grain edges. (a) Large opaque grains are Cr-spinel with blue-coloured chlorite rims. Chlorite is present in the groundmass too. (b) As above, except the development of chlorite rims around serpentinised olivine inclusions in Cr-spinel is evident, as well as the formation of chlorite along narrow intra-Cr-spinel fractures (arrowed). (c) Brittle deformation of Cr-spinel aggregates in the Cliff chromitites, forming fractures infilled with fibrous chrysotile.

Figure 4. (a) Reflected light photomicrograph illustrating the typical texture of the Cliff chromitites. Note the highlighted 'bright' phases, which are Ni-arsenide, which may be located both at grain edges and isolated within the groundmass. (b) Back-scattered electron micrograph of a Cr-spinel grain showing complex resorption features, particularly at grain boundaries and along intra-grain fractures. The presence of several Ni-arsenide phases is highlighted. The dark grey Cr-spinel at image centre is the fresh, unaltered composition. (c) Close-up back-scattered electron micrograph of Cr-spinel grain boundary. Note the development of ferrian chromite rims (lighter grey) at Cr-spinel grain boundaries and along a narrow zone of alteration within the Cr-spinel crystal, and the presence of three sperrylite crystals in this zone. The large bright phase at the edge of the Cr-spinel is maucherite, as is the smaller phase below it. (d) Back-scattered electron micrograph illustrating a narrow intra-Cr-spinel fracture, containing numerous maucherite and sperrylite grains (bright phases). (e) Back-scattered electron micrograph of an area of complex alteration at the edge of a cluster of Cr-spinel crystals. Note the extensive development of sieve-textured ferrian chromite, and close spatial association of numerous Ni-arsenide and PGM at the edges of the Cr-spinels and within the fractures that cross-cut the grains. (f) Close-up back-scattered electron micrograph of a maucherite grain, showing internal compositional heterogeneity, and (arrowed) an irarsite grain.

Figure 5. Plots of (a) S wt.% versus Ni wt.% and (b) Sb wt.% versus As wt.% in maucherite.

Figure 6. Plots of platinum-group mineral compositional variations (a) Rh wt.% versus S wt.%, (b) Ni wt.% versus Fe wt.%, (c) Fe wt.% versus Pt wt.%, (d) Ir wt.% versus As wt.%, (e) Pt wt.% versus Cu wt.%, (f) Sb wt.% versus As wt.%. See text for further discussion.

Figure 7. Plots of trace elements in Cr-spinel as follows: (a) Zn versus Co, (b) As versus Co, (c) Pb versus As.

Figure 8. Plots of trace elements in Ni-arsenide as follows: (a) Au versus Cu, (b) Pt versus Cu, (c) Co versus Rh, (d) Pt versus Pd. See text for further discussion.

Figure 9. (a)-(c) Various time-resolved LA-ICP-MS traverses from Cr-spinel (dark grey) into serpentinised groundmass. The yellow colour represents background levels (i.e., laser off) and the white colour indicates data collection in the serpentinite groundmass. See text for further discussion. Specific features of the data referred to in the text are labeled on the panels.

754 **Figure 10.** X-ray tomography data from the Cliff chromitites. (a) An example of the ~1 mm
755 diameter cores analysed by synchrotron. The dense phase coloured in light brown is Cr-spinel. (b)
756 Hand specimen scanned at the MXIF facility, University of Manchester, using the Nikon Metris
757 225/320 kV Xray CT, showing the apparently random distribution of Ni-arsenides (false-colour
758 purple grains) throughout the sample. (c) Cross section of a ~1 mm diameter core, analysed by
759 synchrotron. The partial outline of the core in plan view is shown by the white dashed line. The
760 false-coloured light-brown phase is Cr-spinel. Note the irregular amorphous grey features in the
761 serpentinised groundmass, interpreted here to be relic (pseudomorphed) olivine grains.

762
763 **Figure 11.** X-ray tomography data from the Cliff chromitites. (a) False-coloured μ CT scan showing
764 Cr-spinel (light-brown) and clusters of Ni-arsenide grains (false-colour purple) sited at the point
765 where a fracture network intersects the boundary between a Cr-spinel grain and the serpentinised
766 groundmass. (b) False-coloured image highlighting the abundance and complexity of fracturing of
767 Cr-spinel grains (coloured green). Note the presence of high-density phases (Ni-arsenides) coloured
768 purple, again sited within or close to intra-Cr-spinel fractures. (c) Close-up image of a Ni-arsenide
769 (maucherite) entirely enclosed in the serpentinite groundmass – such occurrences tend to exhibit the
770 coarsest grainsizes of Ni-arsenide (light-brown is Cr-spinel and purple is Ni-arsenide).

771
772 **Figure 12.** Sketch of the temperature-composition phase diagram for the Ni-As system, reproduced
773 from Yund (1961). The dashed lines indicate work not determined by that study. Mineral phase
774 abbreviations as follows; ma: maucherite, nc: niccolite, rm: rammelsbergite, prm:
775 pararammelsbergite.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12