Diagenetic controls on the reservoir quality of the tight gas Collyhurst sandstone formation, lower Permian, east Irish Sea basin, United Kingdom

DOI:

[10.1016/j.sedgeo.2018.04.006](https://doi.org/10.1016/j.sedgeo.2018.04.006)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Oluwadebi, A., Taylor, K., & Dowey, P. (2018). Diagenetic controls on the reservoir quality of the tight gas Collyhurst sandstone formation, lower Permian, east Irish Sea basin, United Kingdom. *Sedimentary Geology*, 371, 55-74. <https://doi.org/10.1016/j.sedgeo.2018.04.006>

Published in:

Sedimentary Geology

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Accepted Manuscript

Diagenetic controls on the reservoir quality of the tight gas Collyhurst sandstone formation, lower Permian, east Irish Sea basin, United Kingdom

Ayomiposi G. Oluwadebi, Kevin G. Taylor, Patrick J. Dowe

PII: S0037-0738(18)30111-8
DOI: doi:[10.1016/j.sedgeo.2018.04.006](https://doi.org/10.1016/j.sedgeo.2018.04.006)
Reference: SEDGEO 5335

To appear in:

Received date: 14 March 2018
Revised date: 27 April 2018
Accepted date: 29 April 2018

Please cite this article as: Ayomiposi G. Oluwadebi, Kevin G. Taylor, Patrick J. Dowe , Diagenetic controls on the reservoir quality of the tight gas Collyhurst sandstone formation, lower Permian, east Irish Sea basin, United Kingdom. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Sedgeo(2018), doi:[10.1016/j.sedgeo.2018.04.006](https://doi.org/10.1016/j.sedgeo.2018.04.006)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Diagenetic Controls on the Reservoir Quality of the Tight Gas Collyhurst Sandstone
Formation, Lower Permian, East Irish Sea Basin, United Kingdom**

Ayomiposi G. Oluwadebi¹, Kevin G. Taylor¹ and Patrick J. Dowey¹

¹ *School of Earth and Environmental Sciences, University of Manchester, Manchester, M13 9PL, UK.*

Corresponding author: ayomiposi.oluwadebi@manchester.ac.uk

Abstract

The Lower Permian Collyhurst Sandstone Formation is an important gas reservoir both offshore and onshore in the East Irish Sea Basin of north-western UK, and is considered to be a tight gas reservoir. Diagenetic processes have significantly influenced the reservoir quality of the sandstone. This study employs petrographic, mineralogical, and stable isotopic analyses on two different onshore core successions to investigate the composition and texture, the diagenetic history, and the impact of lithology and diagenesis on reservoir quality. The Collyhurst Sandstone Formation consists of fine- to very coarse-grained, poorly to well-sorted, sub-arkosic to lithic arenite sands. The reservoir properties of the sandstone are relatively poor, with average porosity and permeability of 6.6% and 0.387 mD respectively. On the basis of petrographic analyses, five sandstone lithofacies were identified: fine-grained, conglomeratic, carbonate-cemented, quartz-cemented and matrix-free. Contrasting characteristics between the reservoir units suggest the development of divergent diagenetic pathways. Features observed during early diagenesis include compaction, growth of pore-filling early carbonate cements (calcite I, dolomite and siderite I) and vermicular kaolinite. Late diagenetic features include: further carbonate cementations (ferroan dolomite/ankerite, calcite II and siderite II-III), clay mineral cementation (illite and chlorite) with both grain-coating and pore-filling textures. Also common in late diagenesis are syntaxial quartz overgrowths. Albite, barite and anhydrite are minor late diagenetic cement phases. Compaction processes rather than cementation accounts for the major destruction of

the initial sandstone porosity. But cementation still accounts for up to 32% of porosity loss. Carbonate cements mostly fill intergranular pores, thus block pore throat and prevent migration of fluid. Clay minerals bridge and occlude pores, resulting in further destruction of the reservoir quality. The study provides insights into how lithofacies (primary composition, texture and sorting) and diagenesis can influence reservoir qualities of tight gas sandstones.

Keywords

Collyhurst Sandstone, East Irish Sea Basin, diagenesis, reservoir quality, Tight gas.

1. Introduction

The Lower Permian Collyhurst Sandstone Formation is an important gas-bearing reservoir in the East Irish Sea Basin (EISB) of north-western UK. It forms part of the New Red (Permo-Triassic) Sandstone in the British stratigraphic succession (Bjørlykke, 2011). Due to an average permeability of less than 1 mD and porosity below 10%, it is considered to be a tight sandstone (Newman, 1999; Smythe, 2014). Sandstone reservoirs (including tight sandstones) can be extensively altered by diagenetic processes (Primmer *et al.*, 1997; Worden and Burley, 2003; Schmid *et al.*, 2004; Karim *et al.*, 2010; Taylor *et al.*, 2010). Therefore, a key factor in identifying net pay zones in tight sandstones is the ability to understand and predict diagenetic processes and their impact on reservoir quality (Fic and Pedersen, 2013; Stroker *et al.*, 2013).

Research on diagenesis in low permeability reservoirs is becoming increasingly important owing to the development of substantial gas reserves in tight sandstones (Stroker *et al.*, 2013). However, studies on sandstone diagenesis are mostly associated with high permeability examples since they tend to be better reservoirs. Examples of low permeability sandstones (tight sandstone) are valuable as they may identify which factors control reservoir quality. Isolated examples from Chinese and US basins (e.g., Zhang *et al.*, 2009; Tobin *et al.*, 2010;

Stroker *et al.*, 2013; Xi *et al.*, 2015; Lai *et al.*, 2017) do not give a sufficiently broad consideration to the potential geological controls on tight sandstone reservoir quality.

Within the East Irish Sea Basin and adjoining areas, several works have been published on regional geology, stratigraphy, tectonic setting and hydrocarbon potential (e.g., Stuart and Cowan, 1991; Jackson *et al.*, 1997; Cowan *et al.*, 1999; Knott and Hill, 1999; Yaliz and Taylor, 2003; Xi *et al.*, 2015). However, studies on diagenesis and reservoir quality within East Irish Sea Basin reservoirs have previously focused on the well understood, high permeability Triassic Sherwood Sandstone Group (e.g., Burley, 1984; Bushell, 1986; Meadows and Beach, 1993; Greenwood and Habesch, 1997; Bastin *et al.*, 2003). The Collyhurst Sandstone then serves as a good example of the diagenetic evolution of a tight gas sandstone reservoir within the EISB and the UK more widely.

This study aims to improve the understanding of diagenesis and its controls on reservoir quality in tight gas sandstones. The objectives are to utilise petrographic, mineralogical, and stable isotopic techniques to: 1) characterize the composition and texture of the Collyhurst Sandstone Formation, 2) identify lithofacies within the reservoir, 3) outline the diagenetic evolution and reconstruct the paragenetic sequence, and 4) identify how primary depositional and secondary diagenetic processes control reservoir quality in the Collyhurst Sandstone Formation.

2. Geological setting

The study area is located in East Irish Sea Basin (EISB), north-western UK (Fig. 1). The EISB is the largest of five Permo-Triassic basins within the Irish Sea and is an established hydrocarbon province (Floodpage *et al.*, 2001). The stratigraphic succession within the Permo-Triassic of the East Irish Sea Basin consists of five depositional sequences (Fig. 2). The Collyhurst Sandstone Formation which is part of Appleby Group (Lower Permian)

makes up the first sequence, followed by Cumbrian Coast Group (Upper Permian), Sherwood Sandstone Group (Lower Triassic), Mercia Mudstone Group (Upper Triassic) and Penarth Group (Upper Triassic). Each of the depositional sequences is separated by major sequence boundaries or disconformities (Jackson *et al.*, 1997).

The Collyhurst Sandstone Formation developed during a late syn-rift event. The rifting produced normal faults with NNE-SSW and ENE-WSW orientations, which were active through the Middle Permian (Knott and Hill, 1999). The Collyhurst Sandstone Formation is the basal unit of the Permo-Triassic sequence and was unconformably deposited on the underlying Stephanian red bed sandstones of the Kidston Group (Carboniferous). It also locally forms a disconformity with the Early Permian Tormentil Volcanics (Fig. 2, Jackson *et al.*, 1997). The Collyhurst Sandstone Formation consists dominantly of aeolian and fluvial sandstones (red to brownish grey) with interbedded conglomerate and the rare occurrence of mudstone (Jackson *et al.*, 1997; Brandon *et al.*, 1998). The Collyhurst Sandstone Formation is a reservoir in the Elswick gas-field. The Elswick gas-field has limited production from a single well (Knott and Hill, 1999), it was hydraulically fractured due to its low permeability (Warwick, 2001), and directly generates about 1MW of electricity annually.

3. Samples and methodology

Forty-six rock samples were collected within the core depths (1043-1082 m) from the Elswick-1 (436965 N, 342380 E) and Grange Hill-1 well (438939 N, 339179 E) at a regular spacing (approximately 1 m). From the samples, polished thin sections were prepared for petrographic and mineralogical analysis. The thin sections were impregnated with blue dye to enhance the visibility of porosity; lithological parameters such as texture, sorting and mineral composition were described using optical microscopy. All light microscopic analyses were performed using a Nikon Eclipse E200 microscope and a Nikon Japan microscope equipped

with a CL8200-MKS cold cathodoluminescence optical system. Light and CL microscopic analysis of thin section texture and composition were supplemented with mechanical, quantitative point-counting on each of the forty-six thin sections. Three hundred point-counts per thin section sample were completed.

Further detailed analysis of sandstone microstructure and composition was undertaken using a XL30 environmental scanning electron microscope (ESEM) equipped with Energy Dispersive X-ray (EDX) spectrometer. Representative carbon-coated thin section and stub mounted samples were studied with back-scattered electron (BSE) and secondary electron (SE) modes. EDX scans allowed the chemical compositions of minerals to be identified. In addition to energy dispersive X-ray analysis of samples, full quantitative analysis of some representative samples were performed using Cameca Electron Microprobe attached with wave-length-dispersive X-ray to determine their elemental composition. Major elements analysed include Carbon (C), Calcium (Ca), Magnesium (Mg) and Iron (Fe), while trace elements are Strontium (Sr) and Manganese (Mn).

Mineralogical identification of seventeen samples utilised X-ray diffraction (XRD) analysis on whole rock and clay-size fractions ($\leq 2 \mu\text{m}$). Rock samples were dispersed using de-ionised water, and clay-size fractions were separated from the coarse fractions ($> 2 \mu\text{m}$) using the centrifuge method (Ronald, 1967). The clay-size fractions were smeared on to glass slide and analysed using a Bruker D8Advance X-ray diffractometer equipped with a Göbel Mirror a Lynxeye detector. The X-ray tube had a copper source, providing $\text{CuK}_{\alpha 1}$ X-rays with a wavelength of $1.5406 \text{ \AA}$. Samples were scanned from $5-70^\circ 2\theta$, with a step size of 0.02° and a count time of 0.2s per step. The resultant diffraction patterns were evaluated using EVA software (version 4), which compares experimental data to standards from the International Centre for Diffraction Data (ICDD). X-ray diffraction patterns were obtained for both air

dried and ethyl glycolation. Treatment with ethyl glycolation enables easy identification of swelling clay minerals.

The carbon ($^{13}\text{C}/^{12}\text{C}$) and oxygen ($^{18}\text{O}/^{16}\text{O}$) isotope ratios of eight selected dolomite samples were determined following the classical “sealed vessel” procedure described by (McCrea, 1950) and (Swart *et al.*, 1991). Sufficient powdered sample (to yield between 40 and 50 μmol s of CO_2) was reacted at 50 °C for 48 hrs with phosphoric acid (specific gravity 1.92) under vacuum. Product CO_2 was recovered cryogenically and mass ratios were measured against a comparison CO_2 (‘reference’ gas) using a dual-inlet VG SIRA 10 mass spectrometer. Resultant delta values ($\delta_{\text{CO}_2}^{45/44}$ and $\delta_{\text{CO}_2}^{46/44}$) were corrected for ^{17}O effects (Craig, 1957) and adjusted for temperature-dependent kinetic oxygen isotope fractionation associated with the dolomite-phosphoric acid reaction using a fractionation factor (α) of 1.01066 (Rosenbaum and Sheppard, 1986). All data are reported as delta (δ) values with respect to the Vienna Pee Dee Belemnite (VPDB) carbon and oxygen isotope scales (*via* secondary laboratory reference calcite calibrated against NBS 19 and NBS18). Estimated analytical precision (1σ), based on the replicate analysis of an in-house quality control dolomite, is better than 0.1‰ for both carbon and oxygen isotope values.

Associated porosity and permeability (poroperm) datasets for Elswick-1 (EL-1) well were supplied by UK land well record. These data were not available for Grange Hill-1 (GH-1) well. The measurements were collected using standard poroperm measurement techniques (Hornung and Aigner, 1999; Ruf and Aigner, 2004).

4. Results

4.1 Detrital mineralogy

Petrographic analysis of samples from both studied wells indicate that the sandstones consist of very fine-grained to very coarse-grained, poorly to well-sorted, sub-angular to sub-

rounded sands (Fig. 3A and 3B). Long grain contacts and concavo-convex contacts are dominant (Fig. 3C), while suture contacts and stylolites (Fig. 3D) are also present in some samples. Generally, EL-1 well consists of fine to coarse-grained sandstone, whereas GH-1 well is predominantly of very fine to fine-grained sandstone and mudstone.

Compositionally, quartz is the dominant detrital mineral, with monocrystalline quartz ranging from 23.5 to 61.0% (average 48.6%) and polycrystalline quartz ranging from 0.0 to 13.5% (average 6.0%). Total feldspar varies from 2.0 to 10.5% (average 5.3%); K-feldspar is the dominant type of the feldspar, making up to 70.0% of total feldspar. Rock fragments include chert, volcanic and metamorphic rocks and mud intraclasts. The total concentrations of all rock fragments range from 3.0% to 42.5% (average 14.4%). Mica grains (muscovite and biotite) range from 0.0 to 4.5 % (average 1.3%). Heavy minerals including garnet, rutile, ilmenite and glauconite are rare, ranging from 0.0 to 3.5% (average 1.2%).

Lithic rock fragments are more common in samples from EL-1 well than GH-1 well; except mica grains and heavy minerals such as ilmenite and rutile which are more common in GH-1 well. Compositional data from point count analysis of the Collyhurst Sandstone Formation is presented in Fig. 4. Based on the classification scheme of Pettijohn (1957) the sandstones in the two studied wells range from a lithic arenite to a sub-arkosic and sub-lithic arenite to a quartz arenite. Majority of samples from both wells falls within the sublitharenite and fewer samples are subarkose, while only one sample from Elswick-1 well is quartz arenite.

4.2 Lithofacies description

Five lithofacies were identified based on lithological characteristics (including texture, mineral composition, sorting) and dominant diagenetic minerals, these are; (A) fine-grained

sandstone; (B) conglomeratic sandstone; (C) carbonate-cemented sandstone; (D) quartz-cemented sandstone. (E) matrix-free sandstone.

Lithofacies A consists of very fine- to fine-grained sandstone largely consist abundant monocrystalline quartz (Fig. 5A) and it is moderately sorted. Microscopic observations indicate that this lithofacies is extensively/poorly cemented (average of 10.6%). Clay mineral cements are the most common (average of 5.9%) with varying proportions of carbonate (average of 4.3%) and quartz cements (average of 0.6%). The fine-grained sandstone is interbedded with silty mudstone. It is dominant in GH-1 well and lower section of EL-1 well (1075 – 1082 m).

Lithofacies B consists of a medium- to very coarse-grained sandstone. It is poorly sorted sandstone and clast-supported (Fig. 5B). Samples of this lithofacies are highly compacted due to the presence of ductile mud intraclasts and lithic grains (average of 29.0%). Clay minerals cements account for major cement in this lithofacies (average of 2.6%) and it is dominant in EL-1 well.

Lithofacies C is typically a fine to coarse-grained sandstone. It is moderately sorted and heavily cemented by carbonate cement, with an average of 13.1% (Fig. 5C and 5D). This lithofacies is observed in both studied wells; however, there is variation in the distribution of the carbonate cements type. Calcite and dolomite cements are common in EL-1 well, whereas GH-1 well is dominated by dolomite and siderite cement.

Lithofacies D is a medium to coarse-grained, moderately sorted sandstone. Quartz cementation dominates this lithofacies (Fig. 5E). Lithofacies D is primarily identified in samples from the EL-1 well (quartz cement averages 3.5%). A rare occurrence of this lithofacies is also observed in samples from GH-1 well, with an average of 0.6% of quartz cement. Quartz cement occurs mainly as overgrowth on the detrital quartz grains; whereas, in some examples it was noted to completely fill intergranular pore volumes (Fig. 5E).

Lithofacies E consists of medium to coarse-grained sandstone, which is moderately to well-sorted (Fig. 5F). This lithofacies is termed matrix-free sandstone due to its abundant detrital quartz (average of 51.4%), lithic rock fragments (average of 14.4%) and feldspar grains (average of 6.6%) with low concentrations of cement (total cement averages 8.8%). This lithofacies occurs predominantly within the EL-1 well.

4.3 Diagenetic processes

The main diagenetic processes identified in the studied samples include compaction and cementation (carbonate, clay minerals and quartz cements). The vertical distribution of major cements in EL-1 and GH-1 well is shown in Fig. 6. Carbonate (mainly dolomite and calcite) and clay mineral (chlorite and illite) cements are the most common pore-filling cements, followed by authigenic quartz cement. Barite, anhydrite and albite are minor cements.

4.3.1 Carbonate cements

Carbonate cements (mainly dolomite and calcite) are the dominant cement in the studied sandstone (Fig. 6). Carbonate cements within both studied wells are unevenly distributed (Fig. 6A). On the basis of the point count data, dolomite is the dominant carbonate cement and ranges from 0.0 to 21.0% (average 3.2%). It is present in almost all samples and occurs as both ferroan and non-ferroan dolomite cements. Non-ferroan dolomite occurs as monocrystalline euhedral rhombic crystals and ranges in size from $<5\ \mu\text{m}$ to $50\ \mu\text{m}$ (Fig. 7A). Non-ferroan dolomite cement fills intergranular pores. Analysis of thin sections under cold cathodoluminescence microscopy revealed non-ferroan dolomite has an orange luminescence colour (Fig. 7B), filling and occluding primary pores. Ferroan dolomite/ankerite (Fe-D) mostly zoned non-ferroan dolomite (Fig. 7A); however, it occasionally grows as large individual crystals ($\geq 20\ \mu\text{m}$) filling intergranular pores. In addition to petrographic texture, electron probe analysis of Fe concentrations in some analysed samples show a clear contrast between non-ferroan and ferroan dolomite. Non-ferroan dolomite cements have low Fe

contents (0.05 – 4.0 mol %), whereas, ferroan dolomite is Fe-rich (5.0 - 17.0 mol %). Both ferroan and non-ferroan dolomite cements are observed in the two studied wells, but they are more common in the EL-1 well.

Calcite cement is the second most abundant carbonate cement (varying from 0.0 to 15.5%, averages 2.2%). Two successive calcite phases were recognised. Calcite (I) is microcrystalline cement which fills large intergranular pores, about 100 μm long (Fig. 7C). Calcite (I) predates the development of quartz overgrowths, it precipitated around detrital quartz and other grains, no evidence of quartz cement growing prior to the development of calcite (I) cement. Rarely, calcite (I) crystal has a shape similar to that of framework grains (Fig. 7C). Calcite (II) is also a microcrystalline cement that fills intergranular pores, however, it engulfs quartz overgrowths (Fig. 7D), indicating that it post-dates quartz cementation. Calcite (II) is observed under optical microscopy as replacement mineral. EDX microanalysis indicates that both types of calcite are non-ferroan calcite. Calcite II is less distributed than calcite I in the samples analysed. Both generations of calcite were observed only in samples from EL-1 well.

Based on morphology, three types of authigenic siderite are identified. Siderite (I) is a microcrystalline cement which primarily occurs on the rims of detrital grains ($>2\ \mu\text{m}$ thick) and predates quartz overgrowths (Fig. 7E). This morphology is observed only in samples from GH-1 well. Siderite (II) is a massive ($\geq 20\ \mu\text{m}$), subhedral cement; it fills intergranular pores and in some examples, postdates quartz overgrowths (Fig. 7F). Siderite II is rare but observed in both wells. Siderite (III) forms as aggregate cement, sometimes partially dissolved cement (Fig. 7G), overgrown pore filling clay cements and microcrystals of dolomite. Siderite (III) is observed in samples from both wells.

4.3.2 Clay mineral cements

Based on SEM morphological characteristics and XRD mineralogical analysis, three main clay minerals were identified in samples from Collyhurst Sandstone Formation: illite, chlorite and kaolinite (Fig. 8). Each clay mineral has varying textures and concentrations. Point count data identifies an average of 4.2% total clay mineral cement concentration in both wells. In both studied well, clay mineral cement concentrations show a similar trend of increasing with depth (Fig. 6B).

Authigenic illite is common in EL-1 and less abundant in GH-1. Morphologically, the illite cement has flaky and fibrous crystals. The flaky crystal is $\leq 5 \mu\text{m}$ in size (Fig. 9A) occurring as grain coating perpendicular to the grain surface (Fig. 9B) and also sometimes developing into pore-bridging cements (Fig. 9B). Grain coating illite is well developed in medium to coarse-grained sandstone (lithofacies B) of the EL-1 well. Under optical microscopy, illite coats can be up to $50 \mu\text{m}$ thick (Fig. 9C), which are largely absent at grain contacts. Detailed SEM analysis on stubs and thin sections further revealed a concentrically laminated morphology of clay coats (Fig. 9D). This morphology is not pervasive in the sandstones. Energy dispersive X-ray indicates the presence of mixed illite-smectite composition (Fig. 9E) preserved in the coated grains, which complement XRD result with an illite-smectite peak (Fig. 8). Illite grain coating minerals are present in most samples from EL-1 well, but rare in samples from GH-1. Continuous growth of illite cement developed into the fibrous morphology of interlocking mesh-work, bridging and filling intergranular pores (Fig. 9B).

Chlorite cement is observed in samples from both wells, although more common in EL-1 well than GH-1. Authigenic chlorite developed as pore-lining and pore-filling cement. Pore-filling chlorite is mostly observed in samples from GH-1 whereas samples from EL-1 revealed both pore-lining and pore-filling chlorite. SEM observation of stub samples revealed

the chlorite as microcrystalline with rosette morphology (Fig. 9F). Pore-lining chlorite nucleated parallel on detrital grains surfaces (Fig. 9G), it is also observed to grow after illite coats on some grains (Fig. 9E) and appear to inhibit the development of quartz overgrowth (Fig. 9H). Continuous growth of pore-lining chlorite cements developed into pore-filling cements growing in between grains and sometimes completely fills intergranular pores (Fig. 9G). Pore-filling chlorite sometimes shows fan-like habit with individual crystals longer than (up to 20 μm) the earlier formed pore-lining chlorite (Fig. 9G). EDX scans identify the chlorite cement as Fe-rich type (chamosite). Textural and microscopic evidence indicates that chlorite occasionally forms with authigenic illite (Fig. 9F).

A vermicular form of kaolinite is observed, occurring as pseudo-hexagonal, pore-occluding cement with individual crystal up to 5 μm long (Fig. 9I). A stacked aggregate of vermicular kaolinite (>10 μm) can fill intergranular pore. Kaolinite cement was observed to replace detrital feldspar and mica grains. Kaolinite appears to inhibit the growth of chlorite crystals (Fig. 9I), and thus predates the development of chlorite cement. Kaolinite is abundant in samples from GH-1 well and not observed to postdate quartz overgrowth.

4.3.3 Quartz cement

Quartz cement was identified with point-counting and more common in samples from EL-1 than GH-1 well. Quartz cement tends to decrease with depth within the two wells; (Fig. 6C). Quartz cements have pore-lining and pore-filling morphologies. Pore-lining quartz cements form euhedral, syntaxial overgrowths (up to 200 μm long) on detrital quartz grains. Pore-filling quartz cements form anhedral overgrowths, which sometimes completely fill primary pores (Fig. 10A). The pore-lining, syntaxial quartz overgrowths were recognised in the majority of samples from the two wells, while about three samples had pore-filling quartz overgrowths. Quartz overgrowths often developed where clay rims are thin or discontinuous

(Fig. 10A). Quantitative point-count data shows that quartz cement concentrations vary from 0.0 to 5.5% with an average of 2.2%.

4.3.4 Other cements

Barite and anhydrite are minor (<1%) sulphate cements in both studied wells. Barite was observed in both wells and occur as poikilotopic, pore-filling cement (<20 μm). It engulfs quartz overgrowths but replaced by authigenic albite (Fig. 10B). Observation of barite in some samples revealed that it developed after clay mineral cements (illite and chlorite) and Fe-dolomite. It also replaced calcite (II) cement, suggesting it precipitated after late calcite.

Anhydrite cement is common in GH-1 well but was observed in only one sample from the EL-1. Anhydrite occurs as monocrystalline cement filling large intergranular pores (>60 μm across). It engulfs quartz overgrowths, suggesting it postdates quartz cement precipitation (Fig. 10C).

Minor authigenic albite cement (<1%) occurs as overgrowths on detrital plagioclase feldspar and K-feldspars grain (Fig. 10D). It also occurs occasionally as pore-occluding cement (Fig. 10D) and sometimes replaced pore-filling anhydrite (Fig. 10C). Optical microscopy revealed partial albitization of some detrital K-feldspar grains. Authigenic albite is present in both studied wells.

Haematite minerals enclosed almost all sand grains in the studied sandstones. It is present in both wells, but mostly common within the medium to coarse-grained sandstone of EL-1 well. Haematite occurs as a reddish-brown colour on grain rims under an optical microscope and mostly absent at the point of grain contact (Fig. 10E). SEM observation revealed small crystals of haematite (up to 5 μm long) growing from the breakdown of ferromagnesian minerals (Fig. 10F). Authigenic illite/chlorite and dolomite are sometimes

associated with haematite cement. XRD analysis also indicates the presence of haematite on samples from both wells.

4.3.5 Mineral dissolution

Dissolution of feldspars, unstable lithic grains and leaching of authigenic minerals were observed in both studied wells, resulting in secondary porosity. Analysed samples show intense leaching and partial dissolution of K-feldspar and plagioclase feldspar (Fig. 11A and B), contributing to intragranular porosity. The growth of authigenic illite is noticed in some feldspar grains that have undergone almost complete dissolution (Fig. 11B). Alterations of unstable minerals such as mica (muscovite and biotite) were also noticed (Fig. 11C and 11D) as well as the partial dissolution of carbonate cement (Fig. 11E).

4.3.6 Compaction

The extent of compaction during diagenesis of the Collyhurst Sandstone Formation was assessed from petrographic observations and by calculating the degree of porosity loss by compaction and cementation using the method of Houseknecht (1987). Petrographic observations identify mica grain bending (Fig. 11F) and the fracturing of quartz grains. Common straight and concave-convex grain contacts between detrital framework grains also indicate extensive compaction. Sutured grains and stylolite features are observed in some samples (Fig. 3C and 3D).

Calculation of the degree of porosity loss by compaction and cementation (Houseknecht, 1987) is performed by plotting the percentage of original porosity destroyed by compaction (mechanical and chemical) against the percentage of original porosity destroyed by cementation. This approach assumes the original porosity of the sandstones (sand) was 40%. The percentage of original porosity destroyed by compaction is calculated as:

$$\frac{40 - \text{intergranular volume}}{40} \times 100$$

Where, intergranular volume is expressed as the percentage of the whole rock,
And the percentage of the original porosity destroyed by cementation is calculated as:

$$\frac{\text{cement}}{40} \times 100$$

Where, cement refers to the volume of cement expressed as the percentage of the whole rock.

Also,

$$\text{intergranular porosity} = \text{intergranular volume} - \text{cement}.$$

The resulting plot of intergranular volume (IGV) against cement volume for EL-1 and GH-1 is shown in (Fig. 12A and 12B) respectively. The plot indicates that over 60% of the samples from EL-1 well and 100% of GH-1 well falls between the 0% and 10% intergranular porosity lines. On the basis of lithofacies, almost all the samples from lithofacies A-D falls within 0 to 10% intergranular porosity line, while most samples from lithofacies E (matrix-free sandstones) fall mostly on 10-20% intergranular line with two samples slight above this. GH-1 well predominantly consist of lithofacies A (fine-grained sandstone and mudstones) and all samples analysed here are between the 0% and 10% intergranular porosity lines, indicating higher compaction rate than cementation. Generally, an average value for EL-1 data is 20.6% intergranular volume and 13.1% of cement, whereas it is 13.6% intergranular volume and 8.1% of cement for GH-1 samples. Further application of the above equation to obtain the approximate rate (percentage) of compaction and cementation in the studied well is express under the discussion section.

4.4 Porosity and permeability

Comparison of the values obtained from helium porosity measurements and point count porosity for EL-1 well is in a similar range (Fig. 13A). However, some samples have slightly higher point counting data porosities than measured helium porosity. Helium porosity

measurements vary from 1.2 to 16.0% with an average of 6.6%, while point count porosity varies from 0.5 to 23.0% with an average of 8.4%. Both helium and point count porosity decreases with depth (Fig. 12A). Horizontal permeability values range from 0.008-6.707 mD with an average of 0.387 mD for EL-1 well samples. Analysis of core permeability data also shows a decreasing trend with increasing burial depth (Fig. 13B).

Petrographic observations identify two pore types in the Collyhurst Sandstone Formation; intergranular and intragranular. Intergranular pores (Fig. 14A) are more common (<1% to 21%, with an average of 8.4%) than intragranular pores (<1% to 8.5%, with an average of 2.6%). Intragranular porosity is mainly associated with the dissolution of feldspar and other reactive detrital and authigenic minerals (Fig. 14B). Both types of porosity are common in samples from EL-1 well, while micro-fracture porosity with small pore aperture is also noticed in some samples from GH-1 well (Fig. 14C).

4.5 Carbon and oxygen stable isotope data

Results of carbon and oxygen isotope analysis on selected dolomite cements mainly from lithofacies C (Table 1, Fig. 15), show that early dolomite (non-ferroan) have distinctive isotopic values from later dolomite (ferroan). $\delta^{13}\text{C}$ value for non-ferroan dolomite ranges between -0.48 to -2.37‰ (average of -1.45‰) and $\delta^{18}\text{O}$ value of -2.01 to -4.51‰ (average of -2.82‰). For ferroan dolomite cement, $\delta^{13}\text{C}$ value is between -2.18 to -9.22‰ (average of -6.59‰) and $\delta^{18}\text{O}$ value of -4.39 to -7.90‰ with an average of -5.44‰ .

5. Discussion

5.1 Paragenetic sequence

The textural relationships between the diagenetic minerals, combined with the isotopic composition of selected phases were used to reconstruct an overall paragenetic sequence of the studied reservoir (El-Ghali *et al.*, 2009; Karim *et al.*, 2010). The main diagenetic minerals

identified within this paragenesis are cementation by carbonate, authigenic clay minerals and quartz cements. The paragenetic sequence for both wells follows a similar trend with slight differences (Fig. 16). However, diagenetic alteration varies between different lithofacies. The diagenetic pathway through each lithofacies is summarized in (Table 2).

5.1.1 Early diagenesis

Early diagenesis refers to processes that affect sediments after deposition due to their proximity to the surface (Choquette and Pray, 1970), often a regime with a maximum depth limit of 1-2 km and temperature range of 30 °C to 70 °C (Worden and Burley, 2003). Depositional composition has a major control on early cementation within the studied reservoir.

Coatings of haematite around sand grains (Fig. 10E) are interpreted to be early diagenetic in origin with precipitation occurring shortly after deposition. Precipitation of early haematite is common in red-bedded sandstones (e.g., Hubert and Reed, 1978; Ixer *et al.*, 1979; Burley *et al.*, 1985; Morad *et al.*, 1995; Chan *et al.*, 2005), and has been interpreted to be the result of ferrous iron oxidation released during the dissolution of ferromagnesian minerals (Fig. 10F and 11D; Robb, 1949; Walker *et al.*, 1981).

Calcite (I) and non-ferroan dolomite are dominant carbonate cement occurring as pore-filling cement in both studied wells. Petrographic evidence indicates that they developed prior to precipitation of quartz cement, as there is no evidence that the cements has engulfed quartz overgrowth (Fig. 7B and 7C), but mostly has direct contact with detrital framework grains. This suggests that their temperature of the formation is lower than typical lower temperatures (~60 °C) required for quartz cement precipitation (Walderhaug, 1994). Carbon and oxygen stable isotopic data (Table 1, Fig. 15) for early dolomite cement (non-ferroan dolomite) of $\delta^{13}\text{C}$ (−0.48 to −2.37‰ VPDB) and $\delta^{18}\text{O}$ (−2.01 to −4.51‰ VPDB) are compatible with a carbon source dominated by meteoric fluids (Morad, 1998). The oxygen

isotope values in this study fall closely to the negative values (−2.4 to −5.9‰) obtained by Bath *et al.* (1987) for dolomite cement of Permo-Triassic sandstone in South Yorkshire. Meteoric water composition for the study area has not been properly documented. However, previous authors have estimated $\delta^{18}\text{O}$ of −3‰ (SMOW) for Triassic-Permian continental meteoric (e.g., Naylor *et al.*, 1989; Greenwood and Habesch, 1997). With respect to the estimated $\delta^{18}\text{O}$ water composition of −3‰ SMOW, a precipitation temperature of 27 °C to 39 °C is calculated for early dolomite cements in the Collyhurst sandstone (Fig. 17).

The absence of or low Fe content in calcite (I) and non-ferroan dolomite as revealed by EDX attached to SEM analysis and bright cathodoluminescence can further be used to explain the occurrence of the two cements as early carbonate cements (cf., Schmid *et al.*, 2006). It also suggests precipitation in an oxygen-rich environment which characterises vadose zone carbonates (Morad, 1998). Carbonate in a continental environment can either be pedogenic, vadose or phreatic in origin (e.g., Khadkikar *et al.*, 1998; Beckner and Mozley, 2009). The studied samples have no evidence of a pedogenic origin, such as cemented root cast, root trace or nodules. The reaction of Ca^{2+} and Mg^{2+} from the breakdown of unstable minerals and plagioclase feldspars with CO_3^{2-} derived from organic matter in the vadose zone (Morad, 1998) likely precipitated early calcite and dolomite in this study. The organic carbonate could originate from the accumulation of skeletal part of organisms, photosynthesis and transpiration of CO_2 by plant or through elevated CO_2 by bacteria (Schmid *et al.*, 2006). Additionally, petrographic observation shows that these carbonates decreased with depth (Fig. 6A), it can then be proposed that some of the ions required for the precipitation of these carbonates may be derived from overlying evaporites.

Siderite (I) occur as a coating on detrital grains (Fig. 7E), common in GH-1 well. Grain coating siderite can result from marginal replacement of detrital grains (mostly quartz grains in this study) as more Fe^{2+} become present in the pore fluid, whereas Mg^{2+} is reduced (Pye

and Krinsley, 1986). From petrographic observation, distribution of dolomite cement is less than siderite in GH-1 well. Therefore, it can be inferred that occurrence of minerals such as ilmenite and other ferruginous minerals with high iron composition common in the well could release Fe^{2+} during early burial such that siderite (I) is precipitated.

Petrographic observations show that vermicular kaolinite is common in samples from GH-1 well, occurring as pore-occluding and pore-filling cement (Fig. 9I). Vermicular kaolinite is typical of shallow to intermediate burial diagenesis (Giles *et al.*, 1992; Lanson *et al.*, 2002; Schmid *et al.*, 2004) whereas, blocky kaolinite/dickite precipitate upon deeper burial and higher temperature (Gaupp *et al.*, 1993; Beaufort *et al.*, 1998; Rossi *et al.*, 2001). Authigenic kaolinite in sandstones may be from various sources: (i) crystallization of materials introduced from solutions, (ii) recrystallization of detrital kaolinite clay (fine-grained) and (iii) from alteration of parent mineral rocks (Shelton, 1964). Authigenic kaolinite from crystallization of solution materials often precipitates as vug or cavity infilling and along fractures (Shelton, 1964). However, such morphology is not evident in the studied sandstone. The occurrence of detrital kaolinite is also lacking, suggesting that the observed authigenic kaolinite is not from recrystallization. Alteration of parent minerals such as feldspar (K-feldspar and plagioclase) is the likely source of authigenic kaolinite in this study. Two theories can be used to explain the formation of kaolinite from detrital feldspar. The first is termed “meteoric-water flushing” process, associated with early diagenesis. In this process, kaolinite precipitates at shallow depth by meteoric fluid which flushes the formation (Hancock and Taylor, 1978; Lanson *et al.*, 2002) and it is typical of vermicular kaolinite morphology. In addition to meteoric fluids, the second hypothesis considered organic acid-rich fluids as part of the fluid required for feldspar alteration, which subsequently leads to kaolinite precipitation. Such fluids may be derived from organic matter from adjacent mudstones beds. The temperature range estimated for organic-rich fluid expulsion is $\sim 100^\circ\text{C}$ (Platt, 1993),

which is compatible with temperature for the precipitation of blocky kaolinite/dickite (high temperature kaolinite). Presence of vermicular kaolinite morphology in this study indicates precipitation during early diagenesis under the influence of meteoric fluids.

5.1.2 Late Diagenesis

Late diagenesis refers to processes which are isolated from surface-related processes (Schmid *et al.*, 2004), with burial depth from 1-2 km and temperature up to 200-250 °C (Worden and Burley, 2003). Late diagenetic minerals in the Collyhurst sandstone include late carbonate (calcite II, ferroan dolomite, siderite II and III), chlorite, illite, quartz overgrowth, albite, and sulphates (barite and anhydrite).

Fe-dolomite/ankerite often grows as an outer zone on non-ferroan dolomites, implying that it developed at later temperature range during burial diagenesis than non-ferroan dolomite. Late diagenetic carbonate often contains a significant amount of iron (II) composition (Machent *et al.*, 2007). Considering Fe^{2+} content as a proxy for burial cementation, Fe-dolomite/ankerite in the Collyhurst sandstone indicate late carbonate phase. Studies from the adjacent Sherwood Sandstone Formation identified precipitation of ankerite at a temperature range of 110 °C and 165 °C (Schmid *et al.*, 2004). Upon progressive burial, temperature increases while oxygen level becomes depleted, Fe^{2+} is released into the pore water as Ca^{2+} and Mg^{2+} decreases, resulting in precipitation of ferroan dolomite/ankerite. Presence of Fe^{2+} in this carbonate phase suggests a significance of iron (III) reduction rather than sulphate reduction. A thorough examination of samples from the studied wells indicates an absence of pyrite cement which inferred sulphate depletion or low sulphate concentration in the deeper sequence of the sediments. According to Curtis (1987), such an event is dominated by microbial methanogenesis which possibly contributed to the late ferroan carbonate. There is no evidence of late dolomite replacing earlier formed calcite contrary to the findings in some similar sandstones beds (e.g., Glennie *et al.*, 1978).

Carbon and oxygen isotope value obtained for later dolomite (ferroan dolomite) cements are lower than those for the early dolomite; $\delta^{13}\text{C}$ varies from -2.18 to -9.22‰ VPDB and $\delta^{18}\text{O}$ range from -4.39 to -7.90‰ VPDB (Table 1, Fig. 15). The low $\delta^{18}\text{O}$ value for the later dolomite cements still consistent with carbonate precipitation from meteoric fluids (Taylor *et al.*, 2000; Moore, 2001), but precipitated at a slightly higher temperature of $39\text{--}58\text{ }^{\circ}\text{C}$ (Fig. 17) upon continual burial. The isotopic value from one of the samples (sample 56516, Table 1) indicates that some of the ferroan dolomite precipitated during early diagenesis which is in agreement with petrographic textural relationship observed in the analysed samples (Fig. 16).

Petrographic textural observation revealed siderite II and III as late carbonate. Siderite (II) developed after precipitation of quartz overgrowth (Fig. 7F), whereas siderite (III) overgrown pore-filling illite (Fig. 7G). Late siderite possibly precipitated due to further enrichment of Fe^{2+} ion and depletion of Ca^{2+} and Mg^{2+} in the pore water. Purvis (1992), documents carbonate phases of dolomite – ferroan dolomite – siderite in his study where a further increase of Fe^{2+} in the pore water and less Mg^{2+} and Ca^{2+} was responsible for siderite precipitation as a stable and last phase. Additionally, ilmenite is a common mineral in the studied wells (mostly GH-1), alteration of this may account for more Fe^{2+} needed for siderite formation. According to Karim *et al.* (2010), Fe^{2+} in siderite is often derived from iron hydroxides or ilmenite alteration.

Textural relationships revealed that calcite (II) engulf and postdates quartz overgrowths (Fig. 7D), suggesting a temperature of precipitation higher than that of calcite (I) and quartz overgrowths. The study by Greenwood and Habesch (1997) on surrounding wells (Sherwood Sandstone Group) indicates precipitation of late calcite cement at temperature up to $110\text{ }^{\circ}\text{C}$. The result from their isotopic data showed that CO_3^{2-} derived from organic source

was incorporated into the pore water which precipitated late calcite cement. A similar condition might possibly precipitate late calcite cement in Collyhurst Sandstone Formation.

Illite occurs as grain coats on detrital grains and as pore-filling cement (Fig. 9A – 9C). It sometimes has an illite-smectite composition (Fig. 9D – 9E). Petrographic observations indicate that clay coats are largely absent at grain contacts (Fig. 9C), suggesting a post-deposition process. Two mechanisms can account for the development of the observed illite; replacement of kaolinite or replacement of smectite. Illitization of kaolinite often develops at a temperature above 120 °C (usually between 120-140 °C) with a low precipitation rate (Bjørlykke *et al.*, 1995). The fact that grain coating illite precipitated prior to significant quartz cements with a temperature of formation around 60 °C (Bjørlykke and Egeberg, 1993; Walderhaug, 1994) suggests that kaolinite is not the probable precursor mineral for the observed illite. Additionally, petrographic observation indicates that illite is abundant in eastern well, while kaolinite is absent, thereby supporting the first concept. Conversely, the transformation of smectite to illite is recorded to occur below 90 °C (Worden and Morad, 2003), may even form at a very low temperature between 20-30 °C and depth of 500 m (Buatier *et al.*, 1992). This hypothesis support smectite as likely precursor mineral in addition to observed flaky morphology (Fig. 9A), which is usually associated with smectite mixed layer precursor (Pollastro, 1985; Storvoll *et al.*, 2002). Whereas, sheet-like morphology is common with illite from kaolinite source (Pollastro, 1985). Upon progressive burial, modification of earlier formed illite (flaky morphology) occurs, such that illite with fibrous intergrown texture is developed, which mostly bridge and fill the pore spaces (Figure 9B, Pollastro, 1985). Illitization of K-feldspar is another possible reaction in this study, microscopic observation revealed the growth of illite from the breakdown of detrital K-feldspar (Fig. 11B, EL-1 well). Such reaction could occur in the absence of kaolinite but under

acidic condition (eq (1), McAulay *et al.*, 1993), since kaolinite is not observed as present in the eastern well.

Acid fluids needed for feldspar dissolution must have been expelled from the underlain Carboniferous Coal Measures (Platt, 1993).

Chlorite developed as pore-lining and pore-filling cement in both studied wells. Microscopic observation shows that chlorite developed after grain coating illite (Fig. 9E) and occasionally show synchronized texture with illite (Fig. 9F), suggesting that both minerals probably precipitated at a similar time (same temperature range). Different sources of chlorite cement in sandstone have been documented (e.g., Worden and Morad, 2003; Ziegler, 2006; Dowey *et al.*, 2012). Chlorite can develop from precursor clay minerals such as berthierine, odinite, kaolinite, and smectite at a temperature range of 70 °C to 100 °C (Hillier and Velde, 1992; Ehrenberg, 1993; Bloch *et al.*, 2002; Worden and Morad, 2003). Dissolution of iron and magnesium-rich grains, and breakdown of volcanic rock fragments can also lead to chlorite precipitation (Thomson and Stancliffe, 1990; dos Anjos *et al.*, 2000; Worden and Morad, 2003). In this study, there is no evidence of berthierine and odinite clay minerals, suggesting that chlorite cement probably originated from a smectite precursor as applicable to illite cement. Illite coating is often generated from dioctahedral smectite while chlorite is usually from the more iron-rich trioctahedral smectite (Pollastro, 1985; Worden and Morad, 2003). ESEM analysis revealed both forms of chlorite as iron-rich; supporting the fact that Fe^{2+} could have been from trioctahedral smectite. Dissolution of iron and magnesium-rich detrital grains present in analysed samples (Fig. 10F and 11D) can also contribute to chlorite formation in the studied sandstones.

Quartz overgrowth developed during burial of the studied sandstone. Petrographic evidence revealed concave-convex, long grain suture and stylolite contacts (Fig. 3C, 3D and 5E) of detrital grains, indicating that pressure solution possibly accounts for some of the quartz cementation. Grain coating/pore lining chlorite is however observed to somewhat inhibit the precipitation of quartz overgrowth (Fig. 9H). Other possible sources of silica for quartz overgrowth precipitation in this study include dissolution of feldspar and unstable rock fragments (cf., Burley, 1984; Bjorlykke and Egeberg, 1993; Walderhaug, 1996).

Authigenic albite is rare, growing mostly as an overgrowth on detrital K-feldspar and plagioclase feldspar (Fig. 10D). Albitization of feldspar is often a late diagenetic phase (Strong and Milodowski, 1987; Ramseyer *et al.*, 1992; Strong, 1993) precipitating at high temperature up to 120 °C (Schmid *et al.*, 2004). Pressure dissolution of detrital plagioclase feldspar grains with mutual grain contacts (Fig. 10D) can release the Na^+ , Al^{3+} and Si^{4+} for albite reprecipitation.

Both sulphate cements (anhydrite and barite) are late cements. Microscopic observation indicates that anhydrite developed earlier than the growth of barite cement (Fig. 10C). Dissolution or re-precipitation of early gypsum cement can generate late anhydrite precipitation due to its high solubility (Holliday, 1970; Greenwood and Habesch, 1997). However, there is no evidence for early gypsum or anhydrite in the Collyhurst Sandstone Formation. Strong *et al.* (1994), suggested that fluid for anhydrite precipitation can be supplied from evaporates within, above or beneath a sequence. The Collyhurst Sandstone Formation is overlain by the Cumbrian Group (an equivalent of Permian Zechstein), therefore Manchester Marl or St. Bees Evaporites (Fig. 2) could probably supply the fluids needed for sulphate precipitation.

5.2 Diagenetic controls on reservoir quality

Reservoir quality is a function of various depositional and diagenetic controls including sorting, grain size, cementation, dissolution and compaction (Schmid *et al.*, 2004; Bjørlykke and Jahren, 2010; Xi *et al.*, 2015). In the studied sandstones, reservoir quality is mainly controlled by compaction followed by cementation. Petrographic analysis revealed that most rigid grains have concave-convex, sutures and stylolite point contacts (Fig. 3C and 3D), while ductile grains are compacted around more rigid grains (Fig. 5B). These observations support the result that compaction has greatly influenced the sandstones reservoir quality. Petrographic observations and the plot of intergranular volume against cement (Fig. 12) is evidence that the degree of compaction is higher in GH-1 than in EL-1 well. The GH-1 well predominantly comprises of fine-grained sandstones and silty mudstones, thus faster deformation of ductile grains (e.g. mica and clay matrix) occurs, leading to increasing compaction.

A plot of intergranular volume against cement (Fig. 12A and 12B) revealed that over 60% of the samples from EL-1 well and 100% of GH-1 well falls between the 0% and 10% intergranular porosity lines, suggesting that compaction processes rather than cementation account for the major destruction of the initial porosity (cf., Houseknecht, 1987; Lander and Walderhaug, 1999). It can also be deduced from the plot that more compaction occurs within samples from lithofacies A-D than lithofacies E. The average values for the data from EL-1 well are 20.6% intergranular volume and 13.1% of cement. Considering the evaluation method of Houseknecht (1987), this indicates that 48.5% of original porosity has been destroyed by compaction (mechanical and chemical), whereas 32.8% has been destroyed by cementation assuming the original porosity was 40%. For GH-1 well, average data revealed intergranular volume of 13.6% and cement of 8.1%. This indicates that 66.1% of original

porosity has been destroyed by compaction (mechanical and chemical), whereas only 20.2% has been destroyed by cementation.

The assessment of porosity loss (by compaction and cementation) support petrographic evidence between the two studied wells, revealing higher compaction rate in GH-1 well than in EL-1 well, which is largely influenced by lithofacies variation. In this study, compaction contributed to permeability loss within the reservoir, deformation and bending of lithic grains occur, ductile grains get crushed and extruded between rigid sands grains, thus blocking pore throats and reducing the pore connectivity (cf., Islam, 2009).

Other than compaction, cementation also severely reduced the reservoir quality of the studied sandstone through the development of early carbonate cement (mainly early dolomite and calcite) and clay mineral cement (mostly late clay cement). Carbonate cement accounts for almost half of the total observed cements, indicating that carbonate cement has a significant impact on reservoir quality of the studied formation. Dolomite occurs in almost all the samples analysed from the two wells, filling intergranular pores (Fig. 7A and 7B), blocking pore throats and thereby reducing reservoir quality. Early calcite completely filled intergranular pores (Fig. 5C, 5D and 7C) preventing macro-porosity and migration of pore fluid.

Authigenic clay mineral precipitation can significantly influence sandstone reservoir properties. The relative abundance of illite in the eastern well (EL-1) and kaolinite in the western well (GH-1) contribute greatly to the reduction of reservoir quality of the studied sandstone. Flaky illite (Fig. 9A and 9B) tends to sit between grains, occluding pores and blocking pore throats, thereby destroying the reservoir quality. Authigenic illite usually reduces permeability in sandstone reservoir (e.g., Kantorowicz, 1990; Gaupp *et al.*, 1993; Leveille *et al.*, 1997). Vermicular kaolinites also occlude and fill intergranular pores (Fig. 9I), thus reducing the sandstone permeability. Pore-lining chlorite possibly has a slight effect in

inhibiting the development of quartz overgrowth, evidence in some samples (Fig. 9H), likely maintained the reservoir quality to a certain degree (cf., Ehrenberg, 1993; Ziegler, 2006). However, pore-filling chlorite tends to reduce the pore throat diameter, resisting fluid flow.

Quartz cementation has been recognized as one of the main cause of porosity loss in many sandstone reservoirs (e.g., Blatt, 1979; McBride, 1989; Ehrenberg, 1990; Swarbrick, 1999). The quartz cementation in this study occurs in moderate amounts (average 2.2%); it probably has little effect in reducing the reservoir quality due to the dominating effect of carbonate and clay mineral cements. Sulphate cement (barite and anhydrite) and albite occur as minor cement (<1%), and they will have a negligible effect on the reservoir quality.

5.4 Reservoir quality prediction within various lithofacies

Fig. 18 shows the reservoir quality variation (porosity and permeability) within different lithofacies present in the studied sandstone. Lithofacies have a significant impact on the reservoir quality of the studied sandstone. A predictive model for each lithofacies against reservoir properties (Fig. 18 and 19) is focused on EL-1 well due to the heterogeneity of samples in this well and the availability of permeability data. GH-1 well is dominated by homogenous fine-grained sandstone and mudstone. The cross-plot of each lithofacies against the porosity and permeability is useful in understanding the reservoir properties of the studied sandstone (Fig. 18) as well as predicting the reservoir quality to greater depth (Fig. 19). The diagenetic pathway through each lithofacies is illustrated in (Table 2).

Evidence from petrographic data indicates that a greater percentage of reservoir quality is lost through compaction. Generally, the initial porosity for the studied sandstones is estimated to be about 40% as applicable to most sandstones (e.g., Beard and Weyl, 1973; Houseknecht, 1987). However, the point-count porosity of the analysed samples ranges from 0.5% to 23.0% with an average of 8.2%; whereas horizontal permeability measurement ranges from 0.008 mD to 6.707 mD with an average of 0.387 mD for EL-1 well samples (Fig.

18). Porosity of most of the studied samples is below 10% and permeability is less than 1 mD (Fig. 18), compatible with reservoir properties of a tight gas sandstones (Newman, 1999; Holditch, 2006; Smith *et al.*, 2009).

In lithofacies A, mechanical infiltration of detrital clay occurs shortly after deposition, enclosing most grain surfaces or forming an argillaceous matrix. The mud intraclasts are easily compacted, leading to a greater reduction of intergranular porosity. Although the porosity of these mud-supported sands would have been high at deposition, however, it decreases extensively during progressive compaction (cf., Lai *et al.*, 2017). Due to the abundance of detrital matrix and ductile grains such as mica (Fig. 5A), mechanical compaction continued during late burial diagenesis, resulting in further decrease of intergranular porosity (Fig. 18 and 19). Clay mineral cement is common within this lithofacies contributing to a greater reduction in porosity.

Lithofacies B (conglomeratic sandstone) is characterized by poorly-sorted sandstone with dominant lithic grains (Fig. 5B). Mechanical compaction is dominant here, ductile grains tend to squeeze into more stable grains (Fig. 5B) and small grains could fit into intergranular pores due to their poor sorting, therefore the original porosity is not preserved (Fig. 18 and 19). Upon progressive burial, porosity loss is often common with poorly sorted sandstones (cf., Walderhaug *et al.*, 2012).

In Lithofacies C, the vast proportion of early carbonate cements (Fig. 5C, 5D and 7C) led to the deterioration of the reservoir properties, rather than creating a grain-supported framework for the sandstone. The rate of cementation in this lithofacies is higher than compaction. Precipitation of the early carbonate cement within this lithofacies reduces compaction. As diagenesis progress, compaction becomes limited, while precipitation of carbonates cement continued, resulting in later carbonate phase. Evidence from petrographic analysis revealed late calcite precipitation, developing after quartz overgrowth (Fig. 7D) and

late siderite also precipitated after the growth of clay minerals cements (Fig. 7F and 7G), which significantly contribute to porosity reduction.

Within quartz-cemented sandstone (lithofacies D), compaction dominates during early diagenesis. Chemical compaction is common, evident in samples with concave-convex, suture and stylolite contacts (Fig. 3C and 3D). Upon burial, quartz cementation prevails, sometimes quartz overgrowth completely fills intergranular pores (Fig. 5E) thereby reducing the sandstones porosity.

Lithofacies E (matrix-free sandstones) mainly consists of medium to coarse-grained and well-sorted sandstone. The highest percentage of porosity and permeability is recorded in samples from this lithofacies (Fig. 18 and 19). Mechanical compaction and mineral dissolution account for early diagenesis. Limited mechanical compaction is observed due to the abundance of stable grains (Fig. 5F). Grain dissolution is evidence through petrographic analysis where samples show the partial dissolution of feldspar and rock fragments (Fig. 5F, 11B and 14C) leading to secondary porosity and the rigid quartz grains also preserve primary porosity to large extent. Burial depth prediction model indicates that porosity is preserved to a certain extent within this lithofacies (Fig. 19).

In conclusion, it is evident from petrographic analysis and the cross-plotting of each lithofacies against core porosity and permeability data (Fig. 18 and 19) that lithofacies A have the poorest reservoir properties due to extensive compaction and cementation by clay minerals (illite and chlorite). Reservoir quality of lithofacies B (conglomeratic sandstone) appear a bit fairer than that of lithofacies A. Carbonate-cemented, lithofacies C reflects low permeability and low porosity due to extensive distribution of carbonate cement within the samples. Quartz cemented sandstone (Lithofacies D) show a range of reservoir properties which can be linked to varying degree of quartz cementation and compaction. Lithofacies E (matrix-free sandstone) has the highest reservoir properties and serves as the best reservoir

(Fig. 18 and 19). This lithofacies has limited compaction as well as less cementation resulting in higher primary porosity, dissolution of detrital grains also contribute to secondary porosity. Lithofacies E has a minimum porosity of 10% and average permeability is about 1.00 mD (Fig. 18). Lithofacies E has highest values of reservoir properties and therefore serves as a prospective target for exploration survey.

6. Conclusions

Two onshore core successions (Elswick-1 and Grange hill-1) from the Collyhurst Sandstone Formation, Northern England, in the East Irish Sea Basin have been analysed. Petrographic, mineralogical and stable isotopic evidence from the samples indicate that:

1. The tight-gas bearing aeolian and fluvial Collyhurst Sandstone Formation is mostly sub-arkosic to lithic arenite with very fine to very coarse-grained size and poor to moderate sorting. The reservoir properties are relatively poor with low porosity and permeability.
2. Five lithofacies were identified based on lithological characteristics and dominant diagenetic minerals: (A) fine-grained sandstone; (B) conglomeratic sandstone; (C) carbonate-cemented sandstone; (D) quartz-cemented sandstone; (E) matrix-free sandstone.
3. The tight sandstones have undergone significant early and late diagenetic processes. Early diagenetic minerals include: early carbonate cements (calcite I, siderite I, dolomite) and kaolinite are mostly pore-filling and mineral replacive cements. Late diagenetic minerals include: late carbonate cements (calcite II, ferroan dolomite and siderites II-III). Quartz overgrowths, illite, chlorite, albite, barite and anhydrite all developed as late cements.

4. Reservoir quality is greatly reduced by compaction and cementation. Compaction processes rather than cementation account for the major destruction of the initial porosity of the sandstone. For EL-1 well, 48.5% of original porosity has been destroyed by compaction (mechanical and chemical), whereas 32.8% has been destroyed by cementation. For GH-1, 66.1% of original porosity has been destroyed by compaction, whereas only 20.2% has been destroyed by cementation.
5. Carbonate cements and authigenic clay minerals are major pore-filling components; which significantly influenced the sandstone reservoir quality. Dolomite fills pores and block pore throats, thereby reduced reservoir quality. Early calcite completely filled intergranular pores, preventing macro-porosity and migration of pore fluid. Flaky illite sits between grains and block pore throats, it destroyed the reservoir quality. Vermicular kaolinites also occlude and fill intergranular pores, reducing the sandstone permeability. Pore-lining chlorite possibly has a slight effect in inhibiting the development of quartz overgrowth, thereby maintained the reservoir porosity to a certain extent. Dissolution of feldspar and lithic grains play a role in contributing to secondary (intragranular) porosity, enhancing the sandstone reservoir quality.
6. The study assisted in understanding various diagenetic processes and lithofacies variability and their subsequent control on reservoir quality of the Collyhurst tight-gas sandstone. The best reservoir quality is present in lithofacies E (clean sandstone) due to limited compaction and cementation as well as the development of secondary porosity.
7. Findings from this study provide conditioning information useful in predicting reservoir quality in similar tight-gas sandstone settings.

Acknowledgements

The authors would like to appreciate British Geological Survey and Cuadrilla Limited for the core data released. Special thanks to the Nigerian Federal Government through Tertiary Education Trust Fund (TETFund) and Schlumberger Foundation for their financial support of the study. We are grateful to John Waters and Jon Fellows for their assistance during the laboratory works. Steve Stockley is appreciated for thin sections preparation.

REFERENCES

- Bastin, J., Boycott-Brown, T., Sims, A., Woodhouse, R., 2003. The South Morecambe Gas Field, Blocks 110/2a, 110/3a, 110/7a and 110/8a, East Irish Sea. Geological Society of London Memoirs 20, 107-118.
- Bath, A., Milodowski, A., Spiro, B., 1987. Diagenesis of carbonate cements in Permo-Triassic sandstones in the Wessex and East Yorkshire-Lincolnshire Basins, UK: a stable isotope study. Geological Society of London, Special Publications 36, 173-190.
- Beard, D., Weyl, P., 1973. Influence of texture on porosity and permeability of unconsolidated sand. American Association of Petroleum Geologists Bulletin 57, 349-369.
- Beaufort, D. et al., 1998. Kaolinite-to-dickite reaction in sandstone reservoirs. Clay minerals 33, 297-316.
- Beckner, J., Mozley, P., 1998. Origin and spatial distribution of early vadose and phreatic calcite cements in the Zia Formation, Albuquerque Basin, New Mexico, USA. In: Morad, S. (Ed.), Carbonate Cementation in Sandstones. Special Publication, International Association of Sedimentologists 26, pp. 27-51.
- Bjørlykke, K., 2011. Open-system chemical behaviour of Wilcox Group mudstones. How is large scale mass transfer at great burial depth in sedimentary basins possible? A discussion. Marine and Petroleum Geology 28, 1381-1382.
- Bjørlykke, K., Aagaard, P., Egeberg, P.K., Simmons, S.P., 1995. Geochemical constraints from formation water analyses from the North Sea and the Gulf Coast Basins on quartz, feldspar and illite precipitation in reservoir rocks. Geological Society of London, Special Publications 86, 33-50.
- Bjørlykke, K., Egeberg, P., 1993. Quartz cementation in sedimentary basins. American Association of Petroleum Geologists Bulletin 77, 1538-1548.

- Bjørlykke, K., Jahren, J., 2010. Sandstones and sandstone reservoirs. In: K. Bjørlykke (Ed.), *Petroleum Geoscience: from Sedimentary Environments to Rock Physics*. Springer-Verlag Berlin Heidelberg, pp. 113-140.
- Blatt, H., 1979. Diagenetic Processes in Sandstones. *The Society of Economic Paleontologists and Mineralogists* 26, 141-157.
- Bloch, S., Lander, R.H., Bonnell, L., 2002. Anomalously high porosity and permeability in deeply buried sandstone reservoirs: Origin and predictability. *American Association of Petroleum Geologists Bulletin* 86, 301-328.
- Brandon, A. et al., 1998. *Geology of the Country around Lancaster: British Geological Survey. Memoir Sheet 59 (England and Wales)*, pp. 1-200.
- Buatier, M.D., Peacor, D.R., O'Neil, J.R., 1992. Smectite-illite transition in Barbados accretionary wedge sediments: TEM and AEM evidence for dissolution/crystallization at low temperature. *Clays and Clay Minerals* 40, 65-80.
- Burley, S., 1984. Patterns of diagenesis in the Sherwood Sandstone Group (Triassic), United Kingdom. *Clay Minerals* 19, 403-440.
- Burley, S.D., Kantorowicz, J.D., Waugh, B., 1985. *Clastic Diagenesis*. Geological Society of London, Special Publications 18, 189-226.
- Bushell, T., 1986. Reservoir geology of the Morecambe Field. Geological Society of London, Special Publications 23, 189-208.
- Chan, M.A., Bowen, B.B., Parry, W., Ormö, J., 2005. Red rock and red planet diagenesis: comparisons of Earth and Mars concretions. *Geological Society of America Today* 15, 4-10.
- Choquette, P.W., Pray, L.C., 1970. Geologic Nomenclature and Classification of porosity in Sedimentary Carbonates. *American Association of Petroleum Geologists Bulletin* 54, 207-250.

- Craig, H., 1957. Isotopic standards for carbon and oxygen and correction factors for mass-spectrometric analysis of carbon dioxide. *Geochimica et cosmochimica acta* 12, 133-149.
- Cowan, G. et al., 1999. Oil and gas migration in the Sherwood Sandstone of the East Irish Sea Basin, *Petroleum Geology of northwest Europe. Proceedings of the 5th Conference of Geological Society of London*, pp. 1383-1398.
- Curtis, C., 1987. Mineralogical consequences of organic matter degradation in sediments: inorganic/organic diagenesis. In: J.K. Leggett and G.G. Zuffa (Eds.), *Marine Clastic Sedimentology*. Graham and Trotman, London, pp. 108-123.
- Dixon, S., Summers, D., Surdam, R., 1989. Diagenesis and preservation of porosity in Norphlet Formation (Upper Jurassic), southern Alabama. *American Association of Petroleum Geologists Bulletin* 73, 707-728.
- dos Anjos, S.M., De Ros, L.F., de Souza, R.S., de Assis Silva, C.M., Sombra, C.L., 2000. Depositional and diagenetic controls on the reservoir quality of Lower Cretaceous Pendorcia sandstones, Potiguar rift basin, Brazil. *American Association of Petroleum Geologists Bulletin* 84, 1719-1742.
- Dowey, P.J., Hodgson, D.M., Worden, R.H., 2012. Pre-requisites, processes, and prediction of chlorite grain coatings in petroleum reservoirs: A review of subsurface examples. *Marine and Petroleum Geology* 32, 63-75.
- Ehrenberg, S., 1990. Relationship between diagenesis and reservoir quality in sandstones of the Garn formation, Haltenbanken, mid-Norwegian Continental shelf. *American Association of Petroleum Geologists Bulletin* 74, 1538-1558.
- Ehrenberg, S., 1993. Preservation of anomalously high porosity in deeply buried sandstones by grain-coating chlorite: examples from the Norwegian continental shelf. *American Association of Petroleum Geologists Bulletin* 77, 1260-1286.

- El-Ghali, M.A.K. et al., 2009. Distribution of diagenetic alterations within depositional facies and sequence stratigraphic framework of fluvial sandstones: evidence from the Petrohan Terrigenous Group, Lower Triassic, NW Bulgaria. *Marine and Petroleum Geology* 26, 1212-1227.
- Fic, J., Pedersen, P.K., 2013. Reservoir characterization of a “tight” oil reservoir, the middle Jurassic Upper Shaunavon Member in the Whitemud and Eastbrook pools, SW Saskatchewan. *Marine and Petroleum Geology* 44, 41-59.
- Floodpage, J., Newman, P., White, J., 2001. Hydrocarbon prospectivity in the Irish Sea area: insights from recent exploration of the Central Irish Sea, Peel and Solway basins. Geological Society of London, Special Publication 188, 107-134.
- Gaupp, R., Matter, A., Platt, J., Ramseyer, K., Walzebuck, J., 1993. Diagenesis and fluid evolution of deeply buried Permian (Rotliegende) gas reservoirs, northwest Germany. *American Association of Petroleum Geologists Bulletin* 77, 1111-1128.
- Giles, M. et al., 1992. The reservoir properties and diagenesis of the Brent Group: a regional perspective. Geological Society of London, Special Publications 61, 289-327.
- Glennie, K., Mudd, G., Nagtegaal, P., 1978. Depositional environment and diagenesis of Permian Rotliegendes sandstones in Leman Bank and Sole Pit areas of the UK southern North Sea. *Journal of the Geological Society* 135, 25-34.
- Greenwood, P., Habesch, S., 1997. Diagenesis of the Sherwood Sandstone Group in the southern East Irish Sea Basin (Blocks 110/13, 110/14 and 110/15): constraints from preliminary isotopic and fluid inclusion studies. Geological Society of London, Special Publications 124, 353-371.
- Hancock, N., Taylor, A., 1978. Clay mineral diagenesis and oil migration in the Middle Jurassic Brent Sand Formation. *Journal of the Geological Society* 135, 69-72.

- Herries, R.D., Cowan, G., 1997. Challenging the 'sheetflood' myth: the role of water-table-controlled sabkha deposits in redefining the depositional model for the Ormskirk Sandstone Formation (Lower Triassic), East Irish Sea Basin. Geological Society of London, Special Publications 124, 253-276.
- Hillier, S., Velde, B., 1992. Chlorite interstratified with a 7 mineral: an example from offshore Norway and possible implications for the interpretation of the composition of diagenetic chlorites. Clay Minerals 27, 475-475.
- Holditch, S.A., 2006. Tight gas sands. Journal of Petroleum Technology 58, 86-93.
- Holliday, D., 1970. The petrology of secondary gypsum rocks: a review. Journal of Sedimentary Research 40, 734-744.
- Hornung, J., Aigner, T., 1999. Reservoir and aquifer characterization of fluvial architectural elements: Stubensandstein, Upper Triassic, southwest Germany. Sedimentary Geology 129, 215-280.
- Houseknecht, D.W., 1987. Assessing the Relative Importance of Compaction Processes and Cementation to Reduction of Porosity in sandstone. American Association of Petroleum Geologists Bulletin 71, 633-642.
- Hubert, J.F., Reed, A.A., 1978. Red-bed Diageneses in the East Berlin Formation, Newark Group, Connecticut Valley. Journal of Sedimentary Research 48, 175-184.
- Islam, M.A., 2009. Diagenesis and reservoir quality of Bhuban sandstones (Neogene), Titas Gas Field, Bengal Basin, Bangladesh. Journal of Asian Earth Sciences 35, 89-100.
- Ixer, R., Turner, P., Waugh, B., 1979. Authigenic iron and titanium oxides in Triassic red beds:(St. Bees Sandstone), Cumbria, northern England. Geological Journal 14, 179-192.
- Jackson, D., Johnson, H., Smith, N., 1997. Stratigraphical relationships and a revised lithostratigraphical nomenclature for the Carboniferous, Permian and Triassic rocks of

- the offshore East Irish Sea Basin. Geological Society of London, Special Publications 124, 11-32.
- Kantorowicz, J., 1990. The influence of variations in illite morphology on the permeability of Middle Jurassic Brent Group sandstones, Cormorant Field, UK North Sea. *Marine and Petroleum Geology* 7, 66-74.
- Karim, A., Pe-Piper, G., Piper, D.J., 2010. Controls on diagenesis of Lower Cretaceous reservoir sandstones in the western Sable Subbasin, offshore Nova Scotia. *Sedimentary Geology* 224, 65-83.
- Khadkikar, A., S. Merh, J. Malik and L. Chamyal, 1998. Calcretes in semi-arid alluvial systems: Formative pathways and sinks. *Sedimentary Geology* 116, 251-260.
- Knott, D.Q.S.R.I., Hill, J.R.L., 1999. Petroleum geology and future hydrocarbon potential of the Irish Sea. *Journal of Petroleum Geology* 22, 243-260.
- Lai, J. et al., 2017. Diagenesis and reservoir quality in tight gas sandstones: The fourth member of the Upper Triassic Xujiahe Formation, Central Sichuan Basin, Southwest China. *Geological Journal* 53, 629-646.
- Lander, R.H., Walderhaug, O., 1999. Predicting porosity through simulating sandstone compaction and quartz cementation. *American Association of Petroleum Geologists Bulletin* 83, 433-449.
- Lanson, B. et al., 2002. Authigenic kaolin and illitic minerals during burial diagenesis of sandstones: a review. *Clay minerals* 37, 1-22.
- Leveille, G.P., Primmer, T.J., Dudley, G., Ellis, D., Allinson, G.J., 1997. Diagenetic controls on reservoir quality in Permian Rotliegendes sandstones, Jupiter Fields area, southern North Sea. Geological Society of London, Special Publications 123, 105-122.
- Machent, P.G., Taylor, K.G., Macquaker, J.H., Marshall, J.D., 2007. Patterns of early post-depositional and burial cementation in distal shallow-marine sandstones: Upper

- Cretaceous Kenilworth Member, Book Cliffs, Utah, USA. *Sedimentary Geology* 198, 125-145.
- McAulay, G., Burley, S., Johnes, L., 1993. Silicate mineral authigenesis in the Hutton and NW Hutton fields: implications for sub-surface porosity development, Geological Society of London, Petroleum Geology Conference series, pp. 1377-1394.
- McBride, E.F., 1989. Quartz cement in sandstones: a review. *Earth-Science Reviews* 26, 69-112.
- McCrea, J.M., 1950. On the isotopic chemistry of carbonates and a paleotemperature scale. *The Journal of Chemical Physics* 18, 849-857.
- Meadows, N., Beach, A., 1993. Controls on reservoir quality in the Triassic Sherwood Sandstone of the Irish Sea, Geological Society of London, Petroleum Geology Conference series, pp. 823-833.
- Moore, C.H., 2001. Carbonate Reservoirs. Porosity Evolution and Diagenesis in a Sequence Stratigraphic Framework. In: *Developments in Sedimentology* 55, 444 pp.
- Morad, S., 1998. Carbonate cementation in sandstones; distribution patterns and geochemical evolution. In: Morad, S. (Ed.), *Carbonate Cementation in Sandstones*. International Association of Sedimentologists, 26, 1-26.
- Morad, S. et al., 1995. Diagenesis of a mixed siliciclastic/evaporitic sequence of the Middle Muschelkalk (Middle Triassic), the Catalan Coastal Range, NE Spain. *Sedimentology* 42, 749-768.
- Naylor, H., Turner, P., Vaughan, D., Fallick, A., 1989. The Cherty Rock, Elgin: A petrographic and isotopic study of a Permo-Triassic calcrete. *Geological Journal* 24, 205-221.
- Newman, P., 1999. The geology and hydrocarbon potential of the Peel and Solway basins, East Irish Sea. *Journal of Petroleum Geology* 22, 305-324.

- Pettijohn, F.J., 1957. *Sedimentary Rocks*, 2nd ed. Harper and Row, London, 718 pp.
- Platt, J., 1993. Controls on clay mineral distribution and chemistry in the early Permian Rotliegend of Germany. *Clay Minerals* 28, 393-393.
- Pollastro, R.M., 1985. Mineralogical and morphological evidence for the formation of illite at the expense of illite/smectite. *Clays and Clay Minerals* 33, 265-274.
- Primmer, T.J. et al., 1997. Global Patterns in Sandstone Diagenesis: Their Application to Reservoir Quality Prediction for Petroleum Exploration. *American Association of Petroleum Geologists Memoir* 69, 61-77.
- Purvis, K., 1992. Lower Permian Rotliegend sandstones, southern North Sea: a case study of sandstone diagenesis in evaporite-associated sequences. *Sedimentary Geology* 77, 155-171.
- Pye, K., Krinsley, D., 1986. Diagenetic carbonate and evaporite minerals in Rotliegend aeolian sandstones of the southern North Sea: their nature and relationship to secondary porosity development. *Clay Minerals* 21, 44 -457.
- Ramseyer, K., Boles, J.R., Lichtner, P.C., 1992. Mechanism of plagioclase albitization. *Journal of sedimentary research* 62, 349-356.
- Robb, G.L., 1949. Red bed coloration. *Journal of Sedimentary Research* 19, 99-103.
- Ronald, J., 1967. Quantitative X-ray diffraction analysis using clay mineral standards extracted from the samples to be analysed. *Clay Minerals* 7, 79-91.
- Rosenbaum, J., Sheppard, S., 1986. An isotopic study of siderites, dolomites and ankerites at high temperatures. *Geochimica et Cosmochimica Acta* 50, 1147-1150.
- Rossi, C., Marfil, R., Ramseyer, K., Permanyer, A., 2001. Facies-related diagenesis and multiphase siderite cementation and dissolution in the reservoir sandstones of the Khataiba Formation, Egypt's Western Desert. *Journal of Sedimentary Research* 71, 459-472.

- Ruf, M., Aigner, T., 2004. Facies and poroperm characteristics of a carbonate shoal (Muschelkalk, South German Basin): a reservoir analogue investigation. *Journal of Petroleum Geology* 27, 215-239.
- Schmid, S., Worden, R.H., Fisher, Q.J., 2004. Diagenesis and reservoir quality of the Sherwood Sandstone (Triassic), Corrib Field, Slyne Basin, west of Ireland. *Marine and Petroleum Geology* 21, 299-315.
- Schmid, S., R.H. Worden and Q.J. Fisher, 2006. Carbon isotope stratigraphy using carbonate cements in the Triassic Sherwood Sandstone Group: Corrib Field, west of Ireland. *Chemical Geology* 225, 137-155.
- Shelton, J.W., 1964. Authigenic kaolinite in sandstone. *Journal of Sedimentary Research* 34, 102-111.
- Smith, T.M., Sayers, C.M., Sondergeld, C.H., 2009. Rock properties in low-porosity/low-permeability sandstones. *The Leading Edge* 28, 48-59.
- Smythe, D., 2014. Planning application no. LCC/2014/0096 by Cuadrilla Bowland Limited to drill at Preston New Road, Lancashire: Objection on grounds of geology and hydrogeology. Non-technical report, pp 1-42.
- Storvoll, V., Bjørlykke, K., Karlsen, D., Saigal, G., 2002. Porosity preservation in reservoir sandstones due to grain-coating illite: a study of the Jurassic Garn Formation from the Kristin and Lavrans fields, offshore Mid-Norway. *Marine and Petroleum Geology* 19, 767-781.
- Stroker, T.M., Harris, N.B., Elliott, W.C., Wampler, J.M., 2013. Diagenesis of a tight gas sand reservoir: Upper Cretaceous Mesaverde Group, Piceance Basin, Colorado. *Marine and Petroleum Geology* 40, 48-68.

- Strong, G., 1993. Diagenesis of Triassic Sherwood Sandstone Group rocks, Preston, Lancashire, UK: a possible evaporitic cement precursor to secondary porosity? Geological Society of London, Special Publications 73, 279-289.
- Strong, G., Milodowski, A., 1987. Aspects of the diagenesis of the Sherwood Sandstones of the Wessex Basin and their influence on reservoir characteristics. Geological Society of London, Special Publications 36, 325-337.
- Strong, G. et al., 1994. The petrology and diagenesis of Permo-Triassic rocks of the Sellafield area, Cumbria. Proceedings of the Yorkshire Geological and Polytechnic Society. Geological Society of London, pp. 77-89.
- Stuart, I., Cowan, G., 1991. The South Morecambe Field, Blocks 110/2a, 110/3a, 110/8a, UK East Irish Sea. Geological Society of London Memoirs 14, 527-541.
- Swarbrick, R.E., 1999. Diagenesis in North Sea HPHT clastic reservoirs - Consequences for porosity and overpressure prediction. Marine and Petroleum Geology 16, 337-353.
- Swart, P.K., S. Burns and J. Leder, 1991. Fractionation of the stable isotopes of oxygen and carbon in carbon dioxide during the reaction of calcite with phosphoric acid as a function of temperature and technique. Chemical Geology: Isotope Geoscience section 86, 89-96.
- Taylor, K.G., Gawthorpe, R.L., Curtis, C.D., Marshall, J.D., Awwiller, D.N., 2000. Carbonate cementation in a sequence-stratigraphic framework: Upper Cretaceous sandstones, Book Cliffs, Utah-Colorado. Journal of Sedimentary Research 70, 360-372.
- Taylor, T.R. et al., 2010. Sandstone diagenesis and reservoir quality prediction: Models, myths, and reality. American Association of Petroleum Geologists Bulletin 94, 1093-1132.

- Thomson, A., Stancliffe, R.J., 1990. Diagenetic controls on reservoir quality, eolian Norphlet Formation, south State Line field, Mississippi. In: Barwis, J.H., McPherson, J.G., Studlick, R.J. (Eds.), Sandstone Petroleum Reservoirs. Springer-Verlag, New York, pp. 205-224.
- Tobin, R.C., T. McClain, R.B. Lieber, A. Ozkan, L.A. Banfield, A.M. Marchand and L.E. McRae, 2010. Reservoir quality modeling of tight-gas sands in Wamsutter field: Integration of diagenesis, petroleum systems, and production data. American Association of Petroleum Geologists Bulletin, 94, 1229-1266.
- Walderhaug, O., 1994. Temperatures of quartz cementation in Jurassic sandstones from the Norwegian continental shelf: Evidence from fluid inclusions. Journal of Sedimentary Research 64, 311-323.
- Walderhaug, O., 1996. Kinetic modeling of quartz cementation and porosity loss in deeply buried sandstone reservoirs. American Association of Petroleum Geologists Bulletin 80, 731-745.
- Walderhaug, O., Eliassen, A., Aase, N.E., 2012. Prediction of permeability in quartz-rich sandstones: examples from the Norwegian continental shelf and the Fontainebleau sandstone. Journal of Sedimentary Research 82, 899-912.
- Walker, T.R., Larson, E.E., Hoblitt, R.P., 1981. Nature and origin of hematite in the Moenkopi Formation (Triassic), Colorado Plateau: a contribution to the origin of magnetism in red beds. Journal of Geophysical Research: Solid Earth 86, 317-333.
- Warwick, E., 2001. Warwick Energy Resources. Available from <https://archive.is/wE6S2>.
- Worden, R., Burley, S., 2003. Sandstone Diagenesis: The Evolution of Sand to Stone, Sandstone Diagenesis. Blackwell Publishing Ltd, pp. 1-44.
- Worden, R., Morad, S., 2003. Clay minerals in sandstones: controls on formation, distribution and evolution. In: Worden, R.H., Morad, S. (Eds.), Clay Mineral Cements

- in Sandstones. International Association of Sedimentologists, Special Publication 34, pp. 3-41.
- Xi, K. et al., 2015. Diagenesis and reservoir quality of the Lower Cretaceous Quantou Formation tight sandstones in the southern Songliao Basin, China. *Sedimentary Geology* 330, 90-107.
- Yaliz, A., Taylor, P., 2003. The Hamilton and Hamilton North Gas Fields, Block 110/13a, East Irish Sea. *Geological Society of London Memoirs* 20, 77-86.
- Zhang, L. et al., 2009. Diagenetic history of tight sandstones and gas entrapment in the Yulin Gas Field in the central area of the Ordos Basin, China. *Marine and Petroleum Geology* 26, 974-989.
- Ziegler, K., 2006. Clay minerals of the Permian Rotliegend Group in the North Sea and adjacent areas. *Clay minerals* 41, 355-393.

FIGURE CAPTIONS

Fig. 1: Map of East Irish Sea Basin showing study wells and surrounding wells within EISB, the red asterisk indicates studied wells modified after (Herries and Cowan, 1997).

Fig. 2: Stratigraphy Sequence of EISB modified after (Jackson et al., 1997; Knott and Hill, 1999).

Fig. 3: Lithology characteristics of representative samples of Collyhurst sandstone. (A) Photomicrograph of medium-grained, well-sorted, sub-angular to rounded sandstone, showing mostly quartz grains (PPL); (B) Photomicrograph of coarse-grained, poorly-sorted sandstone showing quartz, rock fragments and feldspar grains (PPL); (C) Photomicrograph of medium to coarse-grained sandstone indicating long grain contact (arrow I) and concave-convex contacts (arrow II); (D) Photomicrograph of medium to coarse-grained sandstone showing stylolite contact (arrow I) and suture contact (arrow II) of grains (PPL); [PPL = Plane polarized, Q = quartz, F = Feldspar, RF = Rock fragments].

Fig. 4: Classification of Collyhurst Sandstones Formation after (Pettijohn, 1957)

Fig. 5: Representative sample from each lithofacies. (A) Photomicrograph of fine-grained sandstone (PPL); (B) Photomicrograph of conglomeratic sandstone (poorly-sorted) showing mostly ductile grains; (C) BSE image of strongly carbonate-cemented sandstone; (D) Photomicrograph of carbonate-cemented sandstone (PPL); (E) Photomicrograph of quartz-cemented sandstone (PPL); (F) Photomicrograph of matrix-free sandstone (moderately-

sorted). [DQ = Detrital quartz, AQ = Authigenic quartz RF = Rock fragments, Fsp = Feldspar, MI= Mud intraclasts]

Fig. 6: Vertical distribution of major cements within the studied sandstone [upper charts are EL-1 samples and lower charts are GH-1 samples]. (A) Cross plot of total carbonate cement with depth; (B) Cross plot of clay mineral cement with depth; (C) Cross plot of quartz cement with depth.

Fig. 7: Carbonate cements. (A) BSE image of non-ferroan dolomite cement occluding intergranular pore. Outer zone of ferroan dolomite also visible; (B) CL image of non-ferroan dolomite cement filling intergranular pores; (C) BSE image of calcite (I) cement filling large intergranular pores, calcite cement show texture similar to framework grains crystals; (D) BSE image of calcite (II) cement postdating quartz overgrowth (circle shows where calcite engulfed quartz overgrowth); (E) BSE image of siderite (I) cement coating detrital grains; (F) BSE image of siderite II cement filling intergranular pore volume; (G) BSE image of siderite type (III) cement overgrown filling pores illite cement and dolomite rhombs. [Qtz ovgt = quartz overgrowth, Fe-D= Ferroan dolomite]

Fig. 8: XRD pattern of clay fraction from representative samples of Collyhurst Sandstones Formation. (A) XRD diffractogram showing illite, chlorite and kaolinite; (B) XRD diffractogram showing the presence of smectite indicated by a slight shift in the profile pattern (arrow). [Chl = chlorite; ill = illite; K = kaolinite; I-S = illite-smectite; Fsp= Feldspar; Qtz = Quartz].

Fig. 9: Clay morphologies. (A) BSE image of flaky illite cement filling intergranular pore; (B) BSE image of grain coating illite (arrow) growing to pore-bridging illite; (C) PPL Photomicrograph of coarse-grained sandstone revealing grain-coating illite (clay coating is often absent at the point of grain contact); (D) BSE image of grain with concentrically laminated illite-smectite coat; (E) Enlarge image of rectangular box in Fig. 9D, showing the chemistry of mixed smectite layer on top right, (note the presence of chlorite (arrow) post-dating illite-smectite); (F) BSE image of microcrystalline chlorite with rosette morphology (note the growth of illite within the chlorite crystals); (G) BSE image showing pore-lining chlorite developing into pore-filling chlorite cement (note the fan-like habit crystals of chlorite growing within intergranular pore); (H) BSE image of microcrystalline chlorite cement (stub) coating grains and filling pores, note the presence of a small quartz overgrowth (~20 μm) indicating that chlorite inhibits quartz cement; (I) BSE image of vermicular kaolinite filling intergranular porosity and inhibiting crystals of chlorite cement [Qtz ovgt = Quartz overgrowth].

Fig. 10: Photomicrograph showing quartz cement, arrow (I) indicate euhedral syntaxial, pore-lining quartz overgrowth, (II) show quartz overgrowth completely filling pore and (III) reveals break or discontinuity of clay rim resulting in quartz overgrowth, (PPL); (B) BSE image of pore-filling barite cement, engulfed quartz overgrowth and replaced by albite cement; (C) BSE image of pore-filling anhydrite, replaced by albite and engulfed quartz overgrowth; (D) BSE image of albite overgrowth on detrital plagioclase feldspar; (E) Photomicrograph image of haematite coat on detrital grains (PPL); (F) BSE image showing haematite crystals resulting from the breakdown of ferromagnesian mineral. [ill + chl = illite + chlorite, Fsp = Feldspar, Ab = Albite, K = Kaolinite, Qtz ovgt = Quartz overgrowth].

Fig. 11: (A) BSE image showing partial dissolution of K-feldspars; (B) BSE image showing intense dissolution of feldspars grain leading to illite precipitation (illitization of feldspar), note the partial replacement of ferroan dolomite on the feldspar grain and albite growth; (C) BSE image showing partial dissolution of mica sheets; (D) XPL Photomicrograph image showing alteration of biotite grain; (E) BSE image of ferroan dolomite/ankerite showing partial dissolution; (F) BSE image of distorted mica grains owing to mechanical compaction [ill + chl = illite+chlorite, Fsp = Feldspar, Ab = Albite, K = Kaolinite, Qtz ovgt = Quartz overgrowth].

Fig. 12: Plot of intergranular volume (IGV) against cement volume of Collyhurst sandstone reservoir after Houseknecht (1987); (A) EL-1 well and (B) GH-1 well.

Fig. 13: Porosity and Permeability data of the studied reservoir. (A) Distribution of helium porosity and point count porosity with depth in EL-1 well; (B) Distribution of core analysis permeability with depth in EL-1 well.

Fig. 14: Pore types observed in the studied sandstone. (A) Photomicrograph of coarse-grained sandstone showing intergranular porosity (arrow) in between grains (PPL); (B) Photomicrograph showing intragranular porosity (arrow) due to feldspar and rock fragment dissolution (PPL); (C) Photomicrograph showing micro-fracture porosity (arrow) within silty mudstone (PPL)

Fig. 15: The distribution of carbonate and oxygen isotope of representative samples of Collyhurst sandstone (EL-1 and GH-1 wells).

Fig. 16: Diagenetic sequence of Collyhurst Sandstone Formation

Fig. 17: Temperature versus $\delta^{18}\text{O}$ for dolomite cement after (Ehrenberg et al., 2002). Dotted curve represents SMOW value for Triassic-Permian meteoric pore water (-3‰) drawn from fractionation equation of (Matthews and Katz, 1977). Early dolomite reveals lower temperature of formation than late dolomite.

Fig. 18: Plot of reservoir properties of each lithofacies within the studied sandstone (EL-1 well)

Fig. 19: Porosity versus depth prediction model for each lithofacies within Collyhurst Sandstone Formation

TABLE CAPTIONS

Table 1: Isotopic composition of carbonate cement of Collyhurst sandstones

Table 2: Diagenetic scheme of various lithofacies in Collyhurst sandstone (EL-1 well)

Table 1

Sample No	Carbonate Type	$10^3 \delta^{13}\text{C}$ (VPDB)	$10^3 \delta^{18}\text{O}$ (VPDB)
56589	Dolomite	-2.37	-4.51
56591	Dolomite	-1.73	-2.35
56595	Dolomite	-0.48	-2.01
56596	Dolomite	-1.2	-2.42
	Average	-1.45	-2.82
56615	Dolomite (Fe)	-2.18	-7.9
56618	Dolomite (Fe)	-6.76	-4.07
56620	Dolomite (Fe)	-8.18	-5.38
56624	Dolomite (Fe)	-9.22	-4.39
	Average	-6.59	-5.44

Table 2

Lithofacies Type	Early Diagenesis	Late Diagenesis
Lithofacies A	Infiltration of clays; mechanical compaction	Mechanical and compaction; clay cementation (illite and chlorite); intergranular porosity decreases
Lithofacies B	Mechanical compaction; ductile grains squeeze into more stable grains	Mechanical compaction increases, rare intergranular porosity
Lithofacies C	Mechanical and compaction; early carbonate cementation	Compaction reduces; precipitation of late carbonate cementation (late calcite, ferroan dolomite and late siderite); intergranular porosity decreases
Lithofacies D	Mechanical compaction; grain rearrangement; decrease in intergranular porosity	Mechanical and chemical compaction; precipitation of quartz overgrowth
Lithofacies E	Mechanical compaction; dissolution grains	Compaction reduces; generation of secondary porosity through grain dissolution

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

B

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12

Figure 13

Figure 14

Figure 15

Figure 16

Figure 17

Figure 18

Figure 19