Vegetation change in the eastern Pamir Mountains inferred from Lake Karakul pollen spectra of the last 28 kyr

DOI:

[10.1016/j.palaeo.2018.08.010](https://doi.org/10.1016/j.palaeo.2018.08.010)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Heinecke, L., Fletcher, W., Mischke, S., Tian, F., & Herzsuh, U. (2018). Vegetation change in the eastern Pamir Mountains inferred from Lake Karakul pollen spectra of the last 28 kyr. *Palaeogeography, Palaeoclimatology, Palaeoecology*. <https://doi.org/10.1016/j.palaeo.2018.08.010>

Published in:

Palaeogeography, Palaeoclimatology, Palaeoecology

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Accepted Manuscript

Vegetation change in the eastern Pamir Mountains, Tajikistan,
inferred from Lake Karakul pollen spectra of the last 28 kyr

Liv Heinecke, William J. Fletcher, Steffen Mischke, Fang Tian,
Ulrike Herzschuh

PII: S0031-0182(17)31014-3
DOI: doi:[10.1016/j.palaeo.2018.08.010](https://doi.org/10.1016/j.palaeo.2018.08.010)
Reference: PALAEO 8896

To appear in: *Palaeogeography, Palaeoclimatology, Palaeoecology*

Received date: 3 October 2017
Revised date: 21 August 2018
Accepted date: 22 August 2018

Please cite this article as: Liv Heinecke, William J. Fletcher, Steffen Mischke, Fang Tian, Ulrike Herzschuh, Vegetation change in the eastern Pamir Mountains, Tajikistan, inferred from Lake Karakul pollen spectra of the last 28 kyr. *Palaeo* (2018), doi:[10.1016/j.palaeo.2018.08.010](https://doi.org/10.1016/j.palaeo.2018.08.010)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Vegetation change in the eastern Pamir Mountains, Tajikistan, inferred from Lake Karakul pollen spectra of the last 28 kyr

Liv Heinecke^{1, 2, 3 *} (Liv.Heinecke@uni-potsdam.de), William J. Fletcher³ (will.fletcher@manchester.ac.uk), Steffen Mischke⁴ (smi@hi.is), Fang Tian¹ (Fang.Tian@awi.de), Ulrike Herzschuh^{1, 2, 5 *} (Ulrike.Herzschuh@awi.de)

¹ Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Research Unit Potsdam, Telegrafenberg A43, 14473 Potsdam, Germany

² Institute of Earth and Environmental Science, University of Potsdam, Karl-Liebknecht-Str. 24-25, 14476 Potsdam-Golm, Germany

³ Department of Geography, School of Environment, Education and Development, The University of Manchester, Oxford Road, M13 9PL Manchester, United Kingdom

⁴ Faculty of Earth Sciences, University of Iceland, Sturlugata 7, Askja, 101 Reykjavík, Iceland

⁵ Institute of Biochemistry and Biology, University of Potsdam, Karl-Liebknecht-Str. 24-25, 14476 Potsdam-Golm, Germany

Corresponding authors*: Liv Heinecke, Ulrike Herzschuh

Abstract

We present a pollen record for last 28 cal kyr BP from the eastern basin of Lake Karakul, the largest lake in Tajikistan, located in the eastern Pamir Mountains at 3915 m asl, a geographically complex region. The pollen record is dominated by *Artemisia* and Chenopodiaceae, while other taxa, apart from Poaceae, are present in low quantities and rarely exceed 5% in total. Arboreal pollen occur predominantly from ~28 to ~13 cal kyr BP, but as likely no trees occurred in the high mountain regions of the eastern Pamir during this time due to the high altitude and cold climate, arboreal taxa are attributed to long distance transport, probably by the Westerlies, the dominant atmospheric circulation. Tree pollen influx decreases strongly after ~13 cal kyr BP, allowing the pollen spectra to be interpreted as a regional vegetation signal. We infer that from 27.6 to 19.4 cal kyr BP the eastern Pamir was dominated by dry mountain steppe with low vegetation cover, while from 19.0 to 13.6 cal kyr BP *Artemisia* values increase and

Chenopodiaceae, most herb taxa, and inferred far distant input from arboreal taxa decrease. Between 12.9 and 6.7 cal kyr BP open steppe vegetation dominated with maximum values in *Ephedra*, and while steppe taxa still dominated the spectra from 5.4 to 1 cal kyr BP, meadow taxa start to increase. During the last millennium, alpine steppe and alpine meadows expanded and a weak human influence can be ascertained from the increase of Asteraceae and the occurrence of *Plantago* pollen in the spectra.

Keywords: arid Central Asia; high Asia; palynology; vegetation reconstruction; lake sediments

1. Introduction

The magnitude of temperature and precipitation changes in the course of global climate change is projected to vary regionally (Pachauri et al., 2014). Mountain areas, including the Central Asian mountain regions, are among those which are expected to be relatively strongly affected (Lioubimtseva and Henebry, 2009; Pachauri et al., 2014). Climate change in this region is reflected by the increased melt of glaciers (Immerzeel et al., 2010; Makhmadaliev et al., 2008), increased temperature, changes in evaporation, and shifts in precipitation patterns (Barnett et al., 2005; Lutz et al., 2014). Whether these phenomena entail vegetation changes or whether vegetation is more responsive to human impact is rather uncertain as long-term observational time-series are lacking.

Characterizing vegetation responses to past climate change may help us understand better the ongoing and future vegetation change. Lake sediments are best suited to record environmental change on millennial time-scales as they preserve pollen assemblages, which are known to be a reliable proxy of regional terrestrial vegetation (Birks and Birks, 1980; Cohen, 2003).

Palaeovegetation studies from the Central Asian high mountains, i.e. the Tien Shan, Karakorum, and Pamir Mountains, are few (Huang et al., 2014; Lauterbach et al., 2014; Mathis et al., 2014; Tang et al., 2011; Wang et al., 2013). Mathis et al. (2014), for example, reported a dominance of steppe vegetation with more humid phases and regional tree development from 8.3 to 4.5 cal kyr BP (calibrated thousands of years before 1950 AD), and decreasing arboreal pollen due to drier conditions from 4.5 to 2.0 cal kyr BP. More records are available from the western Tibetan

Plateau, e.g. Bangong Co (Gasse et al., 1996; Van Campo et al., 1996); Sumxi Co (Gasse et al., 1991)), the Central Asian lowlands (e.g. Aral Sea (Sorrel et al., 2007); Lake Balkhash (Feng et al., 2013), Lake Manas (Rhodes et al., 1996); Lake Boston (Huang et al., 2009), Wulungu Lake (Liu et al., 2008), Kashgar oasis (Zhao et al., 2012), or the lowlands to the west of the Central Asian high mountains, e.g. Lake Van (Litt et al., 2009, 2014; Wick et al., 2003), although the vegetation histories there seem to differ from the Central Asian high mountains. For example, on the western Tibetan Plateau in the Bangong Co basin, Van Campo et al. (1996) inferred predominantly desert vegetation from 9.9 to 9.5 kyr BP, more steppe-like vegetation from 9.5 to 6.2 kyr BP interrupted by dry spells, for example around 7.7 kyr BP, and a return to more arid conditions from 6.2 kyr BP onward. At Lake Van in Turkey, Litt et al. (2009) reconstructed semi-desert steppe vegetation from 20 to 14.5 kyr BP and semi-desert with a slight increase in tree pollen from 14.5 until 12.7 kyr BP. During the Younger Dryas, *Artemisia* and *Chenopodiaceae* increased until the beginning of the Holocene, when, due to increased moisture availability, desert taxa rapidly decreased and grass-steppe and arboreal taxa increased until approximately 6 kyr BP. After ~3.8 kyr BP human influence increased, as reflected by increasing *Plantago* and *Cerealia* and decreasing *Quercus* pollen percentages.

Studies in the central Asia reflect a very diverse vegetation history, although the spatial density remains extremely low in the Central Asian high mountains, hindering the spatiotemporal reconstruction of past vegetation and thus changes in past climate. The aim of this study is to establish a vegetation record that reaches back into the late Pleistocene to understand better the vegetation history of the eastern Pamir by investigating the pollen record of a sediment core from Lake Karakul (Tajikistan), covering the last 28 cal kyr BP. Thereby we first assess the indicator value of the pollen record; secondly, reconstruct the regional terrestrial vegetation change over the late Pleistocene and Holocene; and thirdly, compare the inferences with other vegetation and climate records, aiming to refine the palaeoenvironmental and climatic history of the eastern Pamir.

2. Study site

Lake Karakul is located in the eastern Pamir Mountains at 3915 m above sea level (asl) in Tajikistan (Fig. 1). The lake occurs in a tectonic graben structure. A horst structure subdivides the lake into a shallow eastern and a deep western basin (Strecker et al., 1995). Triassic granites, Permian mafic metavolcanics, and Carboniferous and Permian phyllites and metasandstones form

the bedrock of the study region and are partially covered by late Pleistocene glacial till, terrace gravel, Holocene lacustrine sediments and gravel deposits (Nöth et al., 1932; Strecker et al., 1995). The eastern Pamir region is a high mountain plateau characterised by discontinuous permafrost formed in the late Pleistocene (Gorbunov, 1990). High mountains, mainly between 5000 and 6000 m elevation, surround the endorheic basin of Lake Karakul (Komatsu and Tsukamoto, 2015). The surface area of the modern lake is ca. 388 km² (Komatsu and Tsukamoto, 2015). Seasonal rivers, mainly fed by snow and glacier melt, drain the 4464 km² catchment. The Westerlies are the main atmospheric circulation influencing the region; however, they are blocked by the Western Pamir, prohibiting effective moisture transport from the west. Accordingly, precipitation is low in the region with an annual mean of 82 mm (Karakul climate station). High insolation (Mętrak et al., 2015) and low precipitation lead to an arid climate with an aridity index of -45 to -61 (Agakhanyantz and Lopatin, 1978; aridity index after Thornthwaite, 1948). The mean annual temperature is -3.9°C, mean January temperature -18.1°C and mean July temperature 8.5°C as recorded at the Karakul station (data from 1934-1995 and 2004-2007).

Due to the steep relief and climatic conditions, the flora of the high Pamir can be categorized into a subalpine zone (from 3500 to 4200 m asl), an alpine zone (4100 to 4800 m asl), and a nival zone (4700 m asl and above; Mętrak et al., 2015). Mountain desert (3500 to 4600 m asl, ~33% of eastern Pamir) is dominated by Chenopodiaceae (e.g. *Ceratoides*) and Asteraceae (*Artemisia*, *Ajana* etc.); mountain steppe (3500 to 4000 m asl; 10% area) is dominated by *Stipa*; mountain meadows (3500 to 4400 m asl, ~5% area) including floodplains, riverbanks, and *tugai* are dominated by Poaceae and Cyperaceae (*Leymus*, *Kobresia*). *Tugai*, a Central Asian type of forest/shrub habitat along rivers or spring-outlets, is found up to 3900 m asl and includes *Betula*, *Populus*, and *Salix* species amongst others (Breckle and Wucherer, 2006; Mętrak et al., 2015). In general, due to the cold, arid conditions, terrestrial vegetation growth is sparse and vegetation cover reaches 5–20% (Breckle and Wucherer, 2006; Mętrak et al., 2015) and is mainly confined to wetlands, springs, and moister valleys. Although the α -diversity is rather low, the β -diversity is high due to the variety of habitats in the eastern Pamir region (Breckle and Wucherer, 2006). *Ceratoides*, *Artemisia*, *Stipa*, *Festuca*, *Leymus*, *Kobresia*, and cushion plant associations, among others, have been recorded in the direct vicinity of the lake. Many of these taxa are grazed by domestic herds (Breckle and Wucherer, 2006). *Teresken* (*Ceratoides papposa* Botsch. & Ikonn.; Chenopodiaceae) is collected widely as fuel for heating and cooking by the locals, and supplementary fodder for livestock during winter (Mętrak et al., 2015).

3. Material and Methods

3.1. *Sediment cores and chronology*

In April 2012, two sediment cores (KK12-1 and KK12-2) were retrieved from Lake Karakul, of 1086 cm and 1226 cm length, respectively. The cores were recovered using UWITEC coring equipment at 39.0176° N / 73.5327°E, 10 m apart from each other, coring from the ice in 2-m segments. A composite core was created based on the correlation of XRF-measurements (Fig. 2; Mischke et al., 2017). A lake reservoir effect of 1.368 kyr was determined by dating living charophytes from 2008 and 2012 (Heinecke et al., 2017b). Twenty-two radiocarbon ages (from total organic carbon, or water-plant remains; 13 from core KK12-1, 9 from core KK12-2) and 10 optically stimulated luminescence (OSL) dates (all from KK12-2) were used to create a composite age-depth model in OxCal 4.2 (Fig. 2; Mischke et al., 2017). The sediment characteristics of KK12-1, including grain size, XRF, geochemistry and isotope measurements are reported in Heinecke et al. (2017b). The sediments from KK12-1 were aligned with the depth scale of core KK12-2 during the establishment of the composite core. A detailed description of the composite core is given in Mischke et al. (2017). Furthermore, a 1.04 m short sediment core covering the last 4 cal kyr BP, which was retrieved in 2008 and presented in Mischke et al. (2010), was also analysed for pollen.

3.2. *Pollen sample preparation and pollen analyses*

Pollen samples were treated at the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research in Potsdam in a designated pollen preparation laboratory. Two *Lycopodium* spore tablets were added to each sediment sample (1.5 ml). Sample preparation included treatments with HCl (10%) and KOH (10%), sieving (400 µm) and boiling with HF (40-45 %), followed by acetolysis and sieving through a 7 µm mesh in an ultrasonic bath. Pollen samples were stored in water-free glycerol for permanent storage.

Fifty pollen samples were prepared and analysed, of which 46 samples were obtained from KK12-1. Two sediment gaps in KK12-1, which occurred due to coring with a piston corer without sediment overlap, were filled with samples from KK12-2 to create the composite pollen record. At least 300 pollen grains were counted, except in samples dated before 17.1 cal kyr BP, where an average of 170 pollen grains per sample were counted, due to very low pollen concentration. Pollen and non-pollen palynomorphs were investigated with an Axio Scope A1 with a 40x and 100x oil immersion objective. Identification is based on literature by Beug (2004) and Moore et al. (1991) and on reference slide collections stored at the University of Manchester and Alfred Wegener Institute. In the short core TAJ-Kar-08-1B, 37 pollen samples were analysed and, on average, 144 pollen grains were counted per sample (min. 50, max. 202).

3.3. Pollen data treatment

Pollen percentages are based on the terrestrial pollen sum (including Cyperaceae, since this taxon may be an important component of regional vegetation as well as a local indicator). The pollen diagram (Fig. 3) was created in Tilia 1.7.16 (Grimm, 2011) and pollen assemblage zones for the long composite core were generated via stratigraphically constrained incremental sum-of-squares cluster analysis (CONISS; Grimm, 1987). The number of pollen assemblage zones was assessed using bootstrapping in the rioja package (Juggins, 2017) in R (R Core Team, 2012). Pollen data were square-root transformed prior to further statistical analyses and included only taxa which reached at least 0.5% in a minimum of three samples. A principal component analysis (PCA) was performed in CANOCO 5 (ter Braak and Šmilauer, 2012). Only data younger than 13.6 cal kyr BP were included in the PCA to reduce the influence of the samples subject to long distance pollen transport. The non-pollen palynomorphs *Botryococcus*, *Pediastrum* and NPP1, an unknown palynomorph, which was identified as gemmae of a bryophyte (e.g. *Bryum*), as well as chironomid remains were counted in the composite core and are displayed as percentages of the terrestrial pollen sum (Fig. 3).

4. Results

4.1. Composite core (KK12-1/2; 27.6 cal kyr BP to present)

In total, 57 pollen taxa were identified. The pollen record (Fig. 3) is dominated by *Artemisia*, which reaches between 45 and 77% (mean 61%). Apart from Cyperaceae (range: 4–25%, mean 16%) and Poaceae (range: 1–12%, mean 7%), other taxa are mostly recorded with very low percentages and only rarely exceed 5%.

The pollen record (Fig. 3) was subdivided into three pollen assemblage zones (PAZ). PAZ1, (1113–716 cm core depth; 27.6–19.4 cal kyr BP) is characterized by relatively high *Pinus* and *Alnus* values. Furthermore it shows variable *Artemisia* percentages (mean in PAZ1: 57%), medium to high values in Chenopodiaceae (mean PAZ1: 17%) and maximum values in the herb taxa Caryophyllaceae, Lamiaceae, Papaveraceae, and *Thalictrum*. The pollen spectra of PAZ 2 (713–595 cm; 19.0–13.6 cal kyr BP) are characterized by maximum values in *Artemisia* (mean PAZ2: 73%), low values in almost all herb taxa, and a decrease in arboreal taxa. In PAZ 3 (580 cm to core top; 12.9 cal kyr BP to modern day), *Artemisia* decreases (mean PAZ3: 61%), while Chenopodiaceae (mean PAZ3: 17%), Cyperaceae (mean PAZ: 3%), and steppe and meadow taxa (i.e. Asteraceae spp., Brassicaceae, Rosaceae) increase. Arboreal pollen remains sparse in PAZ 3. PAZ 3 was subdivided into three subzones - PAZ 3a–c. Increased Chenopodiaceae, high amounts of Cyperaceae, and maximum values of *Ephedra distachya*-type, *E. fragilis*-type, and *Betula* characterize PAZ 3a (580–486 cm; 12.9–6.7 cal kyr BP). In PAZ 3b (457–90 cm; 5.4–1.0 cal kyr BP), Chenopodiaceae, *Ephedra*, and *Betula* decrease slightly, while Brassicaceae and Asteraceae increase. PAZ 3c (62 cm to top; 725 cal yr BP to modern day) shows high values in Brassicaceae and Asteraceae.

In a PCA including all PAZs (not shown), extra-regional taxa *Pinus* and *Alnus* have high abundances in PAZ 1 and 2 and thus strongly contribute to the variance in the pollen spectra. Accordingly, in the PCA presented here only the pollen spectra of PAZ 3 are included. The first two axes of the PCA capture 39.6% of the total variance of the pollen record starting at 13.6 cal kyr BP (Fig. 4). PCA axis 1 accounts for 21.8% and separates the dry steppe desert taxa at the negative end from the cold meadow taxa at the positive end. PCA axis 2 accounts for 17.8% and separates the *Artemisia* from the Cyperaceae signal. The majority of samples from PAZ 3a older than 10 cal kyr BP plot in the upper left corner of the PCA (Fig. 4), samples from PAZ 3a and between 10 and roughly 3 cal kyr BP plot in the lower right, while samples from 3b younger than 3 cal kyr BP plot in the lower left. All samples younger than 1.0 cal kyr BP plot in the upper left of the PCA.

4.2. Short core TAJ-Kar-08-1B

In the short core, 62 pollen taxa were identified. The pollen spectra (Fig. 5) are dominated by *Artemisia* (range: 61–84%, mean 72%), while *Chenopodiaceae* (range: 7–19%, mean 12%), *Poaceae* (range: 0–8%, mean 4%), and *Cyperaceae* (range: 0–5%, mean 1%) are common. Other taxa are recorded with low percentages (maximum values are generally <3%). The pollen spectra generally display no notable temporal trend during the last 4.0 cal kyr.

5. Discussion

5.1. Interpretation of pollen data

Lake Karakul is the largest lake in the Pamir Mountains, with a surface area of approximately 388 km². We therefore assume that most of the pollen deposited in the sediments does not reflect the local scale but rather a regional to extra-regional vegetation signal (Prentice, 1985; Sugita, 1994). For this reason, we refrain from interpreting the pollen spectra of the short core TAJ-Kar-08-1B in which only little variation is reflected. This might be because the core was taken from the centre of the eastern basin of Lake Karakul where the influence of long distance pollen is stronger. A strong extra-regional pollen load is also indicated by the high percentages of arboreal taxa such as *Pinaceae* and *Alnus*, which are particularly high in the central lake core, as arboreal taxa do not occur in the surroundings of the lake and are hardly found in the eastern Pamir. In the composite core, arboreal pollen reaches high percentages, especially during the late Pleistocene. A long distance transported pollen component generally has a stronger representation when local and regional vegetation cover and pollen production are low, for example during cold and dry periods (Van Campo and Gasse, 1993). This effect has also been reported from other sites world-wide that are characterized by open vegetation with low regional pollen productivity, such as Svalbard (Hyvärinen, 1970) and Rogers Lake in Connecticut (USA; Davis, 1967).

Apart from *Pinaceae* and *Alnus*, other woody taxa such as *Betula*, *Salix*, *Populus*, and *Hippophae* are common in riparian environments in the Pamir (Breckle and Wucherer, 2006;

Mętrak et al., 2015). It is furthermore known that pollen grains, including arboreal, shrub and herb pollen, may be transported upwards from lower elevations into more mountainous areas (Markgraf, 1980). These taxa can thus appear at the same time as taxa that indicate drier conditions as they occur in distinct habitats (although spatially very limited to riparian environments) or indicate far distant input.

All samples are dominated by *Artemisia* pollen, a typical steppe indicator and high pollen producer, which is the main component of pollen records from semi-arid to arid Central Asia (Gasse et al., 1996; Herzschuh et al., 2004; Van Campo et al., 1996). Chenopodiaceae, the second-most common pollen taxon represented throughout the core, is likewise a common vegetation component of Central Asia, particular in montane deserts. An increase in Chenopodiaceae relative to *Artemisia* is generally assumed to reflect an increase of desert relative to steppe elements in the vegetation (El-Moslimany, 1990; Gasse et al., 1996; Liu et al., 2008). Accordingly, *Artemisia*/Chenopodiaceae (A/C) ratios are used to differentiate between desert and steppe (El-Moslimany, 1990; Herzschuh, 2007; Y. Zhao et al., 2012). We assume in accordance with Cour et al. (1999) that pollen spectra with A/C ratios ≥ 2 indicate montane steppe or sub-desert. However, Van Campo and Gasse (1993), investigating pollen spectra from the Sumxi Co Basin (Western Tibet), point out that Chenopodiaceae pollen might partly originate from local halophytic taxa close to lake margins. Although saline vegetation can be found along the present shorelines of Lake Karakul, its relative contribution might be small because this large lake should reflect the regional and extra-regional pollen component rather than the local pollen contribution (Sugita, 1994). Different A/C ratios from the wider region of Central Asia are displayed in Fig. 6. In the Lake Karakul record, high percentages of *Ephedra* (including *E. distachya*-type and *E. fragilis*-type) pollen coincide with increases in Chenopodiaceae, which are likewise common dryland taxa (Herzschuh, 2007; Van Campo and Gasse, 1993). Brassicaceae (mean 1.2%, max. ~5%) and Asteraceae (mean ~2%, max. ~7%) are common alpine steppe and alpine meadow taxa and likely represent increased moisture availability (Herzschuh et al., 2010), although it cannot be ruled out that they include pollen from a number of ephemeral species, which have been recorded in Central Asian desert (De-Yuan and Blackmore, 2015). In comparison to Chenopodiaceae and *Artemisia*, Poaceae (mean 6.7%, max. ~12%) and Cyperaceae (mean 1.8%, max. ~5%) require more humid conditions (Jiang et al., 2013). Again, because of the lake size, the relative contribution of Cyperaceae from wetlands connected to the lake's shorelines (Mętrak et al., 2017) might be small. Herzschuh et al. (2004) investigated recent pollen spectra from the

Alashan Plateau and Qilian Mountains (NW China), and found high values of steppe herbs and shrubs such as Brassicaceae, Asteraceae, Rosaceae, and Papaveraceae and in the same samples high values of Poaceae and Cyperaceae in the alpine setting, agreeing well with our interpretation of alpine steppe vegetation as the dominant vegetation composition type.

Cohen (2003) states that although *Botryococcus* occurs in a wide variety of lakes, it is most common in arid and semi-arid regions and shallow waterbodies, while *Pediastrum* can be found in all lake habitat zones, but is often associated with macrophyte occurrences in the more littoral zone. The abundance of the non-pollen-palynomorph (NPP1) shows a similar picture to the *Botryococcus* and *Pediastrum* distribution. We therefore assume that this NPP originates from shallow shoreline areas, rather than the open water body.

5.2. *Vegetation change before and during the global Last Glacial Maximum (27.6-19.4 cal kyr BP)*

Overall, the pollen spectra from Lake Karakul are strongly dominated by *Artemisia* and Chenopodiaceae. Other taxa, except for Poaceae and Cyperaceae, rarely exceed 5%. The contribution of arboreal taxa, especially Pinaceae (~6.5%) and *Alnus* (~3%), is particularly high from 27.6-19.4 cal kyr BP (PAZ 1). As there were likely no natural forests in the high Pamir Mountains due to the elevation and the dry conditions (Körner, 2012), these results suggest a strong influence of an extra-regional pollen component. This long distance component was likely transported via the Westerlies to the region rather than the Asian Summer Monsoon, which was comparatively weak during this period (Qin et al. 1998). Model simulations show a southward shifted route of the Westerlies due to ice sheets covering the northern latitudes during the last glacial (Otto-Bliesner and Brady 2006). The contribution of extra-regional pollen grains to pollen records deposited during the last glacial is a common feature in semi-arid and arid regions of Central Asia (Kramer et al., 2010; Shen et al., 2005; Van Campo and Gasse, 1993), indicative of the low pollen productivity of the regional vegetation. The inference of low regional pollen productivity during PAZ 1 is also supported by the very low pollen concentrations (min. 380 and mean 1180 grain/ml). The eastern Pamir was thus likely dominated by a dry mountain steppe vegetation with low vegetation cover.

Vegetation records from the Central Asian high mountain region covering the pre- and global Last Glacial Maximum (gLGM) have not hitherto been published, although there are

records from elsewhere in Asia. At Lake Balikun (NW China), pollen spectra indicate the expansion of steppe vegetation from 30.7 to 25.1 cal kyr BP that turned into desert during the gLGM period (An et al., 2013). Likewise, *Chenopodiaceae* and *Artemisia* dominate the pollen spectra of Lake Van (eastern Turkey) indicating that dry steppe vegetation dominated the last glacial even in the lowlands to the west of the Central Asian high mountains (Litt et al., 2009, 2014).

The inference of dry steppe vegetation is in line with the globally recorded cold climate and low atmospheric CO₂ concentration (Barnola et al., 1987). Regional climate signals from the Asian high mountains likewise indicate cold conditions: several glacial advances at Lake Yashikul (southern Pamir) around 27 kyr BP (Zech et al., 2005) and the semi-arid western Himalayan-Tibetan stage (SWHTS) 2F around 30±3 kyr BP (Dortch et al., 2013) were reconstructed based on cosmogenic ¹⁰Be surface exposure dates. While the pollen signals from Lake Karakul indicate no severe changes during the preLGM period, the water level of Lake Karakul was reconstructed as higher than modern from approximately 28 to 26 cal kyr BP while it was lower between 26 and 19 cal kyr BP (Heinecke et al., 2017b).

5.3. Vegetation change during the late Glacial (19.0-13.6 cal kyr BP)

In PAZ 2, covering the Late Glacial (19.0-13.6 cal kyr BP), *Artemisia* reaches its maximum pollen percentages while *Chenopodiaceae* decreases, implying that steppe was the dominant vegetation type. Most herb taxa, except *Thalictrum*, decrease rapidly. Towards the upper end of PAZ 2 *Cyperaceae* starts to increase and suggests larger wetland areas in the vicinity of the lake. Arboreal pollen starts to decrease at the onset of PAZ 2 suggesting a slowly decreasing pollen input from far distant sources, probably as a result of higher regional vegetation cover, which is also reflected by the increasing pollen concentration. Accordingly, pollen spectra of Lake Karakul likely reflect more regional vegetation during the Late Glacial.

At Lake Balikun a transition zone from drier to moister conditions from 17.7 to 12.8 cal kyr BP is likewise proposed by An et al. (2013). At Lake Van (eastern Turkey), in contrast, more open vegetation conditions than during the late pleni-glacial were reflected in the pollen record, with an increase in *Chenopodiaceae* compared to *Artemisia*, indicating a vegetation change towards desert communities (Litt et al., 2009).

Globally, climate started to warm from around 19 cal kyr BP onwards, spanning the time of the Late Glacial (Clark et al., 2009). In the high Asian mountains, phases of increased moisture supply, likely associated with the mid-latitude Westerlies, resulted in a number of glacier advances, for example around 17 and 13.7 kyr BP in the Muztag Ata and Kongur Shan massif (Seong et al., 2009), and the SWHTS 2E, 2D, and 2C (20 ± 0.3 kyr, 16.9 ± 0.7 kyr, and 14.9 ± 0.8 kyr, respectively; Dortch et al. 2013).

Similar to the vegetation record, the Lake Karakul lake-level record suggests increased moisture availability compared with the gLGM as indicated both by low Fe/Mn ratios in Lake Karakul sediments (Heinecke et al. 2017b) and by OSL-dated palaeo-shorelines (Komatsu and Tsukamoto, 2015).

5.4. *Vegetation change during the Pleistocene-Holocene transition to mid-Holocene (12.9 to 6.7 cal kyr BP)*

The Holocene part of the core is covered by PAZ 3, which was subdivided into three sub-zones. Whether the first sub-zone also comprises parts of the Late Glacial (i.e. the Bølling/Allerød period and the Younger Dryas) cannot be resolved by the age-depth model or the pollen record. In our pollen record, arboreal pollen is strongly reduced, leading to the conclusion that the long distance input decreased further at the beginning of the Holocene. At the same time the *Artemisia* percentage decreased while the *Chenopodiaceae* percentage increased indicating less moisture availability (El-Moslimany, 1990). This is also reflected in the increased amount of *Ephedra* pollen. Overall the pollen spectra suggest an expansion of dry, rather open, steppe vegetation, while slight increases in *Betula* might reflect a partly mixed signal from long distance transport and regional sources, which together with slight increases in *Poaceae* and *Cyperaceae* probably reflect an increase in riverine wetland area in the lake's vicinity, the shoreline areas and the lowlands. This is in line with the reconstructed alpine meadow vegetation at Lake Son Kul (western Tien Shan), which is located in a high alpine valley area with relatively moist conditions and surrounded by drier mountain ridges (Mathis et al., 2014). Mathis et al. (2014) nevertheless propose a wet early to middle Holocene in the Tien Shan based on a high aridity index, implying higher tree cover in the region, and a high input of arboreal pollen and *Poaceae*.

The expansion of dryland vegetation as recorded by Lake Karakul pollen spectra is also reflected in the pollen signals of Lake Wulungu from 9.5 to 6.7 cal kyr BP (northern Xinjiang,

north-western China) presented by Liu et al. (2008; Fig. 6) and at Boston Lake (located in the eastern Tien Shan), where low A/C ratios (Xu et al. 1998; Fig. 6) and increased *Ephedra* percentages characterize a desert-steppe vegetation from 8.0-6.0 cal kyr BP (Huang et al., 2009). Similar to our record, Herzschuh et al. (2004) find high values of Chenopodiaceae and *Ephedra*, and infer desert as the dominant vegetation type from 10.7 to 5.4 cal kyr BP around the eastern Juyan palaeo-lake on the Alashan Plateau. Desert vegetation was also reconstructed by Van Campo et al. (1996) at Bangong Co from ~9.9 kyr BP to 9.6 kyr BP, with a sudden shift towards steppe vegetation and moister conditions thereafter. A similar trend was recorded from Sumxi Co, where from 12.7 to 10 kyr BP desert-type vegetation with dominant Chenopodiaceae and *Ephedra* was replaced by *Artemisia*-dominated steppe from ca. 10 to 5 kyr BP (Gasse et al., 1991). In the west, at Lake Van, the climate during the early Holocene seems drier too, as tree pollen remains low, while *Ephedra* and Chenopodiaceae reach maximum values (Landmann et al., 1996; Wick et al., 2003).

Global temperature rose notably at the beginning of the Holocene (Marcott et al., 2013). Diverging inferences are given concerning the moisture changes: while desertification is recorded in Central Asia (Jin et al., 2012), higher moisture availability is inferred from glacier advances in the high Asian mountains, i.e. at Muztag Ata and Kongur Shan Mountains during the early Holocene (Seong et al., 2009). The latter agrees with the high lake levels inferred at Lake Karakul (Heinecke et al., 2017a) and increased freshwater input into Lake Issyk-Kul (Ricketts et al., 2001). However, higher moisture availability during the early Holocene is not reflected in the vegetation signals of Lake Karakul where a rapid drop in *Artemisia* pollen and increase in Chenopodiaceae and *Ephedra* pollen percentages indicate a change from steppe to dry steppe vegetation.

The dissimilarities between the inferred moisture supply to the lake and the vegetation may originate from the higher seasonal evapotranspiration as a result of a higher summer insolation. While the lake continues to be fed by glacial meltwater, moisture availability for vegetation away from wetlands and riverine habitats remains low, leading to maximum *Ephedra* percentages and concomitant increases in Cyperaceae. This phenomenon may be intensified by lower atmospheric CO₂ concentrations during the early Holocene as compared to the late Holocene, which are reflected in lower net primary production due to, for example, inhibited photosynthesis as suggested by Herzschuh et al. (2011). Vegetation growing in close vicinity to

streams and the lake margin as well as wetlands is likely to be less affected by insufficient moisture availability.

5.5. *Vegetation change during the middle to late Holocene (5.4 to 1.0 cal kyr BP)*

In the Karakul pollen spectra, the proportion of *Artemisia* increases compared to the early Holocene and Chenopodiaceae and *Ephedra* decrease slightly suggesting an increase in steppe elements. This agrees with the findings at Boston Lake where higher A/C ratios (Fig. 6) imply the replacement of desert with steppe vegetation from 6 to 1.5 kyr BP (Xu et al. 1998; Huang et al., 2009). A similar trend of an increase in *Artemisia* and slight decrease in Chenopodiaceae was noted for Lake Son Kul in the Tien Shan by Mathis et al. (2014) and Lauterbach et al. (2014) (Fig. 6). However, while Poaceae percentages remain relatively high and meadow taxa such as Rosaceae, Asteraceae, and Brassicaceae increase in our Lake Karakul record, at Lake Son Kul Poaceae and *Juniperus* decrease, while Rosaceae, Asteraceae, and Brassicaceae percentages remain constantly low (Lauterbach et al., 2014; Mathis et al., 2014). These differences lead to diverging interpretations concerning moisture availability, but might simply be due to differences in the environmental setting of the studied regions despite their relatively close proximity. Lake Son Kul in the Tien Shan is located at ~3000 m asl and has a modern-day mean annual precipitation of ~470 mm, compared to an elevation of ~3915 m asl and mean annual precipitation of ~82 mm for Lake Karakul. Accordingly, regional differences may mean that dry areas in the eastern Pamir are more sensitive to moisture changes, while the vegetation in the Tien Shan is more sensitive to temperature change. At Lake Karakul we think the increase in steppe vegetation compared to the early Holocene is a result of increased moisture possibly caused by a temperature decrease and the associated reduction in evaporative stress.

On a global scale, warmer and wetter conditions prevailed during the middle Holocene and shifted towards colder and drier conditions in the late Holocene. Increasing $\delta^{18}\text{O}$ values from the Guliya Ice Core (western Tibetan Plateau) reveal an abrupt decrease in temperature at 5 kyr BP (Yao et al., 1997) and Dortch et al. (2013) identified at least three SWHTS between 3.8 ± 0.6 kyr (1C) and 0.4 ± 0.1 kyr (1A). Lower lake levels are reconstructed at Lake Karakul from 6.7 cal kyr until the present-day from the decreased Fe/Mn ratios and are supported by abundant macrophyte remains preserved in the sediment cores of Lake Karakul. A warmer period, with a few colder intervals, led to an increased meltwater influx since 1.9 cal kyr BP (Mischke et al.,

2010; Taft et al., 2014), which coincides with a warm period proposed by Aichner et al. (2015). At Boston Lake, a number of dry spells throughout the late Holocene have also been recorded (Wünnemann et al., 2006), corresponding with a slight drying trend and mainly negative water balance at Lake Son Kul, with a few, up to centennial-long, reversals (Huang et al., 2014; Lauterbach et al., 2014).

5.6 Vegetation change during the last millennium and potential human impact

PAZ 3c spans the core section from 0.7 cal kyr BP to present-day (62 cm to top). The pollen spectra indicate an expansion of alpine steppe and alpine meadow from the high percentages of Asteraceae, Brassicaceae, Cyperaceae, and Poaceae. Possible human impact might be indicated by the increase of Asteraceae pollen, a taxonomic group including many typical grazing-responsive and nitrophile weeds, in combination with *Plantago* pollen recorded in PAZ 3c. At Lake Kichikol (Alai Range, 100 km north of Lake Karakul), human impact based on the appearance of *Plantago* pollen and an increase in charcoal accumulation rates is first inferred around 2.1 cal kyr BP (Beer et al., 2007). At Lake Son Kul stable steppe vegetation is inferred since 5 cal kyr BP and although human impact during the last millennia in the form of nomadic herders and grazing livestock is documented from excavations of burial sites and petroglyphs (Yablonsky, 1995), no signal was imprinted in the pollen record (Lauterbach et al., 2014). Chen et al. (2010) compared a number of studies from arid Central Asia focusing on the last millennium and found that the majority of records show a rather dry Medieval Warm Period and an increase in moisture availability during the Little Ice Age. However, due to spatial differences and a lack of studies it is difficult to draw more detailed conclusions.

6. Conclusions

The pollen spectra from a composite core at Lake Karakul in the eastern Pamir show:

- high arboreal pollen percentages during the late Pleistocene, revealing long-distance input and suggesting a dominant extra-regional vegetation signal in a sparsely vegetated dry mountain steppe landscape
- continued prevalence of steppe, with high values of *Ephedra* as a response to even drier conditions during the period from 12.9-6.7 cal kyr BP
- vegetation changes after 6.7 cal kyr BP tending towards steppe vegetation with slightly moister conditions
- the development of alpine steppe and meadows during the last millennium, with minor indications of grazing activity in the landscape

A comparison of the vegetation signal from Lake Karakul with a previously published sedimentary record shows that during the late Pleistocene both records have similar cold and dry conditions, but differ during the Holocene. In the early Holocene inferred lake levels remain high, but the pollen record appears to respond to a decrease in available moisture. Around 6.7 cal kyr BP a threshold is crossed, which is likely related to decreasing lake level. While the pollen record indicates moister conditions from 6.7 cal kyr BP until present-day, the sedimentary parameter suggests rather drier conditions. These differences are likely related to fine-scale differences as the lake is fed mainly by streams transporting meltwater from glaciers, while the majority of the vegetation does not grow in the direct vicinity of streams and the lake, as well as variations in evaporation due to insolation changes.

Acknowledgments

We thank Romy Zibulsky for the identification of the NPP – gemmae of a Bryophyte, and Xianyong Cao for support during data collection for Fig. 6. L.H. further thanks the Quaternary Environments and Geoarchaeology working group at the School of Environment, Education and Development at the University of Manchester for support during a Visiting Scholarship research stay. We thank the editors Howard Falcon-Lang and Thomas Algeo and two anonymous reviewer for their comments, which helped to improve this manuscript. Funding: This work was supported by the German Research Foundation [grant numbers Mi 730/15-1 and 15-2, He3622/21]; the DFG Graduate School GK 1364]

Data Availability

The palynological datasets presented in this paper are available from Pangaea (doi: xxx will be added upon acceptance)

References

- Agakhanyantz, O.E., Lopatin, I.K., 1978. Main characteristics of the ecosystems of the Pamirs, USSR. *Arct. Alp. Res.* 10, 397–407. doi:10.2307/1550770
- Aichner, B., Feakins, S.J., Lee, J.E., Herzsuh, U., Liu, X., 2015. High-resolution leaf wax carbon and hydrogen isotopic record of the late Holocene paleoclimate in arid Central Asia. *Clim. Past* 11, 619–633.
- An, C.-B., Tao, S.-C., Zhao, J., Chen, F.-H., Lv, Y., Dong, W., Li, H., Zhao, Y., Jin, M., Wang, Z., 2013. Late Quaternary (30.7–9.0 cal ka BP) vegetation history in Central Asia inferred from pollen records of Lake Balikun, northwest China. *J. Paleolimnol.* 49, 145–154. doi:10.1007/s10933-012-9649-7
- Barnett, T.P., Adam, J.C., Lettenmaier, D.P., 2005. Potential impacts of a warming climate on water availability in snow-dominated regions. *Nature* 438, 303–309. doi:10.1038/nature04141
- Barnola, J.M., Raynaud, D., Korotkevich, Y.S., Lorius, C., 1987. Vostok ice core provides 160,000-year record of atmospheric CO₂. *Nature* 329, 408–414. doi:10.1038/329408a0
- Beer, R., Heiri, O., Tinner, W., 2007. Vegetation history, fire history and lake development recorded for 6300 years by pollen, charcoal, loss on ignition and chironomids at a small lake in southern Kyrgyzstan (Alay Range, Central Asia). *The Holocene* 17, 977–985. doi:10.1177/0959683607082413
- Beug, H.-J., 2004. Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete. Dr. Friedrich Pfeil, München.
- Birks, H.J.B., Birks, H.H., 1980. Quaternary palaeoecology. Edward Arnold London.
- Breckle, S., Wucherer, W., 2006. Vegetation of the Pamir (Tajikistan): Land Use and Desertification Problems, in: Spehn, E., Körner, C., Liberman, M. (Eds.), *Land Use Change and Mountain Biodiversity*. CRC Press, pp. 225–237. doi:10.1201/9781420002874.ch16
- Chen, F.-H., Chen, J.-H., Holmes, J., Boomer, I., Austin, P., Gates, J.B., Wang, N.-L., Brooks, S.J., Zhang, J.-W., 2010. Moisture changes over the last millennium in arid central Asia: a review, synthesis and comparison with monsoon region. *Quat. Sci. Rev.* 29, 1055–1068. doi:10.1016/j.quascirev.2010.01.005
- Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica, J.X., Hostetler, S.W., McCabe, A.M., 2009. The Last Glacial Maximum. *Science* 325, 710–714. doi:10.1126/science.1172873
- Cohen, A.S., 2003. *Paleolimnology: the history and evolution of lake systems*. Oxford University Press, USA.
- Cour, P., Zheng, Z., Duzer, D., Calleja, M., Yao, Z., 1999. Vegetational and climatic significance of modern pollen rain in northwestern Tibet. *Rev. Palaeobot. Palynol.* 104, 183–204. doi:10.1016/S0034-6667(98)00062-1
- Davis, M.B., 1967. Pollen accumulation rates at Rogers Lake, Connecticut, during late- and postglacial time. *Rev. Palaeobot. Palynol.* 2, 219–230. doi:10.1016/0034-6667(67)90150-9

- De-Yuan, H. and Blackmore, S., Eds. 2015. *Plant of China: A Companion to the Flora of China*. Cambridge University Press, Cambridge
- Dortch, J.M., Owen, L.A., Caffee, M.W., 2013. Timing and climatic drivers for glaciation across semi-arid western Himalayan–Tibetan orogen. *Quat. Sci. Rev.* 78, 188–208. doi:10.1016/j.quascirev.2013.07.025
- El-Moslimany, A.P., 1990. Ecological significance of common nonarboreal pollen: examples from drylands of the Middle East. *Rev. Palaeobot. Palynol.*, The Proceedings of the 7th International Palynological Congress (Part I) 64, 343–350. doi:10.1016/0034-6667(90)90150-H
- Feng, Z.-D., Wu, H.N., Zhang, C.J., Ran, M., Sun, A.Z., 2013. Bioclimatic change of the past 2500 years within the Balkhash Basin, eastern Kazakhstan, Central Asia. *Quat. Int.*, Hydrological and Ecological Responses to Climatic Change and to Land-use/land-cover changes in Central Asia 311, 63–70. doi:10.1016/j.quaint.2013.06.032
- Gasse, F., Arnold, M., Fontes, J.C., Fort, M., Gibert, E., Huc, A., Bingyan, L., Yuanfang, L., Qing, L., Mélières, F., Campo, E.V., Fubao, W., Qingsong, Z., 1991. A 13,000-year climate record from western Tibet. *Nature* 353, 742–745. doi:10.1038/353742a0
- Gasse, F., Fontes, J.C., Van Campo, E., Wei, K., 1996. Holocene environmental changes in Bangong Co basin (Western Tibet). Part 4: Discussion and conclusions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, Environmental Changes in the Tibetan Plateau and Surrounding Areas 120, 79–92. doi:10.1016/0031-0182(95)00035-6
- Gorbunov, A.P., 1990. Permafrost in the arid mountains of middle asia—the eastern Pamirs, USSR. *Permafr. Periglac. Process.* 1, 309–312. doi:10.1002/ppp.3430010311
- Grimm, E., 2011. *TILIA 1.7*. 16. Ill. State Mus. Res. Collect. Cent.
- Grimm, E.C., 1987. CONISS: a FORTRAN 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares. *Comput. Geosci.* 13, 13–35.
- Heinecke, L., Epp, L.S., Reschke, M., Stoof-Leichsenring, K.R., Mischke, S., Plessen, B., Herzschuh, U., 2017a. Aquatic macrophyte dynamics in Lake Karakul (Eastern Pamir) over the last 29 cal ka revealed by sedimentary ancient DNA and geochemical analyses of macrofossil remains. *J. Paleolimnol.* 1–15. doi:10.1007/s10933-017-9986-7
- Heinecke, L., Mischke, S., Adler, K., Barth, A., Biskaborn, B.K., Plessen, B., Nitze, I., Kuhn, G., Rajabov, I., Herzschuh, U., 2017b. Climatic and limnological changes at Lake Karakul (Tajikistan) during the last ~29 cal ka. *J. Paleolimnol.* 1–18. doi:10.1007/s10933-017-9980-0
- Herzschuh, U., 2007. Reliability of pollen ratios for environmental reconstructions on the Tibetan Plateau. *J. Biogeogr.* 34, 1265–1273. doi:10.1111/j.1365-2699.2006.01680.x
- Herzschuh, U., Birks, H.J.B., Mischke, S., Zhang, C., Böhner, J., 2010. A modern pollen–climate calibration set based on lake sediments from the Tibetan Plateau and its application to a Late Quaternary pollen record from the Qilian Mountains. *J. Biogeogr.* 37, 752–766. doi:10.1111/j.1365-2699.2009.02245.x
- Herzschuh, U., Ni, J., Birks, H.J.B., Böhner, J., 2011. Driving forces of mid-Holocene vegetation shifts on the upper Tibetan Plateau, with emphasis on changes in atmospheric CO₂ concentrations. *Quat. Sci. Rev.* 30, 1907–1917. doi:10.1016/j.quascirev.2011.03.007
- Herzschuh, U., Tarasov, P., Wünnemann, B., Hartmann, K., 2004. Holocene vegetation and climate of the Alashan Plateau, NW China, reconstructed from pollen data. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 211, 1–17. doi:10.1016/j.palaeo.2004.04.001
- Huang, X., Oberhänsli, H., von Suchodoletz, H., Prasad, S., Sorrel, P., Plessen, B., Mathis, M., Usabaliev, R., 2014. Hydrological changes in western Central Asia (Kyrgyzstan) during

- the Holocene as inferred from a palaeolimnological study in lake Son Kul. *Quat. Sci. Rev.* 103, 134–152. doi:10.1016/j.quascirev.2014.09.012
- Huang, X.Z., Chen, F.H., Fan, Y.X., Yang, M.L., 2009. Dry late-glacial and early Holocene climate in arid central Asia indicated by lithological and palynological evidence from Bosten Lake, China. *Quat. Int.* 194, 19–27. doi:10.1016/j.quaint.2007.10.002
- Hyvärinen, H., 1970. Flandrian pollen diagrams from Svalbard. *Geogr. Ann. Ser. Phys. Geogr.* 52, 213–222. doi:10.2307/520815
- Immerzeel, W.W., Beek, L.P.H. van, Bierkens, M.F.P., 2010. Climate Change Will Affect the Asian Water Towers. *Science* 328, 1382–1385. doi:10.1126/science.1183188
- Jiang, Q., Ji, J., Shen, J., Matsumoto, R., Tong, G., Qian, P., Ren, X., Yan, D., 2013. Holocene vegetational and climatic variation in westerly-dominated areas of Central Asia inferred from the Sayram Lake in northern Xinjiang, China. *Sci. China Earth Sci.* 56, 339–353. doi:10.1007/s11430-012-4550-9
- Jin, L., Chen, F., Morrill, C., Otto-Bliesner, B.L., Rosenbloom, N., 2012. Causes of early Holocene desertification in arid central Asia. *Clim. Dyn.* 38, 1577–1591. doi:10.1007/s00382-011-1086-1
- Juggins, S., 2017. rioja: Analysis of quaternary science data, R package.
- Komatsu, T., Tsukamoto, S., 2015. Late Glacial lake-level changes in the Lake Karakul basin (a closed glacierized-basin), eastern Pamirs, Tajikistan. *Quat. Res.* 83, 137–149. doi:10.1016/j.yqres.2014.09.001
- Körner, C., 2012. *Alpine Treelines: Functional Ecology of the Global High Elevation Tree Limits*. Springer Science & Business Media.
- Kramer, A., Herzschuh, U., Mischke, S., Zhang, C., 2010. Late glacial vegetation and climate oscillations on the southeastern Tibetan Plateau inferred from the Lake Naleng pollen profile. *Quat. Res.* 73, 324–335. doi:10.1016/j.yqres.2009.12.003
- Landmann, G., Reimer, A., Kempe, S., 1996. Climatically induced lake level changes at Lake Van, Turkey, during the Pleistocene/Holocene Transition. *Glob. Biogeochem. Cycles* 10, 797–808. doi:10.1029/96GB02347
- Lauterbach, S., Witt, R., Plessen, B., Dulski, P., Prasad, S., Mingram, J., Gleixner, G., Hettler-Riedel, S., Stebich, M., Schnetger, B., Schwalb, A., Schwarz, A., 2014. Climatic imprint of the mid-latitude Westerlies in the Central Tian Shan of Kyrgyzstan and teleconnections to North Atlantic climate variability during the last 6000 years. *The Holocene* 24, 970–984. doi:10.1177/0959683614534741
- Lioubimtseva, E., Henebry, G.M., 2009. Climate and environmental change in arid Central Asia: Impacts, vulnerability, and adaptations. *J. Arid Environ.* 73, 963–977. doi:10.1016/j.jaridenv.2009.04.022
- Litt, T., Krastel, S., Sturm, M., Kipfer, R., Örcen, S., Heumann, G., Franz, S.O., Ülgen, U.B., Niessen, F., 2009. ‘PALEOVAN’, International Continental Scientific Drilling Program (ICDP): site survey results and perspectives. *Quat. Sci. Rev.* 28, 1555–1567. doi:10.1016/j.quascirev.2009.03.002
- Litt, T., Pickarski, N., Heumann, G., Stockhecke, M., Tzedakis, P.C., 2014. A 600,000 year long continental pollen record from Lake Van, eastern Anatolia (Turkey). *Quat. Sci. Rev.*, Special Issue: Results from the PALEOVAN Drilling Project: a 600,000 year long continental archive in the Near East 104, 30–41. doi:10.1016/j.quascirev.2014.03.017
- Liu, X., Herzschuh, U., Shen, J., Jiang, Q., Xiao, X., 2008. Holocene environmental and climatic changes inferred from Wulungu Lake in northern Xinjiang, China. *Quat. Res.* 70, 412–425. doi:10.1016/j.yqres.2008.06.005

- Lutz, A.F., Immerzeel, W.W., Shrestha, A.B., Bierkens, M.F.P., 2014. Consistent increase in High Asia's runoff due to increasing glacier melt and precipitation. *Nat. Clim. Change* 4, 587.
- Makhmadaliev, B., Kayumov, A., Novikov, V., Mustaeva, N., Rajabov, I., 2008. The second national communication of the Republic of Tajikistan under the United Nations framework convention on climate change. Dushanbe 89.
- Marcott, S.A., Shakun, J.D., Clark, P.U., Mix, A.C., 2013. A reconstruction of regional and global temperature for the past 11,300 years. *Science* 339, 1198–1201. doi:10.1126/science.1228026
- Markgraf, V., 1980. Pollen Dispersal in a Mountain Area. *Grana* 19, 127–146. doi:10.1080/00173138009424995
- Mathis, M., Sorrel, P., Klotz, S., Huang, X., Oberhänsli, H., 2014. Regional vegetation patterns at lake Son Kul reveal Holocene climatic variability in central Tien Shan (Kyrgyzstan, Central Asia). *Quat. Sci. Rev.* 89, 169–185. doi:10.1016/j.quascirev.2014.01.023
- Mętrak, M., Chachulski, Ł., Navruzshoev, D., Pawlikowski, P., Rojan, E., Sulwiński, M., Suska-Malawska, M., 2017. Nature's patchwork: How water sources and soil salinity determine the distribution and structure of halophytic plant communities in arid environments of the Eastern Pamir. *PLOS ONE* 12, e0174496. doi:10.1371/journal.pone.0174496
- Mętrak, M., Sulwinski, M., Chachulski, Ł., Wilk, M., Suska-Malawska, M., 2015. Creeping Environmental Problems in the Pamir Mountains: Landscape Conditions, Climate Change, Wise Use and Threats, in: Öztürk, M., Hakeem, K.R., Faridah-Hanum, I., Efe, R. (Eds.), *Climate Change Impacts on High-Altitude Ecosystems*. Springer International Publishing, pp. 665–694. doi:10.1007/978-3-319-12859-7_28
- Mischke, S., Lai, Z., Aichner, B., Heinecke, L., Mahmoudov, Z., Kuessner, M., Herzsuh, U., 2017. Radiocarbon and optically stimulated luminescence dating of sediments from Lake Karakul, Tajikistan. *Quat. Geochronol.* 41, 51–61. doi:10.1016/j.quageo.2017.05.008
- Mischke, S., Rajabov, I., Mustaeva, N., Zhang, C., Herzsuh, U., Boomer, I., Brown, E.T., Andersen, N., Myrbo, A., Ito, E., Schudack, M.E., 2010. Modern hydrology and late Holocene history of Lake Karakul, eastern Pamirs (Tajikistan): A reconnaissance study. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 289, 10–24. doi:10.1016/j.palaeo.2010.02.004
- Moore, P.D., Webb, J.A., Collison, M.E., 1991. *Pollen analysis*. Blackwell scientific publications.
- Nöth, L., Ficker, H.V., Rickmers, W.R. (Eds), 1932. *Geologische Untersuchungen im Nordwestlichen Pamir-Gebiet und Mittleren Trans-Alai, Wissenschaftliche Ergebnisse der Alai-Pamir Expedition 1928*. Reimer-Vohsen, Berlin.
- Otto-Bliesner, B.L. and Brady, E.C., 2006. Last Glacial Maximum and Holocene Climate in CCSM3. *J. of Climate* 19, 2526–2544. doi: 10.1175/JCLI3748.1
- Pachauri, R.K., Allen, M.R., Barros, V.R., Broome, J., Cramer, W., Christ, R., Church, J.A., Clarke, L., Dahe, Q., Dasgupta, P., Dubash, N.K., Edenhofer, O., Elgizouli, I., Field, C.B., Forster, P., Friedlingstein, P., Fuglestedt, J., Gomez-Echeverri, L., Hallegatte, S., Hegerl, G., Howden, M., Jiang, K., Jimenez Cisneroz, B., Kattsov, V., Lee, H., Mach, K.J., Marotzke, J., Mastrandrea, M.D., Meyer, L., Minx, J., Mulugetta, Y., O'Brien, K., Oppenheimer, M., Pereira, J.J., Pichs-Madruga, R., Plattner, G.-K., Pörtner, H.-O., Power, S.B., Preston, B., Ravindranath, N.H., Reisinger, A., Riahi, K., Rusticucci, M., Scholes, R., Seyboth, K., Sokona, Y., Stavins, R., Stocker, T.F., Tschakert, P., van Vuuren, D., van Ypserle, J.-P., 2014. *Climate Change 2014: Synthesis Report. Contribution of Working*

- Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC, Geneva, Switzerland.
- Prentice, I.C., 1985. Pollen representation, source area, and basin size: Toward a unified theory of pollen analysis. *Quat. Res.* 23, 76–86. doi:10.1016/0033-5894(85)90073-0
- Qin, B., Shi, Y., and YU, G. 1998. The reconstruction and interpretations of lake status at 6 ka and 18 ka B.P. in inland mainland Asia. *Chinese Science Bulletin* 43, Issue 14, 1145–1157. doi: 10.1007/BF02883214
- R Core Team, 2012. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rhodes, T.E., Gasse, F., Lin, R., Fontes, J.-C., Wei, K., Bertrand, P., Gibert, E., Mélières, F., Tucholka, P., Wang, Z., Cheng, Z.-Y., 1996. A Late Pleistocene-Holocene lacustrine record from Lake Manas, Zunggar (northern Xinjiang, western China). *Palaeogeogr. Palaeoclimatol. Palaeoecol., Environmental Changes in the Tibetan Plateau and Surrounding Areas* 120, 105–121. doi:10.1016/0031-0182(95)00037-2
- Ricketts, R.D., Johnson, T.C., Brown, E.T., Rasmussen, K.A., Romanovsky, V.V., 2001. The Holocene paleolimnology of Lake Issyk-Kul, Kyrgyzstan: trace element and stable isotope composition of ostracodes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 176, 207–227. doi:10.1016/S0031-0182(01)00339-X
- Seong, Y.B., Owen, L.A., Yi, C., Finkel, R.C., 2009. Quaternary glaciation of Muztag Ata and Kongur Shan: Evidence for glacier response to rapid climate changes throughout the Late Glacial and Holocene in westernmost Tibet. *Geol. Soc. Am. Bull.* 121, 348–365. doi:10.1130/B26339.1
- Shen, J., Liu, X., Wang, S., Matsumoto, Ryo, 2005. Palaeoclimatic changes in the Qinghai Lake area during the last 18,000 years. *Quat. Int.* 136, 131–140. doi:10.1016/j.quaint.2004.11.014
- Sorrel, P., Popescu, S.-M., Klotz, S., Suc, J.-P., Oberhänsli, H., 2007. Climate variability in the Aral Sea basin (Central Asia) during the late Holocene based on vegetation changes. *Quat. Res., Reconstructing past environments from remnants of human occupation and sedimentary archives in western Eurasia* 67, 357–370. doi:10.1016/j.yqres.2006.11.006
- Strecker, M.R., Frisch, W., Hamburger, M.W., Ratschbacher, L., Semiletkin, S., Zamoruyev, A., Sturchio, N., 1995. Quaternary deformation in the Eastern Pamirs, Tadjikistan and Kyrgyzstan. *Tectonics* 14, 1061–1079. doi:10.1029/95TC00927
- Sugita, S., 1994. Pollen representation of vegetation in quaternary sediments: theory and method in patchy vegetation. *J. Ecol.* 82, 881–897. doi:10.2307/2261452
- Sun, X.J., Du, N.Q., Weng, C.Y., Lin, R.F. and Wei, K.Q. 1994. Paleovegetation and paleoenvironment of Manasi Lake, Xinjiang, N.W. China during the last 14 000 years. *Quaternary Sciences* (3): 239–248. (in Chinese with English abstract)
- Taft, L., Mischke, S., Wiechert, U., Leipe, C., Rajabov, I., Riedel, F., 2014. Sclerochronological oxygen and carbon isotope ratios in *Radix* (Gastropoda) shells indicate changes of glacial meltwater flux and temperature since 4,200 cal yr BP at Lake Karakul, eastern Pamirs (Tajikistan). *J. Paleolimnol.* 52, 27–41. doi:10.1007/s10933-014-9776-4
- Tang, Z., Ding, Z., White, P.D., Dong, X., Ji, J., Jiang, H., Luo, P., Wang, X., 2011. Late Cenozoic central Asian drying inferred from a palynological record from the northern Tian Shan. *Earth Planet. Sci. Lett.* 302, 439–447. doi:10.1016/j.epsl.2010.12.042
- Tarasov, P.E. and Kremenetskii, K.V. 1995. Geochronology and stratigraphy of the Holocene lacustrine-bog deposits in northern and central Kazakhstan. *Stratigraphy and Geological Correlation* 3: 73–80.

- ter Braak, C.J.F., Šmilauer, P., 2012. Canoco reference manual and user's guide: software for ordination, version 5.0. Microcomputer Power, Ithaca USA.
- Van Campo, E., Cour, P., Sixuan, H., 1996. Holocene environmental changes in Bangong Co basin (Western Tibet). Part 2: The pollen record. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, Environmental Changes in the Tibetan Plateau and Surrounding Areas 120, 49–63. doi:10.1016/0031-0182(95)00033-X
- Van Campo, E., Gasse, F., 1993. Pollen- and diatom-inferred climatic and hydrological changes in Sumxi Co Basin (Western Tibet) since 13,000 yr B.P. *Quat. Res.* 39, 300–313. doi:10.1006/qres.1993.1037
- Wang, W., Feng, Z., Ran, M., Zhang, C., 2013. Holocene climate and vegetation changes inferred from pollen records of Lake Aibi, northern Xinjiang, China: A potential contribution to understanding of Holocene climate pattern in East-central Asia. *Quat. Int.*, Hydrological and Ecological Responses to Climatic Change and to Land-use/land-cover changes in Central Asia 311, 54–62. doi:10.1016/j.quaint.2013.07.034
- Wick, L., Lemcke, G., Sturm, M., 2003. Evidence of Lateglacial and Holocene climatic change and human impact in eastern Anatolia: high-resolution pollen, charcoal, isotopic and geochemical records from the laminated sediments of Lake Van, Turkey. *The Holocene* 13, 665–675. doi:10.1191/0959683603hl653rp
- Wünnemann, B., Mischke, S., Chen, F., 2006. A Holocene sedimentary record from Bosten Lake, China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 234, 223–238. doi:10.1016/j.palaeo.2005.10.016
- Xu, Y.Q. 1998. The assemblage of Holocene spore pollen and its environment in Bosten Lake area, Xinjiang. *Arid Land Geography* 21(2): 43-49.(in Chinese with English abstract)
- Yablonsky, L.T., 1995. The material culture of the Saka and historical reconstruction. *Nomads Eurasian Steppe Early Iron Age* Berkeley Zinat Press P 201–35.
- Yao, T., Thompson, L.G., Shi, Y., Qin, D., Jiao, K., Yang, Z., Tian, L., Thompson, E.M., 1997. Climate variation since the Last Interglaciation recorded in the Guliya ice core. *Sci. China Ser. Earth Sci.* 40, 662–668. doi:10.1007/BF02877697
- Zech, R., Abramowski, U., Glaser, B., Sosin, P., Kubik, P.W., Zech, W., 2005. Late Quaternary glacial and climate history of the Pamir Mountains derived from cosmogenic ^{10}Be exposure ages. *Quat. Res.* 64, 212–220. doi:10.1016/j.yqres.2005.06.002
- Zhao, K., Li, X., Dodson, J., Atahan, P., Zhou, X., Bertuch, F., 2012. Climatic variations over the last 4000 cal yr BP in the western margin of the Tarim Basin, Xinjiang, reconstructed from pollen data. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 321, 16–23. doi:10.1016/j.palaeo.2012.01.012
- Zhao, Y., Liu, H., Li, F., Huang, X., Sun, J., Zhao, W., Herzsuh, U., Tang, Y., 2012. Application and limitations of the *Artemisia*/Chenopodiaceae pollen ratio in arid and semi-arid China. *The Holocene* 22, 1385–1392. doi:10.1177/0959683612449762

Figure Captions

Fig.1: Lake Karakul in the Pamir Mountains, arid Central Asia (regional map –Landsat-8 (OLI) image, 22 July 2014). The yellow circle indicates the coring location; A-C refers to photographs of different vegetation communities with A - typical teresken steppe to desert vegetation, B - wetland on saline soils close to the lake shoreline and C - sparse steppe vegetation with riverine habitat dominated by Cyperaceae.

Fig. 2: Sedimentology of cores KK12-1 and KK12-2, including sample depth for radiocarbon and OSL dating and XRF peaks in selected elements used to create the composite core. Age-depth model created in OxCal 4.2 (adapted from Mischke et al. 2017).

Fig. 3: Pollen stratigraphy of the composite core from Lake Karakul. Black filled silhouettes represent original pollen abundance; white silhouettes exaggerate original abundance by five to improve detectability.

Fig. 4: Biplot of axis 1 and axis 2 of a Principal Component Analysis (PCA) performed on pollen from pollen assemblage zone 3 of the composite core from Lake Karakul and age of samples in cal yr BP.

Fig. 5: Pollen spectra of the short core TAJ-Kar-08-1B. Core chronology is presented in Mischke et al. (2010).

Fig. 6. Comparison of *Artemisia*/Chenopodiaceae ratios from lakes from the wider region of Central Asia, from west to east, Lake Karakul (this study), Lake Kichikul (Beer et al. 2007), Lake Sonkul (Lauterbach et al. 2014), Lake Pashennoe (Tarosov et al. 1995), Lake Bangong (Van Campo et al. 1996), Lake Sumxi (Van Campo and Gasse 1993), Lake Aibi (Wang et al. 2013), Lake Sayram (Jiang et al. 2013), Lake Wulung (Liu et al. 2008), Lake Manas (Sun et al. 1994), Lake Boston (Xu 1998).

Highlights:

- First vegetation study covering the late Pleistocene in high arid Central Asian
- Pollen spectra reveal high far distant input before ~13 cal kyr BP
- Regional terrestrial vegetation signal after 13 cal kyr BP for eastern Pamir
- Detection of human influence in the last millennium

Figure 1

Figure 2

Figure 4

Figure 5

Figure 6