Quaternary disappearance of tree taxa from Southern Europe: timing and trends

DOI:

[10.1016/j.quascirev.2017.02.014](https://doi.org/10.1016/j.quascirev.2017.02.014)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Magri, D., Di Rita, F., Arabbari Erkiaga, J., Fletcher, W., & González Sampériz, P. (2017). Quaternary disappearance of tree taxa from Southern Europe: timing and trends. *Quaternary Science Reviews*, 163, 23-55. <https://doi.org/10.1016/j.quascirev.2017.02.014>

Published in:

Quaternary Science Reviews

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Manuscript Details

Manuscript number	JQSR_2016_62
Title	Quaternary disappearance of tree taxa from Southern Europe: timing and trends
Article type	Invited review article

Abstract

A hundred pollen and macrofossil records over the Iberian Peninsula, Southern France, the Italian Peninsula, Greece and the Aegean, and the southwestern Black Sea area have formed the basis for a review on the Quaternary extirpation of tree populations from Southern Europe. Following a discussion of the caveats/challenges about using pollen data, the Quaternary history of tree taxa has been reconstructed with attention to *Taxodium*/*Glyptostrobus*, *Sciadopitys*, *Cathaya*, *Cedrus*, *Tsuga*, *Eucommia*, *Engelhardia*, *Carya*, *Pterocarya*, *Parrotia*, *Liquidambar*, and *Zelkova*. The timing of extinction, distributed over the whole Quaternary, appears very diverse from one region to the other, in agreement with current biodiversity in Southern Europe. The geographical patterns of persistence/disappearance of taxa show unexpected trends and rule out a simple North to South and/or West to East trend in extirpations. In particular, it was possible to detect disjunct populations (*Engelhardia*), long-term persistence of taxa in restricted regions (*Sciadopitys*), distinct populations/species/genera in different geographical areas (*Taxodium* type). Some taxa that are still widespread in Europe have undergone extirpation in Mediterranean areas in the lateglacial period and Holocene (*Buxus*, *Carpinus betulus*, *Picea*); they provide indication on the modes of disappearance of tree populations that may be useful to evaluate correctly the vulnerability of modern fragmented plant populations. The demographic histories of tree taxa obtained by combined palaeobotanical and genetic studies is a most challenging field of research not only to assess species/population differentiation, but also to reach a better understanding of extinction processes, an essential task in the current global change scenario.

Keywords	Quaternary; Southern Europe; vegetation dynamics; <i>Taxodium</i> -type; <i>Sciadopitys</i> ; <i>Cathaya</i> ; <i>Cedrus</i> ; <i>Tsuga</i> ; <i>Eucommia</i> ; <i>Engelhardia</i> ; <i>Carya</i> ; <i>Pterocarya</i> ; <i>Parrotia</i> ; <i>Liquidambar</i> ; <i>Zelkova</i>
-----------------	---

Corresponding Author	Donatella Magri
-----------------------------	-----------------

Corresponding Author's Institution	Sapienza University
---	---------------------

Order of Authors	Donatella Magri, Federico Di Rita, Josu Aranbarri Erkiaga, William Fletcher, Penélope González Sampériz
-------------------------	---

Suggested reviewers	Chronis Tzedakis, Adele Bertini, Suzanne Leroy, Demet Biltekin
----------------------------	--

Submission Files Included in this PDF

File Name [File Type]

Magri et al_ cover letter.docx [Cover Letter]

Magri et al_Quaternary disappearance.docx [Manuscript File]

Magri et al_Quaternary disappearance_Fig 1.jpg [Figure]

Magri et al_Quaternary disappearance_Fig 2.jpg [Figure]

Magri et al_Quaternary disappearance_Fig 3.jpg [Figure]

Magri et al_Quaternary disappearance_Fig 4.jpg [Figure]

Magri et al_Quaternary disappearance_Fig 5.jpg [Figure]

Magri et al_Quaternary disappearance_Fig 6.jpg [Figure]

Magri et al_Quaternary disappearance_Fig 7.jpg [Figure]

Magri et al_Quaternary disappearance_Fig 8.jpg [Figure]

Magri et al_Quaternary disappearance_Table 1.docx [Table]

Magri et al_ Highlights.docx [Highlights]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Roma, 31 October 2016

Dear Editor,

Please find uploaded the manuscript "Quaternary disappearance of tree taxa from Southern Europe: timing and trends" by Donatella Magri, Federico Di Rita, Josu Aranbarri Erkiaga, William Fletcher, Penélope González Sampériz.

All authors have materially participated in the collection of data and in the article preparation. They have seen and approved the final version of the manuscript being submitted. The article is the authors' original work, hasn't received prior publication and isn't under consideration for publication elsewhere.

This manuscript is an invited paper (Editor: Pepe Carrion).

Yours sincerely,

Prof. Donatella Magri

1 **Title:**
2 **Quaternary disappearance of tree taxa from Southern Europe: timing and trends**
3
4 **Authors:**
5 Donatella Magri¹, Federico Di Rita¹, Josu Aranbarri Erkiaga², William Fletcher³, Penélope González Sampériz⁴
6
7
8
9 ¹ Department of Environmental Biology, Sapienza University of Rome, Piazzale Aldo Moro, 5, 00185 Roma,
10 Italy
11 ² Department of Geography, Prehistory and Archaeology, University of Basque Country, C/ Tomás y Valiente
12 s/n, 01006 Vitoria-Gasteiz, Spain
13 ³ Geography, School of Environment, Education and Development, The University of Manchester, Oxford
14 Road, Manchester M13 9PL, UK
15 ⁴ Pyrenean Institute of Ecology (IPE-CSIC), Av/ Montañana 1005, 50059 Zaragoza, Spain
16
17
18 Corresponding author:
19 Donatella Magri
20 e-mail: donatella.magri@uniroma1.it
21
22

23 **Abstract**

24 A hundred pollen and macrofossil records over the Iberian Peninsula, Southern France, the Italian Peninsula,
25 Greece and the Aegean, and the southwestern Black Sea area have formed the basis for a review on the
26 Quaternary extirpation of tree populations from Southern Europe. Following a discussion of the
27 caveats/challenges about using pollen data, the Quaternary history of tree taxa has been reconstructed with
28 attention to *Taxodium/Glyptostrobus*, *Sciadopitys*, *Cathaya*, *Cedrus*, *Tsuga*, *Eucommia*, *Engelhardia*, *Carya*,
29 *Pterocarya*, *Parrotia*, *Liquidambar*, and *Zelkova*. The timing of extinction, distributed over the whole
30 Quaternary, appears very diverse from one region to the other, in agreement with current biodiversity in
31 Southern Europe. The geographical patterns of persistence/disappearance of taxa show unexpected trends
32 and rule out a simple North to South and/or West to East trend in extirpations. In particular, it was possible
33 to detect disjunct populations (*Engelhardia*), long-term persistence of taxa in restricted regions (*Sciadopitys*),
34 distinct populations/species/genera in different geographical areas (*Taxodium* type). Some taxa that are still
35 widespread in Europe have undergone extirpation in Mediterranean areas in the lateglacial period and
36 Holocene (*Buxus*, *Carpinus betulus*, *Picea*); they provide indication on the modes of disappearance of tree
37 populations that may be useful to evaluate correctly the vulnerability of modern fragmented plant
38 populations. The demographic histories of tree taxa obtained by combined palaeobotanical and genetic
39 studies is a most challenging field of research not only to assess species/population differentiation, but also
40 to reach a better understanding of extinction processes, an essential task in the current global change
41 scenario.

42

43

44 **Key-words**

45 Quaternary; Southern Europe; vegetation dynamics; *Taxodium*-type; *Sciadopitys*; *Cathaya*; *Cedrus*;
46 *Tsuga*; *Eucommia*; *Engelhardia*; *Carya*; *Pterocarya*; *Parrotia*; *Liquidambar*; *Zelkova*

47

1. Introduction

The progressive disappearance of tree taxa from Europe in the course of the late Cenozoic, and their persistence in East Asia and/or eastern North America, has long been recognized. The first papers highlighting the question, based on seed floras, date back to more than one century ago ([Reid and Reid, 1907; 1915](#)). Starting in the 1950s, several “fathers” of the European palynology published pioneering studies dealing with Quaternary extinctions of plants, including [van der Hammen \(1951\)](#), [van der Hammen et al. \(1971\)](#), [Zagwijn \(1957, 1963, 1974\)](#), [West \(1962, 1970\)](#), and [Lona \(1950\)](#). Since then, following a considerable increase in the number of published Early and Middle Pleistocene pollen records in Europe and a significant refinement of the chronological setting of the Quaternary stages, our knowledge of the timing of disappearance of tree taxa from Europe has much improved. Recently, a number of synthesis works on the history of vegetation and the Quaternary progression of plant extirpations (or regional extinctions, hereafter called extinctions) in Southern Europe have been published, describing the state of knowledge in different regions: Spain ([González-Sampériz et al., 2010; Postigo-Mijarra et al., 2010](#)), Italy ([Bertini, 2010; Magri and Palombo, 2013; Martinetto, 2015](#)), Greece ([Tzedakis et al., 2006; Velitzelos et al., 2014](#)), and Anatolia ([Biltekin et al., 2015](#)). In addition, several studies have examined the history of taxa currently extinct, or severely reduced in Europe, including *Taxodium* ([Biltekin et al., 2015](#)), *Cathaya* ([Liu and Basinger, 2000](#)), *Cedrus* ([Magri, 2012](#)), *Carya* ([Orain et al., 2013](#)), *Aesculus* ([Ravazzi, 1994; Postigo Mijarra et al., 2008](#)), *Zelkova* ([Follieri et al., 1986](#)), and *Parrotia* ([Bińka et al., 2003](#)). The present review paper is intended to summarize the available information obtained from the Pleistocene pollen records and from the synthesis papers, and to provide a holistic view of timing and trends of tree taxa disappearance in Southern Europe.

Previous works dealing with plant extinctions in Europe have mainly focused on two main problems:

1. Usage of abundances and/or last occurrences of extinct plant taxa as biostratigraphical markers. This practice has been rather common since the beginning of pollen analysis of Plio-Pleistocene deposits. In the absence of, or in support of other chronostratigraphical constraints, the presence of locally extinct taxa in pollen records may provide useful indications about the age of study sites. Many such cases can be quoted, especially before the development of advanced chronostratigraphical tools, but

the presence of extinct taxa in the pollen records is still considered an important biostratigraphical indicator. For example, the pollen record from La Côte in the western French Alps has been ascribed to the temperate Holsteinian interglacial, corresponding to Marine Isotope Stage (MIS) 11, on the basis of the occurrence of *Pterocarya* and the dominance of *Abies* (Field et al., 2000), according to the records of the Velay maar in the French Massif (Reille et al., 2000). Similarly, the record from Torre Mucchia (Central Italy) has been correlated to MIS 17 on the basis of the underlying Bruhnes/Matuyama magnetostratigraphic boundary coupled with a significant abundance of *Tsuga* (Pieruccini et al., 2016), which is only sporadically found in Central and Southern Italy after MIS 16 (Russo Ermolli et al., 2015). On the other hand, this approach may be rather weak and may lead to misleading interpretations. A classic example is the setting of the “Tiberian boundary” between the Gelasian (formerly in the Upper Pliocene) and the Calabrian (Early Pleistocene) with respect to a significant reduction of Taxodiaceae (Lona and Bertoldi, 1972; Suc, 1973). Recent pollen studies have shown that this event is not clear-cut or synchronous over a large region as previously believed, suggesting that biostratigraphical schemes need to be constantly updated and that they may be considered valid only for regions of limited extension. Thoughtful discussions on the stratigraphical potentiality and risks of palynology in the study of late Neogene and Quaternary deposits may be found in Leroy (2007) and Bertini (2010).

2. Relation between plant extinctions and climate changes. This was first raised by Reid and Reid (1915), who considered the presence of barriers to migration in the form of west-east oriented mountain ranges (e.g., Pyrenees, Alps, and Carpathians) the main cause for the extinction of taxa in NW Europe, preventing plant population migrations in response to Quaternary climate oscillations (West, 1970). Huntley (1993) suggested that the enhanced rate of extinction of genera and families in Europe during the Quaternary climate changes may depend on a much smaller area of forest vegetation persisting during glacial stages in Europe compared to eastern North America and East Asia. This reduced area is due, on the one hand, to the considerable extent of glaciated and periglacial surfaces at high latitudes, and, on the other hand, to a trend of increasing seasonality and dryness in the Mediterranean regions since the Pliocene, which according to Suc (1984) is at the origin of the

Mediterranean evergreen vegetation, although with clear latitudinal and longitudinal gradients (Suc and Popescu, 2005). In Central and Northern Europe, most temperate trees are believed to have disappeared during the Quaternary glacial periods and migrated back during interglacial periods, but a number of them were extirpated (see list in Svenning, 2003, Suppl. Materials), and some of them are still living in Southern Europe in a relictual state (e.g., *Zelkova*, *Aesculus*, *Castanea*, *Celtis*, *Cercis*, *Cupressus*, *Styrax*, *Juglans*, *Ostrya*, *Platanus*). Svenning (2003) has found that genera of woody plants that are currently widespread in Europe have greater tolerance to cold growing season and winter temperatures with respect to regionally extinct and relictual genera, while relictual genera are more drought tolerant. This indicates that the persistence (or extinction) of tree populations in Europe is largely controlled by a deterministic ecological sorting process. Bhagwat and Willis (2008) showed that biogeographical traits (geographical distribution, habitat preference and life-history) of woody plants may have determined their ability to survive in inhospitable climates, so influencing their persistence (or extinction) in response to Pleistocene climate change. Eiserhardt et al. (2015) showed that late Cenozoic climate change induced phylogenetically selective regional extinction of northern temperate trees, leading to significant losses of phylogenetic diversity. Studying the relationships between climate oscillations and ecological traits, as well as the genetic characteristics of plants that went extinct in Europe during the Quaternary is especially important in terms of modern biodiversity conservation. This is a hot topic, as a better understanding of the ability of species to survive abrupt warming is indispensable in view of predicted global climate changes (Dawson et al., 2011). Linking studies of modern biodiversity and ecosystem patterns with paleoclimate dynamics may offer important scientific advances in both reconstructing the climatic conditions under which past extinction events happened, and inferring how present-day biodiversity patterns may be affected by climate change (Svenning et al., 2015).

Biostratigraphical and palaeoclimatic aspects have been often considered together, in connection to the question of whether Quaternary climate changes may have determined the extinction of plants in Southern Europe, in particular specific events including the Pliocene/Pleistocene transition (Popescu et al., 2010; Bertini, 2010; Jiménez-Moreno et al., 2010; Martinetto et al., 2015; Biltekin et al., 2015), the

128 Gelasian/Calabrian transition (Combourieu-Nebout, 1993; Bertini, 2010; Martinetto et al., 2015;
129 Combourieu-Nebout et al., 2015), the Middle Pleistocene revolution (Suc and Popescu, 2005; Leroy, 2007;
130 Postigo Mijarra et al., 2009, 2010; Bertini, 2010; Magri and Palombo, 2013), and specific glacial stages
131 corresponding to MIS 22, 16 and 12 (Muttoni et al., 2003; Tzedakis et al., 2006). Moreover, this question
132 implicitly involves the still open problem of whether plant extinctions were step-wise (e.g., Svenning, 2003;
133 Martinetto et al., 2015) or progressive (e.g., Bertini, 2010; Combourieu-Nebout et al., 2015, Biltekin et al.,
134 2015).

135 Other aspects related to the Pleistocene disappearance of tree taxa have not been adequately discussed yet,
136 and will be encompassed in the present paper. Starting from the compilation of schematic regional records
137 of past tree populations spanning the whole Quaternary, we intend to examine:

- 138 • value and limitations of pollen data to reconstruct the Pleistocene extinction of tree taxa,
- 139 • state of the art of the Early and Middle Pleistocene palaeofloristic data available from Southern
140 Europe,
- 141 • Quaternary history of tree taxa no longer present in Southern Europe mainland,
- 142 • timing of tree population reduction and eventual disappearance,
- 143 • geographical patterns of extinction,
- 144 • long-term persistence areas of tree populations in Southern Europe,
- 145 • ongoing reduction trends in Southern European tree taxa populations, in the light of timing and
146 trends of past tree population disappearance.

147

148 2. Recognizing plant extinctions from pollen data: value and limitations

149 Most knowledge of Quaternary floristic change comes from fossil pollen preserved in lacustrine and marine
150 sediments. In particular, continuous and detailed pollen records, stretching over multiple glacial-interglacial
151 cycles, are of incommensurable value to reveal the presence of species outside their present range, to

152 provide direct evidence of the persistence of plant populations in an area, to show the dynamics of species
153 turnover following climate change, and to document the disappearance of taxa. They are particularly
154 valuable when obtained from precisely dated deposits. Short records may also prove useful, especially when
155 they are chronologically well-constrained and fill gaps of knowledge for specific time intervals. The best
156 possible condition is the availability of pollen data from multiple dated sites that can substantiate the
157 interpretation of rare fossil occurrences ([Gavin et al., 2014](#)), which is a main problem when dealing with
158 extinctions.

159 In terms of limitations, one faces the classic problem “absence of evidence is not evidence of absence” – this
160 may particularly be an issue for palynological detection of rare taxa when total count sizes do not typically
161 exceed 300 grains and may be lower in the case of poor pollen concentration or recovery from geological
162 sections. On the other hand, reworking and long-distance transport may contribute to rare observations of a
163 particular pollen type such that its occurrence may not be indicative of contemporaneous regional presence.
164 Application of fluorescence approaches may be informative for discrimination of reworked grains and
165 demonstration of the occurrence of a taxon in an area (e.g. [Biltekin et al., 2015](#)). Long-distance transported
166 pollen is particularly troublesome when the aim is to detect the presence of reduced populations undergoing
167 extirpation, which by their nature are revealed by only rare pollen finds. For this reason, it may be very
168 difficult to assess the disappearance of trees belonging to the Pinaceae family (e.g. *Cedrus*; [Postigo-Mijarra
169 et al., 2010; Magri, 2012](#)) that are often huge producers of wind-transported pollen.

170 In the case of long-distance pollen occurrence, palynological records may post-date the time of disappearance
171 of a taxon from a region. For this reason, it is of great importance to consider the pollen productivity and
172 dispersal of taxa that are now absent from Southern Europe, but still present in neighbouring geographical
173 areas ([Filipova-Marinova et al., 2010](#)). In most cases, the last occurrences of pollen (and plant macrofossils)
174 tend to predate the extinction of plant populations, as it is presumed that the probability of finding pollen or
175 macroremains belonging to the very last individual existing in a region is virtually null. The error will be
176 smaller if long continuous records are available, and larger in case of fragmentary and discontinuous records.
177 Besides, under climatic conditions below the tolerance threshold of a species, plants may switch from

178 primarily sexual to vegetative reproduction and ultimately to a non-regenerative state, such that
179 disequilibrium populations on the verge of extirpation may be palynologically weakly visible or even invisible
180 ([Tallantire, 2002](#)). This may be relevant on centennial timescales for long-lived, stress tolerator species.
181 Therefore, it is important to be aware that the assessment of the exact timing of the extinction of plant
182 population is impossible.

183 The problem of determining the timing of plant extinctions has parallels with the difficulty of identifying
184 glacial refugia in areas far from the current distribution of species ([Gavin et al., 2014](#)), which is the subject of
185 vibrant and still unresolved discussions ([Stewart and Lister, 2001](#); [Tzedakis et al., 2013](#)). In many cases, rare
186 pollen finds distant from the modern species ranges and occurring much later than the last local population
187 expansion can only be considered as circumstantial and need to be confirmed and reinforced by the
188 accumulation of different sources of evidence, for example multiple records from the same region. Besides,
189 extinctions may be masked by taxonomic smoothing in the pollen record, which rarely permits species-level
190 differentiation ([Jackson and Weng, 1999](#)). For this reason, the role of macrofossils is especially important, as
191 they can often be identified with greater taxonomic precision than pollen, and they tend to be much more
192 locally distributed from their source ([Birks and Birks, 2000](#)). Besides, in some cases, macrofossil records
193 provide information on species whose pollen is badly preserved, for example *Laurus*, which we would expect
194 to be abundant in the Mediterranean pollen records ([Rodríguez-Sánchez et al., 2009](#)), but it is only seldom
195 found. However, the macrofossil record is generally very discontinuous in space and time, and the probability
196 of finding macrofossils of plants on the way to extinction is very low.

197 Specific problems affect the study of Pleistocene plant extinctions. One possible matter of confusion is that
198 the Gelasian Stage was included in the Pliocene until 2009, when the IUGS (International Union Geological
199 Sciences) ratified its transfer to the Quaternary System ([Gibbard et al., 2010](#)). This change requires a careful
200 consideration of the chronostratigraphic setting of pollen records. Going back in time to the Early and Middle
201 Pleistocene, the chronological control of pollen sequences may be highly inaccurate, as radiometric dating
202 and tephrochronology cannot always be applied, magnetostratigraphy may not be conclusive, astronomic
203 calibration is limited to sections with continuous sedimentation, direct correlation with oxygen isotope

204 stratigraphy is limited to marine sediments, and other biostratigraphic proxies studied in the same deposits
205 may suffer from the same chronostratigraphical uncertainties as palynology. In some cases, the margin of
206 error may be of hundreds of thousands of years, but even so the pollen evidence may be of great value, when
207 no other data are available.

208 When long continuous pollen records spanning multiple glacial-interglacial cycles are not available from a
209 region, the timing of plant extinctions is necessarily reconstructed through a compilation of different records,
210 thus accumulating chronological and taxonomical uncertainties. This difficulty is increased when the available
211 records come from both continental and marine environments, that are affected by different biases related
212 to pollen transportation, accumulation and preservation ([Hooghiemstra et al, 2006](#)).

213 Taxonomical uncertainties may be severe when dealing with extinct plants. It is likely that many extinct
214 species will never be recognized, as the identification of pollen at the species level is often impossible.
215 Furthermore, genus-level identification may be ambiguous. For example, *Cathaya* is often referred to as
216 *Cathaya* type, or included in *Pinus* haploxylon ([Liu and Basinger, 2000](#)). The distinction between *Taxodium*
217 and *Glyptostrobus* pollen is very difficult, so that it is often reported as *Taxodium* type, or Taxodiaceae.
218 However, many genera formerly included in the old Taxodiaceae family are now assigned to different
219 subfamilies of the Cupressaceae family (Taxodioideae: *Taxodium*, *Glyptostrobus*, and *Cryptomeria*;
220 Sequoioideae: *Sequoia*, *Sequoiadendron*, and *Metasequoia*) ([Farjon, 2005](#)). The correct assignment of
221 Taxodiaceae or *Taxodium*-type pollen to one or the other taxon is not possible unless the distinction is clearly
222 reported on the original paper ([Bertini, 2010](#)). Another common case is that of *Zelkova* pollen, which is often
223 counted together with *Ulmus* and grouped in an *Ulmus/Zelkova* type, so making it difficult to reconstruct the
224 history of this taxon, still included in the European flora.

225 A further complication, affecting several Mediterranean records of Early and Middle Pleistocene age, is that
226 the aim of the original studies was the reconstruction of palaeoenvironmental and palaeoclimatic
227 fluctuations rather than evaluation of plant population dynamics. For this reason, often only summary
228 cumulative diagrams of ecological groups and a few selected taxa are reported in the published pollen
229 records (e.g., [Combourieu-Nebout, 1993](#); [Müllenders et al. 1996](#); [Subally et al., 1999](#); [Sabato et al., 2005](#);

230 [Popescu et al., 2010](#); [Sadori et al., 2016](#)). In some cases, a list of the identified taxa is available ([Klotz et al.,](#)
231 [2006](#); [Degeai et al., 2013](#)), but much uncertainty remains as to the abundance and the age of the finds. It is
232 unfortunate that the full pollen records for these sites are not published and available to the scientific
233 community, after having required much work and effort.

234

235 3. South European records of Quaternary plant extirpations

236 We have reviewed a hundred Quaternary pollen records, showing the history of taxa that have disappeared
237 from Southern Europe mainland ([Fig. 1 and Table 1](#)). These records were dated by different methods,
238 including radiometric dates and tephrochronology, magnetotratigraphy, counting of annual laminated
239 sediments, astronomic calibration, correlation with paleontological records, or direct correlation with oxygen
240 isotope curves in the case of marine sediments ([Table 1](#)). In addition to many records spanning multiple
241 glacial-interglacial cycles, several well-dated shorter records conveniently fill gaps of knowledge for specific
242 time intervals (e.g., Colle Curti: [Bertini, 2000](#); Alleret Maar AL2: [Pastre et al., 2007](#); Cal Guardiola: [Postigo](#)
243 [Mijarra, 2007](#)). In same cases, also sequences that are not chronologically well constrained, but that fill large
244 gaps of information, have been considered, for example the records from Tres Pins ([Leroy, 2007](#)) and Saint
245 Macaire ([Leroy et al. 1994](#)), without which the information on the vegetation history in NE Spain and S France
246 during the late Early Pleistocene would be very limited.

247 We present here a number of regional overviews, necessarily limited to the most important sites, presented
248 in the form of a schematic compilation of selected pollen records from peri-Iberian marine cores ([Fig. 2](#)) and
249 Northeastern Spain-Southern France (<46°N) in [Fig. 2](#), Northern (>44°N; [Fig. 3](#)) and Central Italy (41°-44°N)
250 in [Fig. 3](#), Southern Italy (<41°N), Greece and the Aegean in [Fig. 4](#), and Tenaghi Philippon in Macedonia
251 ([Tzedakis et al., 2006](#)) and site DSDP 380 from the Black Sea ([Biltekin et al., 2015](#)) in [Fig. 5](#). This is at the
252 margin of Southern Europe, but may be considered an important bridge towards the the Anatolian-Caucasus-
253 Hyrcanian region that still hosts several species currently extinct from the European mainland. In some cases,
254 pollen data are complemented by macrofossil data.

255

Fig. 1. Location of the Quaternary records considered in this paper. Numbering of the sites follows Table 1. Red dots represent the peri-Iberian marine sites included in Fig. 2; blue dots correspond to the sites from NW Iberia and S France represented in Fig. 2; red squares include sites from N Italy (Fig. 3); yellow dots correspond to sites from central Italy of Fig. 3; pink dots are sites from S Italy represented in Fig. 4; green dots include Greece and the Aegean (Fig. 4); the blue square corresponds to the site of Tenaghi Philippon (Fig. 5) and the pink square to DSDP site 380 (Fig. 5). White dots indicate the location of sites not represented in Figs 2-5 but cited in the text.

Table 1. List of the vegetation records represented in Fig. 1.

Site number	Site	Age (ka) MIS	Pollen (P) Macrofossil (M)	Marine (M) Continental (C)	Chronological control	References
1	MD95-2043	48-2 ka	P	M	14C, event stratigraphy	Sanchez Goñi et al., 2002; Fletcher and Sanchez Goñi, 2008
2	Padul	100-0 ka	P	C	14C and vegetation history	Pons and Reille, 1988
3	ODP976	MIS 31-23	P	M	Oxygen isotope stratigraphy	Joannin 2007; Joannin et al., 2011; Combourieu Nebout et al., 2009
4	Andalucia G1	Miocene-Gelasian	P	M	Oxygen isotope stratigraphy	Feddi et al., 2011
5	Morgandinho	Early-Middle Pleistocene	P	C	Micromammals	Antunes et al., 1986
6	Site U1385	MIS 38	P	M	Orbital tuning	Tzedakis et al., 2015
7	MD95-2042	MIS 6-2	P	M	14C and Oxygen isotope stratigraphy	Sanchez Goñi et al., 1999
8	MD01-2443	345-180 ka	P	M	Oxygen isotope stratigraphy	Roucoux et al., 2006
9	Monsarros	Plio-Pleistocene	P	C	Stratigraphy	Vieira, 2009
10	MD03-2697	340-270 ka	P	M	Oxygen isotope stratigraphy	Desprat et al., 2009
11	MD01-2447	430-360 ; 250-180 ka	P	M	Oxygen isotope stratigraphy	Desprat et al., 2005
12	MD99-2331	MIS 5-2	P	M	14C, event stratigraphy	Sanchez-Goñi et al., 2005, 2008
13	Santa Maria de Oia	MIS 5-3	P	C-M (coastal)	14C, coastal stratigraphy	Gómez-Orellana et al., 2013
14	Area Longa	MIS 5-3	P	C-M (coastal)	14C, coastal stratigraphy	Gómez-Orellana et al., 2007
15	Pointe de la Négade North	Early Pleistocene	P+M	M	Pollen biostratigraphy	O'Brien and Jones, 2003
16	La Balise	MIS 11	P+M	M	Pollen biostratigraphy	O'Brien and Jones, 2003
17	Lezetxiki	MIS 3	P	C	14C, archaeological data (lithic tools)	Sánchez-Goñi, 1992
18	Atapuerca	900-400 ka	P	C	Luminiscence	García-Antón & Sainz-Ollero, 1991
19	Cañizar de Villaquemado	MIS 5-1	P	C	14C and OSL dating	González-Sampériz et al., 2013; Aranbarri et al., 2014; García-Prieto, 2015
20	Garraf	5300-1750 ka	P	M	Foraminifer biostratigraphy	Suc and Cravatte, 1982
21	Abric Romaní	70-40 ka	P	C	U/Th	Burjachs et al., 2012
22	Cal Guardiola	1200-800 ka	P-M	C	Micromammals and palaeomagnetism	Postigo Mijarra et al., 2007
23	Val d'en Bas	250 ka	P	C	Volcanic crust	Cross et al., 1986
24	Pla de l'Estany-Garrotxa	MIS 5?-1	P	C	Vegetation history and 14C for the upper part	Burjachs, 1990
25	Pla de l'Estany-Incarcal-Crespia	Plio-Pleistocene	M	C	Fauna (Mammals biostratigraphy)	Villalta and Vicente, 1972; Suc and Cravatte, 1982; Roiron, 1983
26	Tres Pins	2600-1200 ka	P	C	Vegetation composition (presence of carya)	Leroy, 1997
27	Mas Grill	Mid-	P	C	Vegetation history (older	Geurts, 1977; 1979

		Pleistocene?			than Bòbila Ordis)	
28	Bòbila-Ordis	MIS 36-33	P	C	Micromammals and palaeomagnetism	Løvlie and Leroy, 1995; Leroy, 2008
29	Banyoles	1000-950 ka	P	C		Julia Bruguès and Suc, 1980
30	Molí Vell	Early Pleistocene	P	C	Vegetation history (younger than Bòbila Ordis)	Geurts, 1977, 1979
31	Mas Miquel	Early Pleistocene	P	C	Fauna (Mammals biostratigraphy)	Leroy, 1990
32	Saint-Macaire	1400-700 ka	P	C	K/Ar dating, magnetostratigraphy	Leroy et al., 1994
33	Nogaret	MIS 75	P	C	K/Ar dating, mammal biostratigraphy	Brugal et al., 1990; Leroy and Seret, 1992
34	Bernasso	2160-1960 ka	P	C	K/Ar dating, magnetostratigraphy	Suc, 1978; Leroy and Roiron, 1996
35	Celleneuve	Early Calabrian	P	C-M	Biostratigraphy	Suc, 1973
36	Alleret	MIS 17-15	P	C	Tephrochronology, K/Ar and Ar/Ar dating	Pastre et al., 2007; Degeai et al., 2013
37	Senèze	MIS 85-76	P	C	Ar/Ar dating, mammal biostratigraphy, magnetostratigraphy	Elhaï, 1969; Roger et al., 2000
38	Velay maars	430-0 ka	P	C	Biostratigraphy, Ar/Ar dating	Reille et al., 2000
39	Ceyssac	1000-900 ka	P	C	K/Ar dating, magnetostratigraphy	Ablin, 1991
40	Saint-Vallier	1900 ka	P	C	Biostratigraphy	Argant, 2004
41	La Côte	MIS 11	P	C	Pollen biostratigraphy	Field et al., 2000
42	Fornaci di Ranica	MIS (36)30 or 28	P,M	C	Magnetostratigraphy, mammal biostratigraphy	Ravazzi et al., 2005
43	Leffe-Fornace Martinelli	MIS 64-22	P,M	C	Magnetostratigraphy, mammal biostratigraphy	Ravazzi and Rossignol-Strick, 1995; Ravazzi, 2003; Muttoni et al., 2007
44	Pianico Sèllere	MIS 21-19	P, M	C	K/Ar and Ar/Ar dating, mammal biostratigraphy	Ravazzi, 2003; Martinetto, 2009
45	Pianengo	1200-700 ka	P	C	Magnetostratigraphy	Muttoni et al., 2003
46	Stirone	5100-2200 ka	P	M	Mollusc and foraminifer biostratigraphy	Lona and Bertoldi, 1972; Bertini, 2001
47	Castell'Arquato	Early Pleistocene	P	M	Mollusc biostratigraphy	Lona and Bertoldi, 1972
48	San Pietro di Ragogna	Gelasian	P, M	C	Lithostratigraphy, biostratigraphy	Martinetto et al., 2012
49	Azzano Decimo	215-20 ka	P	C	14C dating, biostratigraphy	Pini et al., 2009
50	Visogliano	MIS 13 or 11	P	M	Mammal biostratigraphy	Cattani et al., 1991; Abbazzi et al., 2000
51	Venice	2150-110 ka	P	M	Magnetostratigraphy, molluscs biostratigraphy, sapropel stratigraphy	Mullenders et al., 1996; Massari et al., 2004
52	Core 223-S17	130-0 ka	P	C/M	Foraminifer and ostracod biostratigraphy	Amorosi et al., 2004
53	Oriolo (Faenza)	ca. 800 ka	M	C/M	Magnetostratigraphy, molluscs biostratigraphy	Martinetto and Sami, 2001
54	Lamone Valley	MIS 64-46	P	M	Mollusc and ostracod biostratigraphy	Fusco, 2007
55	Monte Poggiolo	1400-1070? ka	P	M	Magnetostratigraphy, molluscs biostratigraphy, ESR dating	Messenger et al., 2011
56	Poggio Rosso	1900-1800 ka	P	C	Magnetostratigraphy	Bertini et al., 2010

57	Gubbio	1600-1300? Ka	P	C	Biostratigraphy	Lona and Ricciardi, 1961
58	Colle Curti	ca. 990 ka	P	C	Magnetostratigraphy	Bertini, 2000
59	Cesi	750-600? ka	P	C	Magnetostratigraphy, Tephrochronology	Bertini, 2000
60	Pietrafitta	1400? ka	P	C	Mammal biostratigraphy	Ricciardi, 1961; Martinetto et al., 2014
61	Fosso Bianco	MIS 100-82	P	C	Magnetostratigraphy, Mammal biostratigraphy	Pontini and Bertini 2000
62	Dunarobba	3500-1500? ka	P, M	C	Biostratigraphy	Martinetto et al., 2014
63	Leonessa	1400-1200? ka	P	C	Biostratigraphy	Ricciardi, 1965
64	Madonna della Strada	ca. 1250 ka	P	C	Mammal biostratigraphy	Magri et al., 2010
65	Pagliare di Sassa	1100 ka	P	C	Magnetostratigraphy, mammal biostratigraphy	Palombo et al., 2010
66	Torre Mucchia (Ortona)	MIS 17	P	M	Magnetostratigraphy	Pieruccini et al., 2016
67	Carsoli	ca. 530	M	C	Ar/Ar dating	Sadori et al., 2010
68	Peperino (Rome)	513±3 ka	M	C	Ar/Ar dating	Follieri, 2010; Marra et al., 2014
69	Riano	450-350? ka	M	C	Mammal biostratigraphy	Follieri, 1958, 1962
70	Valle Ricca	2100-1800 ka	P	M	Magnetostratigraphy, biostratigraphy	Urban et al., 1983; Arias et al., 1990
71	Valle di Castiglione	250-0 ka	P	C	Varve counting, astronomic calibration	Follieri et al., 1988, 1989; Di Rita et al., 2013
72	Coste San Giacomo	ca. 2100 ka	P	C	Magnetostratigraphy, mammal biostratigraphy	Bellucci et al., 2014
73	Fontana Ranuccio	MIS 23	P	C	Magnetostratigraphy	Corrado and Magri, 2011
74	Ceprano	MIS 15-13	P	C	Magnetostratigraphy	Manzi et al., 2010
75	Isernia - La Pineta	900-700 ka	P	C	Magnetostratigraphy, Ar/Ar dating	Lebreton, 2002
76	Sessano	MIS 14-13	P	C	Magnetostratigraphy, Tephrochronology, Ar/Ar dating	Russo Ermolli et al., 2010a, 2015
77	Boiano	MIS 13-9	P	C	Ar/Ar dating, tephrochronology	Orain et al., 2013, 2015; Russo Ermolli et al., 2015
78	Saticula	ca. 1200 ka	P	C	Mammal biostratigraphy	Russo Ermolli et al., 2010b
79	Acerno	MIS 13-12	P	C	tephra, Ar/Ar dating	Munno et al., 2001; Russo Ermolli et al., 2015
80	Lago Grande di Monticchio	130±0	P	C	14C dating, tephrochronology, varve counting	Allen et al., 2000; Allen and Huntley, 2009
81	Vallo di Diano	650-450 ka	P	C	tephra, Ar/Ar dating	Russo Ermolli, 1994; Russo Ermolli et al., 2015
82	Camerota	1750-1600? ka	P	C	Biostratigraphy	Brenac, 1984
83	Mercure	MIS 13	P	C	Tephrochronology, Ar/Ar	Petrosino et al., 2014
84	San Lorenzo	1200-1100 ka	P	C	Magnetostratigraphy, mammal biostratigraphy	Sabato et al., 2005
85	Montalbano Jonico	1250-900 ka	P	M	Ar/Ar dating, Oxygen isotope stratigraphy	Joannin et al., 2008; Bertini et al., 2015
86	Semaforo-Vrica	2460-2110 ka	P	M	Magnetostratigraphy, foraminifer biostratigraphy, K/Ar	Combourieu-Nebout and Vergnaud-Grazzini, 1991;

					dating	Combourieu-Nebout, 1993; Klotz et al., 2006
87	Santa Lucia	MIS 43-40	P	M	Biostratigraphy, Oxygen isotope stratigraphy	Joannin et al., 2007b
88	Valle di Manche	MIS 20-18	P	M	Magnetostratigraphy, Oxygen isotope stratigraphy, Tephrochronology, Biostratigraphy, Ar/Ar dating	Capraro et al., 2005
89	Ohrid	500-0 ka	P	C	¹⁴ C, tephra, stratigraphy	Wagner et al., 2009, 2014; Sadori et al., 2016
90	Ioannina	450-0 ka	P	C	Pollen biostratigraphy	Tzedakis, 1994; Roucoux et al., 2008
91	Acheloos Delta	5-3 ka	P	M	¹⁴ C dating	Fouache et al., 2005
92	Zakynthos	2000-1800 ka	P	M	Magnetostratigraphy, oxygen isotope stratigraphy	Subally et al., 1999
93	Gulf of Corinth	1800-600 ka	P	M	Pollen biostratigraphy, regional stratigraphical correlations	Rohais et al., 2007
94	Kopais	500-2 ka	P	C	Pollen biostratigraphy	Tzedakis, 1999; Okuda et al., 2001
95	Megalopolis Basin	650-300 ka	P	C	Magnetostratigraphy, palaeontology	Okuda et al., 2002
96	Kallithea	Calabrian	M	C	Mollusc biostratigraphy	Mai and Velitzelos, 2007; Velitzelos et al., 2014
97	Kolymia Bay	500 ka	M	M	Magnetostratigraphy	Boyd, 2009
98	Tsampika Bay	MIS 42-1	P	M	Magnetostratigraphy, Oxygen isotope stratigraphy	Joannin et al., 2007a
99	Tenaghi Philippon	1350-0 ka	P	C	Pollen-orbital tuning	Wijmstra, 1969; Wijmstra and Smit, 1976; Van der Wiel and Wijmstra, 1987a,b; Tzedakis et al., 2006; Fletcher et al., 2013; Pross et al., 2015
100	DSDP Site 380	6000-0 ka	P	M	Nannoplankton	Biltekin, 2010; Popescu et al., 2010; Biltekin et al., 2015

268

269

270 3.1 Iberian Peninsula

271 The Iberian Peninsula has always been a region of great physiographical heterogeneity over time, resulting
272 in a unique evolutionary history and particular vegetation behaviour and composition (Carrión and Leroy,
273 2010). This fact has been considered one of the main reasons for survival and/or delayed extinction of many
274 taxa in this region of southern Europe (Postigo-Mijarra et al., 2010). There are many Quaternary sites, both
275 continental and marine sequences (Fig. 2), which provide information from Iberia, although those from Early
276 and Middle Pleistocene are relatively scarce and no long continuous records such as those from France,
277 Greece or Italy are available (see compilations of González-Sampériz et al., 2010 and Postigo-Mijarra et al.,
278 2010).

280 **Fig. 2.** Stratigraphical setting and schematic pollen biostratigraphy of selected taxa of Pleistocene records
281 from the Alboran Sea and off Portugal, and from NW Iberia and S France. Question marks indicate
282 uncertainties in the chronological setting of the records. Asterisks indicate plant macrofossils. Red lines
283 represent regionally extinct taxa. Blue lines represent taxa that are only mentioned in the papers or uncertain
284 determinations (e.g. *Ulmus/Zelkova*).

285 During the Early Pleistocene (ca. 2.6-0.8 Ma, MIS 100-20), the majority of available data come from
 286 Northeastern Spain (Fig. 2), thus presenting a partial image of the whole of Iberia. These include Garraf, Pla
 287 de l'Estany-Incarcal-Crespià (the three names are sometimes interchanged in the literature because the
 288 different outcrops are located in the same site), Mas Miquel, Tres Pins, Bòbila Ordis, Molí Vell and Cal
 289 Guardiola, while ODP 976 and G1 are the only marine cores which cover this period. Monsarros, in Portugal,
 290 is also considered a Plio-Pleistocene site (see location, more details of the different sequences and
 291 compilation of references in both González-Sampériz et al., 2010 and Carrión (Ed.), 2012). In relation to
 292 extinct taxa considered in this work, Garraf (Suc and Cravatte, 1982; Suc and Popescu, 2005) records
 293 Taxodiaceae, *Liquidambar*, *Cathaya* and *Eucommia*; Pla de l'Estany-Incarcal-Crespià shows *Carya*, *Pterocarya*,
 294 *Zelkova*, *Parrotia* and *Carpinus* leaf imprints (Villalta and Vicente, 1972) and pollen presence of both *Carpinus*
 295 and *Parrotiopsis* (Suc and Cravatte, 1982); Mas Miquel pollen data (Geurts, 1979; Leroy, 1990) includes
 296 *Carpinus*, *Liquidambar*, *Carya*, *Pterocarya*, *Nyssa* and even *Acacia* type between other meso-thermophytes;
 297 Tres Pins captures *Cathaya*, *Cedrus*, *Picea*, *Tsuga*, *Carpinus*, *Eucommia*, *Liquidambar*, *Carya*, *Pterocarya*,
 298 *Parrotia* and *Zelkova* (Leroy 1990, 1997), while in the site of Bòbila Ordis, dated at around 1.2 Ma (MIS 36-
 299 33) (Lovlie and Leroy, 1995), *Engelhardtia*, *Eucommia*, *Carpinus*, *Carya*, *Pterocarya*, *Tsuga*, *Zelkova* and
 300 *Parrotia* occur (Leroy 1990, 2008) (Fig. 2). Both Molí Vell (Geurts 1977, 1979) and Cal Guardiola (1.2-0.8 Ma:
 301 Postigo-Mijarra et al. 2007) show *Carya* and *Carpinus* as well as Taxodiaceae in the case of Cal Guardiola,
 302 pointing the last evidence in Iberia and one of the last in Europe for this taxon (Postigo-Mijarra et al., 2010).
 303 Likewise, the Portuguese site of Monsarros (Vieira, 2009) documents the presence of *Liquidambar*,
 304 Taxodiaceae, *Engelhardia*, *Symplocos*, *Nyssa*, and *Cathaya*, while G1 core, in the Alboran Sea, shows evidence
 305 of *Picea*, *Tsuga*, *Cathaya*, Taxodiaceae, *Carpinus*, *Engelhdaria* and *Liquidambar* (Feddi et al., 2011) and ODP
 306 976 *Liquidambar*, *Zelkova*, *Tsuga*, *Pterocarya*, and *Carpinus* (Joannin et al., 2011).
 307 In summary, a complex scenario emerges from the patchy, discontinuous geographical distribution of sites
 308 across Iberia. Long temporal series are generally lacking and vegetation signals are usually dominated by
 309 conifers and/or the herbaceous component. A varied presence of other relict taxa is nevertheless apparent
 310 (González-Sampériz et al., 2010; Postigo-Mijara et al., 2010), at least near the Mediterranean and until the
 311 Early-Middle Pleistocene transition (between MIS 36-18). After this transition, the last occurrences of some

312 taxa which concern the topic of this work such as *Engelhardia*, *Eucommia*, *Liquidambar*, *Sciadopitys*, *Parrotia*,
313 *Taxodium* type and *Tsuga*, can be detected (Leroy, 2008; Postigo-Mijarra et al., 2009, 2010).

314 With respect to the Middle Pleistocene (ca. 0.8-0.126 Ma, MIS 20-6), following the review of Postigo-Mijarra
315 et al. (2010), there is a noticeable increase in information related to last recognised evidence of relicts, but
316 the number of records is similar to the Early Pleistocene and again most of the sites are only fragmentary
317 sequences, with a low number of samples studied and poor chronological control (for more details see table
318 compilation in González-Sampériz et al. (2010) and individual sites description in Carrión (Ed.) (2012).
319 Nevertheless, valuable information emerges from both marine and continental sites, including some
320 archaeological sites and again not only pollen but macro-remains, too.

321 Thus, *Carpinus* is recorded in marine cores ODP 976 (Combourieu-Nebout et al., 1999; Joannin, 2007, 2011),
322 MD99-2331 (Sánchez-Goñi et al., 2005, 2008), MD01-2447 (Desprat et al., 2005, 2007), MD03-2697 (Desprat
323 et al., 2009) and MD01-2443 (Roucoux et al., 2006). *Pterocarya* occurs in both MD01-2447 during MIS 11
324 (Desprat et al., 2005) and MD03-2697 during MIS 9 (at around 298 ka BP: Desprat et al., 2009). However, a
325 greater diversity of extinct taxa can be observed in continental areas sites (Fig. 2). This is the case of *Carya*,
326 for example, which has been recorded after MIS 18 in the Banyoles area but imprecisely dated (Geurts, 1979).
327 Val d'en Bas includes *Carpinus* in the spectra located below a volcanic crust dated at 250 ka ago (Cross et al.,
328 1986). Mas Grill shows the presence of *Fagus*, *Ostrya*, *Juglans* and *Carya* (Geurts, 1979), although only five
329 pollen samples have been studied. *Carpinus* is also recorded at the Northern Iberian Lezetxiki site, attributed
330 to the end of the Middle Pleistocene following the archaeological sequence (Sánchez-Goñi, 1992). The
331 Atapuerca pollen sequence, from both Galería (TG) and Gran Dolina (TD) outcrops (García-Antón and Sainz-
332 Ollero, 1991), records the last evidence of *Cedrus*, *Taxodium*, *Platanus* and *Carya* from inner continental areas
333 at around 0.8-0.6 Ma (TD 5-TD 8) while *Carpinus*, *Picea* or *Juglans* still persist in the sequence and other sites
334 until Holocene or current times in same cases. In Padul, the longest pollen sequence from southern Spain,
335 *Abies*, *Tsuga*, *Fagus* and *Quercus*, amongst other taxa, are recorded in levels probably attributed to the
336 interglacial Mindel-Riss (Holstein) despite controversy between Florschütz et al. (1971) and Pons and Reille
337 (1988) with respect to the chronology of the sequence. All these sites are located in Spain, but data from
338 Portugal is also available such as in Morgandinho (Antunes et al., 1986), where *Cathaya* and *Keteleeria* were

339 present, as well as *Picea* (ca. 3%), in a conifer-dominated landscape (mainly formed by *Pinus*) between,
340 probably, MIS 20-17.

341 Thus, the Middle Pleistocene Iberian sites offer an informative but always incomplete view of the extinction
342 and/or persistence of relicts pointing the key role of southern European latitudes for vegetation refugia in
343 the face of intense global climatic changes such as those associated to glacial/interglacial periods and the
344 consequent adaptation, survival or disappearance of numerous taxa.

345 Finally, Iberian available sites from Late Pleistocene ([González-Sampériz et al., 2010](#)) and Holocene ([Carrión](#)
346 [et al., 2010](#)) are much more numerous than previous periods, geographically well distributed and benefit
347 from generally better chronological control, but information with respect to extinct taxa becomes limited
348 ([Postigo-Mijarra et al., 2008, 2010](#)) because *Engelhardia*, *Eucommia*, *Liquidambar*, *Sciadopitys*, *Parrotia*,
349 *Taxodium* type and *Tsuga*, amongst other relicts already disappeared.

350 Concerning currently extinct conifers in Iberia, *Cedrus* is relatively frequently present in continental
351 palynological records (marine cores from southern Iberia cannot be considered here for its potential African
352 pollen rain capture area; [Fig. 2](#)), both from the Late Pleistocene ([Pons and Reille, 1988](#); [Burjachs and Julià,](#)
353 [1994](#); [Carrión and van Geel, 1999](#); [Yll et al., 2003](#); [Carrión et al., 2008](#); [Cortés-Sánchez et al., 2008](#)) and
354 Holocene ([Pons and Reille, 1988](#); [Carrión and van Geel, 1999](#); [Carrión et al., 2003, 2007](#); [Cortés-Sánchez et](#)
355 [al., 2008](#); [Aranbarri et al., 2014](#)).

356 New Iberian long sequences which extend from, at least, the Middle-Late Pleistocene transition and include
357 the whole Holocene, such as the Cañizar de Villarquemado record ([González-Sampériz et al., 2013](#); [García-](#)
358 [Prieto, 2015](#)), Conquezueta palaeolake ([González-Sampériz, pers. com.](#)), or a new study at the well-known
359 Padul peatbog ([Jiménez-Moreno, pers. comm.](#)), are currently under study and will be published in the next
360 years. Current methodologies usually imply a multidisciplinary context of analyses and an accurate
361 independent chronology. Thus, these future results will be of great interest and could contribute to the
362 understanding of the survival history of rare taxa in the Mediterranean context.

363

364

365 3.2 Southern France

366 Only pollen records at latitudes <46°N have been considered from Southern France, where the paleofloristic
367 record of the Early Pleistocene is rather fragmented (Fig. 2). The pollen record from Senèze, dated to around
368 2.1 Ma (Roger et al., 2000), lists many “exotic trees”, including *Taxodium/Sequoia*, *Sciadopitys*, *Pinus*
369 *haploxylon*, *Nyssa*, *Liquidambar*, *Eleagnus*, *Symplocos*, *Carya*, *Pterocarya*, *Juglans*, *Eucommia*, *Cedrus*, and
370 *Tsuga* (Elhaï, 1969). The site of Bernasso, which provided both pollen and plant macrofossil data, developed
371 during the time interval between ca. 2.16 and 1.96 Ma and documents the presence of a rich flora with *Cedrus*,
372 *Tsuga*, *Eucommia*, *Parrotia*, *Carya*, *Pterocarya*, *Engelhardia*, *Myrica*, *Zelkova*, *Carpinus orientalis*, and
373 *Parthenocissus* (Suc, 1978; Leroy and Roiron, 1996). The Nogaret deposit, dated between 2 Ma and 1.7 Ma
374 shows a substantially similar flora, including also scarce grains of *Cathaya* (Brugal et al., 1990; Leroy and
375 Seret, 1992). The Saint-Macaire maar (NW Mediterranean coastal plain of France), considered to be
376 deposited between 1.4 and about 0.7 Ma, presents many floristic elements typical of the Early Pleistocene,
377 including *Cathaya*, *Cedrus*, *Tsuga*, *Liquidambar*, *Carya*, *Pterocarya*, *Eucommia*, *Parthenocissus*, and *Ulmus*-
378 *Zelkova* (Leroy et al., 1994). The long pollen record from Ceysac depicts the vegetation history of the Massif
379 Central from the beginning of the Pleistocene to approx. 0.9 Ma. The bottom part of the record, substantially
380 confirming the presence of the taxa found at Senèze, Bernasso, Nogaret and Saint-Macaire, is not precisely
381 dated, while pollen zone 8 is bracketed between two K/Ar dates suggesting an age between 1.1 and 0.9 Ma
382 (Ablin, 1991). It still includes a number of taxa that are not currently present in the region, including *Cathaya*
383 type, *Cedrus*, *Tsuga*, *Parrotia*, *Eucommia*, *Carya*, *Pterocarya*, *Parthenocissus*, and *Zelkova*. Other pollen
384 records of Early Pleistocene age were studied from the French Pays Basques (Oldfield, 1968) and from the
385 Médoc region (O’Brien and Jones, 2003). Although they are not set in a precise chronostratigraphic
386 framework, they provide the important information of significant expansions of *Tsuga*, reaching the Atlantic
387 coast in a time period preceding 0.8 Ma.

388 The vegetation history of the early Middle Pleistocene is well studied in the Alleret maar (Massif Central),
389 which comprises two records: the oldest is dated 750 to 650 ka and is still only partially published (Degeai et
390 al., 2013), while the youngest is correlated with MIS 15 (Pastre et al., 2007). The flora appears impoverished
391 with respect to the Early Pleistocene, as taxa such as *Cedrus*, *Eucommia*, *Parrotia*, and *Parthenocissus* are not

392 present. *Carya* and *Liquidambar* are still recorded during MIS 17 (Degeai et al., 2013), but were absent in MIS
 393 15 (Pastre et al., 2007). *Cathaya*, *Tsuga*, *Nyssa*, *Pterocarya* and *Zelkova* were still present in both records.
 394 The records from the Velay (Reille et al., 2000 and references therein) constitute a benchmark for the
 395 vegetation history of southern France across five glacial-interglacial cycles (MIS 11 to the present). *Cathaya*,
 396 *Nyssa*, *Tsuga* and *Zelkova* are not found any more, while *Pterocarya* is recorded for the last time in the
 397 Praclaux interglacial, correlated with MIS 11. The same age is prospected for the finds of *Pterocarya* in the
 398 interglacial deposits from La Côte, in the western French Alps (Field et al., 2000), but a personal
 399 communication of M. Reille reveals that traces of *Pterocarya* were found in the Massif Central in MIS 9
 400 (Tzedakis et al., 2006).

401

402 3.3 Italian Peninsula

403 In the Italian Peninsula, a large number of Pleistocene pollen records have been published over the years,
 404 documenting the presence of taxa nowadays absent from the Italian territory. Several long pollen records,
 405 often spanning multiple glacial-interglacial cycles, have been studied from lacustrine sites, such as Azzano
 406 Decimo (Pini et al., 2009), Leffe (Ravazzi and Rossignol-Strick, 1995; Ravazzi, 2003; Muttoni et al., 2007),
 407 Pianengo (Muttoni et al., 2003), Fornaci di Ranica (Ravazzi et al., 2005), Fosso Bianco (Pontini and Bertini,
 408 2000), Valle di Castiglione (Follieri et al., 1988, 1989), Vallo di Diano (Russo Ermolli, 1994), Acerno (Munno et
 409 al., 2001; Russo Ermolli et al., 2015), Boiano (Orain et al., 2013), Sessano (Russo Ermolli et al., 2010a), and
 410 Camerota (Brenac, 1984). Other long pollen records were obtained from marine sites, including Venice
 411 (Müllenders et al., 1996, Massari et al., 2004), Stirone (Lona and Bertoldi, 1972; Bertini 2001), Lamone Valley
 412 (Fusco, 2007), Valle Ricca, (Urban et al., 1983; Arias et al., 1990), Santa Lucia (Joannin et al., 2007b)
 413 Montalbano Jonico (Joannin et al., 2008; Bertini et al., 2015), and Crotone (Coubourieu Nebout and Vergnaud
 414 Grazzini, 1991; Coubourieu-Nebout, 1993; Klotz et al., 2006; Suc et al., 2010). In addition to these long
 415 sequences, many pollen records span only single glacial-interglacial cycles or shorter time intervals.
 416 Nonetheless, they provide useful information about the presence/absence of plant taxa, especially when
 417 they are set into a sound chronostratigraphic framework. Among them, it is worth mentioning Pianico Sèllere
 418 (Ravazzi et al., 2003; Martinetto, 2009); Valle di Manche (Capraro et al., 2005), Colle Curti (Bertini, 2000),

419 Fontana Ranuccio (Corrado and Magri, 2011), Mercure (Petrosino et al., 2014), and Lago Grande di
420 Monticchio (Allen et al., 2000; Allen and Huntley, 2009). A number of pollen records have been studied from
421 paleontological sites, with mammal fauna providing additional palaeoecological information. They include
422 Coste San Giacomo (Bellucci et al., 2014), Madonna della Strada (Magri et al., 2010), Poggio Rosso (Bertini et
423 al., 2010), Pagliare di Sassa (Palombo et al., 2010), Riano (Follieri, 1958, 1962), Pietrafitta (Ricciardi, 1961;
424 Martinetto et al., 2014), Saticula (Russo Ermolli et al., 2010), and San Lorenzo in the Sant'Arcangelo Basin
425 (Sabato et al., 2005). A few pollen records were obtained from archaeological sites: Visogliano (Cattani et al.,
426 1991; Abbazzi et al., 2000), Isernia-La Pineta (Lebreton et al., 2002), Monte Poggiolo (Messenger et al., 2011),
427 and Ceprano (Manzi et al., 2010). A large number of macrofossil sites complement the information provided
428 by pollen records. They are mostly located in northern and central Italy (Martinetto 2001, 2009, 2015,
429 Martinetto and Sami, 2001, Martinetto et al., 2014, 2015).

430 While no single pollen record in Italy spans the whole (or most) of the Quaternary, the wealth of available
431 data has made it possible to compose a satisfactory picture of the succession of tree taxa disappearances
432 from the Peninsula, which is presented separately for Northern (>44°N; Fig. 3) and Central Italy (44°-41°N) in
433 Fig. 3, and Southern Italy (<41°N) in Fig. 4). These regions do not correspond exactly to the administrative
434 subdivision of Italy, but are appropriate for our purposes.

435 In Northern Italy (>44°N; Fig. 3), the base of the Quaternary is well represented, in continuity with the end of
436 the Pliocene, in the Stirone section, recording the altitudinal forest development in the Northern Apennines
437 (Italy) from 2.7 Ma to approx. 2.2 Ma (Bertini, 2001). In the Pleistocene portion of the record, a number of
438 tree taxa are found that later disappear, including *Taxodium* type, *Sequoia* type, *Sciadopitys*, *Cathaya*, *Tsuga*,
439 *Cedrus*, *Liquidambar*, *Engelhardia*, *Carya*, *Pterocarya*, *Zelkova*, and *Parthenocissus*. These taxa were still
440 present, and in some cases abundant, in the marine sediment of Calabrian age from the Lamone Valley,
441 where also *Nyssa* and cf. *Parrotia* were found (Fusco, 2007). A clear reduction in the recorded taxa appear in
442 the pre-Jaramillo record from Lefte-Fornace Martinelli (Ravazzi and Rossignol-Strick, 1995; Ravazzi, 2003;
443 Muttoni et al., 2007), where *Sciadopitys*, *Cathaya*, *Parrotia*, and *Engelhardia* are not reported, but other most
444 interesting taxa are represented, including *Aesculus* (Ravazzi, 1994), *Styrax*, and *Parthenocissus*. It is
445 important to note that *Taxodiaceae*, still abundant in the Lamone Valley, are only found at the base of the

446 Leffe-Fornace Martinelli record. In the early Middle Pleistocene, a further reduction of tree taxa is observed
 447 in the section of Pianico Sèllere (Ravazzi, 2003; Martinetto, 2009), whose fossil flora does not display taxa
 448 typical of the Early Pleistocene (e.g. Taxodiaceae, *Carya*, *Liquidambar*, and *Eucommia*), but shows the last
 449 sporadic appearance of *Cedrus*, which was abundant at Leffe. *Parrotia* and *Zelkova* are still found in the
 450 macroflora of Oriolo (Martinetto and Sami, 2001), but never again in the mid-Middle Pleistocene site of
 451 Visogliano (350–500 ka), where also *Carya* and *Tsuga* were lost, but *Pterocarya* was still present (Cattani et
 452 al., 1991; Falguères et al., 2008). The Azzano Decimo record (Pini et al., 2009), spanning the last 215 ka, does
 453 not display any tree taxon currently absent from the peninsula. However, pollen records from the Eastern Po
 454 Plain (Amorosi et al., 2004) and from Venice (Müllenders et al., 1996) document the presence of *Zelkova* until
 455 the end of the Eemian interglacial.

456
 457 **Fig. 3.** Stratigraphical setting and schematic pollen biostratigraphy of selected taxa of Pleistocene records
 458 from N Italy (>44°N) and Central Italy (44°-41°N). Note that the sites of Sessano and Boiano are
 459 administratively located in S Italy. Question marks indicate uncertainties in the chronological setting of the
 460 records. Asterisks indicate plant macrofossils. Red lines represent regionally extinct taxa. Blue lines represent
 461 taxa that are only mentioned in the papers or uncertain determinations (e.g. *Ulmus/Zelkova*).

Fig. 4. Stratigraphical setting and schematic pollen biostratigraphy of selected taxa of Pleistocene records from S Italy (<41°N), Greece and the Aegean. Question marks indicate uncertainties in the chronological setting of the records. Asterisks indicate plant macrofossils. Red lines represent regionally extinct taxa. Blue lines represent taxa that are only mentioned in the papers or uncertain determinations (e.g. *Ulmus/Zelkova*).

In Central Italy (44°-41°N; Fig. 3), *Taxodium* type is abundant in all the Gelasian records (Fosso Bianco: Pontini and Bertini, 2000; Poggio Rosso: Bertini et al., 2010; Coste San Giacomo: Bellucci et al., 2014). However, at Fosso Bianco there are still many Pliocene taxa that are not found in the Quaternary records of Central Italy, including *Panax* type (Araliaceae), *Engelhardia*, *Nyssa*, *Symplocos*, and cf. *Parthenocissus*. It is also worth noting the last occurrence of *Aesculus* in Central Italy, recorded at Poggio Rosso. During the Calabrian, a number of sites that are not precisely dated (Pietrafitta: Ricciardi, 1961; Martinetto et al., 2014; Gubbio: Lona and Ricciardi, 1961; Leonessa: Ricciardi, 1965; Madonna della Strada: Magri et al., 2010; and Saticula: Russo Ermolli et al., 2010b), together with other records dated by means of magnetostратigraphy (Colle Curti: Bertini, 2000; Fontana Ranuccio: Corrado and Magri, 2011) show the last occurrences of *Sciadopitys* (Pietrafitta), *Parrotia* (Madonna della Strada), *Liquidambar* (Colle Curti), *Cathaya* type (Fontana Ranuccio) and the last continuous record of *Cedrus* (Fontana Ranuccio). Sparse pollen grains of *Taxodium* type are still

479 found in the Calabrian records. In the early Middle Pleistocene, the Torre Mucchia record, correlated with
 480 MIS 17 (Pieruccini et al., 2016), displays the last significant abundance (>3%) of *Tsuga* and *Carya*, which are
 481 recorded for the last time during interglacial periods corresponding to MIS 13 (Sessano: Russo Ermolli et al.,
 482 2010a, 2015) and MIS 9 (Boiano: Orain et al., 2013; Russo Ermolli et al., 2015), respectively. The last
 483 occurrence of *Pterocarya*, still very abundant in both the pollen and macrofossil records from Riano (Follieri,
 484 1958, 1962), likely correlated to MIS 11, is recorded at Valle di Castiglione around 200 ka (Follieri et al., 1988).
 485 *Zelkova*, a main component of the Eemian vegetation at Valle di Castiglione (Follieri et al., 1986, 1988, 1989)
 486 disappears from several records in Central Italy at the end of MIS 3, while *Picea* is found with percentages
 487 >2% until about 25 ka (Follieri et al., 1998; Magri, 1999; Magri and Sadori, 1999).

488 In Southern Italy (<41°N; Fig. 4), all the records of Early Pleistocene age were studied from marine sediments,
 489 with the exception of Camerota (Brenac, 1984). The long Crotone sequence, including the Semaforo (2.46 to
 490 2.11 Ma) and Vrica (2.2 to 1.36 Ma) sections (Combourieu-Nebout and Vergnaud-Grazzini, 1991;
 491 Combourieu-Nebout, 1993; Klotz et al., 2006; Suc et al., 2010), provides a well-dated and detailed record of
 492 the Gelasian vegetation, including a large number of taxa nowadays absent from Italy, belonging to the
 493 subtropical humid forest: *Cathaya*, *Distylium*, *Sciadopitys*, *Taxodium*, *Engelhardia*, *Myrica*, *Magnolia*, *Nyssa*,
 494 *Symplocos*, and *Sapotaceae*. Besides, there were a number of taxa typical of the deciduous forest
 495 (*Liquidambar*, *Parrotia*, *Carya*, *Pterocarya*, *Eucommia*) and of the altitudinal coniferous belt (*Cedrus*, *Picea*,
 496 *Tsuga*), that are not living any more in Southern Italy. Some of these taxa were not retrieved in the pollen
 497 record from Camerota, dated to the early Calabrian (Brenac, 1984), namely *Magnolia*, *Nyssa*, *Eucommia*, and
 498 *Symplocos*. *Parthenocissus* was found in one level. Other taxa were lost in the Santa Lucia section, correlated
 499 with MIS 43-40: *Engelhardia*, *Sciadopitys*, and *Parrotia* (Joannin et al., 2007b). The floristic list is substantially
 500 the same at Montalbano Jonico during MIS 37 to 23, with *Liquidambar*, *Tsuga*, and *Carya* showing a
 501 progressive decrease (Joannin et al., 2008). In the same site, during MIS 21 to 19 (858 to 745 ka), *Liquidambar*
 502 and *Taxodium* type are represented by sporadic grains, while *Cathaya* is virtually absent (Bertini et al., 2015).
 503 In the coeval site of Valle di Manche (Capraro et al., 2005), *Cathaya* is documented for the last time in
 504 Southern Italy. During the Middle Pleistocene, the long pollen record from Vallo di Diano depicts the
 505 vegetation changes during MIS 16 to 13 (Russo Ermolli, 1994; Russo Ermolli et al., 2015). Only single grains

506 of *Taxodium* type, *Tsuga*, *Pterocarya*, and *Liquidambar* were found; *Cedrus*, *Picea*, and *Carya* were strongly
507 reduced but still present, especially during the interglacial phase corresponding to MIS 13. During the same
508 forest phase, *Zelkova* was rather abundant, reaching percentages >5%. All these taxa disappeared from
509 Southern Italy before the Late Pleistocene, except *Picea* that persisted until the last glacial maximum, as
510 documented by the pollen record from Lago Grande di Monticchio (Allen et al., 2000; Allen and Huntley,
511 2009).

512

513 3.4 Balkan Peninsula and Anatolia

514 In the Balkans, several lines of information contribute to the understanding of regional Quaternary tree taxa
515 persistence and extinctions (Figs 4-5). Pollen records on long core sequences from subsiding terrestrial basins
516 provide continuous records spanning multiple glacial-interglacial cycles, such as Lake Ohrid (Wagner et al.,
517 2009; Sadori et al., 2016), Lake Ioannina (Tzedakis, 1994), Tenaghi Philippon (Tzedakis, 2006; Pross et al.,
518 2015), and Kopais (Tzedakis, 1999, Okuda et al., 2001). Exposed geological sections, such as in the
519 Megalopolis Basin (Okuda et al., 2002) and on Rhodes (Joannin et al., 2007a) provide discontinuous
520 sequences that can be integrated into the wider framework through radiometric dating, palaeomagnetic and
521 isotope stratigraphy, and correlations with astronomical cycles (Fig. 4). Pollen records from continuous
522 marine sediment sequences also represent an important contribution, notably a 6 Ma sequence from DSDP
523 Site 380 in the southwestern Black Sea (Biltekin, 2010; Popescu et al., 2010; Biltekin et al., 2015).
524 Complementary information is also available from plant macrofossil locales, principally in Southwestern
525 Greece and in the Hellenic Arc region (reviewed in Velitzelos et al., 2014).

526 Long-term patterns of persistence and extinction are revealed in the long, continuous pollen records. At
527 Tenaghi Philippon (Wijmstra, 1969; Wijmstra and Smit, 1976; Van der Wiel and Wijmstra, 1987a, 1987b;
528 Tzedakis et al., 2006; Fletcher et al., 2013; Pross et al., 2015), the key patterns of extinctions and their
529 relationship to glacial-interglacial cycles of the last 1.3 Ma is discussed by Tzedakis et al. (2006). The record
530 reveals a progressive broad-scale floristic shift in forest composition, with Early Pleistocene spectra rich in
531 *Castanea*, *Pterocarya*, *Carya* and *Cedrus* replaced by assemblages with *Carpinus*, *Ostrya* and *Corylus*
532 alongside increased *Quercus* (Fig. 5). Within this transition, which occurred between 0.9 and 0.6 Ma (and is

533 associated with the first major glaciations of MIS 22 and MIS 16) a series of individual taxa disappearances is
 534 recorded, with last occurrences of *Liquidambar* and *Parthenocissus* prior to MIS 22, and subsequent
 535 sequential disappearance of *Parrotia*, *Eucommia*, *Tsuga*, *Pterocarya*, *Carya* and *Cedrus* (Tzedakis et al., 2006).
 536 At DSDP site 380 (Popescu et al., 2010; Biltekin et al., 2015) two taxa, *Avicennia* and *Nyssa*, do not appear to
 537 persist into the Middle Pleistocene at all (Fig. 5). These taxa are documented by rare occurrences in the Early
 538 Pleistocene, and they appear more broadly associated with taxa whose last appearances in this sequence are
 539 documented in the late Miocene and Pliocene (Arecaceae, *Parrotia*). Several taxa are consistently recorded
 540 during the early Pleistocene, prior to around 1.2 Ma, specifically *Engelhardia*, *Zelkova*, *Pterocarya* and *Tsuga*.
 541 These taxa all occur more sporadically in the middle Pleistocene sediments, and are generally absent in the
 542 upper part of the sequence, corresponding to the late Middle Pleistocene and Late Pleistocene. *Cathaya*
 543 displays a similar pattern of sporadic occurrences in the early Middle Pleistocene, although the main period
 544 of consistent presence was earlier, during the Pliocene. A small group of taxa (Taxodiaceae, *Carya* and *Cedrus*)
 545 are documented sporadically through the later Middle Pleistocene and Late Pleistocene.
 546 At other sites with pollen records extending into the Middle Pleistocene, occasional occurrences of extinct
 547 or relictual taxa are reported. The presence of *Pterocarya* in mainland Greece during MIS 11 and MIS 9 is
 548 confirmed in the pollen record of Kopais shown by Okuda et al., 2001. In the Ioannina-249 core (Tzedakis,
 549 1994), *Zelkova* is reported in interglacials prior to the last glacial.
 550 Important insights are also derived from the shorter, or discontinuous records. Subally et al. (1999) report
 551 on the Coupe Citadelle, Zakynthos (2.0-1.8 Ma), where oscillations between *Artemisia* and *Cedrus* dominated
 552 assemblages are associated with interglacial/glacial cycles. Alongside *Cedrus*, *Cathaya* and Taxodiaceae are
 553 well represented. Joannin et al. (2007a) report findings from Early to Middle Pleistocene deposits in Rhodes
 554 (Tsampika formation). Pollen diagrams for sequence TS1-3 and TS5-6 correspond to MIS 31-27 (1.1–0.98 Ma)
 555 and MIS 19 (~800-700 ka), respectively, as determined by palaeomagnetic polarity and correlation of glacial-
 556 interglacial cycles to the marine isotope stratigraphy. The pollen record for the earlier interval (MIS 31-27)
 557 documents several mesothermic taxa no longer occurring in the regional vegetation, including *Carya*,
 558 *Zelkova*, *Pterocarya* and *Liquidambar*, as well as mega-mesothermic *Taxodium* type and mid-altitude

575 phases. A small sample set from exposures in the southern coastal area of the Gulf of Corinth also documents
576 the presence of *Carya*, *Tsuga*, *Cedrus*, *Pterocarya*, *Liquidambar* and *Zelkova*, and has been used to support a
577 Lower and Middle Pleistocene age for fluvio-lacustrine and fan-delta syn-rift deposits (Rohais et al., 2007)
578 Other sites have potential to contribute to the issue, but details of rare taxa, including extinct or relictual
579 elements, may not yet be published. Sadori et al. (2016) present a new continuous pollen record spanning
580 the last 500 ka at Lake Ohrid, Albania. The data published to date reveals the good correspondence between
581 forested/non-forested phases and glacial-interglacial cycles, and highlights a progressive shift towards
582 warmer and drier interglacials associated with a diminishing role of *Abies* and *Picea* in the regional
583 vegetation. Future insights regarding extinct taxa for this, and the deeper sequence extending to 1.2 Ma
584 (Wagner et al., 2009) will be important. Ongoing investigation of new boreholes at Tenaghi Philippon at high
585 (centennial-scale) resolution may also contribute new insights into the pattern of regional extinctions in
586 Northeastern Greece over the last 1.35 Ma (Pross et al., 2015).

587

588 4. History of tree taxa extirpated from S Europe

589 The compilation of the pollen records in Figs 2-5, although schematic and in many cases fragmentary,
590 provides the basis for a comprehensive view of the population dynamics of tree taxa that disappeared from
591 Southern Europe during the Pleistocene. In particular, they provide regional overviews of the expansions and
592 last occurrences of the most important tree taxa that went extinct from the Southern European mainland.
593 For this review, we selected a number of taxa that are reliably identified and appreciably represented in all
594 the Quaternary pollen floras of South Europe: *Taxodium*-type, *Sciadopitys*, *Cathaya*, *Cedrus*, *Tsuga*,
595 *Eucommia*, *Engelhardia*, *Carya*, *Pterocarya*, *Parrotia*, *Liquidambar*, and *Zelkova* (Figs 6-8). Other taxa, not
596 discussed in this review, including Sapotaceae, Juglandaceae, Arecaceae, *Avicennia*, *Aesculus*,
597 *Parthenocissus*, *Nyssa*, *Picea omorika*, and *Platanus* may deserve a further work.

598

599

Fig. 6. Schematic stratigraphic distribution of *Taxodium* type, *Sciadopitys*, *Cathaya* type, and *Cedrus*. Blue lines represent taxa that are only mentioned in the original papers.

4.1 *Taxodium* type

As mentioned above, the pollen types indicated as Taxodiaceae (e.g., Ravazzi, 2003; Joannin et al., 2007a,b; Biltekin et al., 2015) and *Taxodium*-type (e.g., Fusco, 2007; Bertini et al., 2010) may belong to different genera of the Taxodiaceae subfamily (Cupressaceae family). According to Fusco (2007) and Bertini (2010), in the Italian pollen records they may include both *Taxodium* cf. *distichum* and *Glyptostrobus*, while Biltekin et al. (2015) ascribe it mostly to *Glyptostrobus*. Both genera were found in the Miocene macrofossil record of Europe (Kovar-Eder et al., 2006), but wood samples analyses from the Early Pleistocene deposit of Leffe (Ravazzi and van der Burgh, 1995) as well as shoots, cones and seeds from the Early Pleistocene deposit of Dunarobba in Central Italy (Martinetto et al., 2014) suggest that *Taxodium*-type pollen might be referred to *Glyptostrobus*.

On the other hand, fossil wood from a pyroclastic deposit near Rome, Ar/Ar dated at 514 ka (Marra et al., 2014), was ascribed to *Taxodium* (Follieri, 2010). Until more conclusive data are collected, the question of the exact attribution of *Taxodium*/*Glyptostrobus*-type pollen to one or the other genus is still open, as well as the possibility that more than one species or genus were present in the Quaternary deposits of Southern Europe. The question is highly relevant considering the present distribution of the two genera: *Taxodium* is limited to the southeastern regions of North America (*T. distichum* (L.) Rich.) and Mexico (*T. huegelii* C.Lawson), while *Glyptostrobus pensilis* (Staunton ex D.Don) K.Koch (the sole extant species of this genus) lives in southern China. Both *Taxodium* and *Glyptostrobus* live in swampy environments and riverbanks.

Bearing in mind these taxonomic uncertainties, the Pleistocene pollen records in South Europe provide a rather consistent picture of the history of *Taxodium*/*Glyptostrobus* (here after named *Taxodium* type) (Fig. 6). During the Gelasian, *Taxodium* type was present only in very low percentages in the peri-Iberian marine cores (core G1: Feddi et al., 2011), while it was rather abundant (5-10%) in Northeastern Spain (Garraf: Suc and Cravatte, 1982), and in Italy (Stirone: Lona and Bertoldi, 1972; Fosso Bianco: Pontini and Bertini, 2000; Coste San Giacomo: Bellucci et al., 2014; Poggio Rosso: Bertini et al., 2010; Semaforo: Combourieu-Nebout, 1993). During the early Calabrian, *Taxodium* type was still significantly represented (>5%) in Northern Italy (Lamone valley: Fusco, 2007; Lefte: Ravazzi, 2003). In Southern Italy it was abundant in the marine sediments of the Vrica section (not shown in Fig. 6) (Combourieu-Nebout and Vergnaud-Grazzini, 1991), where it progressively declined around 1.3 Ma, while it was discontinuously represented in the lacustrine sediments from Camerota (Brenac, 1984). Starting in the middle Calabrian, *Taxodium* type was strongly reduced in the Western and Central Mediterranean regions, but it still shows sporadic appearances before its final disappearance, dated around 600-500 ka both in France (MIS 15 at Alleret maar: Pastre et al., 2007; Degeai et al., 2013) and in Central-Southern Italy (MIS 15-13 at Vallo di Diano: Russo Ermolli, 1994; MIS 13 at Sessano: Russo Ermolli et al., 2010a). Macrofossils of *Taxodium* from near Rome, dated around 500 ka, confirm the age of these occurrences (Follieri, 2010).

In Greece, *Taxodium*-type is continuously found in low amounts at Zakynthos (Subally et al., 1999) before 1.8 Ma; it was sporadic at Rhodes until the early Middle Pleistocene (ca. 700 ka; Joannin et al., 2007a), where a

639 fossil wood of *Glyptostrobus* is dated around 500 ka (Boyd, 2009). No record is reported from Tenaghi
640 Philippon (Wijmstra, 1969; Wijmstra and Smit, 1976; Van der Wiel and Wijmstra, 1987a,b; Tzedakis et al.,
641 2006), Ohrid (Sadori et al., 2016), and Megalopolis (Okuda et al., 2002). In core DSDP 380 (Biltekin et al.,
642 2015) it is abundant throughout the Quaternary until recent, but it underwent two severe declines at about
643 1.2 and 0.5 Ma.

644 A clear pattern appears from the synoptical view of Fig. 6. The scarcity/lack of *Taxodium* type in Greece
645 suggests that two distinct populations/species/genera were probably present in South Europe, which had
646 different fates. The first one, found in Northeastern Spain, Southern France and Italy, underwent a severe
647 reduction in Northeastern Spain-Southern France and Central Italy at the end of the Gelasian, persisted in
648 Northern Italy until the mid-Calabrian (1.4-1.2 Ma), and disappeared definitely from Central and Southern
649 Italy after ca. 500 ka. The second one was distributed in Anatolia-Rhodes; it underwent a first severe
650 reduction around 1.2 Ma and a second reduction around 500 ka (at approximately the same age of the last
651 occurrence in Italy), but was in any case able to survive until recent times.

652

653 4.2 *Sciadopitys*

654 The sole extant species of the family Sciadopityaceae is *Sciadopitys verticillata* (Thunb.) Siebold & Zucc.
655 Molecular dating shows that it represents one of the earliest diverging plant lineages (ca. 220 Ma) survived
656 by a single species in the world alongside other classic 'living fossils' such as *Ginkgo* (Worth et al., 2013). It is
657 currently patchily distributed in high rainfall montane areas of Japan (islands of Honshu, Shikoku and Kyushu)
658 within temperate forests, where mean annual precipitation can exceed 4000 mm/yr (Farjon, 2005). It was
659 found in several Pliocene sites in Northwestern Europe (Kemna and Westerhoff, 2007), France (Suc, 1973),
660 and Northern Italy (Pini et al., 2003). The pollen data from Southern Europe (Fig. 6) indicate that, apart from
661 a single find in the marine core G1 from the Alboran Sea (Feddi et al., 2011), the only country where it is
662 recorded is Italy, where it is found not only during the Gelasian (Stirone: Lona and Bertoldi, 1972; Bertini,
663 2001; Fosso Bianco: Pontini and Bertini, 2000; Semaforo: Combourieu-Nebout, 1993; Klotz et al., 2006), but
664 also during the early Calabrian (Lamone Valley: Fusco, 2007; Pietrafitta: Lona and Bertoldi, 1972; Camerota:

665 [Brenac, 1984](#)). In Southern France, a few pollen grains were found in the Saint-Macaire maar, of Early
666 Calabrian age, but they are considered reworked from underlying Early Pliocene marine clays ([Leroy et al.,](#)
667 [1994](#)).

668 No occurrences of *Sciadopitys* are reported from Greece the Aegean islands and DSDP site 380, indicating
669 that the genus may have disappeared from Southeastern Europe in the course of the Pliocene, as it was
670 present in upper Miocene sediments from Samos and in Lower Pliocene deposits of Ptolemaida, in Western
671 Macedonia, as documented by macrofossils ([Velitzelos et al., 2014](#)).

672 The comparative view of the pollen records from Southern Europe has provided novel information about the
673 past distribution of this taxon, which has considerable evolutionary interest. In particular, Italy appears as a
674 long-term refuge area for *Sciadopitys*, that persisted in several areas of the Italian Peninsula for over one
675 million years after it had disappeared from the rest of Europe. This long-term persistence might be connected
676 with generally high rainfall along the Italian mountain chains.

677

678 4.3 *Cathaya*

679 *Cathaya* is a monotypic genus of the Pinaceae family, presently living only in Southeastern China, in
680 mesophilous forests at 900-1900 m elevation ([Farjon, 2008](#)). It is categorised as a palaeoendemic, with a
681 fossil history dating at least to the Cretaceous ([Liu and Basinger, 2000](#)). Pollen of *Cathaya* has often been
682 misidentified, having been confused with *Pinus* haploxylon-type, or even *Podocarpus* ([Liu and Basinger,](#)
683 [2000](#)). In many European records, it is named *Cathaya* type, leaving a certain margin of error. This is because
684 many determinations of fossil grains are based on published photographs and descriptions ([Caratini and](#)
685 [Sivak, 1972; Sivak, 1976; Liu and Basinger, 2000; Zanni and Ravazzi, 2007](#)), and not on fresh reference
686 collections, due to the rarity of the single extant species, *Cathaya argyrophylla*, which was formally described
687 only in 1958 by Chun and Kuang. In several Early Pleistocene Italian sites its pollen has been often classified
688 as, or joined to *Pinus* haploxylon-type ([Capraro et al., 2005; Fusco, 2007](#); see photographs in [Lona and](#)
689 [Ricciardi, 1961; Ricciardi, 1965](#); and revision by [Liu and Basinger, 2000](#)).

690 *Cathaya* fossils were found in many Miocene and Pliocene sites in Europe as both macrofossils (Kovar-Eder
691 et al., 2006; Velitzelos et al., 2014) and pollen (Jiménez-Moreno and Suc, 2007; Popescu et al., 2010). During
692 the early Gelasian, *Cathaya* was continuously present in the marine record from the Alboran Sea (Feddi et
693 al., 2011), and abundant (even >10%) in the record from Garraf 1 in Northeastern Spain (Suc and Cravatte,
694 1982). All the fragmented records from Northeastern Spain-Southern France of Early Pleistocene age reveal
695 its presence until at least the Jaramillo subchron, although in minimal amounts. In Northern Italy, it was well
696 represented at Stirone, in the Gelasian (Bertini, 2001). It was still present at the beginning of the Calabrian
697 at the Lamone Valley, where it is grouped together with *Pinus* haploxylon (Fusco, 2007), then it was not any
698 more found (Ravazzi, 2005). In Central Italy, *Cathaya* was again well represented during the Gelasian, with
699 percentages in some cases reaching values of 10% (Pontini and Bertini, 2000; Bertini et al., 2010; Bellucci et
700 al., 2014), while in the Calabrian it is recorded (as *Pinus* haploxylon) in two sites (Gubbio and Pietrafitta),
701 whose age – not well constrained – is around 1.4 Ma (Lona and Ricciardi, 1961; Lona and Bertoldi, 1972). Its
702 last occurrence is at Fontana Ranuccio in an interglacial phase dated around 850 ka (Corrado and Magri,
703 2011). In Southern Italy, *Cathaya* was abundant in the Early Pleistocene marine record of Croton especially
704 from ca. 1920 ka to 1740 ka; then it decreased but was still present around 1.3 Ma (Combourieu-Nebout and
705 Vergnaud- Grazzini, 1991). Small amounts of *Cathaya* pollen are found in Southern Italy until approx. 750 ka
706 (Capraro et al., 2005; Joannin et al., 2007b, 2008). In Greece, *Cathaya* is well represented in the Gelasian
707 record from Zakynthos (Subally et al., 1999), it is represented by macrofossils finds of Calabrian age at Rhodes
708 (Mai and Velitzelos, 2007; Velitzelos et al., 2014), while it is not recorded at Tenaghi Philippon, where it could
709 have been counted as *Pinus* haploxylon-type, almost continuously present up to 850 ka (Wijmstra and Smit,
710 1976; Van der Wiel and Wijmstra, 1987a). *Cathaya* pollen is rare in the Early Pleistocene record at the DSDP
711 380 site (Biltekin et al., 2015).

712 During the Middle Pleistocene, only two sites report sporadic occurrences of *Cathaya* in Southern Europe:
713 Alleret in Southern France at 700-550 ka (Pastre et al., 2007; Degeai et al., 2013) and Site 380, discontinuously
714 present until 380 ka (Biltekin et al., 2015).

715 On the whole, in spite of identification difficulties and of nomenclature uncertainties, the Southern European
716 records provide a consistent picture of the history of *Cathaya*, which was present at all sites during the
717 Gelasian. In the Calabrian it was probably more common in Southern Italy ([Combourieu-Nebout and](#)
718 [Vergnaud- Grazzini, 1991](#)) than in other regions, where it was already severely fragmented. However, the
719 sparse occurrences documented in France and Anatolia between 700 ka and 380 ka suggest that very small
720 relict populations may have persisted in isolated location and warn of the possibility to find more fossil
721 remains of Middle Pleistocene age in Southern European deposits.

722

723 4.4 *Cedrus*

724 The genus *Cedrus* includes two extant accepted species, native in the Mediterranean mountains: *Cedrus*
725 *atlantica* (Endl.) Manetti ex Carrière, living in Algeria and Morocco ([Slimani et al., 2014](#)), and *Cedrus libani*
726 A.Rich., distributed in Asia Minor ([Fady et al., 2008](#)), including *Cedrus libani* var. *brevifolia* Hook.f. in Cyprus,
727 sometimes considered as a separate species ([Jasińska et al., 2013](#)). A further species, *Cedrus deodara* (Roxb.
728 ex D.Don) G.Don, lives in Afghanistan and in the southern slopes of the western Himalaya ([Bou Dagher-](#)
729 [Kharrat et al., 2007](#)). The modern Mediterranean populations of *Cedrus* are severely fragmented as a result
730 of the unfavorable climate conditions of the last glacial period ([Cheddadi et al., 2009](#)), of a long history of
731 human activity ([Hajar et al., 2010b](#)), and of increasing aridity in the late Holocene ([Hajar et al., 2010a](#)).

732 Modern pollen data obtained in/near cedar forests ([Hajar et al., 2008](#)) show that percentages above 5-7%
733 indicate its presence near the site ([Hajar et al., 2010a](#); [Bell and Fletcher, 2016](#)). This result is confirmed by
734 the Holocene pollen records from Morocco, Algeria and Lebanon, showing high (>10%) pollen percentages
735 of *Cedrus* in the vicinity of modern cedar forests ([Lamb et al., 1989](#); [Hajar et al., 2008](#); [Cheddadi et al., 2009](#)).
736 For this reason, sparse pollen grains of *Cedrus* that are often found in late Quaternary sedimentary deposits
737 from the north Mediterranean regions, especially during the last glacial period ([Magri and Parra, 2002](#)), have
738 been interpreted as the effect of long-distance transportation from north Africa ([Reille, 1990](#)).

739 Fossil data show that *Cedrus* has a long history in Europe, being already present during the Middle Miocene
740 (16-11 Ma) in Anatolia ([Yavuz-Işık, 2007](#), [Biltekin et al., 2015](#)), Bulgaria ([Ivanov et al., 2002](#)), and Austria

([Jiménez-Moreno et al., 2008](#)). However, in the course of the Pleistocene, *Cedrus* underwent a strong reduction, disappearing from Europe, where it is currently absent. It is worth noticing that a strong reduction of *Cedrus* has been reported in also China since the Upper Pliocene, and ascribed to an intensification of the East Asian winter monsoon ([Su et al., 2013](#)).

The Southern European pollen records ([Fig. 6](#)) document a regionally diverse history of *Cedrus*. In the marine sediment cores from the Alboran sea, *Cedrus* had a discontinuous presence at the end of the Pliocene and during the Gelasian ([Feddi et al., 2011](#)), while continuous and appreciable percentages of *Cedrus* pollen are found during the time intervals ca. 1.09-0.90 Ma (core ODP976: [Joannin et al., 2011](#)), 48-26 ka (core MD95-2043: [Sánchez-Góñi et al., 2002](#)), and the last 25 ka (core ODP976: [Combourieu-Nebout et al., 2009](#)). These records support a continuous presence of cedar throughout the Pleistocene in the regions facing the Alboran sea, with a most likely North African origin of this anemophilous pollen type, also considering the documented presence of *Cedrus* in Morocco during the late Middle Pleistocene (Benzú rockshelter: [Ramos et al., 2008](#)) and the Late Pleistocene ([Lamb et al., 1989](#); [Cheddadi et al., 2009](#)).

The Early Pleistocene record from the Iberian Peninsula documents only sporadic occurrences of *Cedrus*, except for the Gelasian record from Garraf 1 ([Suc and Cravatte, 1982](#)). Pollen of *Cedrus* was very rare or absent at Morgadinho in Algarve ([Antunes et al., 1986](#)), at Tres Pins ([Leroy, 1997](#)), Banyoles ([Julia Bruguès and Suc, 1980](#)), Cal Guardiola D2 ([Postigo Mijarra et al., 2007](#)), and Bòbila Ordis IV ([Leroy, 2008](#)). In Southern France, *Cedrus* was found in various deposits of Pliocene age ([Fauquette et al., 1999](#); [Suc and Drivaliari, 1991](#)), but in the Early Pleistocene it was recorded in very low amounts at Senèze ([Elhaï, 1969](#)), Bernasso ([Leroy and Roiron, 1996](#)), Nogaret ([Leroy and Seret, 1992](#)), Saint Macaire ([Leroy et al., 1994](#)), and Ceyssac-8 ([Ablin, 1991](#)). By contrast, the Early Pleistocene record of *Cedrus* from the Italian Peninsula is very rich. In Northern Italy, where it was abundant already during the Pliocene (Aulla: [Bertoldi, 1988](#); Stirone: [Bertini, 2001](#)), pollen percentages >5% document its local presence at Stirone ([Lona and Bertoldi, 1972](#); [Bertini, 2001](#)), Lamone Valley ([Fusco, 2007](#)), and Leffe ([Ravazzi and Rossignol-Strick, 1995](#); [Ravazzi, 2003](#); [Muttoni et al., 2007](#)). It becomes sporadic in the Prealps at the end of the Early Pleistocene (Fornaci di Ranica: [Ravazzi et al., 2005](#)). In Central Italy, *Cedrus* is well represented in several Pliocene sites (e.g., Upper Valdarno Basin: [Bertini and](#)

767 [Roiron, 1997](#); [Bertini and Martinetto, 2011](#); Borgorose and Marano de' Marsi: [Sadori et al., 2010](#)), and also
 768 many Early Pleistocene sites document abundant pollen of cedar, with values in some levels >10% at Fosso
 769 Bianco ([Pontini and Bertini, 2000](#)), Coste San Giacomo ([Bellucci et al., 2014](#)), Gubbio ([Lona and Ricciardi,](#)
 770 [1961](#)) and Colle Curti ([Bertini, 2000](#)). In some late Early Pleistocene (Calabrian) sites *Cedrus* was present but
 771 sparse (Madonna della Strada: [Magri et al., 2010](#); Pagliare di Sassa: [Palombo et al., 2010](#); Saticula: [Russo](#)
 772 [Ermolli et al., 2010b](#); Fontana Ranuccio: [Corrado and Magri, 2011](#)), suggesting that its distribution was
 773 already rather fragmented in Central Italy. Also in Southern Italy, *Cedrus* was already present in the Pliocene
 774 together with other montane conifers (*Cathaya*, *Tsuga*, *Abies*, and *Picea*) (Punta Piccola: [Combourieu-Nebout](#)
 775 [et al., 2004](#)). In the Early Pleistocene it was abundant in the marine sediments at Semaforo-Vrica
 776 ([Combourieu-Nebout and Vergnaud-Grazzini, 1991](#); [Combourieu-Nebout, 1993](#)), Camerota ([Brenac, 1984](#)),
 777 Santa Lucia ([Joannin et al., 2007b](#)), and Montalbano Jonico ([Joannin et al., 2008](#)). Marine sediments from
 778 Rhodes ([Joannin et al., 2007a](#)) show a continuous presence of *Cedrus* in the island in the Early Pleistocene,
 779 with values much higher than at Tenaghi Philippon ([Tzedakis et al., 2006](#)) and at Site 380 ([Biltekin et al., 2015](#)),
 780 where it was rather discontinuous.

781 During the Middle and late Pleistocene the French sites in the Velay region document a very discontinuous
 782 occurrence of *Cedrus* ([Reille et al., 2000](#), [Pastre et al., 2007](#); [Degeai et al., 2013](#)). In the Iberian Peninsula,
 783 *Cedrus* is relatively frequent in continental pollen records during the Late Pleistocene and Holocene, although
 784 with percentages values never >2% (see compilation by [González-Sampériz et al. \(2010\)](#) and references
 785 therein, as well as the recent pollen record from Cañizar de Villarquemado: [González-Sampériz et al., 2013](#);
 786 [Aranbarri et al., 2014](#); [García-Prieto, 2015](#)). In Italy, *Cedrus* underwent a dramatic reduction during the
 787 Middle Pleistocene, being recorded in significant amounts only in Southern Italy at the Early to Middle
 788 Pleistocene transition ([Bertini et al., 2015](#); [Capraro et al., 2005](#)). In Northern Italy it was absent in the early
 789 Middle Pleistocene (Pianico Sèllere: [Moscariello et al., 2000](#); [Ravazzi, 2003](#)) and was not found until present
 790 (see Azzano Decimo: [Pini et al., 2009](#), Fimon: [Pini et al., 2010](#)). In Central and Southern Italy the record of
 791 *Cedrus* was discontinuous at all sites during the Middle and Late Pleistocene (Cesi: [Bertini, 2000](#); Torre
 792 Mucchia: [Pieruccini et al., 2016](#); La Pineta: [Lebreton, 2002](#); Ceprano: [Manzi et al., 2010](#); Sessano, [Russo](#)
 793 [Ermolli et al., 2010a](#); Vallo di Diano: [Russo Ermolli, 1994](#); Acerno: [Munno et al., 2001](#); Boiano, [Orain et al.,](#)

2015; Valle di Castiglione: [Follieri et al., 1988](#); Lagaccione: [Magri, 1999](#); Lago Grande di Monticchio: [Allen et al., 2000](#)). Unexpectedly, *Cedrus* is well represented (>2%) in the lateglacial record from Pesce Luna ([Di Rita et al., 2015](#)) near Rome. In the Balkan Peninsula, the Middle Pleistocene records from Ohrid ([Sadori et al., 2016](#)), Ioannina ([Tzedakis, 1994](#); [Roucoux et al., 2008](#)), Megalopolis ([Okuda et al., 2002](#)) and Kopais ([Okuda et al., 2001](#)) do not document a presence of *Cedrus* in the region. It is also possible that these diagrams, representing only selected taxa, do not report occasional finds of *Cedrus*, whose presence must have been negligible. At Tenaghi Philippon ([Tzedakis et al., 2006](#)) during the Middle Pleistocene *Cedrus* became sporadic until its last occurrence, dated approx. 0.35 Ma. In the marine record from Rhodes ([Joannin et al., 2007a](#)), which is relatively close to the modern range of *Cedrus* in Turkey, it is found with frequencies >20% in forest phases of the early Middle Pleistocene correlated to MIS 19 and 17. Interestingly, fossil leaves of *Cedrus* were found in Rhodes in sediments with an age of about 0.5 Ma, confirming the local presence of the genus until at least the Middle Pleistocene ([Boyd, 2009](#)). In the Black Sea, *Cedrus* is continuously documented until recent, although with discontinuous occurrences, suggesting its long-time persistence on the mountain areas bordering the Anatolian Plateau and its elimination from the northern and western margins of the Black Sea by human disturbance ([Biltekin et al., 2015](#)).

The interpretation of the Southern European pollen record of *Cedrus* is not totally conclusive: while it is clear that it was widespread in Italy and in the Aegean region during the Early Pleistocene, as documented by high pollen percentage values, it is very difficult to evaluate the discontinuous single grains found in the Iberian, French and Italian records since the Middle Pleistocene. Following the interpretation by [Postigo-Mijarra et al. \(2010\)](#) who consider that even sparse occurrences of *Cedrus* indicate its local presence, *Cedrus* should have been present in Southern Europe until the Holocene. Alternatively, supporting the hypothesis by [Reille et al. \(2000\)](#) and [Magri and Parra \(2002\)](#) that sparse pollen grains are of North African origin, *Cedrus* was never present in continental Iberia during the Quaternary, disappeared from Northeastern Spain-Southern France at the beginning of the Pleistocene, and from the Italian Peninsula at an early stage of the Middle Pleistocene. In Greece, *Cedrus* possibly persisted a little longer, but the continuous record from Tenaghi Philippon indicates that during the last 1.4 Ma it was always rather sparse. The almost continuous presence of *Cedrus* in the Black Sea core is consistent with its relative vicinity in Turkey.

Fig. 7. Schematic stratigraphic distribution of *Tsuga*, *Eucommia*, *Parrotia*, and *Liquidambar*. Blue lines represent taxa that are only mentioned in the original papers.

4.5 *Tsuga*

The genus *Tsuga* (Pinaceae) is an important element of subalpine coniferous and temperate broad-leaved deciduous forests, currently showing a disjunct distribution in Eastern and Western North America, Japan, China and Himalaya. It lives at elevations from sea level to 3800 m, in areas where precipitation ranges from 700 to 4500 mm/year and is distributed throughout the year (Farjon, 2008). An abundant fossil record of *Tsuga* indicates that it was distributed throughout North America and Eurasia from the Late Cretaceous to the Plio-Pleistocene (LePage, 2003). In the Northwestern European records of Early Pleistocene age, *Tsuga* pollen is found during the Tiglian (MIS 79–63), Waalian (MIS 49–37) and Bavelian (MIS 31–22) (Zagwijn, 1985).

833 The last occurrence of *Tsuga* pollen grains in Central Europe is during the Leerdam interglacial, a post-
 834 Jaramillo late Early Pleistocene warm stage (Zagwijn and De Jong, 1984).
 835 During the Gelasian, *Tsuga* is well represented at all sites in Northeastern Spain-Southern France and in Italy
 836 (Fig. 7), including Garraf (Suc and Cravatte, 1982), Bernasso (Leroy and Roiron, 1996), Nogaret (Leroy and
 837 Suc, 1992), Stirone (Lona and Bertoldi, 1972; Bertini, 2001), Fosso Bianco (Pontini and Bertini, 2000), Coste
 838 San Giacomo (Bellucci et al., 2014), Poggio Rosso (Bertini et al., 2010), Semaforo and Vrica (Combourieu-
 839 Nebout, 1993; Combourieu-Nebout and Vergnaud-Grazzini, 1991). In contrast, it is not represented in the
 840 marine records south of 38°N from the Alboran Sea (G1: Feddi et al., 2011) and from Zakynthos (Subally et
 841 al., 2009). In core DSDP 380 it is discontinuously represented.
 842 The Calabrian record of *Tsuga* is generally very rich, showing higher percentages than in the Gelasian, at least
 843 until the Jaramillo subchron. In Northeastern Spain it is particularly abundant at Bòbila-Ordis IV (Leroy, 2008)
 844 and in S France at Ceyssac-8 (Abilin, 1991). It is also present with values >10% in Southwestern France in
 845 the Médoc region (O'Brien and Jones, 2003), reflecting the fact that it was largely distributed over Southern
 846 Europe. In Northern Italy it was found with percentages often >10% at the Lamone Valley (Fusco, 2007), Leffe
 847 (Ravazzi and Rossignol Strick, 1995; Ravazzi, 2003), Pianengo (Muttoni et al., 2003), and in the late Early
 848 Pleistocene lacustrine sediments from Fornaci di Ranica (Ravazzi et al., 2005). It was also found in Slovenia
 849 near Novo Mesto (Šercelj, 1961) and near Ilirska Bistrica (Šercelj, 1965). In Central Italy it was abundant in
 850 many sites, reaching its highest percentages (>30%) at Madonna della Strada (Magri et al., 2010) around 1.3
 851 Ma and at Colle Curti (>10%) during the Jaramillo subchron (Bertini, 2000). It was continuously present also
 852 in the records of Pietrafitta (Lona and Bertoldi, 1972), Gubbio (Lona and Ricciardi, 1961), Leonessa (Ricciardi,
 853 1965), Saticula (Russo Ermolli et al., 2010b), and Fontana Ranuccio (Corrado and Magri, 2011). In Southern
 854 Italy it is very abundant in the marine sediments (>30%) at Santa Lucia during MIS 43-40 (Joannin et al.,
 855 2007b) and is continuously present at Montalbano Jonico during MIS 37 to 23 (Joannin et al., 2008). At
 856 Tenaghi Philippon in Macedonia (Tzedakis et al., 2006), *Tsuga* displays declining percentages in the interval
 857 1.35-0.8 Ma. In Site 308 its declining trend is very evident, as it becomes sporadic after 1.27 Ma (Biltekin et
 858 al., 2015).

859 During the Middle Pleistocene, *Tsuga* underwent a marked general decline, although with different timings
860 from one region to the other. There are very few data from Spain that can help to assess the local presence
861 of *Tsuga*. In Southern France, there were still sporadic occurrences in two interglacial periods corresponding
862 to MIS 17 and MIS 15 at the Alleret maar (Pastre et al., 2007; Degeai et al., 2013). In Northern Italy, the
863 lacustrine succession of Pianico-Sèllere (Moscariello et al., 2000), close to the Early/Middle Pleistocene
864 boundary (Pinti et al., 2001), records only sporadic grains of *Tsuga* (Ravazzi et al., 2005), which correspond
865 to the last occurrences for this taxon in Northern Italy. However, macrofossil finds are reported at the site of
866 Oriolo dated around 800 ka (Martinetto and Sami, 2001). In Central Italy, *Tsuga* was still found with
867 percentages >3% in an early interglacial phase correlated with MIS 17 (Pieruccini et al., 2016), and with single
868 pollen grains in the early Middle Pleistocene (Cesi: Bertini, 2000), and during MIS 13 (Sessano: Russo Ermolli
869 et al., 2010). In Southern Italy, *Tsuga* was still well represented (2-5%) at the Early to Middle Pleistocene
870 transition (Valle di Manche: Capraro et al., 2005; Montalbano Jonico: Bertini et al., 2015), but was dubitably
871 present during MIS 13 (Vallo di Diano Russo Ermolli, 1994). However, it was virtually absent from some sites
872 (Lebreton, 2002), suggesting that its distribution was rather patchy. In the Aegean, the last occurrences of
873 *Tsuga* date back to MIS 17 (Joannin et al., 2007), slightly later than at Tenaghi Philippon, where it disappears
874 near the Brunhes/Matuyama transition (Tzedakis et al. 2006). At Lake Ohrid, located at the border between
875 the Former Yugoslavian Republic of Macedonia and Albania, although the detailed record has not been
876 published yet, the occurrence of *Tsuga* is reported with an age younger than 500 ka (Sadori et al., 2016). At
877 Site 380 *Tsuga* is sporadically found until 230 ka (Biltekin et al., 2015).

878 Considered altogether, the Southern European Quaternary records of *Tsuga* appear rather homogenous (Fig.
879 7), with sparse presence in the most southern sites (Alboran Sea, Rhodes), and significant abundances in
880 Northeastern Spain-Southern France and the Italian Peninsula during the Early Pleistocene, especially during
881 the Calabrian. A general increase in *Tsuga*, recorded between 1.8 Ma and 1 Ma, is often considered indicative
882 of cooler but relatively moist conditions (Bertini, 2010). From the population dynamics point of view, it is
883 noteworthy that this general expansion of *Tsuga* during the Calabrian, corresponding also to a significant
884 presence in central Europe, occurred just before its decline, which was moderate until approx. 700 ka, and
885 then so severe as to disrupt the Southern European range of *Tsuga* and lead to its rapid and definitive

886 disappearance. This is recorded at different times: approx. 750 ka in Northern Italy, 550 ka in Southern France,
887 and 280 ka in Anatolia. In the light of the most recently published data from several Middle Pleistocene sites,
888 and considering the under-representation of *Tsuga* pollen ([Janssen, 1984](#)), the single pollen grains found at
889 Vallo di Diano ([Russo Ermolli, 1994](#)) may be considered representative of local small stands of *Tsuga*, so fixing
890 around 450 ka the age of the last occurrence of *Tsuga* in Southern Italy.
891 [Bertini \(2000\)](#) related the disappearance of *Tsuga* to the Middle Pleistocene shift from 41 to 100 ka cyclicity,
892 leading to increased global aridity and decreased temperatures during the glacial stages, whereas [Ravazzi et](#)
893 [al. \(2005\)](#) ascribed the *Tsuga* decline recorded at Fornaci di Ranica to catastrophic fires triggered by summer
894 drought.

895

896 4.6 *Eucommia*

897 *Eucommia ulmoides* Oliv. is the single extant species of the genus *Eucommia* (Eucommiaceae). It is a small
898 deciduous tree currently living only in a hilly area of South China ([Wang et al., 2003](#)). It is more tolerant of
899 dry climate than many other extinct genera ([Svenning, 2003](#)), although spring temperature is a limiting factor
900 for *Eucommia* ([Wang et al., 2003](#)).

901 It was present in Central Europe during the Pliocene ([Hably and Kvacek, 1998](#)), where it was found until the
902 late Early Pleistocene ([Ber, 2005](#)).

903 With respect to Southern Europe, *Eucommia* pollen was not found in the marine cores from the Alboran Sea,
904 where it was missing also in Pliocene sediments ([Feddi et al., 2011](#)). In contrast, it is found in almost all sites
905 in Northeastern Spain and Southern France, until the end of the Early Pleistocene (ca. 800 ka), suggesting a
906 local persistence in the area. The Middle Pleistocene records from the Alleret Maar (700-550 ka) do not
907 report *Eucommia* any more ([Pastre et al., 2007](#); [Degeai et al., 2013](#)).

908 In Italy, pollen of *Eucommia* was present in the Gelasian marine sediments from Semaforo, but its record is
909 not shown in detail in the original papers ([Combourieu-Nebout, 1993](#); [Klotz et al., 2006](#)). It is missing from all
910 the Pleistocene pollen records from Central Italy. It is discontinuously documented in Northern Italy during
911 the early Calabrian ([Fusco, 2007](#)), but increased in the course of time, reaching its highest values at the

912 Jaramillo subchron (Ravazzi, 2003; Muttoni et al., 2007). Macrofossils of *Eucommia* were found in the
913 Castelletto Cervo II site, of possible Gelasian age (Cavallo and Martinetto, 2001), at Arda-Castell'Arquato both
914 in Gelasian and in late Calabrian layers (Martinetto, 2015; Martinetto et al., 2015), as well as in the latest
915 Calabrian (Stirone-Laurano site: Gregor, 1986). The consistency of pollen and macrofossil data supports the
916 hypothesis advanced by Martinetto (2015) of an Early Pleistocene immigration or increase of *Eucommia* in
917 Northern Italy. In Italy, *Eucommia* is not recorded after the Calabrian.

918 *Eucommia* is not present in the Aegean records, while it is recorded at Tenaghi Philippon from the base of
919 the record up to approx. 600 ka (Tzedakis et al., 2006). In Site 380 it is found only in Pliocene sediments
920 (Popescu et al., 2010).

921 The history of *Eucommia* is clearly distinct from the other taxa that disappeared from Europe during the
922 Quaternary, as this temperate tree persisted in central Europe at least as long as in Southern France, until
923 the late Calabrian. It expanded in Northern Italy during the Early Pleistocene, without reaching Central Italy,
924 where it was never found. The population at Tenaghi Philippon was possibly a separate one, as it is not
925 matched by any other data from Greece and around the Black Sea.

926

927 4.7 *Parrotia*

928 The genus *Parrotia* (family Hamamelidaceae) includes two species: *P. persica* C.A. Mey., endemic to the
929 Alborz Mountains of northern Iran (Sefidi et al., 2011), and *P. subaequalis* (Hung T. Chang) R.M. Hao & H.T.
930 Wei, endemic to a restricted area in Eastern China (Li and Del Tredici, 2008). The disjunct distribution and the
931 relictual state of the genus reflect the dramatic reduction it underwent during the Pleistocene.

932 The European records of Miocene and Pliocene age indicate that *Parrotia* was widespread, together with
933 other genera currently restricted in the Hyrcanian region, including *Pterocarya* and *Zelkova* (Jiménez-Moreno
934 and Suc 2007; Jiménez-Moreno et al., 2008, 2010).

935 The Pleistocene records of *Parrotia* in Southern Europe are sparse (Fig. 7), probably because its pollen is
936 markedly under-represented, as shown in the Early Pleistocene site of Bernasso, where a large number of

937 fossil leaves (at least 130 specimens; 15% of total finds) are matched by only low pollen percentages (max
938 4%) (Leroy and Roiron, 1996). Pollen morphology is illustrated in Leroy (2007) and Bińka et al. (2003).
939 During the Early Pleistocene, *Parrotia* is recorded in all Southern European regions (Spain; Southern France;
940 North, Central and Southern Italy; and Northeastern Greece at Tenaghi Philippon), and even in IODP Site
941 U1385 off Portugal, dated around 1.26 Ma (Tzedakis et al., 2015). Macrofossils were found in the Iberian
942 Peninsula at Crespià-Incarcal (Villalta and Vicente, 1972; Roiron, 1983) and in France at Bernasso (Leroy and
943 Roiron, 1996).
944 The last record of *Parrotia* in Southern Europe comes from Northern Italy at the onset of the Middle
945 Pleistocene, where it is represented by fossil leaves (Martinetto and Sami, 2001). The record by Bottema
946 (1974), who reported up to 10% *Parrotia* in Ioannina-II core, is very uncertain, as *Parrotia* was not found
947 again at Ioannina in any part of the 423 kyr I-249 sequence (Tzedakis, 1994), nor in the stratigraphically longer
948 I-284 sequence (Lawson et al., 2004), nor in the Ohrid core, spanning the last 500 ka (Sadori et al., 2016).
949 In Southern Europe, the lack of fossil remains of *Parrotia* younger than ca. 800 ka is surprising, also in the
950 light of its continuous record during an interglacial phase attributed to the Holsteinian in Eastern Poland
951 (Bińka et al., 2003). Various explanations may be suggested to solve this puzzle: either it had a long-term
952 refuge in Central Europe, or it migrated there from Southern Europe but it was never found there due to its
953 poor pollen dispersal, or it reached Eastern Poland from the Caspian region, where it is still documented by
954 Holocene pollen records (Leroy et al., 2013), passing north of the Caucasus Range. The latter hypothesis is
955 also supported by the absence of *Parrotia* from the Southwestern side of the Black Sea since the Pliocene
956 (Biltekin et al., 2015).

957

958 4.8 *Liquidambar*

959 The genus *Liquidambar* (family Altingiaceae) is distributed over a wide geographical range extending from
960 North America to East Asia. In the Mediterranean basin, it is represented only by the species *L. orientalis*
961 Mill., naturally distributed in the temperate parts of Western and Southern Anatolia and on the island of

962 Rhodes ([Öztürk et al., 2008](#)). It prefers the wet-mild bioclimate of moist bottomlands, occasionally flooded
 963 swampy plains, riparian habitats and humid areas.

964 It was a rather common component of mixed deciduous and evergreen forests in Europe during the Miocene
 965 and Pliocene, being often an element of a riparian vegetation, accompanied by *Salix*, *Alnus*, *Carya*,
 966 *Pterocarya*, *Carpinus*, *Zelkova*, and *Ulmus* ([Jiménez-Moreno and Suc 2007](#); [Jiménez-Moreno et al., 2008](#),
 967 [2010](#)).

968 *Liquidambar* is recorded in the Early Pleistocene marine cores from the Alboran Sea up to 900 ka ([Feddi et](#)
 969 [al., 2011](#); [Joannin et al., 2011](#)) and in the records from Spain until the Jaramillo subchrone ([Fig. 7](#)). In the
 970 Massif central it was present during MIS 17 ([Degeai et al., 2013](#)), while it was no longer found in the sediments
 971 corresponding to MIS 15 ([Pastre et al., 2007](#)).

972 In Northern and Central Italy, *Liquidambar* is documented in the Calabrian sediments of the Lamone valley
 973 ([Fusco, 2007](#)), Lefte ([Ravazzi and Rossignol-Strick, 1995](#); [Ravazzi, 2003](#)), Gubbio ([Lona and Ricciardi, 1961](#)),
 974 Pietrafitta ([Lona and Bertoldi, 1972](#)), and Colle Curti ([Bertini, 2000](#)), while it is missing in all the Middle
 975 Pleistocene sites. In Southern Italy, *Liquidambar* was almost continuously present until 1.3 Ma ([Joannin et](#)
 976 [al., 2007](#)), was rather discontinuous until 900 ka ([Joannin et al., 2008](#)), and represented by sparse pollen
 977 grains at the Early to Middle Pleistocene transition ([Capraro et al., 2005](#); [Bertini et al., 2015](#)). The pollen
 978 record from Vallo di Diano documents *Liquidambar* around 600 ka ([Russo Ermolli, 1994](#)). This find was
 979 considered reworked, but in the light of the most recent papers recording *Liquidambar* around 750 ka in
 980 Southern Italy ([Bertini et al., 2015](#)) and 700 ka in France ([Degeai et al., 2013](#)), as well as the last occurrence
 981 of *Liquidambar* at Tenaghi Philippon around 480 ka ([Wijmstra and Smit, 1976](#); [Van der Wiel and Wijmstra,](#)
 982 [1987a, 1987b](#)), it may be reasonably regarded as valid.

983 A significant presence in the Middle Pleistocene pollen records ([Joannin et al., 2007](#)), as well as macrofossil
 984 finds ([Boyd, 2009](#); [Velitzelos et al., 2014](#)) from Rhodes are not surprising, considering the small population of
 985 *Liquidambar* currently living in the island. At Site 380, an almost continuous presence until 1.28 Ma and
 986 disappearance at 270 ka are revealed by the pollen record.

987 The recent pollen studies from Southern Europe and Anatolia allow the definition of a marked decline of
 988 *Liquidambar* populations in Europe towards the end of the Early Pleistocene. During the Middle Pleistocene

its remnant populations were scattered in various regions (Southern France, Southern Italy, coasts of the Black Sea), but still surviving for a few glacial/interglacial cycles before being reduced to the small populations currently living in Rhodes and Southeastern Turkey.

Fig. 8 Schematic stratigraphic distribution of *Engelhardia*, *Carya*, *Pterocarya*, and *Zelkova*. Blue lines represent taxa that are only mentioned in the original papers.

4.9 *Engelhardia*

Engelhardia is a genus of the family Juglandaceae currently widespread in tropical to subtropical forests of South China, Indochina, Malaysia, Philippines and New Guinea, with mean annual precipitation between 740 and 3500 mm (Fang et al., 2011; Meng et al., 2015). It was very common in European Miocene and Pliocene

deposits, where it was found both as macrofossils and pollen (e.g., [Ivanov et al., 2002](#); [Kovar-Eder et al., 2006](#); [Jiménez-Moreno and Suc, 2007](#); [Jiménez-Moreno et al., 2008](#); [Velitzelos et al., 2014](#)).

The genus shows an extinction pattern during the Pliocene with decreasing ages from North to South Europe, in relation with the Northern Hemisphere temperature evolution, with a longer persistence in the Mediterranean regions where it probably disappeared because of increasing dryness ([Popescu et al., 2010](#); [Biltekin et al., 2015](#)).

During the Gelasian, it was recorded in low amounts in the marine core G1 from the Alboran Sea ([Feddi et al., 2011](#)), and abundantly (up to 10%) in the site of Garraf ([Suc and Cravatte, 1982](#); [Fauquette et al., 1998](#), [González-Sampériz et al., 2010](#)), in continuity with its Pliocene distribution ([Fig. 8](#)). At Fosso Bianco in Southern Italy ([Pontini and Bertini, 2000](#)), it is also in continuity with the Pliocene record, but it shows much lower percentages (<5%). In Northern Italy it only shows a sporadic occurrence in the Stirone section, where it was continuously present during the Pliocene ([Bertini, 2001](#)). *Engelhardia* is reported in Southern Italy at Semaforo, but its presence was probably modest, as it is not shown separately from other elements of the Subtropical humid forest, altogether showing percentages always <5% ([Combourieu-Nebout, 1993](#); [Klotz et al., 2006](#)). It is reported from Rhodes up to 1.7 Ma ([Biltekin et al., 2015](#)), and is present in Site 380 ([Biltekin et al., 2015](#)), although discontinuously.

In the Calabrian, *Engelhardia* appears strongly reduced in Spain and Southern France, where it was anyway found in several sites until approx. 1 Ma ([Ablin, 1991](#); [Leroy et al., 1994](#); [Leroy, 2007, 2008](#)). In Italy it is recorded both in Northern Italy (Lamone Valley: [Fusco, 2007](#)) and in Southern Italy (Camerota: [Brenac, 1984](#)). It was possibly present also in the Vrica section: detailed data have not been published ([Combourieu-Nebout and Vergnaud-Grazzini, 1991](#)), but its last occurrence at Vrica is reported at 1.35 Ma by [Biltekin et al. \(2015\)](#).

Interestingly, *Engelhardia* was not found in any site of Central Italy, nor in the records from Greece and the Aegean, while at Site 380 it was relatively abundant until 1.25 Ma, then it is sporadic ([Biltekin et al., 2015](#)).

In the Middle Pleistocene, *Engelhardia* was not found in Spain, France and Italy. Sporadic finds at the Early to Middle Pleistocene transition in Southern Italy ([Capraro et al., 2005](#); [Bertini et al., 2015](#)) are considered as reworked ([Bertini, 2015](#)). However, it is reported from a Mid-Pleistocene core from the North Adriatic Sea

(unpublished report: [Biltekin et al., 2015](#)), and it is represented by sporadic occurrences at Site 380 until ca. 200 ka ([Biltekin et al., 2015](#)). Interestingly, it was found further East in the Western Caucasus in the early Middle Pleistocene ([Shatilova et al., 2011](#)).

The Quaternary history of *Engelhardia* in Southern Europe shows a marked fragmentation pattern since the end of the Pliocene, when *Engelhardia* was distributed over large parts of the European territory. It persisted longer in Northeastern Spain/Southern France (until ca. 1 Ma) than in the Italian Peninsula, where it was probably split into two separate populations: one in Southern Italy, persisting until approx. 1.35 Ma (or possibly later if the finds at the Early to Middle Pleistocene transition will be confirmed), and one in Northern Italy, lasting until the early Calabrian, separated by a large region in Central Italy where *Engelhardia* was absent. A separate population, lasting until ca. 200 ka, is documented by the long DSDP 380 record in the Black Sea.

1037

1038 4.10 *Carya*

The genus *Carya* (family Juglandaceae) is currently distributed in subtropical and temperate continental biomes over North America in the United States, Canada and Mexico, and Southeast Asia in China, Viet Nam, Laos and India, with rainfall values between 1000 and 1500 mm/year ([Manchester, 1987](#); [Fang et al., 2011](#); [Orain et al., 2013](#)). It was rather common in central Europe during the Miocene and Pliocene ([Pontini and Bertini, 2000](#); [Jiménez-Moreno et al., 2008, 2010](#); [Kovar-Eder et al., 2006](#)).

In the Gelasian Southern European record ([Fig. 8](#)), *Carya* is well represented at all sites, except the Iberian marine cores, where it is absent. In Site 380 it is almost continuously represented, although in very low frequencies.

Towards the end of the Gelasian and during the Calabrian, there was a general increase in pollen percentage values of *Carya*, in some cases surpassing 30%. In the late Gelasian French sites of Nogaret ([Brugal et al., 1990](#); [Leroy and Suc, 1992](#)) and Bernasso ([Leroy and Roiron, 1996](#)), high percentages of *Carya* pollen (up to 35%) and macrofossils document its dominance in the local vegetation. A large number of macrofossils were found by [Martinetto \(2015\)](#) in Italy. At all sites in Italy, *Carya* is one of the main elements of the forest

1052 vegetation during the early Calabrian ([Russo Ermolli, 2010b](#)). Apparently, this considerable expansion of
 1053 *Carya* forests ended first in Southern Italy, with percentages being always <5% since ca. 1.25 Ma ([Joannin et](#)
 1054 [al., 2008](#)), then in Central Italy, where low frequencies are found since approx. 1.1 ka at the beginning of the
 1055 Jaramillo subchrone ([Bertini, 2000](#)), and soon after in Northern Italy ([Ravazzi, 2003](#); [Muttoni et al., 2007](#)). In
 1056 Northeastern Iberia, *Carya* was still well represented at the Early to Middle Pleistocene transition (Banyoles:
 1057 [Geurts, 1979](#)). At Tenaghi Philippon and at Site 380, *Carya* was always present in low amounts during the
 1058 Early Pleistocene.

1059 The last occurrences of *Carya* in S Europe are recorded during the Middle Pleistocene. In France, it was
 1060 recorded at the Alleret maar between 700 and 550 ka ([Pastre et al., 2007](#); [Degeai et al., 2013](#)), but it was not
 1061 any more present in the Praclaux interglacial, correlated with MIS 11 ([Reille et al., 2000](#)). An unpublished
 1062 record of *Carya* (Suc et al., in progress) identified in Southern France up to the Last Glacial ([Biltekin et al.,](#)
 1063 [2015](#)) needs to be better documented.

1064 Macrofossils of *Carya*, reported from the site of Oriolo in N Italy (ca. 800 ka; [Martinetto and Sami, 2001](#)),
 1065 more or less coeval to the last finds of *Carya* pollen at Pianico Sèllere ([Rossi, 2003](#); [Ravazzi et al., 2005](#); [2008](#)),
 1066 agree with pollen data from the magnetostratigraphically dated Venice core sediments ([Müllenders et al.,](#)
 1067 [1996](#)). In Central and Southern Italy, *Carya* was still locally present at Torre Mucchia ([Pieruccini et al., 2016](#))
 1068 at the onset of the Middle Pleistocene, as well as at various sites during the forest phase corresponding to
 1069 MIS 15 and 13 (Vallo di Diano: [Russo Ermolli, 1994](#); Ceprano: [Manzi et al., 2010](#)), as also confirmed by a
 1070 macrofossil find at Carsoli ([Sadori et al., 2010](#)). However at the same time it was absent from a number of
 1071 other sites, for example at La Pineta before MIS 15 ([Lebreton, 2002](#)) and at Sessano before MIS 13 ([Russo](#)
 1072 [Ermolli et al., 2010a,b](#)), as well as at Acerno during MIS 13 ([Petrosino et al., 2014](#)), probably because its
 1073 distribution was already rather fragmented. Its last occurrence in the Italian Peninsula is recorded at Boiano
 1074 ([Orain et al., 2013](#)), a site that is administratively located in Southern Italy, but in this paper included Central
 1075 Italy because it located at a latitude > 41°N ([Figs 4 and 7](#)). Its persistence in the Boiano basin may be linked
 1076 to specific local physiographical characters enhancing humid climate conditions, that may have favored the
 1077 establishment of a vegetation refuge ([Orain et al., 2013](#)). At Tenaghi Philippon, *Carya* shows an almost

1078 continuous record until the Early to Middle Pleistocene transition, then it presents only sporadic occurrences
1079 until 200 ka (Tzedakis et al., 2006). A Middle Pleistocene presence of *Carya* in the Balkan peninsula is also
1080 confirmed by its occurrence at in the 500,000 years old record from Lake Ohrid, for which detailed pollen
1081 data have not been published yet (Sadori et al., 2016). In the Black Sea core DSDP 380, *Carya* is
1082 discontinuously recorded until the Holocene (ca. 2 ka), which is the most recent occurrence in Europe and
1083 the Near East (Biltekin et al., 2015).

1084 A few remarks can be made on the Pleistocene history of *Carya* in Southern Europe. First of all, the studies
1085 carried out in the last few years have considerably delayed the age of the last occurrence of *Carya*, both in
1086 the Italian Peninsula (MIS 9) and in Anatolia (Holocene). New published data are expected for Southern
1087 France and Northeastern Spain to define the age of its disappearance. A second observation is that, as also
1088 shown for *Tsuga*, *Carya* populations were very vigorous before their final decline in the Middle Pleistocene
1089 and did not show any sign preluding to the impending extirpation. Interestingly, dense populations lasted
1090 longer in regions (Southern France and Northern Italy) where they disappeared first. In Central and Southern
1091 Italy, as well as at Tenaghi Philippon, they decreased sooner, but lasted longer. So, while *Carya*-dominated
1092 forests ended in the eastern and southern regions of Southern Europe earlier than the Northwestern
1093 Mediterranean areas, the opposite is true for the last occurrences of *Carya* pollen.

1094

1095 4.11 *Pterocarya*

1096 The genus *Pterocarya* (family Juglandaceae) has at present a disjunct distribution, with several species living
1097 in both the western and eastern regions of the Asiatic continent. The species with the distribution closest to
1098 Europe is *Pterocarya fraxinifolia* (Poir.) Spach, living in Caucasus and in the Hyrcanian forests and extending
1099 further west along the Black Sea, with extrazonal populations in S Turkey and in the Zagros Mts (Akhani and
1100 Salimian, 2003). It currently populates flooded forests and valleys with running water, often in association
1101 with *Alnus glutinosa* (Akhani et al., 2010; Sheykhislami and Ahmadi, 2009).

1102 *Pterocarya* was a constituent of evergreen and deciduous mixed forest in Europe, with other riverine taxa,
1103 during the Miocene and Pliocene (Jiménez-Moreno et al., 2008, 2010; Popescu et al., 2010). During the

1104 Pleistocene it was found in Northwestern and Central Europe in interglacial phases, until the Holsteinian
1105 (generally correlated with MIS 11), for which period *Pterocarya* is often considered a characteristic taxon
1106 (Bińka et al., 2003).

1107 In Southern Europe, it is found in most records (Fig. 8), generally in very low frequencies (<2%), but in some
1108 sites it is very abundant, attaining >25% at the Gelasian site of Coste San Giacomo (Bellucci et al., 2014), >10%
1109 at Lefte (Ravazzi, 2003), >60% at Fontana Ranuccio (Corrado and Magri, 2011) in the Calabrian, and >50% in
1110 the Middle Pleistocene site of Riano (Follieri, 1962), where also fossil fruits and leaves were recovered
1111 (Follieri, 1958). This uneven behavior suggests that the abundance of *Pterocarya* was more related to local
1112 hydrological regimes (e.g., flooding), than to general climatic conditions, according to its modern ecological
1113 requirements.

1114 As to the timing of its disappearance, *Pterocarya* became rare in Anatolia by 1.26 Ma and was recorded
1115 around 400 ka for the last time (Biltekin et al., 2015). In Greece, it was present at Tenaghi Philippon (Tzedakis
1116 et al., 2006) and Kopais (Okuda et al., 2001) until MIS 9 (approx. 320 ka). In Central Italy it was found at Valle
1117 di Castiglione (Follieri et al. 1988) until MIS 7 (ca. 200 ka). In the French Massif Central it is reported until MIS
1118 11 (Reille et al. 2000), but traces are found also in MIS 9 (M. Reille, pers. commun., in Tzedakis et al., 2006).
1119 The same age is assigned to a pollen record from the French Alps, where *Pterocarya* attains 10% (Field et al.,
1120 2003). *Pterocarya* is recorded in two marine cores off Portugal: MD01-2447 during MIS 11 (ca. 400 ka Desprat
1121 et al., 2005) and MD03-2697 during MIS 9 (ca. 298 ka: Desprat et al., 2009). In continental Spain, isolated
1122 grains of *Pterocarya* were observed in the lowest part of the Cañizar de Villarquemado sequence, attributed
1123 to MIS 5 (González-Sampériz et al., 2013; García-Prieto, 2015). This is the first time that *Pterocarya* pollen
1124 grains are found in European sequences during the Late Pleistocene. However, since the study is still in
1125 progress, it must be taken with caution.

1126 Summing up this information, *Pterocarya* was more or less present at all European sites with different
1127 densities, depending on the local situation, until 400 ka, after which time its distribution was severely
1128 fragmented causing its disappearance at different times until its last recorded occurrence during, probably,
1129 the Eemian (ca. 120 ka).

1130

1131 4.12. *Zelkova*

1132 The genus *Zelkova* (family Ulmaceae) includes three extant species living in East Asia, one in the Caucasus
1133 and nearby mountain ranges in Turkey and Iran (*Z. carpinifolia* (Pall.) K. Koch), one in Crete (*Z. abelicea* (Lam.)
1134 Boiss.), and one in Sicily (*Z. sicula* Di Pasq., Garfi & Quézel) (Denk and Grimm, 2005). Its modern distribution
1135 is therefore disjunct and is subject to a progressive reduction (Kozłowski et al., 2012), above all in its south-
1136 western Eurasian species, which have haplotypes that are distinctly clustered according to species
1137 assignment (Christie et al., 2014).

1138 One major problem concerning the fossil record of *Zelkova* is the difficulty of identifying its pollen.
1139 Morphological differences with *Ulmus* are found in the ectexine, especially for its marked sculpture and the
1140 thickening around the pores, in the number of pores (mostly 4), and in the sub-quadrangular to square (if 4-
1141 porate) shape of pollen grains (Kuprianova, 1965; Follieri et al., 1986; Nakagawa et al., 1998; Kvavadze and
1142 Connor, 2005). In many publications, *Ulmus* and *Zelkova* pollen types have been lumped together.
1143 Nonetheless, we have taken these records into consideration and represented them (in blue) in Figs 2-5 and
1144 8.

1145 The Pleistocene records from Spain are not conclusive about the regional history of *Zelkova*, as in most cases
1146 its pollen was not counted separately from *Ulmus*. However, the local presence of *Zelkova* in Northeastern
1147 Iberia is documented in the leaf flora of the Early Pleistocene site of Incarcal at Crespià (Roiron, 1983). During
1148 the Middle and Late Pleistocene it is not recorded in Spain any more. In Southern France, during the Gelasian
1149 *Zelkova* is found in the pollen record from Senèze in the Massif Central (Elhaï, 1969), and in the leaf flora
1150 from Nogaret and Bernasso (Leroy and Seret, 1992; Leroy and Roiron, 1996). It is documented in the late
1151 Early Pleistocene record of Ceyssac-8 (Ablin, 1991) and in the Middle Pleistocene sediments from the Alleret
1152 maar (Pastre et al., 2007; Degeai et al., 2013), which attest the last occurrence of *Zelkova* in France during
1153 MIS 15.

1154 In Italy, *Zelkova* shows discontinuous presence in Northern Italy, where well preserved macrofossil finds are
1155 reported from Oriolo around 800 ka (Martinetto and Sami, 2001). The last pollen occurrences in Northern

Italy are reported from the Southeastern Po Plain (Amorosi et al., 2004) and Venice (Müllenders et al., 1996) during the Eemian interglacial (ca. 120 ka). In Central and Southern Italy, modest but continuous presence of *Zelkova* is documented throughout the Pleistocene, until its disappearance. Only a few expansions show frequencies around 10% (Fig. 8): at the end of the Calabrian at Santa Lucia (Joannin et al., 2008), during MIS 13 at Vallo di Diano (Russo Ermolli et al., 2015), at Riano in a forest phase probably corresponding to MIS 11, where also macrofossils of *Zelkova* were found (Follieri, 1958, 1962), and during the Eemian interglacial at Valle di Castiglione near Rome (Follieri et al., 1986, 1988). The last occurrence of *Zelkova* in the Italian Peninsula is documented in the open forest phases corresponding to MIS 3 (somewhat older than 30 ka) at three sites of the Lazio region: Valle di Castiglione (Follieri et al., 1986, 1988), Lago di Vico (Magri and Sadori, 1999), and Lagaccione (Magri, 1999). Such late occurrence of *Zelkova* is supported by the relatively near location of two modern populations of *Z. sicula* in Sicily (Garfi et al., 2011).

Zelkova is recorded in Rhodes by both pollen and macrofossils until approx. 500 ka (Joannin et al., 2007a; Boyd, 2009). After this time there are no other pollen records available for the area that can attest the age of its disappearance. At Tenaghi Philippon, *Ulmus/Zelkova* is documented until 420 ka (Van der Wiel and Wijmstra, 1987a). It is also mentioned as present in the 500 ka-old record from Lake Ohrid (Sadori et al., 2016). *Zelkova* was found during the Eemian interglacial (ca. 120 ka) at Ioannina (Tzedakis et al., 2003). In Southern Greece *Ulmus/Zelkova* is recorded between 650 and 300 ka at Megalopolis (Okuda et al., 2002), and between 500 ka to recent at Kopais (Okuda et al., 2001). This late record is corroborated by the occurrence of *Zelkova* between 4 and 3 ka in the Gulf of Corinth (Fouache et al., 2005) and by the populations of *Z. abelicea* still living in Crete (Kozłowski et al., 2014).

In Site 380, *Zelkova* is continuously recorded up to 1.22 Ma and sporadically up to 150 ka (Biltekin et al., 2015), an age that is consistent with the modern presence of *Z. carpinifolia* in the nearby Western Transcaucasia (Kozłowski et al., 2012).

On the whole, the fossil records from Southern Europe, together with the biogeographical and genetic data available for the extant species, form a fairly accurate reconstruction of the history of *Zelkova* in S Europe, except for the Iberian Peninsula, where either *Zelkova* disappeared very early compared to the other

European regions, or the scanty fossil record available for the Middle Pleistocene has failed to trace its presence. In Italy, in Greece and in the Black Sea, the existence of relatively close population of relict species strongly support the pollen finds dating back to the Late Pleistocene/Holocene.

1185

1186 **5. Timing and trends of tree population extinctions**

1187 The wealth of data available from the Early and Middle Pleistocene pollen records of S Europe allows
1188 discussion of several aspects of long-term dynamics of tree populations, related to the reduction and
1189 extinction of species.

1190 A first observation concerns the general changes of the vegetation in which the extinction processes are
1191 framed. **Figs 2-5** show that in the course of the Pleistocene there was a progressive decrease in forest
1192 vegetation and a corresponding increase in herbaceous and steppe environments. This is especially evident
1193 in the peri-Iberian marine records, in Northeastern Spain-Southern France and in Central-Southern Italy. In
1194 Northern Italy, apart from a time interval of a few hundred thousand years during the Calabrian with
1195 relatively high values of herbs, herbaceous vegetation was never dominant. Even during the glacial periods,
1196 *Pinus* was widespread. In Greece and the Aegean, including Tenaghi Philippon, the general decrease in
1197 forestation is less evident, as the vegetation appears rather open already in the Early Pleistocene. At Site 380,
1198 a progressive decline of trees appears related to the decline of *Taxodium* populations, as *Quercus* does not
1199 show significant changes in abundance along the Pleistocene.

1200 This overall decline of forest vegetation in the Mediterranean region, paralleling the amplitude of Pleistocene
1201 climatic cycles, has been observed and discussed by several authors ([Suc and Popescu, 2005](#); [Fusco, 2007](#);
1202 [Popescu et al., 2010](#), [Bertini, 2010](#), [Magri and Palombo, 2013](#)). Since the onset of the Pleistocene, 40 ka
1203 obliquity-forced glacial-interglacial cycles led to the alternation of forest vegetation and steppes in many
1204 sites of the Mediterranean Basin, although the direct correlation of steppe phases with glacial periods is a
1205 controversial matter ([Capraro et al., 2005](#); [Bertini, 2010](#)). This climate cyclicity determined a strong reduction
1206 of the Pliocene sub-tropical forests, which required high temperatures and precipitation throughout the year,
1207 and led to a corresponding increase in oak-dominated vegetation, with conifer forests at high altitudes

1208 ([Combourieu-Nebout et al., 2015](#)). A further shift toward a more extended periodicity and amplitude of
1209 glacial interglacial cycles, from 41 ka to approximately 100 ka, provoked a further decline of forests and an
1210 overall spread of steppe conditions during the so-called Mid-Pleistocene Revolution (MPR), which occurred -
1211 in its broadest definition - between 1.2 and 0.5 Ma ([Head and Gibbard, 2005](#)). During this transition,
1212 important changes were observed in the European vegetation ([Suc and Popescu, 2005](#); [Tzedakis et al., 2006](#);
1213 [González-Sampériz et al., 2010](#); [Leroy et al., 2011](#); [Combourieu-Nebout et al., 2015](#)), also driving changes in
1214 faunal communities of both large ([Kahlke et al., 2011](#); [Magri and Palombo, 2013](#)) and small mammals ([van](#)
1215 [Kolfschoten and Markova, 2005](#)).

1216 Climate reconstructions based on selected pollen records from southern Italy suggest a general decline in
1217 winter temperature and annual precipitation from the Early Pleistocene to the Holocene, with main changes
1218 around 2 Ma, 1.4–1.3 Ma and 0.5 Ma ([Combourieu-Nebout et al., 2015](#)). Latitudinal and longitudinal
1219 gradients were observed by [Suc and Popescu \(2005\)](#) and [Combourieu-Nebout et al. \(2015\)](#), who detected a
1220 progressive disappearance of thermophilous vegetation mostly from North to South. In addition, the Asiatic
1221 monsoon enhanced a longitudinal temperature gradient favouring the preservation of thermophilous
1222 elements in the eastern Mediterranean areas ([Suc and Popescu, 2005](#)). This important reduction in moisture
1223 availability and/or the increase in seasonal drought intensity may also be a major factor causing the general
1224 reduction of a type of habitat that was originally widely distributed in the Mediterranean, namely flooded
1225 ground habitats and damp woodland, populated by swamp vegetation, which are currently restricted to
1226 limited areas, also following the extensive reclamation works undertaken during the last centuries.

1227 Despite these general vegetational trends in relation to longitudinal and latitudinal gradients, the
1228 composition of vegetation in S Europe during the Pleistocene shows significant regional differences,
1229 reflecting a mosaic of geographic and climatic situations and partly anticipating the modern complexity of
1230 vegetation patterns ([Figs 2-5](#)).

1231 In the marine cores off Portugal and Spain the dominant element was always *Pinus*, usually associated to its
1232 over-representation in marine sediments, despite inner continental sequences also show important values
1233 of this conifer ([Carrión \(Ed.\), 2012](#)). In the early Pleistocene, the abundance of oaks together with *Olea* is

1234 remarkable, as well as the surprising abundance of *Fagus* since the late Early Pleistocene (Joannin et al.,
 1235 2011).

1236 In Northeastern Spain-Southern France, two clear turnovers of dominant vegetation types can be observed:
 1237 during the Gelasian, forests dominated by subtropical elements, including *Symplocos*, *Engelhardia*, *Taxodium*
 1238 type and *Cathaya* type, were replaced by a vegetation dominated by *Tsuga*, *Picea*, *Carya*, *Pterocarya*, *Quercus*
 1239 and *Carpinus*, and increasing frequencies of *Abies*, in turn substituted by *Quercus*, *Picea*, *Abies*, *Carpinus* and
 1240 increasing *Fagus* during the Early to Middle Pleistocene transition. *Pinus* was abundant throughout.

1241 In Northern Italy, conifers were dominant during the Gelasian, including *Sciadopitys*, *Taxodium* type, *Tsuga*,
 1242 *Cedrus*, *Picea*, and *Abies*, together with Juglandaceae (*Carya*, *Pterocarya*, and *Juglans*). This forest
 1243 composition persisted with little changes throughout most of the Calabrian, with only one important
 1244 turnover at the onset of the Middle Pleistocene. This apparently rapid change may also be an effect of the
 1245 fragmentary character of the available records. The Middle Pleistocene vegetation appears much
 1246 impoverished, and dominated by *Pinus*, *Picea*, *Abies*, *Quercus*, *Carpinus* and *Fagus*.

1247 In Central Italy, a sequence of four main vegetation types is evident: (i) during the first half of the Gelasian,
 1248 forests were dominated by *Taxodium* type, *Cedrus*, *Picea*, *Quercus*, and *Carya*, accompanied by typical
 1249 Pliocene taxa, such as *Engelhardia*, *Nyssa* and *Sciadopitys*; (ii) in the second half of the Gelasian, the
 1250 vegetational landscape was dominated by *Cathaya*, *Taxodium* type, *Cedrus*, *Carya*, *Tsuga* and *Quercus*; (iii)
 1251 in the Calabrian, first *Carya* and then *Tsuga* were the dominant trees, in forests with abundant *Cedrus*, *Picea*,
 1252 and *Quercus*; (iv) starting in the Middle Pleistocene, *Abies*, *Quercus*, *Carpinus*, and *Fagus* increased and
 1253 alternately characterized the forest phases. *Pinus* was always abundant.

1254 In Southern Italy, the succession of vegetation types is similar to Central Italy, but with a clear reduction of
 1255 *Liquidambar*, *Tsuga* and *Carya* during the mid-Calabrian, when forests become dominated by *Cedrus*,
 1256 *Quercus* and *Zelkova*. Besides, *Picea* was much less abundant than in Central Italy. During the Middle
 1257 Pleistocene, forests were characterized by *Quercus*, *Abies*, and *Carpinus*, and then also by *Fagus*, increasing
 1258 around 0.5 Ma.

1259 In Greece and the Aegean, the dominant elements of forest vegetation were *Pinus*, *Quercus* and *Abies*
1260 throughout the Quaternary, with significant amount of *Cedrus* and *Liquidambar* during the Calabrian.

1261 At Tenaghi Philippon, *Quercus* was largely dominant since the Early Pleistocene, accompanied by *Pinus* and
1262 significant percentages only of *Tsuga*, *Pterocarya* and *Cedrus* in the Early Pleistocene, and *Picea* and *Abies* in
1263 the Middle Pleistocene.

1264 In Site 380, Taxodiaceae was predominant, declining through time, while *Quercus* dominated the Quaternary
1265 vegetation.

1266 It clearly appears that it is not possible to identify dominant vegetation types and events of vegetation
1267 turnover valid across Southern Europe that can characterize the single chronostratigraphic units of the
1268 Quaternary. The composite pollen diagrams of [Figs 2-5](#) may constitute reference biostratigraphical schemes
1269 only at a regional scale.

1270 Another important aspect emerging from the long composite pollen records is that several tree genera
1271 persisted regionally from the beginning of the Quaternary until the late Pleistocene. They include: *Quercus*
1272 and *Pinus* in all regions; *Picea*, *Carpinus* and *Fagus* in all regions except Greece and Site 380, where they are
1273 rare; *Taxodium* and *Carya* only in site 380. Recognizing the long-term persistence of tree populations and
1274 their demographic history through multiple glacial-interglacial cycles may be especially important to
1275 determine the timing of the genetic diversification of modern tree populations ([Magri, 2010](#)). Interestingly,
1276 recent papers have advanced hypotheses on the timing of genetic divergence in tree populations that are
1277 consistent with the timing of possible bottlenecks, as reconstructed by long pollen records. [Bagnoli et al.](#)
1278 ([2016](#)), exploring the genetic structure of *Quercus cerris*, suggest that the divergence of its Anatolian and
1279 Balkan populations occurred at the beginning of the Gelasian, at the same time as a dramatic reduction in
1280 oaks, followed by a new population increase, observed at Site 380. The same correspondence between the
1281 timing of genetic differentiation and demographic history of plant populations may be recognized in the
1282 populations of *Abies* in Central-Southern Italy, which were found to have diverged from the northern
1283 populations around 600 ka ([Piotti et al., 2016](#)). It is intriguing that the fossil record indicates that the
1284 populations of *Abies* in Central-Southern Italy show a vigorous expansion at approx. 600 ka, after a period of

1285 over one million years with only modest percentages (Figs 3 and 4), suggesting that the observed expansion
1286 was possibly related to the spread of a new, probably genetically different, population.

1287 The demographic history of tree populations at the timescale of the whole Quaternary may open new
1288 perspectives on the extinction of tree taxa. In fact, rather unexpectedly, several trees that underwent
1289 extinction in Southern Europe show widespread diffusion and considerable abundance before their
1290 disappearance. For example, *Carya* and *Tsuga* markedly expanded during the late Early Pleistocene, to a
1291 degree that provides no indication of their impending extirpation during the Middle Pleistocene. *Zelkova*
1292 displays frequencies around 10% during the Eemian interglacial in Central Italy, only a few tens of thousands
1293 of years before its last occurrence during the last glacial period. However, other taxa persist in small and
1294 sparse populations before their final decline. This is the case of *Cathaya* type, *Liquidambar*, and
1295 *Taxodium/Glyptostrobus* that are found in many pollen records with single occurrences for hundreds of
1296 thousands of years before their last occurrence. In synthesis, it appears that the duration of the extinction
1297 process is very variable from one taxon to the other, or from one population to the other.

1298 The timing of extinction of tree taxa in Southern Europe (Figs 6-8) appears very diverse from one region to
1299 the other. In some cases, there are differences of hundreds of thousands of years even between regions close
1300 to each other. For example, *Cathaya* is not found in Northern Italy after 1.5 Ma, while it persists in Southern
1301 France until approx. 600 ka. *Eucommia* lived in Northern Italy until 1.1 Ma, but it probably disappeared from
1302 Central Italy already at the beginning of the Pleistocene. *Engelhardia* lasted in Site 380 until the Middle
1303 Pleistocene, but was not found in any Greek record during the Pleistocene.

1304 Similarly, the regional distribution of tree taxa that went extinct in Southern Europe is very diversified. For
1305 the first time we have highlighted that the persistence of *Sciadopitys* was much longer in Italy than anywhere
1306 else in Europe (Fig. 6). *Eucommia* had its last refugium near Tenaghi Philippon (Fig. 7), *Cathaya* in Southern
1307 France (Fig. 6), *Taxodium* type along the Anatolian coasts (Fig. 6), *Cedrus* in Rhodes (Fig. 6), probably
1308 *Pterocarya* in inner Northern Spain (Fig. 8). The recent advances in phylogeographical studies, showing that
1309 the Southern European peninsulas often host genetically different populations of the same species (e.g.,
1310 Magri et al., 2006; Bagnoli et al., 2009, 2016; Liepelt et al., 2009), suggest that the different timings of

1311 extinction of tree populations may be strongly connected to their genetic characters, enhancing or reducing
1312 their ability to cope with unfavorable climate. This is particularly evident when the behavior of different
1313 populations is regionally very diverse. For example, *Engelhardia* shows very different behaviours in Northern
1314 Spain and Italy, compared to Anatolia. In the case of *Taxodium*, the Eastern population recorded at Site 380
1315 and the western population in Italy and Spain, separated by a region (Greece) where *Taxodium* type was
1316 almost absent, may even indicate the presence of different species/genera.

1317 Figs 6-8 clearly indicate that it is not always possible to detect geographical trends in the extinction of tree
1318 populations and confute the commonplace view that tree populations disappeared first at higher latitudes
1319 and then in the southern regions. For example, *Engelhardia*, *Eucommia*, and *Cathaya* appear to persist longer
1320 in Southern France than in peninsular Italy and Greece. The case of *Parrotia* is even more unexpected, as its
1321 last occurrence in Europe would date back to MIS 11 in Poland (Bińka et al., 2003). This complexity may reflect
1322 the bioclimatic heterogeneity of southern Europe and its mountain ranges, and also highlights the refugial
1323 role of not only Southern Europe but Eastern Europe and connections to Western Asia.

1324 In many cases, the distribution of tree populations on the way to extinction appears fragmented, as
1325 demonstrated by scattered finds in different coeval records. Interestingly, the last occurrences of taxa that
1326 are still living close to continental Europe (e.g., *Zelkova*, *Liquidambar* and *Cedrus*) are found relatively near
1327 to their modern core distribution ranges. *Zelkova*, still living in Sicily, Crete and in the Hyrcanian region, was
1328 last found in the pollen records from Central Italy, Greece, and Black Sea, respectively. *Liquidambar*, still living
1329 in Southwestern Turkey, had its last appearances in Rhodes and Site 380. Detecting the last fragmented
1330 populations of *Cedrus* is very difficult, but again the macrofossil find from Rhodes (Boyd, 2009) documents
1331 with certainty its late presence close to its extant range in the Eastern Mediterranean. However, if the most
1332 recent record of *Pterocarya* in Northeastern Spain will be confirmed, in a location very far from the present
1333 distribution of *Pterocarya* in the Hyrcanian region, it is also possible that in some cases disjunct populations
1334 may have persisted completely isolated from their distribution center.

1335 The very difficult question of whether the disappearance of tree populations was gradual or step-wise, in
1336 relation to specific climatic events, is probably far from being solved. In many cases, it appears that the

1337 contemporary extinction of more than one taxon in an area is due to the fragmentary nature of the fossil
1338 record. For example, in Southern Italy, the lack of data between MIS 13 and MIS 5 inevitably produces the
1339 effect of several taxa disappearing at the end of MIS 13. Thus, the availability of long continuous sequences
1340 is absolutely necessary to disentangle this issue. The long record from Tenaghi Philippon (Fig. 5) and the
1341 composite, but rather complete, record from C Italy (Fig. 3) do not show any specific time interval in which
1342 more than one last occurrence is found. Extirpations appear distributed over time, even though broad
1343 changes in dominant forest taxa affect the abundance of multiple species. These patterns appear to reflect
1344 the individualistic species potential to survive beyond the most favourable climatic intervals, through
1345 physiological plasticity, evolutionary adaptation and availability of microclimatic refugia.

1346 It would seem obvious to relate plant extinctions to the harsh dry and cold climate conditions of the glacial
1347 periods (Tzedakis et al., 2006), but the exact timing of extirpation is very difficult to be assessed (see
1348 discussion above) such that we cannot easily reconstruct the climate conditions at the time of extirpation.
1349 While the largest glaciations, such as MIS 22 and MIS 16, seem to play a role in conditioning the change in
1350 forest composition (Tzedakis et al., 2006; Pieruccini et al., 2016), and probably enhanced a general
1351 impoverishment of the flora, individual extinctions are not directly linked to these. One possible way to solve
1352 the problem is to investigate the extirpation modes during the last glacial-interglacial cycle, a time period for
1353 which numerous detailed and well-dated records are available.

1354 A number of tree taxa can be found that have strongly reduced their distribution range during the last tens
1355 of thousands of years in Southern Europe:

- 1356 • *Zelkova* was last recorded in Central Italy during one of the final fluctuations of the pleniglacial
1357 interstadials (MIS 3), suggesting its possible extirpation during the last glacial phase (MIS 2).
1358 However, *Zelkova* was last recorded in Greece 4-3 ka ago, when climate conditions were possibly not
1359 unfavourable to its survival (Fouache et al., 2005).
- 1360 • *Glyptostrobus* (*Taxodium* type) and *Carya* may have persisted in northern Anatolia until the Holocene
1361 and disappeared in the last few thousand of years (Biltekin et al., 2015).

- *Chamaerops humilis* was last recorded at Santorini (macrofossil evidence, [Velitzelos et al., 2014](#)) during the Late Pleistocene, pointing to a recent demise of this taxon, which currently has a highly fragmented pan-Mediterranean distribution.
- *Buxus* (including *B. balearica* and *B. sempervirens*) retreated from the southern regions of the Iberian, Italian and Balkan peninsulas during the late Holocene, at the same time of its exponential increase in temperate Europe, thus challenging the common view that the Mediterranean regions are the exclusive and most important refuge areas for evergreen broadleaved trees. During the last 4500 years, populations located in Minorca, Ibiza, Sardinia, Sicily, Apulia, Basilicata, Croatia, Thessaly, W Greece, and Peloponnese have been completely extirpated ([Di Domenico et al., 2012](#)). In other areas in Southern Europe a process of severe reduction is still going on, resulting in the modern highly fragmented distribution area, which is a natural residue of a wider distribution. This reduction is even more striking considering that in these regions *Buxus* persisted throughout the last glacial period.
- *Carpinus betulus* was commonly traced in the Iberian Peninsula during the Middle Pleistocene ([Desprat et al., 2005, 2007](#)). Fossil evidence for *Carpinus* is still quite common in the Late Pleistocene ([Postigo-Mijarra et al., 2008, 2009, 2010; Carrión \(Ed.\), 2012](#)), followed by extreme reduction during the Holocene ([Postigo-Mijarra et al., 2008](#)). The marine sequence MD95-2042 from the Portuguese margin ([Sánchez-Góñi et al. 1999, 2008](#)) records *Carpinus* during the last interglacial Eemian (MIS 5e), and the Area Longa sequence ([Gómez-Orellana et al., 2007](#)) during MIS 5c and MIS 4, as well as Sta. Maria de Oia ([Gómez-Orellana et al., 2012](#)) and Abric Romaní ([Burjachs et al., 2012](#)) during MIS 3. The presence of *Carpinus* in Pla de l'Estany-Garrotxa ([Burjachs, 1990](#)), despite weak chronological control, is also attributed by the author to the Eemian. In the Cañizar de Villarquemado sequence ([González-Sampériz et al., 2013; García-Prieto, 2015](#)), *Carpinus* is recorded in the lower part of the sequence, attributed to MIS 5 ([García-Prieto, 2015](#)). During the Holocene, *Carpinus* populations revealed a progressive depletion, leading to its near disappearance at the Iberian scale: it is sporadically found in records confined to the Cantabrian Mountains (e.g., Lagoa de Lucenza, [Muñoz-Sobrinho et al., 2001](#); Chan do Lamoso, [Ramil-Rego et al., 1998](#); La Roya, [Allen et al., 1996](#); Bidasoa,

Sánchez-Goñi, 1996; Urdaibai, Iriarte-Chiapusso et al., 2016), to the pre-Pyrenean Range (Estanya Lake: González-Sampériz et al., in press), along the Iberian Range (e.g., Las Pardillas, Sánchez-Goñi and Hannon; Moncayo Massif, Aranbarri et al., 2016), as well as in the Central System (Abel-Schaad et al., 2014). In the Mediterranean realm, sequences from the Balearic Islands (e.g., Cala Galdana and Cala'n Porter, Yll et al., 1997), together with those located from the continental environments (e.g., Espinosa del Cerrato, Franco-Mugica et al., 2001), reveal the latest presences of *Carpinus* from inland Iberia between 7 and 4 ka BP. The increased vulnerability of *Carpinus* communities through glacial-interglacial cycles, together with ecological factors (e.g. competition), may have caused the demise of this taxon, whose extirpation from the Iberian Peninsula (excepting a small area in the western Pyrenees, Aizpuru and Catalán, 1984) is still poorly understood.

- *Picea*, widespread in Southern Europe during the Early Pleistocene (Figs 2-5), underwent a marked reduction in most southern regions. In the western Balkans at Lake Ohrid, *Picea* is recorded in low, discontinuous abundances during the last glacial, and generally absent in the Holocene (Sadori et al., 2016). *Picea* is scarcely recorded in Greece from the Middle Pleistocene onwards, being absent in the Middle Pleistocene diagrams of Ioannina, Kopais and Megalopolis Basin (Tzedakis, 1994, 1999; Okuda et al., 2001, 2002), and similarly not recorded in the late Upper Pleistocene and Holocene diagrams of Ioannina (Lawson et al., 2004), Nisi Fen (Lawson et al., 2005) or Northern Aegean marine core GeoTü SL152 (Kotthoff et al., 2008). In Southern Italy, the last significant expansion of *Picea* (>5%) is found at the onset of the Middle Pleistocene (Capraro et al., 2005), but frequent occurrences of single grains are recorded until the Last Glacial Maximum (Allen et al., 2000). In Central Italy, *Picea* was abundant in several periods throughout the Pleistocene, its last important expansion (>20%) being recorded at Boiano (administratively in Southern Italy) during MIS 9 (Orain et al., 2015). Then it decreased and was last found during the Last Glacial Maximum (Follieri et al., 1998). In Northern Italy, *Picea* is widespread in the Alps, but its populations located in the northern Apennines were strongly reduced during the postglacial warming and were not able to compete with more vigorous tree species (Ravazzi et al., 2006; Magri et al., 2015). In France, the pollen records from the Massif central (Reille et al., 2000) show abundant presence of *Picea* during all interglacial phases, except the

Holocene, when it failed to expand after the glacial period. In Spain, *Picea* is recorded both in the Area Longa pollen spectra during MIS 4 (Gómez-Orellana et al., 2007), and in charcoal remains from La Pila, Santa Catalina and Peña del Perro archaeological sites until the early Holocene (Uzquiano, 1995). On the whole, these data indicate that a declining trend, started in the southern regions of Southern Europe during the Middle Pleistocene, was intensified during the last glacial period and especially at the onset of the Holocene, a phase of global warming, which the most southern populations of *Picea* in Europe were not able to withstand.

These documented cases of extirpations that occurred during the Holocene prevent ascribing plant extinctions directly to the adverse climate conditions of the glacial periods, although it cannot be excluded that in some cases a legacy of glacial bottlenecks and range contractions may ultimately lead to eventual extirpations during interglacial periods, with a considerable delay (Tzedakis et al., 2006). On the other hand, these examples of recent contraction of distribution ranges need to be interpreted in the light of long-term population dynamics (10^5 - 10^6 years), as recognized from fossil data. This may be important to evaluate correctly the vulnerability of modern fragmented plant populations, especially in view of conservation actions, and to reach a better understanding of the causes and mechanisms that drive the reduction of tree populations and their eventual extirpation or even extinction.

6. Conclusions

This review on the extinction of tree populations in Southern Europe, based on the collection, organization and interpretation of a large number of pollen and macrofossil records over the Iberian Peninsula, Southern France, the Italian Peninsula, Greece and the Aegean, and the southwestern Black Sea area, has raised a number of methodological issues and highlighted an overall complexity in the timing and modes of plant extinctions.

From the methodological point of view, although fossil records have many inherent uncertainties that are accentuated going back in time to the Middle and Early Pleistocene (e.g., fragmentation, weak chronological setting, reworking and long-distance transport of pollen, lack of modern analogues, taxonomic uncertainties,

1441 bad preservation state, impossibility to determine the exact timing of extirpation), the overall history of tree
1442 populations is strikingly consistent. This demonstrates that in spite all uncertainties, pollen records provide
1443 excellent information even on fragmentary populations on the way to extinction.

1444 Concerning Pleistocene pollen records from the Mediterranean, two main shortcomings have emerged that
1445 may need to be taken in charge by the scientific community. The first one is the habit of publishing pollen
1446 diagrams including only selected taxa, thus omitting important information on rare occurrences of plant
1447 populations undergoing extinction. A key message to the community is the vital importance of archiving full
1448 pollen data in public repositories, especially in this era of generally concise publication of only summary or
1449 partial pollen datasets. The second one is the need for better photographic atlases and pollen descriptions
1450 that may help palynologists recognizing pollen types of plants that are extinct or living far away. This would
1451 help to align pollen identification by different palynologists.

1452 In many cases, the Quaternary distribution of taxa currently absent from the European mainland,
1453 reconstructed through a detailed analysis of the published data, showed unexpected patterns, indicating that
1454 although their extinction in Europe during the Quaternary was common knowledge, the timing and
1455 geographical trends of their disappearance was far from fully known. The main novel aspects highlighted by
1456 this review are the following:

- 1457 • *Taxodium/Glyptostrobus* probably had at least two distinct populations/species/genera in Southern
1458 Europe, which had different fates. One population disappeared from Northern Spain-Southern France
1459 and Central Italy at the end of the Gelasian, persisted in Northern and Southern Italy until the mid-
1460 Calabrian, and disappeared definitely from Central and Southern Italy after ca. 500 ka. The second one
1461 was distributed in Anatolia-Rhodes; it underwent a first severe reduction around 1.2 Ma, and a second
1462 reduction around 500 ka, but was able to survive until the Holocene.
- 1463 • *Sciadopitys* had a long-term Quaternary refuge in the Italian Peninsula, where it persisted for over one
1464 million years after it had disappeared from the rest of Europe.
- 1465 • *Cathaya*, present at all sites during the Gelasian, during the Calabrian was more common in Southern
1466 Italy than in other regions. However, its last occurrence in Italy dates to ca. 850 ka, while small relict

1467 populations may have persisted in isolated locations in France and Anatolia until 700 ka and 380 ka,
 1468 respectively.

- 1469 • *Cedrus* was widespread in Italy and in the Aegean region during the Early Pleistocene. The interpretation
 1470 of discontinuous single pollen grains in the Iberian Peninsula and Southern France during the Early
 1471 Pleistocene, and in Southern-Central Italy during the Middle Pleistocene, is very uncertain. The age of
 1472 the disappearance of *Cedrus* from southern Europe necessarily requires the support of new macrofossil
 1473 data.
- 1474 • *Tsuga* has a rather homogenous history in different regions of Southern Europe, with sparse presence in
 1475 the most southern sites (Alboran Sea, Rhodes), and significant abundances in Northeastern Spain-
 1476 Southern France and the Italian Peninsula during the Early Pleistocene, especially during the Calabrian.
 1477 Its rapid and definitive disappearance is recorded around 750 ka in Northern Italy, around 500 ka in
 1478 Southern France and Central Southern Italy, and 280 ka in Anatolia.
- 1479 • *Eucommia* persisted in central Europe at least as long as in Southern France, until the late Calabrian. It
 1480 expanded in Northern Italy during the Early Pleistocene, without reaching Central Italy, where it was
 1481 never found.
- 1482 • *Parrotia* is not found in Southern Europe after ca. 800 ka, which is surprising considering its continuous
 1483 record during the Holsteinian in Eastern Poland. This result may suggest different hypotheses on its
 1484 persistence areas and Mid-Pleistocene migration routes into Eastern Central Europe.
- 1485 • *Liquidambar*, showing scattered populations in various regions of Southern Europe until the end of the
 1486 Early Pleistocene, survived longer in Southern France, Southern Italy, and the coasts of the Black Sea,
 1487 before being reduced to the small populations currently living in Rhodes and Southeastern Turkey.
- 1488 • *Engelhardia* persisted longer in Northeastern Spain/Southern France (until ca. 1 Ma) than in the Italian
 1489 Peninsula, where it was probably split into two separate populations: one in Southern Italy, persisting
 1490 until approx. 1.35 Ma, and one in Northern Italy, lasting until the early Calabrian, separated by a large
 1491 region in Central Italy where *Engelhardia* was absent. A separate population, lasting until ca. 200 ka, was
 1492 located in the Black Sea region.

- 1493 • *Carya* populations were very vigorous in Italy and France before their decline in the Middle Pleistocene,
1494 and did not show any sign prelude to the impending extirpation. The studies carried out in the last few
1495 years have considerably delayed the age of the last occurrence of *Carya*, both in the Italian Peninsula
1496 (MIS 9) and in Anatolia (Holocene).
- 1497 • *Pterocarya* was more or less present at all European sites with different densities, depending on local
1498 situations, until ca. 400 ka, after which time its distribution was severely fragmented. The disappearance
1499 of *Pterocarya* occurred at different times in different regions. Its last recorded occurrence was possibly
1500 during the Eemian (ca. 120 ka) in Spain.
- 1501 • *Zelkova* pollen was not always distinguished from *Ulmus*. Even so, the fossil records from Southern
1502 Europe, together with the biogeographical and genetic data available for the extant species, provide a
1503 fairly accurate reconstruction of the history of *Zelkova* in Southern Europe, except for the Iberian
1504 Peninsula, where either *Zelkova* disappeared very early compared to the other European regions, or the
1505 scanty fossil record available for the Middle Pleistocene has failed to trace its presence.
- 1506 In addition to these “classical” extinct trees in Europe, a number of taxa have been considered that are still
1507 widespread in Europe, but have undergone fragmentation/reduction/extirpation in wide areas of Southern
1508 Europe during the late Pleistocene-Holocene. They may provide useful insights into the patterns and timing
1509 of range contraction of plant populations, given the good chronological control of their last occurrences and
1510 the large number of pollen records involved. Three such cases appear of particular interest:
- 1511 • *Buxus* retreated from the southern regions of the Iberian, Italian and Balkan peninsulas during the last
1512 4.5 ka, following a general aridification of climate in the Mediterranean Basin. Considering the very low
1513 dispersal ability and growth rate of *Buxus*, and the current extent of human impact, there is very little
1514 chance that the southern territories of Europe will be naturally recolonized in the future, so the southern
1515 extirpation of *Buxus* can be considered permanent.
- 1516 • *Carpinus betulus*, commonly traced in the Iberian Peninsula during the Middle Pleistocene, was still quite
1517 common in the Late Pleistocene. During the Holocene, *Carpinus* populations revealed a progressive
1518 depletion in its southwestern range, leading to its virtual absence at the Iberian scale, excepting a small
1519 area in the western Pyrenean valleys.

• *Picea*, widespread in Southern Europe during the Early Pleistocene, underwent a marked reduction leading to its disappearance in the most southern regions: in Greece, it was absent in many sites since the Late Pleistocene, in Central-Southern Italy it disappeared during the last glacial maximum, in Northern Spain it was last found in the lateglacial period.

These recent extirpation processes are quite unexpected for three main reasons. Firstly, species like *Buxus sempervirens*, *Carpinus betulus*, and *Picea abies* are not at all considered at risk in Europe. Secondly, they contradict the common view that plant populations persist longer in the southern regions of Europe and, together with indications of complex spatial patterns, rule out a simple North to South trend in extirpations. Thirdly, they occurred at the onset or during the current interglacial, while extinctions are generally considered to be enhanced by the adverse climate conditions of the glacial. This might suggest human impact as a possible cause for the reduction of these taxa. On the other hand, extirpations appear distributed over time during the Quaternary, while it is apparent that significant turnovers of the dominant species have occurred in relation to important climatic/geological transitions, such as at the Gelasian-Calabrian transition, and the so-called Middle Pleistocene Revolution.

Future studies are expected to better define the timing and trends of plant extirpation through the study of new long continuous records, especially from regions that are scarcely known (e.g., the northern Balkan Peninsula and inner Iberia), the re-analysis of some “classical” data with the aim to improve/provide robust chronologies for some undated deposits, so providing better chronological constraints on extinctions, and the publication of all the available data in public repositories. Dealing with extirpations, the most challenging field of research to be explored is certainly the combination of palaeobotanical and genetic studies in relation to the demographic histories of taxa, aimed to assess not only the timing and modes of species/population differentiation, but also the mechanisms of plant extinction.

7. Acknowledgments

This paper was in part supported by the projects CGL2012-33063 and CGL2015-69160-R, provided by the Spanish Inter-Ministry Commission of Science and Technology (CICYT).

8. References

- Abbazzi, L., Fanfani, F., Ferretti, M.P., Rook, L., Cattani, L., Masini, F., Mallegni, F., Negrino, F., Tozzi, C., 2000. New human remains of archaic *Homo sapiens* and Lower Palaeolithic industries from Visogliano (Duino Aurisina, Trieste, Italy). *Journal of Archaeological Science* 27, 1173–1186. doi:10.1006/jasc.1999.0541
- Abel-Schaad, D., Pulido, F., López Sáez, J.A., Alba Sánchez, F., Nieto Lugilde, D., Franco Múgica, F., Pérez Díaz, S., Ruiz Zapata, M.B., Gil García, M.J., Dorado Valiño, M., 2014. Persistence of tree relicts in the Spanish Central System through the Holocene. *Lazaroa* 35. doi:10.5209/rev_LAZA.2014.v35.41932
- Ablin, D., 1991. Analyse pollinique des dépôts lacustres de Ceysac, Plio-Pléistocène du Velay (Massif central, France). *Cahiers de micropaléontologie* 6, 21–38.
- Aizpuru, I., Catalán P., 1984. The presence of the bombean in the Iberian Península. *Anales Jard. Bot. Madrid* 41(I), 143–146.
- Akhani, H., Djamali, M., Ghorbanalizadeh, A., Ramezani, E., 2010. Plant biodiversity of Hyrcanian relict forests, N Iran: an overview of the flora, vegetation, palaeoecology and conservation. *Pakistan Journal of Botany* 42, 231–258.
- Akhani, H., Salimian, M., 2003. An extant disjunct stand of *Pterocarya fraxinifolia* (Juglandaceae) in the central Zagros Mountains, W Iran. *Willdenowia* 33, 113–120. doi:10.3372/wi.33.33111
- Allen, J.R., Huntley, B., Watts, W.A., 1996. The vegetation and climate of northwest Iberia over the last 14,000 years. *Journal of Quaternary Science* 11, 125–147.
- Allen, J.R., Watts, W.A., Huntley, B., 2000. Weichselian palynostratigraphy, palaeovegetation and palaeoenvironment; the record from Lago Grande di Monticchio, southern Italy. *Quaternary International* 73, 91–110.
- Allen, J.R.M., Huntley, B., 2009. Last Interglacial palaeovegetation, palaeoenvironments and chronology: a new record from Lago Grande di Monticchio, southern Italy. *Quaternary Science Reviews* 28, 1521–1538. doi:10.1016/j.quascirev.2009.02.013
- Amorosi, A., Colalongo, M., Fiorini, F., Fusco, F., Pasini, G., Vaiani, S., Sarti, G., 2004. Palaeogeographic and palaeoclimatic evolution of the Po Plain from 150-ky core records. *Global and Planetary Change* 40, 55–78. doi:10.1016/S0921-8181(03)00098-5
- Antunes, M.T., Mein, P., Nascimento, A., Pais, J., 1986. Le gisement pleistocène de Morgadinho, en Algarve. *Ciências da Terra* 8, 9–22.
- Aranbarri, J., Bartolomé, M., Alcolea, M., Sancho, C., Celant, A., González-Sampériz, P., Arenas, C., Magri, D., Rodríguez-Lázaro, J., 2016. Palaeobotanical insights from Early-Mid Holocene fluvial tufas in the Moncayo Natural Park (Iberian Range, NE Spain): Regional correlations and biogeographic implications. *Review of Palaeobotany and Palynology* 234, 31–43.
- Aranbarri, J., González-Sampériz, P., Valero-Garcés, B., Moreno, A., Gil-Romera, G., Sevilla-Callejo, M., García-Prieto, E., Di Rita, F., Mata, M.P., Morellón, M., Magri, D., Rodríguez-Lázaro, J., Carrión, J.S., 2014. Rapid climatic changes and resilient vegetation during the Lateglacial and Holocene in a continental region of south-western Europe. *Global and Planetary Change* 114, 50–65. doi:10.1016/j.gloplacha.2014.01.003
- Argant, J., 2004. Le gisement pliocène final de Saint-Vallier (Drôme, France): palynologie. *Geobios* 37, S81–S90.

1590 Arias, C., Bigazzi, G., Bonadonna, F., Iaccarino, S., Urban, B., Dal Molin, M., Dal Monte, L., Martolini,
1591 M., 1990. Valle Ricca Late Neogene stratigraphy (Lazio region, Central Italy). *Paléobiologie*
1592 *continentale* 17, 61–80.

1593 Bagnoli, F., Tsuda, Y., Fineschi, S., Bruschi, P., Magri, D., Zhelev, P., Paule, L., Simeone, M.C.,
1594 González-Martínez, S.C., Vendramin, G.G., 2016. Combining molecular and fossil data to
1595 infer demographic history of *Quercus cerris*: insights on European eastern glacial refugia. *J.*
1596 *Biogeogr.* 43, 679–690. doi:10.1111/jbi.12673

1597 Bagnoli, F., Vendramin, G.G., Buonamici, A., Doulis, A.G., González-Martínez, S.C., La Porta, N.,
1598 Magri, D., Raddi, P., Sebastiani, F., Fineschi, S., 2009. Is *Cupressus sempervirens* native in
1599 Italy? An answer from genetic and palaeobotanical data. *Molecular Ecology* 18, 2276–2286.
1600 doi:10.1111/j.1365-294X.2009.04182.x

1601 Bell, B.A., Fletcher, W.J., 2016. Modern surface pollen assemblages from the Middle and High Atlas,
1602 Morocco: insights into pollen representation and transport. *Grana* 1–16.

1603 Bellucci, L., Bona, F., Corrado, P., Magri, D., Mazzini, I., Parenti, F., Scardia, G., Sardella, R., 2014.
1604 Evidence of late Gelasian dispersal of African fauna at Coste San Giacomo (Anagni Basin,
1605 central Italy): Early Pleistocene environments and the background of early human
1606 occupation in Europe. *Quaternary Science Reviews* 96, 72–85.
1607 doi:10.1016/j.quascirev.2013.10.011

1608 Ber, A., 2005. Polish Pleistocene stratigraphy-A review of interglacial stratotypes. *Netherlands*
1609 *Journal of Geosciences-Geologie en Mijnbouw* 84, 61–76.

1610 Bertini, A., 2010. Pliocene to Pleistocene palynoflora and vegetation in Italy: State of the art.
1611 *Quaternary International* 225, 5–24. doi:10.1016/j.quaint.2010.04.025

1612 Bertini, A., 2001. Pliocene climatic cycles and altitudinal forest development from 2.7 Ma in the
1613 northern Apennines (Italy): evidence from the pollen record of the Stirone section (~ 5.1
1614 to~ 2.2 Ma). *Geobios* 34, 253–265.

1615 Bertini, A., 2000. Pollen record from Colle Curti and Cesi: Early and Middle Pleistocene mammal sites
1616 in the Umbro–Marchean Apennine mountains (central Italy). *Journal of Quaternary Science*
1617 15, 825–840.

1618 Bertini, A., Magi, M., Mazza, P.P.A., Fauquette, S., 2010. Impact of short-term climatic events on
1619 latest Pliocene land settings and communities in Central Italy (Upper Valdarno basin).
1620 *Quaternary International* 225, 92–105. doi:10.1016/j.quaint.2009.05.011

1621 Bertini, A., Martinetto, E., 2011. Reconstruction of vegetation transects for the Messinian–
1622 Piacenzian of Italy by means of comparative analysis of pollen, leaf and carpological
1623 records. *Palaeogeography, Palaeoclimatology, Palaeoecology* 304, 230–246.

1624 Bertini, A., Roiron, P., 1997. Evolution de la végétation et du climat pendant le Pliocène moyen, en
1625 Italie centrale: apport de la palynologie et de la macroflore à l'étude du bassin du Valdarno
1626 supérieur (coupe de Santa Barbara). *Comptes rendus de l'Académie des sciences. Série 2.*
1627 *Sciences de la terre et des planètes* 324, 763–771.

1628 Bertini, A., Toti, F., Marino, M., Ciaranfi, N., 2015. Vegetation and climate across the Early–Middle
1629 Pleistocene transition at Montalbano Jonico, southern Italy. *Quaternary International* 383,
1630 74–88. doi:10.1016/j.quaint.2015.01.003

- 1631 Bertoldi, R., 1988. Una sequenza palinologica di età rusciniiana nei sedimenti lacustri basali del
1632 bacino di Aulla-Olivola (Val di Magra). *Rivista Italiana di Paleontologia e Stratigrafia* 94,
1633 105–138.
- 1634 Bhagwat, S.A., Willis, K.J., 2008. Species persistence in northerly glacial refugia of Europe: a matter
1635 of chance or biogeographical traits? *Journal of Biogeography* 35, 464–482.
1636 doi:10.1111/j.1365-2699.2007.01861.x
- 1637 Biltekin, D., 2010. Vegetation and climate of north Anatolian and north Aegean region since 7 Ma
1638 according to pollen analysis. PhD Thesis, Université Claude Bernard-Lyon I; Université
1639 Technique d'Istanbul.
- 1640 Biltekin, D., Popescu, S.-M., Suc, J.-P., Quézel, P., Jiménez-Moreno, G., Yavuz, N., Çağatay, M.N.,
1641 2015. Anatolia: A long-time plant refuge area documented by pollen records over the last
1642 23million years. *Review of Palaeobotany and Palynology* 215, 1–22.
1643 doi:10.1016/j.revpalbo.2014.12.004
- 1644 Bińka, K., Nitychoruk, J., Dzierżek, J., 2003. *Parrotia persica* C.A.M. (Persian witch hazel, Persian
1645 ironwood) in the Mazovian (Holsteinian) Interglacial of Poland. *Grana* 42, 227–233.
1646 doi:10.1080/00173130310016220
- 1647 Birks, H.H., Birks, H.J.B., 2000. Future uses of pollen analysis must include plant macrofossils. *Journal*
1648 *of biogeography* 27, 31–35.
- 1649 Bottema, S., 1974. Late Quaternary vegetation history of northwestern Greece. Ph.D. Thesis,
1650 University of Groningen.
- 1651 Bou Dagher-Kharat, M., Mariette, S., Lefèvre, F., Fady, B., Grenier-de March, G., Plomion, C.,
1652 Savouré, A., 2007. Geographical diversity and genetic relationships among *Cedrus* species
1653 estimated by AFLP. *Tree Genetics & Genomes* 3, 275–285. doi:10.1007/s11295-006-0065-
1654 x
- 1655 Boyd, A., 2009. Relict conifers from the mid-Pleistocene of Rhodes, Greece. *Historical Biology* 21, 1–
1656 15. doi:10.1080/08912960903033301
- 1657 Brenac, P., 1984. Végétation et climat de la Campanie du Sud (Italie) au Pliocène final d'après
1658 l'analyse pollinique des dépôts de Camerota. *Ecologia mediterranea* 10, 207–216.
- 1659 Brugal, J.-P., Ambert, P., Bandet, Y., Leroy, S., Roiron, P., Suc, J.-P., Vernet, J.-L., 1990. Mammifères
1660 et végétaux du maar Pliocène final de Nogaret (Escandorgue: Hérault, France). *Geobios* 23,
1661 231–247.
- 1662 Burjachs, F., 1990. Palinologia dels dòlmens de l'Alt Empordà i dels dipòsits quaternaris de la cova
1663 de l'Arbreda (Serinyà, Pla de l'Estany) i del Pla de l'Estany (Olot, Garrotxa). *Evolució del*
1664 *paisatge vegetal i del clima des de fa més de 140.000 anys al NE de la Península Ibèrica*.
1665 Ph.D. Thesis, Universitat Autònoma de Barcelona, Bellaterra, 324 pp.
- 1666 Burjachs, F., Julià, R., 1994. Abrupt climatic changes during the last glaciation based on pollen
1667 analysis of the Abric Romani, Catalonia, Spain. *Quaternary Research* 42, 308–315.
- 1668 Burjachs, F., López-García, J.M., Allué, E., Blain, H.-A., Rivals, F., Bennàsar, M., Expósito, I., 2012.
1669 Palaeoecology of Neanderthals during Dansgaard-Oeschger cycles in northeastern Iberia
1670 (Abric Romani): from regional to global scale. *Quaternary International* 247, 26–37.
- 1671 Capraro, L., Asioli, A., Backman, J., Bertoldi, R., Channell, J.E.T., Massari, F., Rio, D., 2005. Climatic
1672 patterns revealed by pollen and oxygen isotope records across the Matuyama-Brunhes

- 1673 Boundary in the central Mediterranean (southern Italy). Geological Society, London, Special
1674 Publications 247, 159–182.
- 1675 Caratini, C., M., Sivak, J., 1972. Pollen de *Cathaya* (Abietaceae) au tertiaire en France. Pollen et
1676 spores 14, 169–172.
- 1677 Carrión, J., Finlayson, C., Fernandez, S., Finlayson, G., Allue, E., López-Sáez, J.A., López-García, P., Gil-
1678 Romera, G., Bailey, G., González-Sampériz, P., 2008. A coastal reservoir of biodiversity for
1679 Upper Pleistocene human populations: palaeoecological investigations in Gorham's Cave
1680 (Gibraltar) in the context of the Iberian Peninsula. Quaternary Science Reviews 27, 2118–
1681 2135. doi:10.1016/j.quascirev.2008.08.016
- 1682 Carrión, J.S., editor, 2012. Paleoflora y Paleovegetación de la Península Ibérica e Islas Baleares:
1683 Plioceno-Cuaternario, 1st edition. ed. Murcia.
- 1684 Carrión, J.S., Fernández, S., González-Sampériz, P., Gil-Romera, G., Badal, E., Carrión-Marco, Y.,
1685 López-Merino, L., López-Sáez, J.A., Fierro, E., Burjachs, F., 2010. Expected trends and
1686 surprises in the Lateglacial and Holocene vegetation history of the Iberian Peninsula and
1687 Balearic Islands. Review of Palaeobotany and Palynology 162, 458–475.
1688 doi:10.1016/j.revpalbo.2009.12.007
- 1689 Carrión, J.S., Fuentes, N., González-Sampériz, P., Sánchez Quirante, L., Finlayson, J.C., Fernández, S.,
1690 Andrade, A., 2007. Holocene environmental change in a montane region of southern
1691 Europe with a long history of human settlement. Quaternary Science Reviews 26, 1455–
1692 1475. doi:10.1016/j.quascirev.2007.03.013
- 1693 Carrión, J.S., Leroy, S.A., 2010. Iberian floras through time: Land of diversity and survival. Review of
1694 Palaeobotany and Palynology 162, 227–230.
- 1695 Carrión, J.S., Sánchez-Gómez, P., Mota, J.F., Yll, R., Chaín, C., 2003. Holocene vegetation dynamics,
1696 fire and grazing in the Sierra de Gádor, southern Spain. The Holocene 13, 839–849.
1697 doi:10.1191/0959683603hl662rp
- 1698 Carrión, J.S., van Geel, B., 1999. Fine-resolution Upper Weichselian and Holocene palynological
1699 record from Navarrés (Valencia, Spain) and a discussion about factors of Mediterranean
1700 forest succession. Review of Palaeobotany and Palynology 106, 209–236.
- 1701 Cattani, L., Cremaschi, M., Ferraris, M., Mallegni, F., Masini, F., Scola, V., Tozzi, C., 1991. Les
1702 gisements du Pléistocène moyen de Visogliano (Trieste): restes humains, industries,
1703 environnement. L'Anthropologie 95, 9–35.
- 1704 Cavallo, P., Martinetto, E., 2001. Flore carpologica del Pliocene di Castelletto Cervo (Biella).
1705 Bollettino-Museo Regionale di Scienze naturali 18, 277–343.
- 1706 Cheddadi, R., Fady, B., François, L., Hajar, L., Suc, J.-P., Huang, K., Demarteau, M., Vendramin, G.G.,
1707 Ortu, E., 2009. Putative glacial refugia of *Cedrus atlantica* deduced from Quaternary pollen
1708 records and modern genetic diversity. Journal of Biogeography 36, 1361–1371.
1709 doi:10.1111/j.1365-2699.2008.02063.x
- 1710 Christe, C., Gregor Kozłowski, Frey, D., Bétrisey, S., Maharramova, E., Garfi, G., Pirintsos, S., Naciri,
1711 Y., 2014. Footprints of past intensive diversification and structuring in the genus *Zelkova*
1712 (Ulmaceae) in south-western Eurasia. Journal of Biogeography 41, 1081–1093.
1713 doi:10.1111/jbi.12276
- 1714 Combourieu Nebout, N., Londeix, L., Baudin, F., Turon, J.-L., Von Grafenstein, R., Zahn, R., 1999.
1715 Quaternary marine and continental paleoenvironments in the western Mediterranean (Site

- 1716 976, Alboran Sea): palynological evidence. Proceedings of the Ocean Drilling Program.
1717 Scientific Results, pp. 457–468.
- 1718 Combourieu-Nebout, N., 1993. Vegetation response to Upper Pliocene glacial/interglacial cyclicity
1719 in the Central Mediterranean. Quaternary Research 40, 228–236.
- 1720 Combourieu-Nebout, N., Bertini, A., Russo-Ermolli, E., Peyron, O., Klotz, S., Montade, V., Fauquette,
1721 S., Allen, J., Fusco, F., Goring, S., Huntley, B., Joannin, S., Lebreton, V., Magri, D., Martinetto,
1722 E., Orain, R., Sadori, L., 2015. Climate changes in the central Mediterranean and Italian
1723 vegetation dynamics since the Pliocene. Review of Palaeobotany and Palynology 218, 127–
1724 147. doi:10.1016/j.revpalbo.2015.03.001
- 1725 Combourieu-Nebout, N., Foucault, A., Mélières, F., 2004. Vegetation markers of palaeoclimate
1726 cyclical changes in the Pliocene of Punta Piccola (Sicily, Italy). Palaeogeography,
1727 Palaeoclimatology, Palaeoecology 214, 55–66. doi:10.1016/j.palaeo.2004.07.021
- 1728 Combourieu-Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U., Marret, F.,
1729 2009. Rapid climatic variability in the west Mediterranean during the last 25 000 years from
1730 high resolution pollen data. Climate of the Past 5, 503–521.
- 1731 Combourieu-Nebout, N., Vergnaud-Grazzini, C., 1991. Late Pliocene northern hemisphere
1732 glaciations: the continental and marine responses in the central Mediterranean.
1733 Quaternary Science Reviews 10, 319–334.
- 1734 Corrado, P., Magri, D., 2011. A late Early Pleistocene pollen record from Fontana Ranuccio (central
1735 Italy). Journal of Quaternary Science 26, 335–344. doi:10.1002/jqs.1459
- 1736 Cortés-Sánchez, M., Morales-Muñiz, A., Simón-Vallejo, M.D., Bergadá-Zapata, M.M., Delgado-
1737 Huertas, A., López-García, P., López-Sáez, J.A., Lozano-Francisco, M.C., Riquelme-Cantal,
1738 J.A., Roselló-Izquierdo, E., others, 2008. Palaeoenvironmental and cultural dynamics of the
1739 coast of Málaga (Andalusia, Spain) during the Upper Pleistocene and early Holocene.
1740 Quaternary Science Reviews 27, 2176–2193.
- 1741 Cross, J., Pérez Obiol, R., Roure, J.M., 1986. Primeres dades sobre la vegetació i el clima del
1742 Quaternari mitjà a Olot. Collectanea botanica 16, 365–369.
- 1743 Dawson, T.P., Jackson, S.T., House, J.I., Prentice, I.C., Mace, G.M., 2011. Beyond predictions:
1744 biodiversity conservation in a changing climate. science 332, 53–58.
- 1745 Degeai, J.-P., Pastre, J.-F., Gauthier, A., Robert, V., Nomade, S., Bout-Roumazeilles, V., Guillou, H.,
1746 2013. La séquence lacustre du maar d'Alleret (Massif Central, France): téphrochronologie
1747 et évolution paléoenvironnementale en Europe occidentale au début du Pléistocène
1748 moyen. Quaternaire 443–459. doi:10.4000/quaternaire.6808
- 1749 Denk, T., Grimm, G.W., 2005. Phylogeny and biogeography of *Zelkova* (Ulmaceae sensu stricto) as
1750 inferred from leaf morphology, ITS sequence data and the fossil record. Botanical Journal
1751 of the Linnean Society 147, 129–157. doi:10.1111/j.1095-8339.2005.00354.x
- 1752 Desprat, S., Sánchez Goni, M.F., McManus, J.F., Duprat, J., Cortijo, E., 2009. Millennial-scale climatic
1753 variability between 340000 and 270000 years ago in SW Europe: evidence from a NW
1754 Iberian margin pollen sequence. Climate of the Past 5, 53–72.
- 1755 Desprat, S., Sánchez Goñi, M.F., Naughton, F., Turon, J.-L., Duprat, J., Malaizé, B., Cortijo, E.,
1756 Peypouquet, J.-P., 2007. Climate variability of the last five isotopic interglacials: Direct land-
1757 sea-ice correlation from the multiproxy analysis of North-Western Iberian margin deep-sea
1758 cores. Developments in Quaternary Sciences 7, 375–386.

- Desprat, S., Sánchez Goñi, M.F., Turon, J.-L., McManus, J.F., Loutre, M.F., Duprat, J., Malaizé, B., Peyron, O., Peypouquet, J.-P., 2005. Is vegetation responsible for glacial inception during periods of muted insolation changes? *Quaternary Science Reviews* 24, 1361–1374. doi:10.1016/j.quascirev.2005.01.005
- Di Domenico, F., Lucchese, F., Magri, D., 2012. *Buxus* in Europe: Late Quaternary dynamics and modern vulnerability. *Perspectives in Plant Ecology, Evolution and Systematics* 14, 354–362. doi:10.1016/j.ppees.2012.07.001
- Di Rita, F., Anzidei, A.P., Magri, D., 2013. A Lateglacial and early Holocene pollen record from Valle di Castiglione (Rome): Vegetation dynamics and climate implications. *Quaternary International* 288, 73–80. doi:10.1016/j.quaint.2011.11.011
- Di Rita, F., Celant, A., Milli, S., Magri, D., 2015. Lateglacial–early Holocene vegetation history of the Tiber delta (Rome, Italy) under the influence of climate change and sea level rise. *Review of Palaeobotany and Palynology* 218, 204–206. doi:10.1016/j.revpalbo.2014.12.005
- Eiserhardt, W.L., Borchsenius, F., Plum, C.M., Ordonez, A., Svenning, J.-C., 2015. Climate-driven extinctions shape the phylogenetic structure of temperate tree floras. *Ecology Letters* 18, 263–272. doi:10.1111/ele.12409
- Elhaï, H., 1969. La flore sporo-pollinique du gisement Villafranchien de Senèze (Massif-Central-France). *Pollen et Spores* 11, 127–139.
- Fady, B., Lefèvre, F., Vendramin, G.G., Ambert, A., Régnier, C., Bariteau, M., 2008. Genetic consequences of past climate and human impact on eastern Mediterranean *Cedrus libani* forests. Implications for their conservation. *Conservation Genetics* 9, 85–95. doi:10.1007/s10592-007-9310-6
- Falguères, C., Bahain, J.-J., Tozzi, C., Boschian, G., Dolo, J.-M., Mercier, N., Valladas, H., Yokoyama, Y., 2008. ESR/U-series chronology of the Lower Palaeolithic palaeoanthropological site of Visogliano, Trieste, Italy. *Quaternary Geochronology* 3, 390–398. doi:10.1016/j.quageo.2008.01.007
- Fang, J., Wang, Z., Tang, Z., 2011. Atlas of woody plants in China: distribution and climate. Springer Science & Business Media.
- Farjon, A., 2008. A natural history of conifers. Timber Press.
- Farjon, A., 2005. A monograph of Cupressaceae and *Sciadopitys*. Royal Botanic Gardens, Kew:
- Fauquette, S., Clauzon, G., Suc, J.-P., Zheng, Z., 1999. A new approach for palaeoaltitude estimates based on pollen records: example of the Mercantour Massif (southeastern France) at the earliest Pliocene. *Earth and Planetary Science Letters* 170, 35–47.
- Feddi, N., Fauquette, S., Suc, J.-P., 2011. Histoire plio-pléistocène des écosystèmes végétaux de Méditerranée sud-occidentale: apport de l'analyse pollinique de deux sondages en mer d'Alboran. *Geobios* 44, 57–69. doi:10.1016/j.geobios.2010.03.007
- Field, M., de Beaulieu, J.-L., Guiot, J., Ponce, P., 2000. Middle Pleistocene deposits at La Côte, Val-de-Lans, Isère department, France: plant macrofossil, palynological and fossil insect investigations. *Palaeogeography, Palaeoclimatology, Palaeoecology* 159, 53–83. doi:10.1016/S0031-0182(00)00032-8
- Filipova-Marionova, M.V., Kvavadze, E.V., Connor, S.E., Sjögren, P., 2010. Estimating absolute pollen productivity for some European Tertiary-relict taxa. *Vegetation History and Archaeobotany* 19, 351–364.

- 1802 Fletcher, W.J., Müller, U.C., Koutsodendris, A., Christanis, K., Pross, J., 2013. A centennial-scale
1803 record of vegetation and climate variability from 312 to 240 ka (Marine Isotope Stages 9c–
1804 a, 8 and 7e) from Tenaghi Philippon, NE Greece. *Quaternary Science Reviews* 78, 108–125.
1805 doi:10.1016/j.quascirev.2013.08.005
- 1806 Fletcher, W.J., Sánchez Goñi, M.F., 2008. Orbital- and sub-orbital-scale climate impacts on
1807 vegetation of the western Mediterranean basin over the last 48,000 yr. *Quaternary*
1808 *Research* 70, 451–464. doi:10.1016/j.yqres.2008.07.002
- 1809 Florschütz, F., Amor, J.M., Wijmstra, T., 1971. Palynology of a thick Quaternary succession in
1810 southern Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology* 10, 233–264.
- 1811 Follieri, M., 2010. Conifer extinction in Quaternary Italian records. *Quaternary International* 225,
1812 37–43. doi:10.1016/j.quaint.2010.02.001
- 1813 Follieri, M., 1962. La foresta colchica fossile di Riano Romano. II. Analisi polliniche. *Annali di Botanica*
1814 27, 245–280.
- 1815 Follieri, M., 1958. La foresta colchica fossile di Riano Romano. I. Studio dei fossili vegetali
1816 macroscopici. *Annali di Botanica* 26, 129–142.
- 1817 Follieri, M., Magri, D., Sadori, L., 1989. Pollen stratigraphical synthesis from Valle di Castiglione
1818 (Roma). *Quaternary International* 3, 81–84.
- 1819 Follieri, M., Magri, D., Sadori, L., 1988. 250000-year pollen record from Valle di Castiglione (Roma).
1820 *Pollen et Spores* 30, 329–356.
- 1821 Follieri, M., Magri, D., Sadori, L., 1986. Late Pleistocene *Zelkova* extinction in central Italy. *New*
1822 *Phytologist* 103, 269–273.
- 1823 Fouache, É., Dalongeville, R., Kunesch, S., Suc, J., Subally, D., Prieur, A., Lozouet, P., 2005. The
1824 environmental setting of the harbor of the classical site of Oeniades on the Acheloos delta,
1825 Greece. *Geoarchaeology* 20, 285–302.
- 1826 Franco Múgica, F., García Antón, M., Maldonado Ruiz, J., Morla Juaristi, C., Sainz-Ollero, H., 2001.
1827 The Holocene history of *Pinus* forests in the Spanish Northern Meseta. *The Holocene* 11,
1828 343–358. doi:10.1191/095968301669474913
- 1829 Fusco, F., 2007. Vegetation response to early Pleistocene climatic cycles in the Lamone valley
1830 (Northern Apennines, Italy). *Review of Palaeobotany and Palynology* 145, 1–23.
1831 doi:10.1016/j.revpalbo.2006.08.005
- 1832 García-Antón, M., Sainz-Ollero, H., 1991. Pollen records from the middle Pleistocene Atapuerca site
1833 (Burgos, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* 85, 199–206.
- 1834 García-Prieto, E., 2015. Dinámica paleoambiental durante los últimos 135.000 años en el Alto Jiloca:
1835 el registro lacustre de El Cañizar (PhD Thesis). Universidad de Zaragoza, Spain.
- 1836 Garfi, G., Carimi, F., Pasta, S., Rühl, J., Trigila, S., 2011. Additional insights on the ecology of the relic
1837 tree *Zelkova sicula* di Pasquale, Garfi et Quézel (Ulmaceae) after the finding of a new
1838 population. *Flora - Morphology, Distribution, Functional Ecology of Plants* 206, 407–417.
1839 doi:10.1016/j.flora.2010.11.004
- 1840 Gavin, D.G., Fitzpatrick, M.C., Guggen, P.F., Heath, K.D., Rodríguez-Sánchez, F., Dobrowski, S.Z.,
1841 Hampe, A., Hu, F.S., Ashcroft, M.B., Bartlein, P.J., Blois, J.L., Carstens, B.C., Davis, E.B., de
1842 Lafontaine, G., Edwards, M.E., Fernandez, M., Henne, P.D., Herring, E.M., Holden, Z.A.,
1843 Kong, W., Liu, J., Magri, D., Matzke, N.J., McGlone, M.S., Saltré, F., Stigall, A.L., Tsai, Y.-H.E.,
1844 Williams, J.W., 2014. Climate refugia: joint inference from fossil records, species

- 1845 distribution models and phylogeography. *New Phytologist* 204, 37–54.
1846 doi:10.1111/nph.12929
- 1847 Geurts, M., 1979. Approche palynostratigraphique des depots calcaireux quaternaires dans la region
1848 de Banyoles-Besalc (Catalogne), in: *Actas de La IV Reunión Del Grupo de Trabajo Del*
1849 *Cuaternario*. pp. 106–115.
- 1850 Geurts, M., 1977. Premières données a l'étude paplynologique des dépôts calcaireux quaternaires
1851 en Catalogne. *Acta geológica hispánica* 12, 86–89.
- 1852 Gibbard, P.L., Head, M.J., Walker, M.J.C., the Subcommission on Quaternary Stratigraphy, 2010.
1853 Formal ratification of the Quaternary System/Period and the Pleistocene Series/Epoch with
1854 a base at 2.58 Ma. *Journal of Quaternary Science* 25, 96–102. doi:10.1002/jqs.1338
- 1855 Gómez-Orellana, L., Ramil-Rego, P., Muñoz Sobrino, C., 2013. The response of vegetation at the end
1856 of the last glacial period (MIS 3 and MIS 2) in littoral areas of NW Iberia. *Boreas* 42, 729–
1857 744.
- 1858 Gómez-Orellana, L., Ramil-Rego, P., Muñoz Sobrino, C., 2007. The Würm in NW Iberia, a pollen
1859 record from Area Longa (Galicia). *Quaternary Research* 67, 438–452.
1860 doi:10.1016/j.yqres.2007.01.003
- 1861 González-Sampériz, P., Aranbarri, J., Pérez-Sanz, A., Gil-Romera, G., Moreno, A., Leunda, M., Sevilla-
1862 Callejo, M., Corella, J.P., Morellón, M., Oliva, B., Valero-Garcés, 2016. Environmental and
1863 climate change in the southern Central Pyrenees since the Last Glacial Maximum: A view
1864 from the lake records. *Catena*.
- 1865 González-Sampériz, P., García-Prieto, E., Aranbarri, J., Valero Garcés, B.L., Moreno, A., G. Gil-
1866 Romera, Sevilla Callejo, M., Santos, L., Morellón, M., Mata, P., Andrade, A., Carrión, J., 2013.
1867 Reconstrucción paleoambiental del último ciclo glacial-interglacial en la Iberia continental:
1868 la secuencia del Cañizar de Villarquemado (Teruel). *Cuadernos de investigación geográfica*
1869 49–76.
- 1870 González-Sampériz, P., Leroy, S.A.G., Carrión, J.S., Fernández, S., García-Antón, M., Gil-García, M.J.,
1871 Uzquiano, P., Valero-Garcés, B., Figueiral, I., 2010. Steppes, savannahs, forests and
1872 phytodiversity reservoirs during the Pleistocene in the Iberian Peninsula. *Review of*
1873 *Palaeobotany and Palynology* 162, 427–457. doi:10.1016/j.revpalbo.2010.03.009
- 1874 Gregor, H.-J., 1986. Preliminary report about the macrofloral history of the Stirone-river sequence
1875 (Zanclean, Piacenzian, "Calabrian"). *Memorie della Società Geologica Italiana* 31, 219–227.
- 1876 Hably, L., Kvacek, Z., 1998. Pliocene mesophytic forests surrounding crater lakes in western Hungary.
1877 *Review of Palaeobotany and Palynology* 101, 257–269.
- 1878 Hajar, L., François, L., Khater, C., Jomaa, I., Déqué, M., Cheddadi, R., 2010. *Cedrus libani* (A. Rich)
1879 distribution in Lebanon: Past, present and future. *Comptes Rendus Biologies* 333, 622–630.
1880 doi:10.1016/j.crv.2010.05.003
- 1881 Hajar, L., Haïdar-Boustani, M., Khater, C., Cheddadi, R., 2010. Environmental changes in Lebanon
1882 during the Holocene: Man vs. climate impacts. *Journal of Arid Environments* 74, 746–755.
1883 doi:10.1016/j.jaridenv.2008.11.002
- 1884 Hajar, L., Khater, C., Cheddadi, R., 2008. Vegetation changes during the late Pleistocene and
1885 Holocene in Lebanon: a pollen record from the Bekaa Valley. *The Holocene* 18, 1089–1099.
1886 doi:10.1177/0959683608095580

- 1887 Hammen, T. van der, 1951. A contribution to the palaeobotany of the Tiglian. *Geologie Mijnbouw*,
1888 ns 13, 242–250.
- 1889 Hammen, T. van der, Wijmstra, T.A., Zagwijn, W., 1971. The floral record of the late Cenozoic of
1890 Europe, in: Turekian, K.K. (Ed.). *The Late Cenozoic Glacial Ages*. Yale University Press, New
1891 Haven, pp. 391–424.
- 1892 Head, M.J., Gibbard, P.L., 2005. Early-Middle Pleistocene transitions: an overview and
1893 recommendation for the defining boundary. *Geological Society, London, Special*
1894 *Publications* 247, 1–18.
- 1895 Hooghiemstra, H., Lézine, A.-M., Leroy, S.A.G., Dupont, L., Marret, F., 2006. Late Quaternary
1896 palynology in marine sediments: A synthesis of the understanding of pollen distribution
1897 patterns in the NW African setting. *Quaternary International* 148, 29–44.
1898 doi:10.1016/j.quaint.2005.11.005
- 1899 Huntley, B., 1993. Species-richness in north-temperate zone forests. *Journal of Biogeography* 20,
1900 163–180.
- 1901 Iriarte-Chiapusso, M., Muñoz Sobrino, C., Gómez-Orellana, L., Hernández-Beloqui, B., García-
1902 Moreiras, I., Fernández Rodríguez, C., Heiri, O., Lotter, A., Ramil-Rego, P., 2016. Reviewing
1903 the Lateglacial–Holocene transition in NW Iberia: A palaeoecological approach based on
1904 the comparison between dissimilar regions. *Quaternary International* 403, 211–236.
- 1905 Ivanov, D., Ashraf, A.R., Mosbrugger, V., Palamarev, E., 2002. Palynological evidence for Miocene
1906 climate change in the Forecarpathian Basin (Central Paratethys, NW Bulgaria).
1907 *Palaeogeography, Palaeoclimatology, Palaeoecology* 178, 19–37. doi:10.1016/S0031-
1908 0182(01)00365-0
- 1909 Jackson, S.T., Weng, C., 1999. Late Quaternary extinction of a tree species in eastern North America.
1910 *Proceedings of the National Academy of Sciences* 96, 13847–13852.
- 1911 Janssen, C., 1984. Modern pollen assemblages and vegetation in the Myrtle Lake peatland,
1912 Minnesota. *Ecological Monographs* 54, 213–252.
- 1913 Jasińska, A.K., Boratyńska, K., Sobierajska, K., Romo, A., Ok, T., Kharat, M.B.D., Boratyński, A., 2013.
1914 Relationships among *Cedrus libani*, *C. brevifolia* and *C. atlantica* as revealed by the
1915 morphological and anatomical needle characters. *Plant Systematics and Evolution* 299, 35–
1916 48. doi:10.1007/s00606-012-0700-y
- 1917 Jiménez-Moreno, G., Fauquette, S., Suc, J., 2008. Vegetation, climate and palaeoaltitude
1918 reconstructions of the Eastern Alps during the Miocene based on pollen records from
1919 Austria, Central Europe. *Journal of Biogeography* 35, 1638–1649.
- 1920 Jiménez-Moreno, G., Fauquette, S., Suc, J.-P., 2010. Miocene to Pliocene vegetation reconstruction
1921 and climate estimates in the Iberian Peninsula from pollen data. *Review of Palaeobotany*
1922 *and Palynology* 162, 403–415. doi:10.1016/j.revpalbo.2009.08.001
- 1923 Jiménez-Moreno, G., Suc, J.-P., 2007. Middle Miocene latitudinal climatic gradient in Western
1924 Europe: Evidence from pollen records. *Palaeogeography, Palaeoclimatology,*
1925 *Palaeoecology* 253, 208–225. doi:10.1016/j.palaeo.2007.03.040
- 1926 Joannin, S., 2007. Changements climatiques en méditerranée à la transition Pléistocène inférieur-
1927 moyen: pollens, isotopes stables et cyclostratigraphie. Université Claude Bernard-Lyon I.
- 1928 Joannin, S., Bassinot, F., Nebout, N.C., Peyron, O., Beaudouin, C., 2011. Vegetation response to
1929 obliquity and precession forcing during the Mid-Pleistocene Transition in Western

- 1930 Mediterranean region (ODP site 976). *Quaternary Science Reviews* 30, 280–297.
1931 doi:10.1016/j.quascirev.2010.11.009
- 1932 Joannin, S., Ciaranfi, N., Stefanelli, S., 2008. Vegetation changes during the late Early Pleistocene at
1933 Montalbano Jonico (Province of Matera, southern Italy) based on pollen analysis.
1934 *Palaeogeography, Palaeoclimatology, Palaeoecology* 270, 92–101.
1935 doi:10.1016/j.palaeo.2008.08.017
- 1936 Joannin, S., Cornée, J.-J., Moissette, P., Suc, J.-P., Koskeridou, E., Lécuyer, C., Buisine, C., Kouli, K.,
1937 Ferry, S., 2007a. Changes in vegetation and marine environments in the eastern
1938 Mediterranean (Rhodes, Greece) during the Early and Middle Pleistocene. *Journal of the*
1939 *Geological Society* 164, 1119–1131.
- 1940 Joannin, S., Quillévéré, F., Suc, J.-P., Lécuyer, C., Martineau, F., 2007b. Early Pleistocene climate
1941 changes in the central Mediterranean region as inferred from integrated pollen and
1942 planktonic foraminiferal stable isotope analyses. *Quaternary Research* 67, 264–274.
1943 doi:10.1016/j.yqres.2006.11.001
- 1944 Julia Bruguès, R., Suc, J.-P., 1980. Analyse pollinique des dépôts lacustres du Pléistocène inférieur
1945 de Banyoles (Bañolas, site de la Bòbila Ordis-Espagne): un élément nouveau dans la
1946 reconstitution de l'histoire paléoclimatique des régions méditerranéennes d'Europe
1947 occidentale. *Geobios* 13, 5–19.
- 1948 Kahlke, R.-D., García, N., Kostopoulos, D.S., Lacombat, F., Lister, A.M., Mazza, P.P., Spassov, N., Titov,
1949 V.V., 2011. Western Palaeoarctic palaeoenvironmental conditions during the Early and early
1950 Middle Pleistocene inferred from large mammal communities, and implications for
1951 hominin dispersal in Europe. *Quaternary Science Reviews* 30, 1368–1395.
- 1952 Kemna, H.A., Westerhoff, W.E., 2007. Remarks on the palynology-based chronostratigraphical
1953 subdivision of Pliocene terrestrial deposits in NW-Europe. *Quaternary international* 164,
1954 184–196.
- 1955 Klotz, S., Fauquette, S., Combourieu-Nebout, N., Uhl, D., Suc, J.-P., Mosbrugger, V., 2006. Seasonality
1956 intensification and long-term winter cooling as a part of the Late Pliocene climate
1957 development. *Earth and Planetary Science Letters* 241, 174–187.
1958 doi:10.1016/j.epsl.2005.10.005
- 1959 Kotthoff, U., Muller, U.C., Pross, J., Schmiedl, G., Lawson, I.T., van de Schootbrugge, B., Schulz, H.,
1960 2008. Lateglacial and Holocene vegetation dynamics in the Aegean region: an integrated
1961 view based on pollen data from marine and terrestrial archives. *The Holocene* 18, 1019–
1962 1032. doi:10.1177/0959683608095573
- 1963 Kovar-Eder, J., Kvaček, Z., Martinetto, E., Roiron, P., 2006. Late Miocene to Early Pliocene vegetation
1964 of southern Europe (7–4 Ma) as reflected in the megafossil plant record. *Palaeogeography,*
1965 *Palaeoclimatology, Palaeoecology* 238, 321–339. doi:10.1016/j.palaeo.2006.03.031
- 1966 Kozłowski, G., Frey, D., Fazan, L., Egli, B., Bétrisey, S., Gratzfeld, J., Garfi, G., Pirintzos, S., 2014. The
1967 Tertiary relict tree *Zelkova abelicea* (Ulmaceae): distribution, population structure and
1968 conservation status on Crete. *Oryx* 48, 80–87. doi:10.1017/S0030605312001275
- 1969 Kozłowski, G., Gibbs, D., Huan, F., Frey, D., Gratzfeld, J., 2012. Conservation of threatened relict trees
1970 through living ex situ collections: lessons from the global survey of the genus *Zelkova*
1971 (Ulmaceae). *Biodiversity and Conservation* 21, 671–685.
- 1972 Kuprianova, L., 1965. The palynology of the Amentiferae. The Academy of Sciences of the USSR,
1973 Moscow 214.

- 1974 Kvavadze, E.V., Connor, S.E., 2005. *Zelkova carpinifolia* (Pallas) K. Koch in Holocene sediments of
1975 Georgia—an indicator of climatic optima. Review of Palaeobotany and Palynology 133, 69–
1976 89. doi:10.1016/j.revpalbo.2004.09.002
- 1977 Lamb, H.F., Eicher, U., Switsur, V.R., 1989. An 18,000-Year Record of Vegetation, Lake-Level and
1978 Climatic Change from Tigalmamine, Middle Atlas, Morocco. Journal of Biogeography 16,
1979 65. doi:10.2307/2845311
- 1980 Lawson, I., Frogley, M., Bryant, C., Preece, R., Tzedakis, P., 2004. The Lateglacial and Holocene
1981 environmental history of the Ioannina basin, north-west Greece. Quaternary Science
1982 Reviews 23, 1599–1625. doi:10.1016/j.quascirev.2004.02.003
- 1983 Lawson, I.T., Al-Omari, S., Tzedakis, P.C., Bryant, C.L., Christanis, K., 2005. Lateglacial and Holocene
1984 vegetation history at Nisi Fen and the Boras mountains, northern Greece. The Holocene 15,
1985 873–887. doi:10.1191/0959683605hl860ra
- 1986 Lebreton, V., 2002. Végétation et climat au Pléistocène inférieur et moyen à La Pineta (Isernia,
1987 Italie). Comptes Rendus Palevol 1, 11–17.
- 1988 LePage, B.A., 2003. A new species of *Tsuga* (Pinaceae) from the middle Eocene of Axel Heiberg
1989 Island, Canada, and an assessment of the evolution and biogeographical history of the
1990 genus. Botanical Journal of the Linnean Society 141, 257–296. doi:10.1046/j.1095-
1991 8339.2003.00131.x
- 1992 Leroy, S., Ambert, P., Suc, J.-P., 1994. Pollen record of the Saint-Macaire maar (Hérault, southern
1993 France): a Lower Pleistocene glacial phase in the Languedoc coastal plain. Review of
1994 Palaeobotany and Palynology 80, 149–157.
- 1995 Leroy, S.A., Tudryn, A., Chalié, F., López-Merino, L., Gasse, F., 2013. From the Allerød to the mid-
1996 Holocene: palynological evidence from the south basin of the Caspian Sea. Quaternary
1997 Science Reviews 78, 77–97.
- 1998 Leroy, S.A.G., 2008. Vegetation cycles in a disturbed sequence around the Cobb-Mountain subchron
1999 in Catalonia (Spain). Journal of Paleolimnology 40, 851–868. doi:10.1007/s10933-008-
2000 9203-9
- 2001 Leroy, S.A.G., 2007. Progress in palynology of the Gelasian–Calabrian Stages in Europe: Ten
2002 messages. Revue de Micropaléontologie 50, 293–308. doi:10.1016/j.revmic.2006.08.001
- 2003 Leroy, S.A.G., 1997. Climatic and non-climatic lake-level changes inferred from a Plio-Pleistocene
2004 lacustrine complex of Catalonia (Spain): palynology of the Tres Pins sequences. Journal of
2005 Paleolimnology 17, 347–367.
- 2006 Leroy, S.A.G., 1990. Paléoclimats plio-pléistocènes en Catalogne et Languedoc d'après la palynologie
2007 de formations lacustres. These de l'Université Catholique de Louvain, Belgique 319.
- 2008 Leroy, S.A.G., Arpe, K., Mikolajewicz, U., 2011. Vegetation context and climatic limits of the Early
2009 Pleistocene hominin dispersal in Europe. Quaternary Science Reviews 30, 1448–1463.
2010 doi:10.1016/j.quascirev.2010.01.017
- 2011 Leroy, S.A.G., Roiron, P., 1996. Latest Pliocene pollen and leaf floras from Bernasso palaeolake
2012 (Escandorgue Massif, Hérault, France). Review of Palaeobotany and Palynology 94, 295–
2013 328.
- 2014 Leroy, S., Seret, G., 1992. Duration and Vegetation Dynamic of the Nogaret Interglacial (-1.9 Ma, S.
2015 of France): Tentative Correlation with Stage 75, in: Start of a Glacial. Springer, pp. 113–125.

2016 Li, J., Del Tredici, P., 2008. The Chinese *Parrotia*: a sibling species of the Persian *Parrotia*. *Arnoldia*
2017 66, 2–9.

2018 Liepelt, S., Cheddadi, R., de Beaulieu, J.-L., Fady, B., Gömöry, D., Hussendörfer, E., Konnert, M., Litt,
2019 T., Longauer, R., Terhürne-Berson, R., Ziegenhagen, B., 2009. Postglacial range expansion
2020 and its genetic imprints in *Abies alba* (Mill.) — A synthesis from palaeobotanic and genetic
2021 data. *Review of Palaeobotany and Palynology* 153, 139–149.
2022 doi:10.1016/j.revpalbo.2008.07.007

2023 Liu, Y.-S., Basinger, J.F., 2000. Fossil *Cathaya* (Pinaceae) pollen from the Canadian high arctic.
2024 *International Journal of Plant Sciences* 161, 829–847.

2025 Lona, F., 1950. Contributi alla storia della vegetazione e del clima nella Val Padana: analisi pollinica
2026 del giacimento villafranchiano di Leffe (Bergamo). *Atti della Società Italiana di Scienze*
2027 *Naturali* 89, 123–178.

2028 Lona, F., Bertoldi, R., 1972. La storia del Plio-Pleistocene italiano in alcune sequenze vegetazionali
2029 lacustri e marine. *Atti della Accademia Nazionale dei Lincei* 11, 1–45.

2030 Lona, F., Ricciardi, E., 1961. Studio pollinologico stratigrafico su una serie lacustre pleistocenica
2031 dell'Italia centrale (Bacino di Gubbio, Perugia). *Pollen et Spores* 3, 93–100.

2032 Løvlie, R., Leroy, S., 1995. Magnetostratigraphy of Lower Pleistocene Banyoles palaeolake carbonate
2033 sediments from Catalonia, NE Spain: evidence for relocation of the Cobb Mountain sub-
2034 chron. *Quaternary Science Reviews* 14, 473–485.

2035 Magri, D., 2012. Quaternary history of *Cedrus* in Southern Europe. *Annali di Botanica* 2, 57–66.

2036 Magri, D., 2010. Persistence of tree taxa in Europe and Quaternary climate changes. *Quaternary*
2037 *International* 219, 145–151. doi:10.1016/j.quaint.2009.10.032

2038 Magri, D., 1999. Late Quaternary vegetation history at Lagaccione near Lago di Bolsena (central
2039 Italy). *Review of Palaeobotany and Palynology* 106, 171–208.

2040 Magri, D., Agrillo, E., Di Rita, F., Furlanetto, G., Pini, R., Ravazzi, C., Spada, F., 2015. Holocene
2041 dynamics of tree taxa populations in Italy. *Review of Palaeobotany and Palynology* 218,
2042 267–284. doi:10.1016/j.revpalbo.2014.08.012

2043 Magri, D., Palombo, M.R., 2013. Early to Middle Pleistocene dynamics of plant and mammal
2044 communities in South West Europe. *Quaternary International* 288, 63–72.
2045 doi:10.1016/j.quaint.2012.02.028

2046 Magri, D., Parra, I., 2002. Late Quaternary western Mediterranean pollen records and African winds.
2047 *Earth and Planetary Science Letters* 200, 401–408.

2048 Magri, D., Rita, F.D., Palombo, M.R., 2010. An Early Pleistocene interglacial record from an
2049 intermontane basin of central Italy (Scoppito, L'Aquila). *Quaternary International* 225, 106–
2050 113. doi:10.1016/j.quaint.2009.04.005

2051 Magri, D., Sadori, L., 1999. Late Pleistocene and Holocene pollen stratigraphy at Lago di Vico, central
2052 Italy. *Vegetation history and archaeobotany* 8, 247–260.

2053 Magri, D., Vendramin, G.G., Comps, B., Dupanloup, I., Geburek, T., Gomory, D., Latalowa, M., Litt,
2054 T., Paule, L., Roure, J.M., Tantau, I., van der Knaap, W.O., Petit, R.J., de Beaulieu, J.-L., 2006.
2055 A new scenario for the Quaternary history of European beech populations: palaeobotanical
2056 evidence and genetic consequences. *New Phytologist* 171, 199–221. doi:10.1111/j.1469-
2057 8137.2006.01740.x

2058 Mai, D.H., Velitzelos, E., 2007. The fossil flora of Kallithea (Rhodes, Greece) at the
2059 Pliocene/Pleistocene boundary. *Palaeontographica Abteilung B* 75–99.

2060 Manchester, S.R., 1987. The fossil history of the Juglandaceae. *Monographs in Systematic Botany*,
2061 *Missouri bot. Gard* 21, 1–137.

2062 Manzi, G., Magri, D., Milli, S., Palombo, M.R., Margari, V., Celiberti, V., Barbieri, M., Barbieri, M.,
2063 Melis, R.T., Rubini, M., Ruffo, M., Saracino, B., Tzedakis, P.C., Zarattini, A., Biddittu, I., 2010.
2064 The new chronology of the Ceprano calvarium (Italy). *Journal of Human Evolution* 59, 580–
2065 585. doi:10.1016/j.jhevol.2010.06.010

2066 Marra, F., Sottili, G., Gaeta, M., Giaccio, B., Jicha, B., Masotta, M., Palladino, D.M., Deocampo, D.M.,
2067 2014. Major explosive activity in the Monti Sabatini Volcanic District (central Italy) over the
2068 800–390 ka interval: geochronological-geochemical overview and tephrostratigraphic
2069 implications. *Quaternary Science Reviews* 94, 74–101.
2070 doi:10.1016/j.quascirev.2014.04.010

2071 Martinetto, E., 2015. Monographing the Pliocene and early Pleistocene carpoﬂoras of Italy:
2072 methodological challenges and current progress. *Palaeontographica Abteilung B* 57–99.

2073 Martinetto, E., 2009. Palaeoenvironmental significance of plant macrofossils from the Pliocene
2074 Formation, Middle Pleistocene of Lombardy, North Italy. *Quaternary International* 204, 20–
2075 30. doi:10.1016/j.quaint.2008.11.014

2076 Martinetto, E., 2001. The role of central Italy as a centre of refuge for thermophilous plants in the
2077 late Cenozoic. *Acta Palaeobotanica* 41, 299–319.

2078 Martinetto, E., Bertini, A., Basilici, G., Baldanza, A., Bizzarri, R., Cherin, M., Gentili, S., Pontini, M.R.,
2079 2014. The plant record of the Dunarobba and Pietrafitta sites in the Plio-Pleistocene
2080 palaeoenvironmental context of Central Italy. *Alpine and Mediterranean Quaternary* 27,
2081 29–72.

2082 Martinetto, E., Monegato, G., Irace, A., Vaiani, S.C., Vassio, E., 2015. Pliocene and Early Pleistocene
2083 carpological records of terrestrial plants from the southern border of the Po Plain (northern
2084 Italy). *Review of Palaeobotany and Palynology* 218, 148–166.
2085 doi:10.1016/j.revpalbo.2014.10.007

2086 Martinetto, E., Monegato, G., Vassio, E., 2012. An Early Pleistocene plant assemblage with East
2087 European affinity in the Venetian-Friulian basin (NE Italy). *Alpine and Mediterranean*
2088 *Quaternary* 25, 91–104.

2089 Martinetto, E., Sami, M., 2001. Paleoflora delle “sabbie gialle” mediopleistoceniche di Oriolo, presso
2090 Faenza (Ravenna). *Quaderno di Studi e notizie di storia Naturale della Romagna* 14, 1–28.

2091 Massari, F., Rio, D., Serandrei Barbero, R., Asioli, A., Capraro, L., Fornaciari, E., Vergerio, P., 2004.
2092 The environment of Venice area in the past two million years. *Palaeogeography*,
2093 *Palaeoclimatology, Palaeoecology* 202, 273–308. doi:10.1016/S0031-0182(03)00640-0

2094 Meng, H.-H., Su, T., Huang, Y.-J., Zhu, H., Zhou, Z.-K., 2015. Late Miocene *Palaeocarya*
2095 (Engelhardieae: Juglandaceae) from Southwest China and its biogeographic implications.
2096 *Journal of Systematics and Evolution* 53, 499–511.

2097 Messenger, E., Lebreton, V., Marquer, L., Russo-Ermolli, E., Orain, R., Renault-Miskovsky, J.,
2098 Lordkipanidze, D., Desprieé, J., Peretto, C., Arzarello, M., 2011. Palaeoenvironments of
2099 early hominins in temperate and Mediterranean Eurasia: new palaeobotanical data from

2100 Palaeolithic key-sites and synchronous natural sequences. *Quaternary Science Reviews* 30,
2101 1439–1447. doi:10.1016/j.quascirev.2010.09.008

2102 Moscariello, A., Ravazzi, C., Brauer, A., Mangili, C., Chiesa, S., Rossi, S., de Beaulieu, J.-L., Reille, M.,
2103 2000. A long lacustrine record from the Piànico-Sèlere basin (Middle-Late Pleistocene,
2104 Northern Italy). *Quaternary International* 73, 47–68.

2105 Müllenders, W., Favero, V., Coremans, M., Dirickx, M., 1996. Analyses polliniques des sondages à
2106 Venise (VE1, VE1bis, VE2). *Aardkundige Mededelingen* 7, 87–117.

2107 Munno, R., Petrosino, P., Romano, P., Russo Ermolli, E., Juvigné, É., 2001. A late Middle Pleistocene
2108 climatic cycle in southern Italy inferred from pollen analysis and tephrostratigraphy of the
2109 Acerno lacustrine succession. *Géographie physique et Quaternaire* 55, 87–99.

2110 Muñoz Sobrino, C.M., Ramil-Rego, P., Guitián, M.A.R., 2001. Vegetation in the mountains of
2111 northwest Iberia during the last glacial-interglacial transition. *Vegetation History and*
2112 *Archaeobotany* 7–21.

2113 Muttoni, G., Carcano, C., Garzanti, E., Ghielmi, M., Piccin, A., Pini, R., Rogledi, S., Sciunnach, D., 2003.
2114 Onset of major Pleistocene glaciations in the Alps. *Geology* 31, 989–992.

2115 Muttoni, G., Ravazzi, C., Breda, M., Pini, R., Laj, C., Kissel, C., Mazaud, A., Garzanti, E., 2007.
2116 Magnetostratigraphic dating of an intensification of glacial activity in the southern Italian
2117 Alps during Marine Isotope Stage 22. *Quaternary Research* 67, 161–173.
2118 doi:10.1016/j.yqres.2006.07.006

2119 Nakagawa, T., Garfi, G., Reille, M., Verlaque, R., 1998. Pollen morphology of *Zelkova sicula*
2120 (Ulmaceae), a recently discovered relic species of the European Tertiary flora: description,
2121 chromosomal relevance, and palaeobotanical significance. *Review of Palaeobotany and*
2122 *Palynology* 100, 27–37.

2123 O'Brien, C.E., Jones, R.L., 2003. Early and Middle Pleistocene vegetation history of the Médoc region,
2124 southwest France. *Journal of Quaternary Science* 18, 557–579. doi:10.1002/jqs.779

2125 Okuda, M., Van Vugt, N., Nakagawa, T., Ikeya, M., Hayashida, A., Yasuda, Y., Setoguchi, T., 2002.
2126 Palynological evidence for the astronomical origin of lignite–detritus sequence in the
2127 Middle Pleistocene Marathousa Member, Megalopolis, SW Greece. *Earth and Planetary*
2128 *Science Letters* 201, 143–157.

2129 Okuda, M., Yasuda, Y., Setoguchi, T., 2001. Middle to Late Pleistocene vegetation history and
2130 climatic changes at Lake Kopais, Southeast Greece. *Boreas* 30, 73–82.

2131 Oldfield, F., 1968. The Quaternary Vegetational History of the French Pays Basque. I. Stratigraphy
2132 and pollen analysis. *New Phytologist* 67, 677–731. doi:10.1111/j.1469-
2133 8137.1968.tb05494.x

2134 Orain, R., Lebreton, V., Ermolli, E.R., Combourieu-Nebout, N., Sémah, A.-M., 2013. *Carya* as marker
2135 for tree refuges in southern Italy (Boiano basin) at the Middle Pleistocene.
2136 *Palaeogeography, Palaeoclimatology, Palaeoecology* 369, 295–302.
2137 doi:10.1016/j.palaeo.2012.10.037

2138 Orain, R., Russo Ermolli, E., Lebreton, V., Di Donato, V., Bahain, J.-J., Sémah, A.-M., 2015. Vegetation
2139 sensitivity to local environmental factors and global climate changes during the Middle
2140 Pleistocene in southern Italy—A case study from the Molise Apennines. *Review of*
2141 *Palaeobotany and Palynology* 220, 69–77. doi:10.1016/j.revpalbo.2015.05.002

2142 Öztürk, M., Çelik, A., Güvensen, A., Hamzaoglu, E., 2008. Ecology of tertiary relict endemic
2143 *Liquidambar orientalis* Mill. forests. Forest Ecology and Management 256, 510–518.
2144 doi:10.1016/j.foreco.2008.01.027

2145 Palombo, M.R., Mussi, M., Agostini, S., Barbieri, M., Di Canzio, E., Di Rita, F., Fiore, I., Iacumin, P.,
2146 Magri, D., Speranza, F., Tagliacozzo, A., 2010. Human peopling of Italian intramontane
2147 basins: The early Middle Pleistocene site of Pagliare di Sassa (L'Aquila, central Italy).
2148 Quaternary International 223–224, 170–178. doi:10.1016/j.quaint.2009.10.038

2149 Pastre, J.-F., Gauthier, A., Nomade, S., Orth, P., Andrieu, A., Goupille, F., Guillou, H., Kunesch, S.,
2150 Scaillet, S., Renne, P.R., 2007. The Alleret maar (Massif Central, France): A new lacustrine
2151 sequence of the early Middle Pleistocene in western Europe. Comptes Rendus Geoscience
2152 339, 987–997. doi:10.1016/j.crte.2007.09.019

2153 Petrosino, P., Ermolli, E.R., Donato, P., Jicha, B., Robustelli, G., Sardella, R., 2014. Using
2154 Tephrochronology and palynology to date the MIS 13 lacustrine sediments of the Mercure
2155 basin (Southern Apennines–Italy). Italian Journal of Geosciences 133, 169–186.

2156 Pieruccini, P., Di Celma, C., Di Rita, F., Magri, D., Carnevale, G., Farabollini, P., Ragaini, L., Caffau, M.,
2157 2016. Sedimentology, faunal content and pollen record of Middle Pleistocene palustrine
2158 and lagoonal sediments from the Peri-Adriatic basin (Abruzzi, eastern central Italy).
2159 Quaternary Research in press.

2160 Pini, R., Ravazzi, C., Donegana, M., 2009. Pollen stratigraphy, vegetation and climate history of the
2161 last 215 ka in the Azzano Decimo core (plain of Friuli, north-eastern Italy). Quaternary
2162 Science Reviews 28, 1268–1290. doi:10.1016/j.quascirev.2008.12.017

2163 Pini, R., Ravazzi, C., Orombelli, G., 2003. Le ``Argille sotto il Ceppo'' di Paderno d'Adda: nuovi dati
2164 palinologici e biostratigrafici. Istituto Lombardo Rendiconti di Scienze B 136, 93–122.

2165 Pini, R., Ravazzi, C., Reimer, P.J., 2010. The vegetation and climate history of the last glacial cycle in
2166 a new pollen record from Lake Fimon (southern Alpine foreland, N-Italy). Quaternary
2167 Science Reviews 29, 3115–3137. doi:10.1016/j.quascirev.2010.06.040

2168 Pinti, D.L., Quidelleur, X., Chiesa, S., Ravazzi, C., Gillot, P.-Y., 2001. K–Ar dating of an early Middle
2169 Pleistocene distal tephra in the interglacial varved succession of Piànico-Sèllere (Southern
2170 Alps, Italy). Earth and Planetary Science Letters 188, 1–7. doi:10.1016/S0012-
2171 821X(01)00305-3

2172 Piotti, A., Leonarduzzi, C., Bagnoli, F., Spanu, I., Brousseau, L., Urbinati, C., Leonardi, S., Vendramin,
2173 G.G., 2016. Unexpected scenarios from Mediterranean refugial areas: disentangling
2174 complex demographic dynamics along the Apennine distribution of silver fir. Journal of
2175 Biogeography in press.

2176 Pons, A., Reille, M., 1988. The Holocene and Upper Pleistocene pollen record from Padul (Granada,
2177 Spain): a new study. Palaeogeography, Palaeoclimatology, Palaeoecology 66, 243–263.

2178 Pontini, M.R., Bertini, A., 2000. Late Pliocene vegetation and climate in Central Italy: high-resolution
2179 pollen analysis from the Fosso Bianco succession (Tiberino Basin). Geobios 33, 519–526.

2180 Popescu, S.-M., Biltekin, D., Winter, H., Suc, J.-P., Melinte-Dobrinescu, M.C., Klotz, S., Rabineau, M.,
2181 Combourieu-Nebout, N., Clauzon, G., Deaconu, F., 2010. Pliocene and Lower Pleistocene
2182 vegetation and climate changes at the European scale: Long pollen records and
2183 climatostratigraphy. Quaternary International 219, 152–167.
2184 doi:10.1016/j.quaint.2010.03.013

- 2185 Postigo Mijarra, J.M., Barrón, E., Gómez Manzaneque, F., Morla, C., 2009. Floristic changes in the
2186 Iberian Peninsula and Balearic Islands (south-west Europe) during the Cenozoic. *Journal of*
2187 *Biogeography* 36, 2025–2043. doi:10.1111/j.1365-2699.2009.02142.x
- 2188 Postigo Mijarra, J.M., Burjachs, F., Gómez Manzaneque, F., Morla, C., 2007. A palaeoecological
2189 interpretation of the lower-middle Pleistocene Cal Guardiola site (Terrassa, Barcelona, NE
2190 Spain) from the comparative study of wood and pollen samples. *Review of Palaeobotany*
2191 *and Palynology* 146, 247–264. doi:10.1016/j.revpalbo.2007.05.003
- 2192 Postigo Mijarra, J.M., Gómez Manzaneque, F., Morla, C., 2008. Survival and long-term maintenance
2193 of tertiary trees in the Iberian Peninsula during the Pleistocene: first record of *Aesculus* L.
2194 (Hippocastanaceae) in Spain. *Vegetation History and Archaeobotany* 17, 351–364.
2195 doi:10.1007/s00334-007-0130-x
- 2196 Postigo-Mijarra, J.M., Morla, C., Barrón, E., Morales-Molino, C., García, S., 2010. Patterns of
2197 extinction and persistence of Arctotertiary flora in Iberia during the Quaternary. *Review of*
2198 *Palaeobotany and Palynology* 162, 416–426. doi:10.1016/j.revpalbo.2010.02.015
- 2199 Pross, J., Koutsodendris, A., Christanis, K., Fischer, T., Fletcher, W.J., Hardiman, M., Kalaitzidis, S.,
2200 Knipping, M., Kotthoff, U., Milner, A.M., Müller, U.C., Schmiedl, G., Siavalas, G., Tzedakis,
2201 P.C., Wulf, S., 2015. The 1.35-Ma-long terrestrial climate archive of Tenaghi Philippon,
2202 northeastern Greece: Evolution, exploration, and perspectives for future research.
2203 *Newsletters on Stratigraphy* 48, 253–276. doi:10.1127/nos/2015/0063
- 2204 Ramil-Rego, P., Muñoz-Sobrino, C., Rodríguez-Gutián, M., Gómez-Orellana, L., 1998. Differences in
2205 the vegetation of the North Iberian Peninsula during the last 16,000 years. *Plant Ecology*
2206 138, 41–62.
- 2207 Ramos, J., Bernal, D., Dominguezbella, S., Calado, D., Ruiz, B., Gil, M., Clemente, I., Duran, J., Vijande,
2208 E., Chamorro, S., 2008. The Benzú rockshelter: a Middle Palaeolithic site on the North
2209 African coast. *Quaternary Science Reviews* 27, 2210–2218.
2210 doi:10.1016/j.quascirev.2008.08.030
- 2211 Ravazzi, C., 2003. Gli antichi bacini lacustri e i fossili di Leffe, Ranica e Pianico-Sèllere (Prealpi
2212 Lombarde). CNR Consiglio Nazionale delle Ricerche, Istituto per la dinamica dei processi
2213 ambientali.
- 2214 Ravazzi, C., 1994. Polline fossile di *Aesculus* aff. *hippocastanum* L. nel Bacino di Leffe (Pleistocene
2215 inferiore). Posizione sistematica e significato paleoecologico. *Giornale botanico italiano*
2216 128, 751–770. doi:10.1080/11263509409430319
- 2217 Ravazzi, C., Burg, van der, 1995. Coniferous woods in the Early Pleistocene brown coals of the Leffe
2218 Basin (Lombardy, Italy). *Rivista Italiana di Paleontologia e di Stratigrafia* 100, 597–620.
- 2219 Ravazzi, C., Donegana, M., Vescovi, E., Arpentini, E., Caccianiga, M., Kaltenrieder, P., Londeix, L.,
2220 Marabini, S., Mariani, S., Pini, R., Vai, G.B., Wick, L., 2006. A new Late-glacial site with *Picea*
2221 *abies* in the northern Apennine foothills: an exception to the model of glacial refugia of
2222 trees. *Vegetation History and Archaeobotany* 15, 357–371. doi:10.1007/s00334-006-0055-
2223 9
- 2224 Ravazzi, C., Pini, R., Breda, M., Martinetto, E., Muttoni, G., Chiesa, S., Confortini, F., Egli, R., 2005.
2225 The lacustrine deposits of Fornaci di Ranica (late Early Pleistocene, Italian Pre-Alps):
2226 stratigraphy, palaeoenvironment and geological evolution. *Quaternary International* 131,
2227 35–58. doi:10.1016/j.quaint.2004.07.021

- 2228 Ravazzi, C., Rossignol Strick, M., 1995. Vegetation change in a climatic cycle of Early Pleistocene age
2229 in the Leffe Basin (Northern Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology*
2230 117, 105–122.
- 2231 Reid, C., Reid, E.M., 1915. The Pliocene floras of the Dutch-Prussian border. M. Nijhoff, Gravenhage.
- 2232 Reid, C., Reid, E.M., 1907. Fossil flora of Tegelen-sur-Meuse, near Venloo, in the Province of Limburg,
2233 J. Müller. ed, *Verhandelingen der Koninklijke Akademie van Wetenschappen te*
2234 *Amsterdam, Afdeeling Natuurkunde. Tweede sectie, d. 13, no. 6. Amsterdam.*
- 2235 Reille, M., 1990. La tourbière de La Borde (Pyrénées orientales, France): un site clé pour l'étude du
2236 Tardiglaciaire sud-européen. *Comptes rendus de l'Académie des sciences. Série 2,*
2237 *Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la Terre* 310, 823–829.
- 2238 Reille, M., Beaulieu, J.D., Svobodova, H., Andrieu-Ponel, V., Goeury, C., 2000. Pollen analytical
2239 biostratigraphy of the last five climatic cycles from a long continental sequence from the
2240 Velay region (Massif Central, France). *Journal of Quaternary Science* 15, 665–685.
- 2241 Ricciardi, E., 1965. Analisi polliniche di una serie stratigrafica dei sedimenti lacustri del Pleistocene
2242 inferiore nel bacino di Leonessa (Rieti-Italia Centrale). *Giornale Botanico Italiano* 72, 62–
2243 82.
- 2244 Ricciardi, E., 1961. Analisi pollinologica dei sedimenti lacustri lignitiferi di Pietrafitta (Perugia, Italia).
2245 *Pollen et Spores* 3, 325–328.
- 2246 Rodríguez-Sánchez, F., Guzmán, B., Valido, A., Vargas, P., Arroyo, J., 2009. Late Neogene history of
2247 the laurel tree (*Laurus* L., Lauraceae) based on phylogeographical analyses of
2248 Mediterranean and Macaronesian populations. *Journal of Biogeography* 36, 1270–1281.
- 2249 Roger, S., Coulon, C., Thouveny, N., Féraud, G., Van Velzen, A., Fauquette, S., Cocheme, J.J., Prevot,
2250 M., Verosub, K.L., 2000. $^{40}\text{Ar}/^{39}\text{Ar}$ dating of a tephra layer in the Pliocene Senèze maar
2251 lacustrine sequence (French Massif Central): constraint on the age of the Réunion–
2252 Matuyama transition and implications on paleoenvironmental archives. *Earth and*
2253 *Planetary Science Letters* 183, 431–440.
- 2254 Rohais, S., Joannin, S., Colin, J.-P., Suc, J.-P., Guillocheau, F., Eschard, R., 2007. Age and
2255 environmental evolution of the syn-rift fill of the southern coast of the Gulf of Corinth
2256 (Akrata-Derveni region, Greece). *Bulletin de la Société géologique de France* 178, 231–243.
- 2257 Roiron, P., 1983. Nouvelle étude de la macroflore Plio-Pléistocène de Crespià (Catalogne, Espagne).
2258 *Geobios* 16, 687–715.
- 2259 Roucoux, K.H., Tzedakis, P.C., de Abreu, L., Shackleton, N.J., 2006. Climate and vegetation changes
2260 180,000 to 345,000 years ago recorded in a deep-sea core off Portugal. *Earth and Planetary*
2261 *Science Letters* 249, 307–325. doi:10.1016/j.epsl.2006.07.005
- 2262 Roucoux, K.H., Tzedakis, P.C., Frogley, M.R., Lawson, I.T., Preece, R.C., 2008. Vegetation history of
2263 the marine isotope stage 7 interglacial complex at Ioannina, NW Greece. *Quaternary*
2264 *Science Reviews* 27, 1378–1395. doi:10.1016/j.quascirev.2008.04.002
- 2265 Russo Ermolli, E., 1994. Analyse pollinique de la succession lacustre pléistocène du Vallo di Diano
2266 (Campanie, Italie). *Annales de la Société géologique de Belgique* 117, 333–354.
- 2267 Russo Ermolli, E., Aucelli, P.P.C., Di Rollo, A., Mattei, M., Petrosino, P., Porreca, M., Roskopf, C.M.,
2268 2010a. An integrated stratigraphical approach to the Middle Pleistocene succession of the
2269 Sessano basin (Molise, Italy). *Quaternary International* 225, 114–127.
2270 doi:10.1016/j.quaint.2009.04.008

- 2271 Russo Ermolli, E., Di Donato, V., Martín-Fernández, J.A., Orain, R., Lebreton, V., Piovesan, G., 2015.
2272 Vegetation patterns in the Southern Apennines (Italy) during MIS 13: Deciphering pollen
2273 variability along a NW-SE transect. *Review of Palaeobotany and Palynology* 218, 167–183.
2274 doi:10.1016/j.revpalbo.2014.11.004
- 2275 Russo Ermolli, E., Sardella, R., Di Maio, G., Petronio, C., Santangelo, N., 2010b. Pollen and mammals
2276 from the late Early Pleistocene site of Saticula (Sant'Agata de' Goti, Benevento, Italy).
2277 *Quaternary International* 225, 128–137. doi:10.1016/j.quaint.2009.06.013
- 2278 Sabato, L., Bertini, A., Masini, F., Albianelli, A., Napoleone, G., Pieri, P., 2005. The lower and middle
2279 Pleistocene geological record of the San Lorenzo lacustrine succession in the
2280 Sant'Arcangelo Basin (Southern Apennines, Italy). *Quaternary International* 131, 59–69.
2281 doi:10.1016/j.quaint.2004.07.001
- 2282 Sadori, L., Giardini, M., Chiarini, E., Mattei, M., Papasodaro, F., Porreca, M., 2010. Pollen and
2283 macrofossil analyses of Pliocene lacustrine sediments (Salto river valley, Central Italy).
2284 *Quaternary International* 225, 44–57. doi:10.1016/j.quaint.2009.05.008
- 2285 Sadori, L., Koutsodendris, A., Panagiotopoulos, K., Masi, A., Bertini, A., Combourieu-Nebout, N.,
2286 Francke, A., Kouli, K., Joannin, S., Mercuri, A.M., Peyron, O., Torri, P., Wagner, B., Zanchetta,
2287 G., Sinopoli, G., Donders, T.H., 2016. Pollen-based paleoenvironmental and paleoclimatic
2288 change at Lake Ohrid (south-eastern Europe) during the past 500 ka. *Biogeosciences* 13,
2289 1423–1437. doi:10.5194/bg-13-1423-2016
- 2290 Sánchez Goñi, M.F., Hannon, G.E., 1999. High-altitude vegetational pattern on the Iberian Mountain
2291 Chain (north-central Spain) during the Holocene. *The Holocene* 9, 39–57.
- 2292 Sánchez Goñi, M.F., Landais, A., Fletcher, W.J., Naughton, F., Desprat, S., Duprat, J., 2008.
2293 Contrasting impacts of Dansgaard-Oeschger events over a western European latitudinal
2294 transect modulated by orbital parameters. *Quaternary Science Reviews* 27, 1136–1151.
2295 doi:10.1016/j.quascirev.2008.03.003
- 2296 Sánchez Goñi, M.F., Loutre, M.F., Crucifix, M., Peyron, O., Santos, L., Duprat, J., Malaizé, B., Turon,
2297 J.-L., Peypouquet, J.-P., 2005. Increasing vegetation and climate gradient in Western Europe
2298 over the Last Glacial Inception (122–110 ka): data-model comparison. *Earth and Planetary
2299 Science Letters* 231, 111–130. doi:10.1016/j.epsl.2004.12.010
- 2300 Sanchez Goñi, M.F.S., 1996. Vegetation and sea level changes during the Holocene in the estuary of
2301 the Bidasoa (Southern part of the bay of Biscay). *Quaternaire* 7, 207–219.
- 2302 Sánchez Goñi, M.S., Eynaud, F., Turon, J., Shackleton, N., 1999. High resolution palynological record
2303 off the Iberian margin: direct land-sea correlation for the Last Interglacial complex. *Earth
2304 and Planetary Science Letters* 171, 123–137.
- 2305 Sánchez-Goñi, M.F., 1992. Analyse palynologique de sites préhistoriques de Pays Basque: premiers
2306 résultats pour les grottes de Lezetxiki et Urtiaga. The late Quaternary in the western
2307 Pyrenean region. *Universidad del País Vasco* 207–233.
- 2308 Sánchez-Goñi, M.-F., Cacho, I., Turon, J.L., Guiot, J., Sierro, F.J., Peypouquet, J.-P., Grimalt, J.O.,
2309 Shackleton, N.J., 2002. Synchronicity between marine and terrestrial responses to
2310 millennial scale climatic variability during the last glacial period in the Mediterranean
2311 region. *Climate Dynamics* 19, 95–105. doi:10.1007/s00382-001-0212-x
- 2312 Sefidi, K., Marvie Mohadjer, M.R., Etemad, V., Copenheaver, C.A., 2011. Stand characteristics and
2313 distribution of a relict population of Persian ironwood (*Parrotia persica* C.A. Meyer) in

- 2314 northern Iran. Flora - Morphology, Distribution, Functional Ecology of Plants 206, 418–422.
2315 doi:10.1016/j.flora.2010.11.005
- 2316 Šercelj, A., 1965. Lower Pleistocene flora from Bukovica near Ilirska Bistrica. Slovenska Akademija
2317 Znanosti in Umetnosti 8, 439–470.
- 2318 Šercelj, A., 1961. The lower Pleistocene vegetation from Zalog near Novo Mesto. Slovenska
2319 Akademija Znanosti in Umetnosti 6, 417–434.
- 2320 Shatilova, I., Mchedlishvili, N., Rukhadze, L., Kvavadze, E., 2011. The history of the flora and
2321 vegetation of Georgia (South Caucasus). Georgian national museum, Tbilisi.
- 2322 Sheykhoulislami, A., Ahmadi, T., 2009. The Study of Caucasian Walnut (*Pterocarya frexinifolia* (Lam.)
2323 Spach.) in Forests of Mashelak (Noshahr, Iran). Botany Research Journal 2, 28–33.
- 2324 Sivak, J., 1976. Nouvelles espèces du genre *Cathaya* d'après leurs grains de pollen dans le tertiaire
2325 du sud de la France. Pollen et spores 18, 243–288.
- 2326 Slimani, S., Derridj, A., Gutierrez, E., 2014. Ecological response of *Cedrus atlantica* to climate
2327 variability in the Massif of Guetiane (Algeria). Forest Systems 23, 448.
2328 doi:10.5424/fs/2014233-05175
- 2329 Stewart, J.R., Lister, A.M., 2001. Cryptic northern refugia and the origins of the modern biota. Trends
2330 in Ecology & Evolution 16, 608–613.
- 2331 Su, T., Liu, Y.-S. (Christopher), Jacques, F.M.B., Huang, Y.-J., Xing, Y.-W., Zhou, Z.-K., 2013. The
2332 intensification of the East Asian winter monsoon contributed to the disappearance of
2333 *Cedrus* (Pinaceae) in southwestern China. Quaternary Research 80, 316–325.
2334 doi:10.1016/j.yqres.2013.07.001
- 2335 Subally, D., Bilodeau, G., Hillaire-Marcel, C., Tamrat, E., Ferry, S., Debard, E., 1999. Cyclic climatic
2336 records during the Olduvai subchron (Uppermost Pliocene) on Zakynthos Island (Ionian
2337 Sea). Geobios 32, 793–803.
- 2338 Suc, J.-P., 1984. Origin and evolution of the Mediterranean vegetation and climate in Europe. Nature
2339 307, 429–432. doi:10.1038/307429a0
- 2340 Suc, J.P., 1978. Analyse pollinique de dépôts plio-pléistocènes du sud du massif basaltique de
2341 l'Escandorgue (site de Bernasso, Lunas, Hérault, France). Pollen et Spores 20, 497–512.
- 2342 Suc, J.P., 1973. Etude palynologique des marnes de Celleneuve (Pléistocène inférieur). Hérault.
2343 Bulletin de l'Association française pour l'étude du quaternaire 10, 13–24.
2344 doi:10.3406/quate.1973.1214
- 2345 Suc, J.-P., Combourieu-Nebout, N., Seret, G., Popescu, S.-M., Klotz, S., Gautier, F., Clauzon, G.,
2346 Westgate, J., Insinga, D., Sandhu, A.S., 2010. The Croton series: A synthesis and new data.
2347 Quaternary International 219, 121–133. doi:10.1016/j.quaint.2010.01.008
- 2348 Suc, J.-P., Cravatte, J., 1982. Étude palynologique du Pliocène de Catalogne (nord-est de l'Espagne):
2349 apports à la connaissance de l'histoire climatique de la Méditerranée occidentale et
2350 implications chronostratigraphiques. Paléobiologie continentale 13, 1–31.
- 2351 Suc, J.-P., Drivaliari, A., 1991. Transport of bisaccate coniferous fossil pollen grains to coastal
2352 sediments: An example from the earliest Pliocene Orbria (Languedoc, southern France).
2353 Review of Palaeobotany and Palynology 70, 247–253. doi:10.1016/0034-6667(91)90006-O

- 2354 Suc, J.-P., Popescu, S.-M., 2005. Pollen records and climatic cycles in the North Mediterranean region
2355 since 2.7 Ma. Geological Society, London, Special Publications 247, 147–158.
2356 doi:10.1144/GSL.SP.2005.247.01.08
- 2357 Svenning, J.-C., 2003. Deterministic Plio-Pleistocene extinctions in the European cool-temperate
2358 tree flora. Ecology Letters 6, 646–653.
- 2359 Svenning, J.-C., Eiserhardt, W.L., Normand, S., Ordonez, A., Sandel, B., 2015. The Influence of
2360 Paleoclimate on Present-Day Patterns in Biodiversity and Ecosystems. Annual Review of
2361 Ecology, Evolution, and Systematics 46, 551–572. doi:10.1146/annurev-ecolsys-112414-
2362 054314
- 2363 Tallantire, P.A., 2002. The early-Holocene spread of hazel (*Corylus avellana* L.) in Europe north and
2364 west of the Alps: an ecological hypothesis. The Holocene 12, 81–96.
2365 doi:10.1191/0959683602hl523rr
- 2366 Tzedakis, P., Frogley, M., Heaton, T., 2003. Last Interglacial conditions in southern Europe: evidence
2367 from Ioannina, northwest Greece. Global and Planetary Change 36, 157–170.
- 2368 Tzedakis, P.C., 1999. The last climatic cycle at Kopais, central Greece. Journal of the Geological
2369 Society 156, 425–434.
- 2370 Tzedakis, P.C., 1994. Vegetation change through glacial-interglacial cycles: a long pollen sequence
2371 perspective. Philosophical Transactions of the Royal Society of London B: Biological
2372 Sciences 345, 403–432.
- 2373 Tzedakis, P.C., Emerson, B.C., Hewitt, G.M., 2013. Cryptic or mystic? Glacial tree refugia in northern
2374 Europe. Trends in Ecology & Evolution 28, 696–704. doi:10.1016/j.tree.2013.09.001
- 2375 Tzedakis, P.C., Hooghiemstra, H., Pälike, H., 2006. The last 1.35 million years at Tenaghi Philippon:
2376 revised chronostratigraphy and long-term vegetation trends. Quaternary Science Reviews
2377 25, 3416–3430. doi:10.1016/j.quascirev.2006.09.002
- 2378 Tzedakis, P.C., Margari, V., Hodell, D.A., 2015. Coupled ocean–land millennial-scale changes
2379 1.26millionyears ago, recorded at Site U1385 off Portugal. Global and Planetary Change
2380 135, 83–88. doi:10.1016/j.gloplacha.2015.10.008
- 2381 Urban, B., Arias, C., Bigazzi, G., Bonadonna, F., 1983. Early Pleistocene Palynostratigraphy of Fornace
2382 Tini, Valle Ricca (central Italy). Palaeogeography, Palaeoclimatology, Palaeoecology 41,
2383 153–164.
- 2384 Uzquiano, P., 1995. La disparition de *Picea* à la fin du Pléistocène supérieur en région cantabrique
2385 d'après l'anthrocoanalyse: déterminisme climatique et anthropique. Comptes rendus de
2386 l'Académie des sciences. Série 2. Sciences de la terre et des planètes 321, 545–551.
- 2387 Van der Wiel, A., Wijmstra, T., 1987a. Palynology of the 112.8–197.8 m interval of the core Tenaghi
2388 Philippon III, Middle Pleistocene of Macedonia. Review of Palaeobotany and Palynology 52,
2389 89–117.
- 2390 Van der Wiel, A., Wijmstra, T., 1987b. Palynology of the lower part (78–120 m) of the core Tenaghi
2391 Philippon II, Middle Pleistocene of Macedonia, Greece. Review of Palaeobotany and
2392 Palynology 52, 73–88.
- 2393 van Kolfschoten, T., Markova, A.K., 2005. Response of the European mammalian fauna to the mid-
2394 Pleistocene transition. Geological Society, London, Special Publications 247, 221–229.
- 2395 Velitzelos, D., Bouchal, J.M., Denk, T., 2014. Review of the Cenozoic floras and vegetation of Greece.
2396 Review of Palaeobotany and Palynology 204, 56–117. doi:10.1016/j.revpalbo.2014.02.006

- 2397 Vieira, M.C.M., 2009. Palinologia do Pliocénico da Orla Ocidental Norte e Centro de Portugal:
2398 Contributo para a compreensão da cronostratigrafia e da evolução paleoambiental (PhD
2399 Thesis). Universidad do Minho, Braga, Portugal.
- 2400 Villalta, J. de, Vicente, J., 1972. Una flora del Cuaternario antiguo en las cercanías de Crespià. Acta
2401 geológica hispánica 7, 120–128.
- 2402 Wagner, B., Lotter, A.F., Nowaczyk, N., Reed, J.M., Schwalb, A., Sulpizio, R., Valsecchi, V., Wessels,
2403 M., Zanchetta, G., 2009. A 40,000-year record of environmental change from ancient Lake
2404 Ohrid (Albania and Macedonia). Journal of Paleolimnology 41, 407–430.
2405 doi:10.1007/s10933-008-9234-2
- 2406 Wagner, B., Wilke, T., Krastel, S., Zanchetta, G., Sulpizio, R., Reicherter, K., Leng, M.J., Grazhdani, A.,
2407 Trajanovski, S., Francke, A., Lindhorst, K., Levkov, Z., Cvetkoska, A., Reed, J.M., Zhang, X.,
2408 Lacey, J.H., Wonik, T., Baumgarten, H., Vogel, H., 2014. The SCOPSCO drilling project
2409 recovers more than 1.2 million years of history from Lake Ohrid. Scientific Drilling 17, 19–
2410 29. doi:10.5194/sd-17-19-2014
- 2411 Wang, Y.-F., Li, C.-S., Collinson, M.E., Lin, J., Sun, Q.-G., 2003. *Eucommia* (Eucommiaceae), a potential
2412 biothermometer for the reconstruction of paleoenvironments. American journal of botany
2413 90, 1–7.
- 2414 West, R., 1970. Pleistocene history of the British flora, in: D. Walker and R.G. West (Eds) Studies in
2415 the Vegetational History of the British Isles. Cambridge University Press, pp. 1–11.
- 2416 West, R., 1962. Vegetational history of the Early Pleistocene of the Royal Society borehole at
2417 Ludham, Norfolk. Proceedings of the Royal Society of London B: Biological Sciences 155,
2418 437–453.
- 2419 Wijmstra, T., 1969. Palynology of the first 30 metres of a 120 m deep section in northern Greece.
2420 Acta Botanica Neerlandica 18, 511–527.
- 2421 Wijmstra, T., Smit, A., 1976. Palynology of the middle part (30–78 metres) of the 120 m deep section
2422 in northern Greece (Macedonia). Acta Botanica Neerlandica 25, 297–312.
- 2423 Worth, J.R., Sakaguchi, S., Tanaka, N., Yamasaki, M., Isagi, Y., 2013. Northern richness and southern
2424 poverty: contrasting genetic footprints of glacial refugia in the relictual tree *Sciadopitys*
2425 *verticillata* (Coniferales: Sciadopityaceae). Biological Journal of the Linnean Society 108,
2426 263–277.
- 2427 Yavuz-İşık, N., 2007. Pollen analysis of coal-bearing Miocene sedimentary rocks from the Seyitömer
2428 Basin (Kütahya), Western Anatolia. Geobios 40, 701–708.
- 2429 Yll, E.I., Carrión, J.S., Pantaleón-Cano, J., Dupré, M., La Roca, N., Roure, J.M., Pérez-Obiol, R., 2003.
2430 Palinología del Cuaternario reciente en la Laguna de Villena (Alicante). Anales de Biología
2431 25, 65–72.
- 2432 Yll, E.-I., Perez-Obiol, R., Pantaleon-Cano, J., Roure, J.M., 1997. Palynological evidence for climatic
2433 change and human activity during the Holocene on Minorca (Balearic Islands). Quaternary
2434 Research 48, 339–347.
- 2435 Zagwijn, W., 1985. An outline of the Quaternary stratigraphy of the Netherlands. Geologie en
2436 Mijnbouw 64, 17–24.
- 2437 Zagwijn, W., De Jong, J., 1984. Die Interglaziale von Bavel und Leerdam und ihre stratigraphische
2438 Stellung im niederländischen Früh-Pleistozän. Mededelingen Rijks Geologische Dienst 37,
2439 155–169.

- 2440 Zagwijn, W.H., 1974. The Pliocene-Pleistocene boundary in western and southern Europe. *Boreas* 3,
2441 75–97.
- 2442 Zagwijn, W.H., 1963. Pleistocene stratigraphy in the Netherlands, based on changes in vegetation
2443 and climate. *Verh. Kon. Ned. Geol. Mijnbouwk. Gen., Geol. Serie* 21–2.
- 2444 Zagwijn, W.H., 1957. Vegetation, climate and time-correlations in the Early Pleistocene of Europe.
2445 *Geologie en Mijnbouw* 19, 233–244.
- 2446 Zanni, M., Ravazzi, C., 2007. Description and differentiation of *Pseudolarix amabilis* pollen
2447 Palaeoecological implications and new identification key to fresh bisaccate pollen. *Review*
2448 *of Palaeobotany and Palynology* 145, 35–75. doi:10.1016/j.revpalbo.2006.08.004
2449

Figure captions:

Fig. 1. Location of the Quaternary records considered in this paper. Numbering of the sites follows Table 1. Red dots represent the peri-Iberian marine sites included in Fig. 2; blue dots correspond to the sites from NW Iberia and S France represented in Fig. 2; red squares include sites from N Italy (Fig. 3); yellow dots correspond to sites from central Italy of Fig. 4; pink dots are sites from S Italy represented in Fig. 4; green dots include Greece and the Aegean (Fig. 4); the blue square corresponds to the site of Tenaghi Philippon (Fig. 5) and the pink square to DSDP site 380 (Fig. 5). White dots indicate the location of sites not represented in Figs 2-5 but cited in the text.

Fig. 2. Stratigraphical setting and schematic pollen biostratigraphy of selected taxa of Pleistocene records from the Alboran Sea and off Portugal, and from NW Iberia and S France. Question marks indicate uncertainties in the chronological setting of the records. Asterisks indicate plant macrofossils. Red lines represent regionally extinct taxa. Blue lines represent taxa that are only mentioned in the papers or uncertain determinations (e.g. *Ulmus/Zelkova*).

Fig. 3. Stratigraphical setting and schematic pollen biostratigraphy of selected taxa of Pleistocene records from N Italy (>44°N) and Central Italy (44°-41°N). Note that the sites of Sessano and Boiano are administratively located in S Italy. Question marks indicate uncertainties in the chronological setting of the records. Asterisks indicate plant macrofossils. Red lines represent regionally extinct taxa. Blue lines represent taxa that are only mentioned in the papers or uncertain determinations (e.g. *Ulmus/Zelkova*).

Fig. 4. Stratigraphical setting and schematic pollen biostratigraphy of selected taxa of Pleistocene records from S Italy (<41°N), Greece and the Aegean. Question marks indicate uncertainties in the chronological setting of the records. Asterisks indicate plant macrofossils. Red lines represent regionally extinct taxa. Blue lines represent taxa that are only mentioned in the papers or uncertain determinations (e.g. *Ulmus/Zelkova*).

Fig. 5. Stratigraphical setting and schematic pollen biostratigraphy of selected taxa of Pleistocene records from Tenaghi Philippon ([Tzedakis et al., 2006](#)) and DSDP site 380 ([Biltekin et al. 2015](#)). Red lines represent regionally extinct taxa. Blue lines represent taxa that are only mentioned in the papers or uncertain determinations (e.g. *Ulmus/Zelkova*).

Fig. 6. Schematic stratigraphic distribution of *Taxodium* type, *Sciadopitys*, *Cathaya* type, and *Cedrus*. Blue lines represent taxa that are only mentioned in the original papers.

Fig. 7. Schematic stratigraphic distribution of *Tsuga*, *Eucommia*, *Parrotia*, and *Liquidambar*. Blue lines represent taxa that are only mentioned in the original papers.

Fig. 8 Schematic stratigraphic distribution of *Engelhardia*, *Carya*, *Pterocarya*, and *Zelkova*. Blue lines represent taxa that are only mentioned in the original papers.

Table 1. List of the vegetation records represented in Fig. 1.

Site number	Site	Age (ka) MIS	Pollen (P) Macrofossil (M)	Marine (M) Continental (C)	Chronological control	References
1	MD95-2043	48-2 ka	P	M	14C, event stratigraphy	Sanchez Goñi et al., 2002; Fletcher and Sanchez Goñi, 2008
2	Padul	100-0 ka	P	C	14C and vegetation history	Pons and Reille, 1988
3	ODP976	MIS 31-23	P	M	Oxygen isotope stratigraphy	Joannin 2007; Joannin et al., 2011; Combourieu Nebout et al., 2009
4	Andalucia G1	Miocene-Gelasian	P	M	Oxygen isotope stratigraphy	Feddi et al., 2011
5	Morgandinho	Early-Middle Pleistocene	P	C	Micromammals	Antunes et al., 1986
6	Site U1385	MIS 38	P	M	Orbital tuning	Tzedakis et al., 2015
7	MD95-2042	MIS 6-2	P	M	14C and Oxygen isotope stratigraphy	Sanchez Goñi et al., 1999
8	MD01-2443	345-180 ka	P	M	Oxygen isotope stratigraphy	Roucoux et al., 2006
9	Monsarros	Plio-Pleistocene	P	C	Stratigraphy	Vieira, 2009
10	MD03-2697	340-270 ka	P	M	Oxygen isotope stratigraphy	Desprat et al., 2009
11	MD01-2447	430-360 ; 250-180 ka	P	M	Oxygen isotope stratigraphy	Desprat et al., 2005
12	MD99-2331	MIS 5-2	P	M	14C, event stratigraphy	Sanchez-Goñi et al., 2005, 2008
13	Santa Maria de Oia	MIS 5-3	P	C-M (coastal)	14C, coastal stratigraphy	Gómez-Orellana et al., 2013
14	Area Longa	MIS 5-3	P	C-M (coastal)	14C, coastal stratigraphy	Gómez-Orellana et al., 2007
15	Pointe de la Négade North	Early Pleistocene	P+M	M	Pollen biostratigraphy	O'Brien and Jones, 2003
16	La Balise	MIS 11	P+M	M	Pollen biostratigraphy	O'Brien and Jones, 2003
17	Lezetxiki	MIS 3	P	C	14C, archaeological data (lithic tools)	Sánchez-Goñi, 1992
18	Atapuerca	900-400 ka	P	C	Luminiscence	García-Antón & Sainz-Ollero, 1991
19	Cañizar de Villaquemado	MIS 5-1	P	C	14C and OSL dating	González-Sampériz et al., 2013; Aranbarri et al., 2014; García-Prieto, 2015

20	Garraf	5300-1750 ka	P	M	Foraminifer biostratigraphy	Suc and Cravatte, 1982
21	Abric Romaní	70-40 ka	P	C	U/Th	Burjachs et al., 2012
22	Cal Guardiola	1200-800 ka	P-M	C	Micromammals and palaeomagnetism	Postigo Mijarra et al., 2007
23	Val d'en Bas	250 ka	P	C	Volcanic crust	Cross et al., 1986
24	Pla de l'Estany-Garrotxa	MIS 5?-1	P	C	Vegetation history and 14C for the upper part	Burjachs, 1990
25	Pla de l'Estany-Incarcal-Crespià	Plio-Pleistocene	M	C	Fauna (Mammals biostratigraphy)	Villalta and Vicente, 1972; Suc and Cravatte, 1982; Roiron, 1983
26	Tres Pins	2600-1200 ka	P	C	Vegetation composition (presence of carya)	Leroy, 1997
27	Mas Grill	Mid-Pleistocene?	P	C	Vegetation history (older than Bòbila Ordis)	Geurts, 1977; 1979
28	Bòbila-Ordis	MIS 36-33	P	C	Micromammals and palaeomagnetism	Løvlie and Leroy, 1995; Leroy, 2008
29	Banyoles	1000-950 ka	P	C		Julia Bruguès and Suc, 1980
30	Molí Vell	Early Pleistocene	P	C	Vegetation history (younger than Bòbila Ordis)	Geurts, 1977, 1979
31	Mas Miquel	Early Pleistocene	P	C	Fauna (Mammals biostratigraphy)	Leroy, 1990
32	Saint-Macaire	1400-700 ka	P	C	K/Ar dating, magnetostratigraphy	Leroy et al., 1994
33	Nogaret	MIS 75	P	C	K/Ar dating, mammal biostratigraphy	Brugal et al., 1990; Leroy and Seret, 1992
34	Bernasso	2160-1960 ka	P	C	K/Ar dating, magnetostratigraphy	Suc, 1978; Leroy and Roiron, 1996
35	Celleneuve	Early Calabrian	P	C-M	Biostratigraphy	Suc, 1973
36	Alleret	MIS 17-15	P	C	Tephrochronology, K/Ar and Ar/Ar dating	Pastre et al., 2007; Degeai et al., 2013
37	Senèze	MIS 85-76	P	C	Ar/Ar dating, mammal biostratigraphy, magnetostratigraphy	Elhaï, 1969; Roger et al., 2000
38	Velay maars	430-0 ka	P	C	Biostratigraphy, Ar/Ar dating	Reille et al., 2000
39	Ceyssac	1000-900 ka	P	C	K/Ar dating, magnetostratigraphy	Ablin, 1991
40	Saint-Vallier	1900 ka	P	C	Biostratigraphy	Argant, 2004
41	La Côte	MIS 11	P	C	Pollen biostratigraphy	Field et al., 2000
42	Fornaci di Ranica	MIS (36)30 or 28	P,M	C	Magnetostratigraphy, mammal biostratigraphy	Ravazzi et al., 2005
43	Leffe-Fornace Martinelli	MIS 64-22	P,M	C	Magnetostratigraphy, mammal biostratigraphy	Ravazzi and Rossignol-Strick, 1995; Ravazzi, 2003; Muttoni et al.,

						2007
44	Pianico Sèllere	MIS 21-19	P, M	C	K/Ar and Ar/Ar dating, mammal biostratigraphy	Ravazzi, 2003; Martinetto, 2009
45	Pianengo	1200-700 ka	P	C	Magnetostratigraphy	Muttoni et al., 2003
46	Stirone	5100-2200 ka	P	M	Mollusc and foraminifer biostratigraphy	Lona and Bertoldi, 1972; Bertini, 2001
47	Castell'Arquato	Early Pleistocene	P	M	Mollusc biostratigraphy	Lona and Bertoldi, 1972
48	San Pietro di Ragogna	Gelasian	P, M	C	Lithostratigraphy, biostratigraphy	Martinetto et al., 2012
49	Azzano Decimo	215-20 ka	P	C	¹⁴ C dating, biostratigraphy	Pini et al., 2009
50	Visogliano	MIS 13 or 11	P	M	Mammal biostratigraphy	Cattani et al., 1991; Abbazzi et al., 2000
51	Venice	2150-110 ka	P	M	Magnetostratigraphy, molluscs biostratigraphy, sapropel stratigraphy	Mullenders et al., 1996, Massari et al., 2004
52	Core 223-S17	130-0 ka	P	C/M	Foraminifer and ostracod biostratigraphy	Amorosi et al., 2004
53	Oriolo (Faenza)	ca. 800 ka	M	C/M	Magnetostratigraphy, molluscs biostratigraphy	Martinetto and Sami, 2001
54	Lamone Valley	MIS 64-46	P	M	Mollusc and ostracod biostratigraphy	Fusco, 2007
55	Monte Poggiolo	1400-1070? ka	P	M	Magnetostratigraphy, molluscs biostratigraphy, ESR dating	Messenger et al., 2011
56	Poggio Rosso	1900-1800 ka	P	C	Magnetostratigraphy	Bertini et al., 2010
57	Gubbio	1600-1300? Ka	P	C	Biostratigraphy	Lona and Ricciardi, 1961
58	Colle Curti	ca. 990 ka	P	C	Magnetostratigraphy	Bertini, 2000
59	Cesi	750-600? ka	P	C	Magnetostratigraphy, Tephrochronology	Bertini, 2000
60	Pietrafitta	1400? ka	P	C	Mammal biostratigraphy	Ricciardi, 1961; Martinetto et al., 2014
61	Fosso Bianco	MIS 100-82	P	C	Magnetostratigraphy, Mammal biostratigraphy	Pontini and Bertini 2000
62	Dunarobba	3500-1500? ka	P, M	C	Biostratigraphy	Martinetto et al., 2014
63	Leonessa	1400-1200? ka	P	C	Biostratigraphy	Ricciardi, 1965
64	Madonna della Strada	ca. 1250 ka	P	C	Mammal biostratigraphy	Magri et al., 2010
65	Pagliare di Sassa	1100 ka	P	C	Magnetostratigraphy, mammal	Palombo et al., 2010

					biostratigraphy	
66	Torre Mucchia (Ortona)	MIS 17	P	M	Magnetostratigraphy	Pieruccini et al., 2016
67	Carsoli	ca. 530	M	C	Ar/Ar dating	Sadori et al., 2010
68	Peperino (Rome)	513±3 ka	M	C	Ar/Ar dating	Follieri, 2010; Marra et al., 2014
69	Riano	450-350? ka	M	C	Mammal biostratigraphy	Follieri, 1958, 1962
70	Valle Ricca	2100-1800 ka	P	M	Magnetostratigraphy, biostratigraphy	Urban et al., 1983; Arias et al., 1990
71	Valle di Castiglione	250-0 ka	P	C	Varve counting, astronomic calibration	Follieri et al., 1988, 1989; Di Rita et al., 2013
72	Coste San Giacomo	ca. 2100 ka	P	C	Magnetostratigraphy, mammal biostratigraphy	Bellucci et al., 2014
73	Fontana Ranuccio	MIS 23	P	C	Magnetostratigraphy	Corrado and Magri, 2011
74	Ceprano	MIS 15-13	P	C	Magnetostratigraphy	Manzi et al., 2010
75	Isernia - La Pineta	900-700 ka	P	C	Magnetostratigraphy, Ar/Ar dating	Lebreton, 2002
76	Sessano	MIS 14-13	P	C	Magnetostratigraphy, Tephrochronology, Ar/Ar dating	Russo Ermolli et al., 2010a, 2015
77	Boiano	MIS 13-9	P	C	Ar/Ar dating, tephrochronology	Orain et al., 2013, 2015; Russo Ermolli et al., 2015
78	Saticula	ca. 1200 ka	P	C	Mammal biostratigraphy	Russo Ermolli et al., 2010b
79	Acerno	MIS 13-12	P	C	tephra, Ar/Ar dating	Munno et al., 2001; Russo Ermolli et al., 2015
80	Lago Grande di Monticchio	130÷0	P	C	14C dating, tephrochronology, varve counting	Allen et al., 2000; Allen and Huntley, 2009
81	Vallo di Diano	650-450 ka	P	C	tephra, Ar/Ar dating	Russo Ermolli, 1994; Russo Ermolli et al., 2015
82	Camerota	1750-1600? ka	P	C	Biostratigraphy	Brenac, 1984
83	Mercure	MIS 13	P	C	Tephrochronology, Ar/Ar	Petrosino et al., 2014
84	San Lorenzo	1200-1100 ka	P	C	Magnetostratigraphy, mammal biostratigraphy	Sabato et al., 2005
85	Montalbano Jonico	1250-900 ka	P	M	Ar/Ar dating, Oxygen isotope stratigraphy	Joannin et al., 2008; Bertini et al., 2015
86	Semaforo-Vrica	2460-2110 ka	P	M	Magnetostratigraphy, foraminifer biostratigraphy, K/Ar dating	Combourieu-Nebout and Vergnaud-Grazzini, 1991; Combourieu-Nebout, 1993; Klotz et al., 2006
87	Santa Lucia	MIS 43-40	P	M	Biostratigraphy, Oxygen isotope stratigraphy	Joannin et al., 2007b

88	Valle di Manche	MIS 20-18	P	M	Magnetostratigraphy, Oxygen isotope stratigraphy, Tephrochronology, Biostratigraphy, Ar/Ar dating	Capraro et al., 2005
89	Ohrid	500-0 ka	P	C	¹⁴ C, tephra, stratigraphy	Wagner et al., 2009, 2014; Sadori et al., 2016
90	Ioannina	450-0 ka	P	C	Pollen biostratigraphy	Tzedakis, 1994; Roucoux et al., 2008
91	Achelooos Delta	5-3 ka	P	M	¹⁴ C dating	Fouache et al., 2005
92	Zakynthos	2000-1800 ka	P	M	Magnetostratigraphy, oxygen isotope stratigraphy	Subally et al., 1999
93	Gulf of Corinth	1800-600 ka	P	M	Pollen biostratigraphy, regional stratigraphical correlations	Rohais et al., 2007
94	Kopais	500-2 ka	P	C	Pollen biostratigraphy	Tzedakis, 1999; Okuda et al., 2001
95	Megalopolis Basin	650-300 ka	P	C	Magnetostratigraphy, palaeontology	Okuda et al., 2002
96	Kallithea	Calabrian	M	C	Mollusc biostratigraphy	Mai and Velitzelos, 2007; Velitzelos et al., 2014
97	Kolymia Bay	500 ka	M	M	Magnetostratigraphy	Boyd, 2009
98	Tsampika Bay	MIS 42-1	P	M	Magnetostratigraphy, Oxygen isotope stratigraphy	Joannin et al., 2007a
99	Tenaghi Philippon	1350-0 ka	P	C	Pollen-orbital tuning	Wijmstra, 1969; Wijmstra and Smit, 1976; Van der Wiel and Wijmstra, 1987a,b; Tzedakis et al., 2006; Fletcher et al., 2013; Pross et al., 2015
100	DSDP Site 380	6000-0 ka	P	M	Nannoplankton	Biltekin, 2010; Popescu et al., 2010; Biltekin et al., 2015

Highlights:

- The Quaternary disappearance of tree taxa in S Europe appears distributed over time
- Simple North to South and/or West to East trend in extirpations cannot be recognized
- Some taxa disappeared from wide areas of S Europe in the Lateglacial and Holocene