Paternal contributions to large-for-gestational-age term babies

DOI:

[10.1017/S2040174419000035](https://doi.org/10.1017/S2040174419000035)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Derraik, J. G. B., Pasupathy, D., McCowan, L. M. E., Poston, L., Taylor, R. S., Simpson, N. A. B., Dekker, G. A., Myers, J., Vieira, M. C., Cutfield, W. S., & Ahlsson, F. (2019). Paternal contributions to large-for-gestational-age term babies: Findings from a multicenter prospective cohort study. *Journal of Developmental Origins of Health and Disease*. <https://doi.org/10.1017/S2040174419000035>

Published in:

Journal of Developmental Origins of Health and Disease

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.

Paternal contributions to large-for-gestational-age term babies: findings from a multicenter prospective cohort study

José G B Derraik^{1,2,3}, Dharmintra Pasupathy⁴, Lesley M E McCowan^{5,6}, Lucilla Poston⁴,
Rennae S Taylor⁵, Nigel A B Simpson⁷, Gustaaf A Dekker⁸, Jenny Myers⁹, Matias Costa
Vieira⁴, Wayne S Cutfield^{1,2}, Fredrik Ahlsson^{3,*}; on behalf of the SCOPE consortium.

¹ Liggins Institute, University of Auckland, Auckland, New Zealand.

² A Better Start – National Science Challenge, University of Auckland, Auckland, New Zealand.

³ Department of Women's and Children's Health, Uppsala University, Uppsala, Sweden.

⁴ Department of Women and Children's Health, School of Life Course Sciences, King's College London, UK.

⁵ Department of Obstetrics and Gynaecology, University of Auckland, Auckland, New Zealand.

⁶ National Women's Hospital, Auckland District Health Board, Auckland, New Zealand.

⁷ Section of Obstetrics and Gynaecology, Leeds Institute of Biomedical & Clinical Sciences, University of Leeds, Leeds, United Kingdom

⁸ Discipline of Obstetrics and Gynaecology, Adelaide Medical School, Robinson Research Institute, University of Adelaide, Adelaide, Australia.

⁹ Maternal and Fetal Health Research Centre, University of Manchester, Manchester, United Kingdom.

* Author for correspondence: Assoc Prof Fredrik Ahlsson, Department of Women's and Children's Health, Uppsala University, Uppsala, Sweden. Email: fredrik.ahlsson@kbh.uu.se

Short title: Paternal factors and LGA infants

Manuscript word count: 2,343

ABSTRACT

We assessed whether paternal demographic, anthropometric and clinical factors influence the risk of an infant being born large-for-gestational-age (LGA). We examined data on 3,659 fathers of term offspring (including 662 LGA infants) born to primiparous women from SCOPE. LGA was defined as birthweight >90th centile as per INTERGROWTH 21st standards, with reference group being infants ≤90th centile. Associations between paternal factors and likelihood of an LGA infant were examined using univariable and multivariable models. Men who fathered LGA babies were 180 grams heavier at birth ($p<0.001$) and were more likely to have been born macrosomic ($p<0.001$) than those whose infants were not LGA. Fathers of LGA infants were 2.1 cm taller ($p<0.001$), 2.8 kg heavier ($p<0.001$), but had similar BMI. In multivariable models, increasing paternal birthweight and height were independently associated with greater odds of having an LGA infant, irrespective of maternal factors. One unit increase in paternal BMI was associated with 2.9% greater odds of having a LGA boy but not girl; however, this association disappeared after adjustment for maternal BMI. There was no association between paternal demographic factors or clinical history and infant LGA. In conclusion, fathers who were heavier at birth and were taller were more likely to have a LGA infant, but maternal BMI had a dominant influence on LGA.

Keywords: LGA, father, mother, birth weight, body mass index, BMI

Abbreviations: BMI, body mass index; LGA, large-for-gestational-age; SGA, small-for-gestational-age.

INTRODUCTION

Birth weight steadily increased worldwide throughout the late 20th century, with a resulting increase in the prevalence of babies that are born large-for-gestational-age (LGA, generally defined as birth weight >90th centile) ¹. Being born either too small (small-for-gestational-age [SGA] – birth weight <10th centile), LGA, or with macrosomia (birth weight >4.0 kg) is associated with adverse long-term health outcomes ². While the evidence for long-term health outcomes in those born LGA is not always consistent ², the association between birth weight and risk of adulthood disease appears to be U-shaped ³. Nonetheless, adults born LGA have an increased risk of obesity ⁴, cardiovascular disease ⁴, diabetes ⁵, and breast cancer ⁶.

Greater maternal body mass index (BMI) is associated with an increased risk of delivering an LGA infant ^{2,7}. In addition, intergenerational studies have shown that women who were born LGA themselves are at increased risk of giving birth to LGA infants ⁸. There is also some evidence of a paternal effect, with father's birth weight reported in a number of studies as an independent predictor of offspring birth weight ⁹⁻¹¹, including a large population study in Sweden ¹². While a positive association between paternal height

and offspring birth weight has been consistently shown^{9,13-17}, the evidence for paternal BMI is conflicting. Studies have reported that greater paternal BMI was associated with increased offspring weight^{10,11,15}. One investigation observed a J-shaped curve, where greater paternal BMI was associated with both low and high birth weight in the offspring¹⁴. However, the evidence is inconclusive as a number of studies have reported no relationship between paternal BMI and offspring birth weight^{9,13,16-18}. The conflicting reports may be explained (at least in part) by sex-specific effects on the offspring. For example, Chen et al. showed that paternal BMI was positively associated with birth weight, abdominal circumference, and other anthropometric parameters in male offspring, but there were no observed associations among females¹⁹.

In addition, there is evidence that the father's metabolic health may affect the offspring. Notably, while maternal diabetes is associated with an increased risk of delivering an LGA baby, there is a consistent reduction in offspring birth weight with paternal type 2 diabetes²⁰⁻²³, as well as paternal insulin resistance^{23,24} or increased glucose levels¹⁴.

Most studies to date have focused on the associations between paternal birth weight or BMI and low birth weight in the offspring, with fewer studies investigating paternal factors associated with the likelihood of having an LGA infant. Thus, we aimed to examine the associations between paternal factors (including anthropometry and metabolic parameters) and the likelihood of having an infant born LGA, while accounting for important maternal confounders.

METHODS

Study participants

Participants were healthy nulliparous women with singleton pregnancies recruited in 2004–2011 to the SCOPE (Screening for Pregnancy Endpoints) study across four countries: Australia (Adelaide), Ireland (Cork), New Zealand (Auckland), and United Kingdom (Manchester, Leeds, and London)²⁵. Ethics approval was provided by the relevant institutional ethic committees²⁵. All women provided written informed consent. If a participant was certain of the identity of the infant's father and she consented, the father was invited to participate and included in this study if he also provided written informed consent.

Study parameters

Paternal characteristics were recorded by 20 weeks of gestation. Paternal height, weight, abdominal circumference, and blood pressure were measured by a research midwife. Data collected included age, race/ethnicity, socioeconomic index²⁶, birth weight, as well as previous diagnosis of diabetes,

hypertension, or heart disease. Birth weight was self-reported, but fathers were asked to confirm the weight from their health records whenever possible. Maternal height and weight were similarly recorded at 14-16 weeks of gestation, with birth weight information obtained as for the fathers'.

Gestational age was determined based on the date of the last menstrual period. However, gestational age was adjusted if one of the following discrepancies between the age from the scan and that calculated by the last menstrual period was observed:

- a) ≥ 7 days based on a scan performed prior to 16 weeks of gestation; or
- b) ≥ 10 days based on the 20-week scan.

If either condition was met, the scan dates were used to determine the estimated date of delivery.

Maternal smoking status and physical activity levels were recorded, with the latter dichotomously divided into whether or not the mother exercised at least moderately (defined as recreational walking ≥ 4 times per week). Infant anthropometric measurements were made soon after birth, usually within 72 hours.

Birth weight centiles were obtained as per INTERGROWTH-21st standards ²⁷, and LGA was defined as birth weight $>90^{\text{th}}$ centile. Macrosomia was defined as a birth weight ≥ 4.0 kg, and it was only adopted as a predictor due to the absence of gestational age data on the fathers. LGA is a more robust indicator of oversize at birth as it takes into account the infant's gestational age at birth, so that the LGA classification can be applied across the gestational age spectrum. Overweight/obesity was defined as BMI ≥ 25 kg/m², obesity as BMI ≥ 30 kg/m², and central adiposity as waist circumference >102 cm for fathers and >88 cm for mothers ²⁸. Paternal hypertension was defined as systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg.

Statistical analyses

Paternal and maternal characteristics in LGA and non-LGA infants were compared using one-way ANOVA, Chi-square tests, or Fisher's exact tests, as appropriate. Univariable logistic regressions were performed to examine associations between paternal factors and the risk of fathering an LGA infant. Continuous predictors were paternal age, socioeconomic status, birth weight, height, weight, BMI, abdominal circumference, systolic and diastolic blood pressures; categorical predictors were race/ethnicity, macrosomia obesity, central adiposity, hypertension at study visit, pre-diagnosed hypertension, and diabetes, as well as fetal sex. The continuous factors significantly associated with LGA birth (at $p < 0.05$) were compared with pairwise linear correlations. A conservative approach was adopted to avoid issues with multicollinearity, and potential predictors were excluded if $|r| > 0.25$ as suggested by Vatcheva et al. ²⁹. Where two parameters were highly correlated, the most clinically logical parameter was selected; where both the continuous parameter and the equivalent categorical variable were found to

be significantly associated with the likelihood of having a LGA infant (e.g. birth weight vs macrosomia), the continuous parameter was chosen for inclusion. All selected paternal parameters were included in a generalized linear regression model incorporating all selected paternal parameters.

The above process was repeated for the corresponding maternal predictors, with the addition of pre-eclampsia, maternal smoking, and moderate-to-vigorous exercise during pregnancy. The final maternal parameters selected were added into a generalized linear regression model including also the selected paternal factors. The above procedures were repeated separately for male and female infants

Each final model discrimination was estimated using the area under the receiver operating characteristic curve (AUROC). Model calibration (i.e. the extent to which the model correctly estimated the absolute risk) was determined using the Hosmer-Lemeshow test ³⁰, with satisfactory calibration determined by $p > 0.05$. Lastly, standardized coefficients were obtained for multivariable models as per King 2007 ³¹.

Statistical analyses were carried out in Minitab v16 (Pennsylvania State University, State College, PA, USA) and SPSS v25 (IBM Corp, Armonk, NY, USA). All statistical tests were two-tailed, and a significance level of 5% was maintained. No adjustments were made for multiple comparisons, and there was no imputation of missing values.

RESULTS

A total of 5,628 pregnant women were recruited into SCOPE; 491 of whom were excluded due to fetal loss at <20 weeks of gestation, perinatal death, major congenital abnormalities, preterm birth, or use of donor sperm or oocyte. A further 13 pregnancies were excluded as the biological father was unknown (Supplementary Figure S1). Of the remaining 5,124 pregnancies, 1,113 biological fathers (21.7%) declined to be in the study, and 352 pregnancies were excluded as paternal birth weight data could not be supplied (Supplementary Figure S1). As a result, we examined data on 3,659 fathers, mothers, and newborns, including 342 infants who were born LGA (18.1%) (Supplementary Figure S1). Rates of LGA birth amongst the offspring of included and excluded fathers were similar (17.2% vs 18.1%, respectively; $p = 0.41$).

LGA

Fathers of LGA infants were demographically similar to those of infants who were not born LGA, as were their respective clinical characteristics (Table 1). However, men who fathered LGA babies were 180 g heavier at birth ($p < 0.001$) and were more likely to have been born macrosomic (Table 1). On average, the fathers of LGA infants were 2.1 cm taller ($p < 0.001$), 2.8 kg heavier ($p < 0.001$), but had similar

BMI (Table 1). While the rate of overweight/obesity was 4.9 percentage points greater among fathers of LGA babies, obesity rates were similar in the two groups (Table 1). The observed differences in paternal characteristics were largely mirrored by maternal characteristics, except that mothers of LGA infants displayed a more marked increase in measures of adiposity compared to those giving birth to non-LGA infants (Table 1).

When individually assessed, demographic and clinical characteristics of the father were not associated with the odds of having a child born LGA, except the odds of the latter were lower among non-Caucasian fathers (Table 2). In contrast, increasing paternal birth weight and paternal macrosomia at birth, as well as greater adult height and weight were associated with greater odds of having an LGA infant (Table 2). Further, every 5-cm increase in abdominal circumference was associated with a 4.3% increase in the odds of LGA, but no association was observed with paternal BMI (Table 2).

When paternal factors were considered together, only increasing paternal birth weight and height remained associated with greater odds of fathering a LGA infant across the whole cohort (Table 3). These associations persisted following the inclusion of maternal predictors (Table 3).

A nearly identical pattern was observed among baby girls, with persistent associations between increasing paternal birth weight and height and the odds of LGA (Table 3). Further, increasing paternal age was associated with lower odds of a LGA girl irrespective of maternal predictors, but standardized coefficients indicated it was a comparatively weaker predictor (Table 3).

Among boys, paternal birth weight and height were also positively associated with the odds of LGA, irrespective of the inclusion of maternal parameters (Table 3). Notably, increasing paternal BMI was associated with greater odds of LGA boys (+2.9% for every 1 kg/m² increment), but this association disappeared when maternal parameters were accounted for (Table 3).

DISCUSSION

We observed that increasing paternal birth weight and adult height were the primary paternal predictors of an increased risk of LGA in the offspring, irrespective of adjustment for maternal factors or the baby's sex. In spite of using the INTERGROWTH 21st standard (which has high rates of LGA when applied to Western settings³²), our findings corroborate previous observations that paternal birth weight is an important independent predictor of offspring birth weight^{10,11,33}.

Increasing paternal BMI was associated with greater odds of having an LGA infant, is in line with other studies showing an association with offspring birth weight^{10,11,34}. However, this association was confined

to boys, and the positive association between paternal BMI and LGA was no longer significant after adjustment for maternal BMI. Thus, paternal BMI was not an independent predictor of offspring birth weight as suggested previously ^{13,17,35}. In addition, other paternal factors related to metabolism (blood pressure, diabetes, and heart disease), body composition (waist circumference), and environment (socioeconomic status) were not associated with the likelihood of fathering an LGA baby. The absence of any association between these parameters recorded at mid pregnancy indicate that the most important paternal factors influencing offspring birth weight are likely to be of genetic origin, as previously suggested ¹³. As a large proportion of an individual's height is genetic ³⁶, our findings are not surprising as greater paternal stature is associated with faster limb growth in the fetus, resulting in longer and heavier infants more likely to be LGA ¹³. Importantly, the genetic paternal contribution contrasts to the maternal influences on offspring size at birth, which are not only genetically driven but also affected by the intrauterine environment ³⁷, as observed among infants born to mothers with gestational diabetes or who are obese ³⁸⁻⁴⁰.

Previous studies have described an association between paternal hypertension and offspring low birth weight ^{24,41}. It has been suggested that the underpinning mechanisms were related to for example, maternal passive smoking ⁴². However, there is a paucity of data regarding offspring born at the upper end of the birth weight spectrum. Here, we observed no association between offspring birth weight (or the likelihood of being born LGA) with paternal blood pressure or central adiposity, irrespective of maternal factors (including smoking during pregnancy).

The limitations of our study include the potential imprecision of maternal and paternal birth weight data, which in many cases could not be verified and relied on recollection. Paternal blood pressure was based on measurement from a single clinic and thus potentially unreliable, but due to the size of the cohort it was not possible to perform repeated measurements over time. As the vast majority of study participants were Caucasian and all infants were born to nulliparous mothers, our findings may not be readily extrapolated to other populations. Further, infants may be LGA by weight and/or length. The paternal influences, such as greater height, are more likely to be associated with a longer rather than a fatter baby, so that the long-term health risks are likely to be greater in the former. Thus, the paternal influences on long-term risk of obesity and related comorbidities in LGA babies are probably minor. The strengths of our study include a large sample size and the robust parental anthropometric data that were measured by a research midwife rather than estimated or recalled by participants.

In conclusion, fathers who are taller as adults and were heavier at birth were more likely to have LGA infants. However, paternal BMI was not found to be an independent predictor of offspring birth weight, suggesting that fathers who are overweight and/or obese do not seem more likely to have an LGA baby

when maternal factors are accounted for. Studies in larger cohorts are necessary to define more accurately the association of paternal clinical factors and offspring birth weight.

Acknowledgements: We would like to thank the pregnant women who participated in the SCOPE Study, the SCOPE Country Project Managers Mrs D Healy, University of Adelaide, Dr A Briley, Kings College London, and Mrs N Murphy and Mrs E Snapes, University College Cork, Miss Eliza Chan for development of the databases, Dr Mik Black for creating the training and validation databases.

Funding: This work was supported by the following funders: New Zealand: New Enterprise Research Fund, Foundation for Research Science and Technology (EM 04–05/03); Health Research Council (04/198); Evelyn Bond Fund, Auckland District Health Board Charitable Trust; Australia: Premier’s Science and Research Fund, South Australian Government; London: Guy’s and St Thomas’ Charity, United Kingdom, Tommy’s the Baby Charity; Manchester: UK Biotechnology and Biological Sciences Research Council, UK National Health Services NEAT Grant, University of Manchester Proof of Concept Funding, Tommy’s the Baby Charity, NIHR; Leeds: Cerebra, UK; and Cork, Ireland: Health Research Board, Ireland; Sweden: the Gillberg Foundation (Uppsala), and Her Royal Highness the Crown Princess Lovisas Society for Child Health Care (Stockholm).

Declaration of interest: The funders had no role in study design, data collection and analysis, decision to publish, or preparation of this manuscript. The authors have no financial conflicts of interest to disclose that may be relevant to this work.

REFERENCES

1. Oken E. Secular trends in birthweight. *Nestle Nutrition Institute workshop series* 2013; 71: 103-14.
2. Chiavaroli V, Derraik JGB, Hofman PL, Cutfield WS. Born large for gestational age: bigger is not always better. *J Pediatr* 2016; 170: 307-11.
3. Wei JN, Sung FC, Li CY, et al. Low birth weight and high birth weight infants are both at an increased risk to have type 2 diabetes among schoolchildren in Taiwan. *Diabetes Care* 2003; 26: 343-8.
4. Skilton MR, Siitonen N, Wurtz P, et al. High birth weight is associated with obesity and increased carotid wall thickness in young adults: the cardiovascular risk in young Finns study. *Arterioscler Thromb Vasc Biol* 2014; 34: 1064-8.
5. Johnsson IW, Haglund B, Ahlsson F, Gustafsson J. A high birth weight is associated with increased risk of type 2 diabetes and obesity. *Pediatr Obes* 2015; 10: 77-83.
6. Innes K, Byers T, Schymura M. Birth characteristics and subsequent risk for breast cancer in very young women. *Am J Epidemiol* 2000; 152: 1121-8.

7. Kramer MS, Morin I, Yang H, et al. Why are babies getting bigger? Temporal trends in fetal growth and its determinants. *J Pediatr* 2002; 141: 538-42.
8. Ahlsson F, Gustafsson J, Tuvemo T, Lundgren M. Females born large for gestational age have a doubled risk of giving birth to large for gestational age infants. *Acta Paediatr* 2007; 96: 358-62.
9. Magnus P, Gjessing HK, Skrondal A, Skjærven R. Paternal contribution to birth weight. *J Epidemiol Community Health* 2001; 55: 873-7.
10. L'Abée C, Vrieze I, Kluck T, Erwich JJ, Stolk RP, Sauer PJ. Parental factors affecting the weights of the placenta and the offspring. *J Perinat Med* 2011; 39: 27-34.
11. Klebanoff MA, Mednick BR, Schulsinger C, Secher NJ, Shiono PH. Father's effect on infant birth weight. *Am J Obstet Gynecol* 1998; 178: 1022-6.
12. Mattsson K, Rylander L. Influence of maternal and paternal birthweight on offspring birthweight – a population-based intergenerational study. *Paediatr Perinat Epidemiol* 2013; 27: 138-44.
13. Pomeroy E, Wells JC, Cole TJ, O'Callaghan M, Stock JT. Relationships of maternal and paternal anthropometry with neonatal body size, proportions and adiposity in an Australian cohort. *Am J Phys Anthropol* 2015; 156: 625-36.
14. Mykkestad K, Vatten LJ, Magnussen EB, Salvesen KÅ, Smith GD, Romundstad PR. Offspring birth weight and cardiovascular risk in parents—a population-based HUNT 2 study. *Am J Epidemiol* 2012; 175: 546-55.
15. Leary S, Fall C, Osmond C, et al. Geographical variation in relationships between parental body size and offspring phenotype at birth. *Acta Obstet Gynecol Scand* 2006; 85: 1066-79.
16. Sletner L, Nakstad B, Yajnik CS, et al. Ethnic differences in neonatal body composition in a multi-ethnic population and the impact of parental factors: a population-based cohort study. *PLoS ONE* 2013; 8: e73058.
17. Nahum GG, Stanislaw H. Relationship of paternal factors to birth weight. *J Reprod Med* 2003; 48: 963-8.
18. Morrison J, Williams GM, Najman JM, Andersen MJ. The influence of paternal height and weight on birth-weight. *Aust N Z J Obstet Gynaecol* 1991; 31: 114-6.
19. Chen Y-P, Xiao X-M, Li J, Reichetzeder C, Wang Z-N, Hoher B. Paternal body mass index (BMI) is associated with offspring intrauterine growth in a gender dependent manner. *PLoS ONE* 2012; 7: e36329.
20. Tyrrell JS, Yaghootkar H, Freathy RM, Hattersley AT, Frayling TM. Parental diabetes and birthweight in 236 030 individuals in the UK Biobank Study. *Int J Epidemiol* 2013; 42: 1714-23.
21. Hypponen E, Smith GD, Power C. Parental diabetes and birth weight of offspring: intergenerational cohort study. *BMJ* 2003; 326: 19-20.
22. Lindsay RS, Dabelea D, Roumain J, Hanson RL, Bennett PH, Knowler WC. Type 2 diabetes and low birth weight: the role of paternal inheritance in the association of low birth weight and diabetes. *Diabetes* 2000; 49: 445-9.

23. Wannamethee SG, Lawlor DA, Whincup PH, Walker M, Ebrahim S, Davey-Smith G. Birthweight of offspring and paternal insulin resistance and paternal diabetes in late adulthood: cross sectional survey. *Diabetologia* 2004; 47: 12-8.
24. Hillman S, Peebles DM, Williams DJ. Paternal metabolic and cardiovascular risk factors for fetal growth restriction: a case-control study. *Diabetes Care* 2013; 36: 1675-80.
25. McCowan LM, Thompson JM, Taylor RS, et al. Clinical prediction in early pregnancy of infants small for gestational age by customised birthweight centiles: findings from a healthy nulliparous cohort. *PLoS ONE* 2013; 8: e70917.
26. Galbraith C, Jenkin G, Davis P, Coope P. *New Zealand Socio-economic Index 1996 – Users' Guide*. Wellington: Statistics New Zealand; 2003.
27. Villar J, Ismail LC, Victora CG, et al. International standards for newborn weight, length, and head circumference by gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-21st Project. *Lancet* 2014; 384: 857-68.
28. Lean MEJ, Han TS, Morrison CE. Waist circumference as a measure for indicating need for weight management. *BMJ* 1995; 311: 158-61.
29. Vatcheva KP, Lee M, McCormick JB, Rahbar MH. Multicollinearity in regression analyses conducted in epidemiologic studies. *Epidemiology* 2016; 6: 227.
30. Hosmer DW, Lemeshow S. Assessing the fit of the model. In: Shewhart WA, Wilks SS, Hosmer DW, Lemeshow S, editors. *Applied Logistic Regression*. 2nd ed: John Wiley & Sons Inc.; 2005. p. 143-202.
31. King JE. Standardized coefficients in logistic regression. *Annual Meeting of the Southwest Educational Research Association*. San Antonio, TX, USA; 2007.
32. Liu S, Metcalfe A, Leon JA, et al. Evaluation of the INTERGROWTH-21st project newborn standard for use in Canada. *PLoS ONE* 2017; 12: e0172910.
33. Magnus P, Gjessing HK, Skrondal A, Skjaerven R. Paternal contribution to birth weight. *J Epidemiol Community Health* 2001; 55: 873-7.
34. Leary S, Fall C, Osmond C, et al. Geographical variation in relationships between parental body size and offspring phenotype at birth. *Acta Obstet Gynecol Scand* 2006; 85: 1066-79.
35. Sletner L, Nakstad B, Yajnik CS, et al. Ethnic differences in neonatal body composition in a multi-ethnic population and the impact of parental factors: a population-based cohort study. *PLoS ONE* 2013; 8: e73058.
36. Wood AR, Esko T, Yang J, et al. Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat Genet* 2014; 46: 1173-86.
37. Derraik JGB, Ahlsson F, Diderholm B, Lundgren M. Obesity rates in two generations of Swedish women entering pregnancy, and associated obesity risk among adult daughters. *Sci Rep* 2015; 5: 16692.

38. Olmos PR, Borzone GR, Olmos RI, et al. Gestational diabetes and pre-pregnancy overweight: possible factors involved in newborn macrosomia. *J Obstet Gynaecol Res* 2012; 38: 208-14.
39. Kc K, Shakya S, Zhang H. Gestational diabetes mellitus and macrosomia: a literature review. *Ann Nutr Metab* 2015; 66: 14-20.
40. Catalano PM, Shankar K. Obesity and pregnancy: mechanisms of short term and long term adverse consequences for mother and child. *BMJ* 2017; 356: j1.
41. Mykkestad K, Vatten LJ, Magnussen EB, Salvesen KA, Smith GD, Romundstad PR. Offspring birth weight and cardiovascular risk in parents: a population-based HUNT 2 study. *Am J Epidemiol* 2012; 175: 546-55.
42. Suter M, Abramovici A, Showalter L, et al. In utero tobacco exposure epigenetically modifies placental CYP1A1 expression. *Metabolism* 2010; 59: 1481-90.

Table 1. Demographic, anthropometric, and clinical characteristics of fathers and mothers of infants who were born large-for-gestational-age (LGA) or not. Where appropriate, data are means $\pm$ standard deviations. *P*-values are from one-way ANOVA or Chi-square tests.

		FATHER			MOTHER		
		Not LGA	LGA	<i>p</i>	Not LGA	LGA	<i>p</i>
n		2,997	662		2,997	662	
Demography	Age (years)	31.0 $\pm$ 5.8	31.0 $\pm$ 5.6	0.79	28.7 $\pm$ 5.2	29.0 $\pm$ 5.2	0.19
	Race/ethnicity (Caucasian)	91.4%	94.0%	0.028	91.4%	93.7%	0.06
	Socioeconomic index	43.1 $\pm$ 15.9	43.3 $\pm$ 15.2	0.76	42.4 $\pm$ 16.7	42.0 $\pm$ 15.9	0.62
At birth	Weight (kg)	3.44 $\pm$ 0.59	3.62 $\pm$ 0.59	<0.001	3.28 $\pm$ 0.52	3.45 $\pm$ 0.56	<0.001
	Macrosomia ^A	15.0%	22.1%	<0.001	7.3%	13.5%	<0.001
Adult anthropometry	Height (cm)	178.6 $\pm$ 6.6	180.7 $\pm$ 6.8	<0.001	164.6 $\pm$ 6.3	167.0 $\pm$ 6.4	<0.001
	Weight (kg)	85.5 $\pm$ 14.6	88.3 $\pm$ 14.2	<0.001	67.8 $\pm$ 13.4	73.5 $\pm$ 15.7	<0.001
	BMI (kg/m ²)	26.77 $\pm$ 4.16	27.03 $\pm$ 3.90	0.15	25.01 $\pm$ 4.67	26.39 $\pm$ 5.49	<0.001
	Overweight/obese (BMI >25 kg/m ²)	65.2%	70.1%	0.016	40.6%	51.8%	<0.001
	Obese (BMI >30 kg/m ²)	17.7%	18.9%	0.47	13.5%	19.9%	<0.001
	Abdominal circumference (cm)	93.1 $\pm$ 11.0	94.1 $\pm$ 10.7	0.027	83.0 $\pm$ 12.3	86.8 $\pm$ 12.5	<0.001
	Central adiposity ^B	16.2%	18.4%	0.19	25.8%	37.0%	<0.001
Clinical features	Systolic blood pressure (mmHg)	123.4 $\pm$ 13.7	123.9 $\pm$ 13.3	0.35	106.8 $\pm$ 10.3	107.9 $\pm$ 10.2	0.009
	Diastolic blood pressure (mmHg)	76.8 $\pm$ 9.7	77.0 $\pm$ 9.5	0.67	64.9 $\pm$ 7.6	65.4 $\pm$ 7.3	0.09
	Hypertension at visit ^C	3.2%	2.7%	0.62	–	–	
	Hypertension ^D	17.2%	16.6%	0.73	–	–	
	Diabetes ^E	0.6%	0.3%	0.40	–	–	

^A Defined as birth weight $\geq$ 4.0 kg

^B Abdominal circumference >102 cm for fathers and >88 cm for mothers.

^C Systolic blood pressure $\geq$ 140 mmHg or diastolic blood pressure $\geq$ 90 mmHg at the study's partner assessment. Note that presence of essential hypertension was an exclusion criteria for mothers.

^D This was defined as a history of hypertension diagnosed by a health practitioner and is taking antihypertensive treatment now or has done in the past. Note that presence of essential hypertension was an exclusion criteria for mothers.

^E Pre-existing diabetes was an exclusion criteria for mothers.

Table 2. Parental demographic, anthropometric, and clinical characteristics and their associated odds ratios (95% confidence intervals) of having a child born large-for-gestational-age (LGA). Statistically significant associations (at $p < 0.05$) are shown in bold.

		Father	Mother
Demography	Age (years)	0.998 (0.984, 1.013)	1.011 (0.995, 1.027)
	Race/ethnicity ^A	0.680 (0.482, 0.959)	0.721 (0.514, 1.010)
	Socioeconomic index	1.001 (0.996, 1.006)	0.999 (0.994, 1.004)
At birth	Birth weight (per 100 g)	1.054 (1.039, 1.070)	1.068 (1.050, 1.087)
	Macrosomia ^B	1.597 (1.296, 1.969)	1.990 (1.526, 2.595)
	Infant sex	0.989 (0.836, 1.171)	0.989 (0.836, 1.171)
Adult anthropometry	Height (cm)	1.047 (1.034, 1.061)	1.061 (1.047, 1.075)
	Weight (kg)	1.013 (1.007, 1.018)	1.026 (1.021, 1.032)
	BMI (kg/m ²)	1.015 (0.995, 1.035)	1.054 (1.037, 1.071)
	Obesity (BMI >30 kg/m ²)	1.084 (0.873, 1.345)	1.594 (1.282, 1.981)
	Abdominal circumference (per 5 cm)	1.043 (1.005, 1.083)	1.44 (1.104, 1.185)
	Central adiposity ^C	1.164 (0.935, 1.450)	1.693 (1.418, 2.023)
Clinical features	Systolic blood pressure (mmHg)	1.003 (0.997, 1.009)	1.011 (1.003, 1.019)
	Diastolic blood pressure (mmHg)	1.002 (0.993, 1.011)	1.009 (0.998, 1.021)
	Hypertension at visit ^D	0.958 (0.765, 1.201)	–
	Hypertension ^E	0.854 (0.512, 1.423)	–
	Diabetes ^F	0.475 (0.110, 2.044)	–
	Preeclampsia	–	1.064 (0.708, 1.598)
Lifestyle	Smoking during pregnancy	–	0.765 (0.618, 0.946)
	Moderate to vigorous exercise in pregnancy	–	0.886 (0.698, 1.124)

^A Non-Caucasian vs Caucasian.

^B Defined as birth weight ≥ 4.0 kg

^C Abdominal circumference >102 cm for fathers and >88 cm for mothers.

^D Systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg at the study's partner assessment. Note that presence of essential hypertension was an exclusion criteria for mothers.

^E This was defined as a history of hypertension diagnosed by a health practitioner and is taking antihypertensive treatment now or has done in the past. Note that presence of essential hypertension was an exclusion criteria for mothers.

^F Pre-existing diabetes was an exclusion criteria for mothers.

Table 3. Paternal characteristics and the respective adjusted odds ratio (aOR) of fathering a large-for-gestational-age (LGA) baby.

Population ^A	Paternal characteristics	Paternal predictors only					Paternal and maternal predictors ^B				
		aOR (95% CI)	Std coef	p-value	H-L	AUROC	aOR (95% CI)	St coef	p-value	H-L	AUROC
All babies	Birth weight (per 100 g)	1.043 (1.028, 1.059)	0.037	<0.001	0.21	0.62 (0.59, 0.64)	1.048 (1.031, 1.065)	0.041	<0.001	0.30	0.68 (0.65, 0.70)
	Abdominal circumference (per 5 cm)	1.019 (0.980, 1.060)	0.019	0.35			0.990 (0.948, 1.033)	-0.003	0.64		
	Height (cm)	1.037 (1.023, 1.051)	0.036	<0.001			1.040 (1.025, 1.054)	0.039	<0.001		
	Ethnicity (Non-Caucasian vs Caucasian)	0.763 (0.538, 1.081)	-0.011	0.13			0.812 (0.559, 1.181)	-0.008	0.28		
Boys	Birth weight (per 100 g)	1.040 (1.017, 1.063)	0.033	<0.001	0.42	0.62 (0.58, 0.65)	1.045 (1.021, 1.070)	0.037	<0.001	0.32	0.70 (0.67, 73)
	BMI (kg/m ²)	1.029 (1.001, 1.058)	0.018	0.041			1.005 (0.973, 1.038)	0.003	0.76		
	Height (cm)	1.042 (1.023, 1.062)	0.041	<0.001			1.032 (1.012, 1.054)	0.031	0.002		
Girls	Birth weight (per 100 g)	1.045 (1.023, 1.067)	0.040	<0.001	0.26	0.62 (0.59, 0.66)	1.054 (1.031, 1.077)	0.047	<0.001	0.19	0.68 (0.65, 0.71)
	Height (cm)	1.037 (1.018, 1.056)	0.036	<0.001			1.031 (1.011, 1.051)	0.037	0.002		
	Age (years)	0.972 (0.951, 0.993)	-0.024	0.009			0.968 (0.946, 0.990)	-0.027	0.005		
	Ethnicity (Non-Caucasian vs Caucasian)	0.627 (0.375, 1.047)	-0.020	0.07			0.791 (0.466, 1.342)	-0.010	0.38		

Data were analysed using multivariable binary logistic regression models.

AUROC, area under the receiver operating characteristic curve; BMI, body mass index; H-L, Hosmer and Lemeshow Test; Std coef, standardized logistic regression coefficient.

^A For all babies n=3,659 (342 LGA), for boys n=1,832 (170 LGA), and for girls n=1,827 (172 LGA).

^B The following maternal predictors were included in the respective models:

- All babies: birth weight, BMI, random glucose at first visit, random glucose at second visit, smoking during pregnancy.
- For boys: age, birth weight, BMI, height, random glucose at first visit, and smoking during pregnancy.
- For girls: birth weight, BMI, and height.

Supplementary Figure S1

Recruitment of participants for the study.